

ECHA note: In appendix V: Overall reference list of the initially uploaded CLH report some of the data owners were reported incorrectly. In this version of the CLH report, the data owner information has been corrected by the Dossier Submitter.

**REGULATION (EC) NO 1272/2008 (CLP REGULATION),
ANNEX VI, PART 2**

**Proposal for Harmonised Classification and Labelling for a biocidal
active substance**

CLH REPORT

***Chrysanthemum cinerariaefolium*, extract from open and mature
flowers of *Tanacetum cinerariifolium* obtained with supercritical
CO₂ (Redefined from Pyrethrins and Pyrethroids and
Chrysanthemum cinerariaefolium, extract)**

EC Number: 289-699-3

CAS Number: 89997-63-7

Index Number:

Contact details of dossier submitter: Ministerio de Sanidad, S. G. de Sanidad, D.G.
Salud Pública, Calidad e Innovación,
Subdirección General de Sanidad Ambiental y
Salud Laboral, Paseo del Prado 18-20, E -
28071 Madrid

Version number: 05 Date: 28 February 2022

Table of Contents

CLH REPORT	4
Summary.....	4
1. Presentation of the Active Substance	4
1.1 Identity of the active substance	4
1.2 Intended Uses and Effectiveness	7
2. Proposed harmonised classification and labelling of the active substance according to the CLP criteria	8
2.1 Proposed harmonised classification and labelling for the active substance.....	8
2.2 History of the previous classification and labelling	11
2.3 Data sources	11
3. Summary of the Human Health Risk Assessment	12
3.1 Summary of the assessment of effects on human health	12
3.2 Reference values.....	15
4. Summary of the Environmental Risk Assessment.....	16
4.1 Fate and behaviour in the environment.....	16
A. Assessment of intrinsic properties and effects of the active substance.....	20
A.1. General substance information	20
A.1.1. Identity of the substance	20
A.1.2. Composition of the substance (reference specifications).....	24
A.1.3. Physical and chemical properties of the active substance.....	26
A.1.4. Physical hazards and respective characteristics	40
A.1.5. Assessment of physical hazards according to the CLP criteria	47
A.2. Assessment of effects on Human Health	48
A.2.1. Toxicokinetics	48
A.2.2. Acute toxicity / STOT SE	53
A.2.3. Skin corrosion and irritation	67
A.2.4. Serious eye damage and Eye irritation.....	69
A.2.5. Skin sensitisation.....	72
A.2.6. Respiratory sensitisation	78
A.2.7. Repeated dose toxicity/STOT RE.....	79
A.2.8. Genotoxicity / Germ cell mutagenicity.....	107
A.2.9. Carcinogenicity	122
A.2.10. Reproductive toxicity	138
A.2.11. Aspiration hazard.....	150
A.2.12. Neurotoxicity	152
A.2.13. Immunotoxicity	156
A.2.14. Endocrine disruption	156
A.2.15. Further Human data.....	160
A.2.16. Other data.....	163
A.3. Environmental effects assessment.....	164

A.3.1. Fate and distribution in the environment	164
A.3.2. Effects on environmental organisms	192
A.3.3. Overall summary of acute and chronic aquatic toxicity data and Comparison with the CLP criteria	208
A.4. Assessment of additional hazards	211
A.4.1. Hazardous to the ozone layer	211
A.5. Additional Labelling	211
A.6. Assessment of exclusion criteria, substitution criteria and POP	212
A.6.1. Exclusion criteria	212
A.6.2. Substitution criteria	215
A.6.3. Assessment of long-range environmental transportation and impact on environmental compartments	216
B. Appendices.....	1
Appendix V: Overall reference list (including data owner and confidentiality claim)	1
Appendix VII: Study summaries.....	1

CLH REPORT

SUMMARY

1. PRESENTATION OF THE ACTIVE SUBSTANCE

Two separate CARs were submitted in support of the active substances.

Pyrethrins were notified as an existing active substance, by Botanical Resources Australia Pty Ltd. (BRA), McLaughlin Gormley King Company (MGK) and SC Johnson & Son Inc., and a CAR was issued in July 2010, by RMS Spain.

A second CAR was issued in July 2010 by RMS Spain in support of the active substance *Chrysanthemum cinerariaefolium*, Extract which was notified as an existing active substance, by Kenya Pyrethrum Information Centre (KPIC).

It was subsequently decided that the substances were technically equivalent and that a combined CAR should be produced.

In accordance with the REACH guidance¹ the extraction method used to prepare the active substance was included in the substance name resulting in two active substances:

- *Chrysanthemum cinerariaefolium*, extract from open and mature flowers of *Tanacetum cinerariifolium* obtained with hydrocarbon solvents (Redefined from Pyrethrins and Pyrethroids and *Chrysanthemum cinerariaefolium*, ext.)
- *Chrysanthemum cinerariaefolium*, extract from open and mature flowers of *Tanacetum cinerariifolium* obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and *Chrysanthemum cinerariaefolium*, ext.)

This CLH report has been prepared for the active substance, "*Chrysanthemum cinerariaefolium*, extract from open and mature flowers of *Tanacetum cinerariifolium* obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and *Chrysanthemum cinerariaefolium*, ext.)". Due to the length of the name the substance will be referred to by the names used in the study reports and in general overview sections as "*Chrysanthemum cinerariaefolium* extract from supercritical CO₂".

1.1 IDENTITY OF THE ACTIVE SUBSTANCE


Table 1.1 Constituents

Constituent(s)	
ISO name	-
IUPAC or EC name	<p><i>Chrysanthemum cinerariaefolium</i>, extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i>, ext.)</p> <p>Pyrethrin 1: (Z)-(S)-2-methyl-4-oxo-3-(penta-2,4-dienyl)cyclopent-2-enyl (1R)-trans-2,2-dimethyl-3-(2-methylprop-1-enyl) cyclopropane carboxylate</p> <p>Cinerin 1: (Z)-(S)-3-(but-2-enyl)-2-methyl-4-oxocyclopent-2-enyl (1R)-trans-2,2-dimethyl-3-(2-methylprop-1-enyl) cyclopropane carboxylate</p>


¹ Guidance for identification and naming of substances under REACH and CLP, Version 2.1 - May 2017.

	<p>Jasmolin 1: (Z)-(S)-2-methyl-4-oxo-3-(pent-2-enyl)cyclopent-2-enyl (1R)-trans-2,2-dimethyl-3-(2-methylprop-1-enyl) cyclopropane carboxylate</p> <p>Pyrethrin 2: (Z)-(S)-2-methyl-4-oxo-3-(penta-2,4-dienyl)cyclopent-2-enyl (E)-(1R)-trans-3-(2-methoxycarbonylprop-1-enyl)-2,2-dimethylcyclopropane carboxylate</p> <p>Cinerin 2: (Z)-(S)-3-(but-2-enyl)-2-methyl-4-oxocyclopent-2-enyl (E)-(1R)-trans-3-(2-methoxycarbonylprop-1-enyl)-2,2-dimethylcyclopropane carboxylate</p> <p>Jasmolin 2: (Z)-(S)-2-methyl-4-oxo-3-(pent-2-enyl)cyclopent-2-enyl (E)-(1R)-trans-3-(2-methoxycarbonylprop-1-enyl)-2,2-dimethylcyclopropane carboxylate</p>
EC number	<p><i>Chrysanthemum cinerariaefolium</i>, extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i>, ext.): 289-699-3</p> <p><i>Chrysanthemum cinerariaefolium</i>: 289-699-3 Pyrethrins: 232-319-8 Pyrethrin 1: 204-455-8 Pyrethrin 2: 204-462-6 Cinerin 1: 246-948-0 Cinerin 2: 204-454-2 Jasmolin 1: not available Jasmolin 2: not available</p>
CAS number	<p><i>Chrysanthemum cinerariaefolium</i>, extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i>, ext.): 89997-63-7</p> <p><i>Chrysanthemum cinerariaefolium</i>: 89997-63-7 Total Pyrethrins: 8003-34-7 Pyrethrin 1: 121-21-1 Pyrethrin 2: 121-29-9 Cinerin 1: 246-948-0 Cinerin 2: 121-20-0 Jasmolin 1: 4466-14-2 Jasmolin 2: 1172-63-0</p>
Index number in Annex VI of CLP	-
Minimum purity / content	100%


Structural formula




pyrethrin 1




cinerin 1




jasmolin 1



pyrethrin 2



cinerin 2



jasmolin 2

The active substance *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ (synonym: Pyrethrum Extract) subject to the CLH proposal is an extract of the flower heads of *Chrysanthemum cinerariaefolium*. It contains Pyrethrins, which may be divided into the two groups Pyrethrins I (consisting of pyrethrin 1, cinerin 1, and jasmolin 1) and

Pyrethrins II (consisting of pyrethrin 2, cinerin 2 and jasmolin 2). It also contains plant material, BHT and water.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ has a minimum purity of 100% w/w (UVCB substance). It is placed in the market as a solution, technical concentrate (c.a. 50% pyrethrins).

Table 1.2 Relevant impurities and additives

For information about the identity of impurities of the active substance, please refer to Confidential data in Appendix VI of this dossier (confidential information). For sake of completeness, information on the solvent used in all (eco)toxicology studies has been included in Appendix VI even though the active substance is stable without it and, therefore, the solvent should not be considered part of the active substance. This solvent (CAS 64742-47-8; EC 265-149-8) is present at a concentration q.s. 100% (solvent range 42.43-50.65%). The solvent has a harmonised classification as Asp. Tox. 1 (H304) and according to REACH registration dossier has no acute toxicity (oral LD₅₀ > 5000 mg/kg bw, dermal LD₅₀ > 2000 mg/kg/bw, and inhalation LC₅₀ > 5.28 mg/L), skin sensitization (one key GPMT, nine supportive studies and three additional support data did not elicit a positive response) and aquatic toxicity properties that would influence the tests results or assessment of the hazard classes to be harmonised.

The following terms are used throughout the CLH report:

- "Total pyrethrins" is a synonym to the active substance and the substance subject to CLH, i.e. it includes pyrethrins, plant material, BHT and water.
- "Extract" is the test substance, which includes in addition to total pyrethrins, also the solvent.

1.2 INTENDED USES AND EFFECTIVENESS

Table 1.3 Use of the active substance

Product type	MG03: Pest control PT18: Insecticides, acaricides and products to control other arthropods
Intended use pattern(s)	<i>Chrysanthemum cinerariaefolium</i> , extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO ₂ , is intended to be used as insecticide against a wide range of flying and crawling pests except those that are plant parasitic, in various applications sites in- and outdoor. Within this dossier the use against flies and mosquitoes is intended.
Users	Professionals and non-professionals

2. PROPOSED HARMONISED CLASSIFICATION AND LABELLING OF THE ACTIVE SUBSTANCE ACCORDING TO THE CLP CRITERIA

2.1 PROPOSED HARMONISED CLASSIFICATION AND LABELLING FOR THE ACTIVE SUBSTANCE

Table 2.1 Proposed harmonised classification and labelling of the substance

	Index No	Chemical name	EC No	CAS No	Classification		Labelling			Specific Conc. Limits, M-factors and ATEs	Notes
					Hazard Class and Category Code(s)	Hazard statement Code(s)	Pictogram, Signal Word Code(s)	Hazard statement Code(s)	Suppl. Hazard statement Code(s)		
Current Annex VI entry	No current Annex VI entry										
Dossier submitters proposal	TBD	<i>Chrysanthemum cinerariaefolium</i> , extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO ₂	289-699-3	89997-63-7	Acute Tox. 4 Acute Tox. 4 Skin Sens. 1B Aquatic Acute 1 Aquatic Chronic 1	H302 H332 H317 H400 H410	GHS07 GHS09 Wng	H302 H332 H317 H410		Acute, M=100 Chronic, M=10 ATE oral = 700 mg/kg bw ATE inhalation = 2.5 mg/L (dusts and mists)	

A warning statement should be included in the summary of the product characteristics for the biocidal products containing *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ indicating that the product contains an active substance which is dangerous or toxic to bees and that the product should only be applied in early morning or late evening when pollinators are unlikely to be foraging.

Table 2.2 Reason for not proposing harmonised classification and labelling and the status under CLH public consultation.

Hazard class	Reason for not proposing classification and labelling	Within the scope of public consultation
Explosives	data conclusive but not sufficient for classification;	Yes
Flammable gases (including chemically unstable gases)	hazard class not applicable	No
Oxidising gases	hazard class not applicable	No
Gases under pressure	hazard class not applicable	No
Flammable liquids	data conclusive but not sufficient for classification;	Yes
Flammable solids	hazard class not applicable	No
Self-reactive substances and mixtures	data conclusive but not sufficient for classification;	Yes
Pyrophoric liquids	data conclusive but not sufficient for classification;	Yes
Pyrophoric solids	hazard class not applicable	No
Self-heating substances and mixtures	hazard class not applicable	No
Substances which in contact with water emit flammable gases	data conclusive but not sufficient for classification;	Yes
Oxidising liquids	data conclusive but not sufficient for classification;	Yes
Oxidising solids	hazard class not applicable	No
Organic peroxides	data conclusive but not sufficient for classification;	Yes
Corrosive to metals	data conclusive but not sufficient for classification;	Yes
Acute toxicity via oral route	Acute Tox. 4, H302: Harmful if swallowed.	Yes
Acute toxicity via dermal route	data conclusive but not sufficient for classification;	Yes
Acute toxicity via inhalation route	Acute Tox. 4, H332: Harmful if inhaled.	Yes
Skin corrosion/irritation	data conclusive but not sufficient for classification;	Yes
Serious eye damage/eye irritation	data conclusive but not sufficient for classification;	Yes
Respiratory sensitisation	data lacking;	No
Skin sensitisation	Skin Sens. 1B, H317: May cause an allergic skin reaction.	Yes
Germ cell mutagenicity	data conclusive but not sufficient for classification;	Yes

Carcinogenicity	data conclusive but not sufficient for classification;	Yes
Reproductive toxicity	data conclusive but not sufficient for classification;	Yes
Specific target organ toxicity-single exposure	data conclusive but not sufficient for classification;	Yes
Specific target organ toxicity-repeated exposure	data conclusive but not sufficient for classification;	Yes
Aspiration hazard	data conclusive but not sufficient for classification;	Yes
Hazardous to the aquatic environment	Aquatic Acute 1, H400: M=100 Aquatic Chronic 1, H410: M=10	Yes
Hazardous to the ozone layer	data conclusive but not sufficient for classification;	Yes

NOTE eCA: Tests on physical hazards were conducted on the active substance as manufactured (TK, ca. 50% pyrethrins), that is, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ has been tested considering the solvent as part of the active substance. eCA Spain, ECHA and MSs have agreed that solvent should not be part of the active substance composition, therefore tests on physical hazards have been repeated on the purified active substance. However, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ does not meet any physical hazard class using the purified active substance. The details of the physical hazard tests conducted on the purified active substance (without the solvent) are described in section A.1.4 and A.1.5.

This plant extract has different components, which are classified:

Plant extract contains different components: pyrethrin 1, pyrethrin 2, cinerin 1, cinerin 2, jasmolin 1 and jasmolin 2. Pyrethrin 1, pyrethrin 2, cinerin 1 and cinerin 2 are included in Annex VI to CLP with the following classification:

Pyrethrin 1 – Index No. 613-023-00-1 – H302, H312, H332, H400, H410

Pyrethrin 2 – Index No. 613-024-00-7 – H302, H312, H332, H400, H410

Cinerin 1 – Index No. 613-025-00-2 – H302, H400, H410

Cinerin 2 – Index No. 613-026-00-8 – H302, H400, H410

Jasmolin 1 and Jasmolin 2 are not included in the C&L inventory, so self-classification is not available

Other components of the extract (excluding the solvent), which are not included in Annex VI of CLP, have the following proposed classifications:

BHT – CAS 128-37-0 – H410 (REACH registration C&L)

Water – CAS 7732-18-5 – Not classified (Notified C&L)

Plant material:

Fatty acids:

Myristic acid – CAS 544-63-8 – Not classified (Notified C&L)

Palmitic acid – CAS 57-10-3 – Not classified (REACH registration C&L)

Stearic acid – CAS 57-11-4 – Not classified (REACH registration C&L)

Myristoleic acid – CAS 544-64-9 – Not classified (Notified C&L)

Palmitoleic acid – CAS 373-49-9 – H315, H319, H335 (Notified C&L)

Oleic acid – CAS 112-80-1 – Not classified (Notified C&L)

Linoleic acid – CAS 60-33-3 – Not classified (Notified C&L)

Terpenoids:

trans- β-farnesene – CAS 18794-84-8 – H304 (REACH registration C&L)

δ-cadinene CAS 483-76-1 – H315, H304 (Notified C&L)

trans-nerolidol - CAS 40716-66-3 – H400, H410 (REACH registration C&L)

Hexahydrofarnesyl acetone - CAS 502-69-2 – H400, H410 (REACH registration C&L)

Sesamin – CAS 607-80-7 – Not classified (Notified C&L)

β-sitosterol – CAS 83-46-5 – Not classified (Notified C&L)

α-amyrin – CAS 638-95-9 – H302 (Notified C&L)

β-amyrin – CAS 559-70-6 – H302 (Notified C&L)

Lupeol – CAS 545-47-1 – H302 (Notified C&L)

β-cubebene (CAS 13744-15-5), *cis*-Z-α-bisabolene epoxide (CAS not available), taxasterol (CAS 1059-14-9), pyrethrosin (CAS not available), γ-cadinene (CAS 39029-41-9), lupeyl acetate (CAS 1617-68-1), and aromadendrene (CAS 498-39-4) are not included in the C&L inventory, so self-classification is not available.

2.2 HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING

Not applicable.

2.3 DATA SOURCES

Please refer to Appendix V for the reference lists and to Appendix VII for the studies included in DAR not in CAR.

3. SUMMARY OF THE HUMAN HEALTH RISK ASSESSMENT

3.1 SUMMARY OF THE ASSESSMENT OF EFFECTS ON HUMAN HEALTH

Table 3.1 Summary of the assessment of effects on human health

Endpoint	Brief description
Toxicokinetics	<p>¹⁴C-Pyrethrin was administered orally to male and female Sprague-Dawley rats. In a preliminary study time peak blood level and blood half-life were determined. In three definitive experiments, absorption, distribution, excretion and metabolism were examined (Selim, 1995). The definitive studies were conducted using three groups of 5 rats/sex/group as follows: a single low-dose (Group 1: 10 mg/kg bw), a single high-dose (Group 2: 50 mg/kg bw for females and 100 mg/kg for males) and a repeated low-dose (Group 3: non-radiolabelled Pyrethrin 1 at 10 mg/kg bw for 14 days prior to administration of a single oral low-dose of ¹⁴C-Pyrethrin 1 on day 15). Two additional groups of 5 rats/sex/group were dosed exclusively for the purposes of isolation and identification of metabolites. In this ¹⁴C-Pyrethrin 1 was administered orally at a single oral low-dose (Group 4: 10 mg/kg bw) and at a single oral high dose (Group 5: 50 mg/kg bw for females and 100 mg/kg for males).</p> <p>All rats survived the studies and no signs of toxicity were observed. The preliminary study showed that less of the administered ¹⁴C-Pyrethrin 1 was systemically absorbed by the male rats than by the female rats with peaks in blood after 5-6 and 6-8 hours, respectively. Male rats excrete the ¹⁴C-Pyrethrin 1 derived radioactivity faster than females, since the elimination half-time in males and females was 5.3 and 6.7 hours, respectively.</p> <p>In all groups of the definitive studies most of the radioactivity was excreted in the urine and faeces during the first 72 hours following administration. The levels of radioactivity expired as ¹⁴CO₂ were very low. Regardless of dosage regimen, males excrete a majority of the administered radioactivity in the faeces via the enterohepatic circulation while females excrete approximately equal amounts in the urine and faeces. Repeated administration of Pyrethrin 1 increased the rate of elimination of the ¹⁴C-Pyrethrin 1-derived radioactivity from the body in both males and females thus suggesting that repeated dose of ¹⁴C-Pyrethrin 1 results in an induction of the liver microsomal enzyme system.</p> <p>Analysis of tissues at 7 days showed that in all dosing regimens, concentrations of radioactivity more than twice that of whole blood were found in liver, ovaries, carcass and fat. The radioactive residues in the fat were consistently higher than the residues in all other tissues analysed. The elimination of substantial amounts of ¹⁴C-Pyrethrin 1-derived radioactivity in the faeces over and extended length of time indicated that enterohepatic circulation played a role in the elimination of the compound and/or its metabolites from the body.</p>
Acute toxicity	<p><i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ showed acute toxicity by oral and inhalation routes, with an oral LD₅₀ in rats for females (the most sensitive sex) of 700 mg/kg bw total pyrethrins (1073 mg/kg bw extract) (Acute Tox. 4 – H302), a dermal LD₅₀ in rabbits for both sexes higher than 2000 mg/kg bw</p>

	total pyrethrins (3064 mg/kg bw extract) (without classification for this route of exposure), and an inhalation LC ₅₀ in rats of 2.5 and 3.9 mg/L for females and males respectively (Acute Tox. 4 – H332).
Corrosion and irritation	The compound was minimally irritating to the skin and eyes not requiring classification.
Sensitisation	A study conducted using the Buehler method did not indicate that <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ is sensitising, however, three LLNA tests indicate sensitisation and, therefore, <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ has been classified as Skin sensitisation, Category 1B, H317: May cause an allergic skin reaction.
Repeated dose toxicity	In sub-chronic tests for toxicity in mice, rats and dogs, the lowest relevant NOAEL after oral administration were 47, 57 and 18 mg/kg bw/d total pyrethrins (72, 87 and 28 mg/kg bw/d extract), respectively. The liver was the main target and increased liver weight was frequently accompanied with increased activity of transaminases. In addition, anaemia was observed in rats and dogs. Dermal administration of pyrethrins at doses up to 300 mg/kg bw/d total pyrethrins (460 mg/kg bw/d extract) for 21 days caused no systemic toxicity in rabbits. In a 13 weeks study in rats exposed by inhalation, the NOAEL for systemic toxicity was 11 mg/m ³ . The effects described in the above paragraph for the oral application was observed.
Genotoxicity	Pyrethrins are non-genotoxic according to the available <i>in vitro</i> and <i>in vivo</i> studies.
Carcinogenicity	In a two-year study of toxicity and carcinogenicity in rats and an 18-month study of carcinogenicity in mice, the NOAEL were 4.4 and 14 mg/kg bw/d total pyrethrins (6 and 22 mg/kg bw/d extract), respectively. Rats showed at the highest dose (130-173 mg pyrethrins/kg bw/d total pyrethrins (199-265 mg/kg bw/d extract)) an increased incidence of adenoma of the liver in females, follicular cell adenomas of thyroid in both sexes and increased numbers of keratoacanthomas in males. Mice exhibited discoloured liver at the highest dose (690-830 mg pyrethrins/kg/d total pyrethrins (1051-1278 mg/kg bw/d extract)) together with vacuolar fatty change in the liver. No oncogenic effects could be detected in mice. Mechanistic studies indicate that tumours found in rat are species specific and caused by induction of CYP enzymes and the subsequent alteration of the metabolism of thyroid hormones. Therefore, an epigenetic mechanism is induced, and a threshold concentration must be assumed.
Reproductive toxicity	Pyrethrins did not show toxic effects on reproduction in a two-generation study on rats at dietary doses equivalent to 360 mg actual pyrethrins/kg bw/d total pyrethrins (552 mg/kg bw/d extract). In teratogenicity studies, the NOAEL for maternal toxicity and for fetotoxicity in rats was 75 mg/kg bw/d total pyrethrins (115 mg/kg bw/d extract). The NOAEL values for reproduction and teratogenicity in rabbits were 25 and 250 mg/kg bw/d total pyrethrins (38 and 383 mg/kg bw/d extract), respectively. The NOAEL for development of F1 and F2 in a two-generation study was 12 mg/kg bw/d total pyrethrins (18 mg/kg bw/d extract).
Neurotoxicity	In a study of neurotoxicity in rats given single oral doses, acute neurological disorders and behavioural effects were noted, with a

	NOAEL of 20 mg/kg bw total pyrethrins (31 mg/kg bw extract).
Immunotoxicity	-
Disruption of the endocrine system	<p>The available data package for <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ has been assessed according to the specified guidance. The data show no effects of <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ on the estrogen, androgen or steroidogenesis modalities. Although some findings related to the thyroid are noted in the rat with chronic and/or high dose exposure, no adverse effects on the thyroid are observed in other species (mouse and dog). Further, based on the available mode of action data, it can be concluded that the thyroid response in the rat is mediated secondary to hepatic enzyme induction and general systemic effects and is not relevant to humans.</p> <p>According to ECHA/EFSA "Guidance for the identification of Endocrine disruptors":</p> <ul style="list-style-type: none"> - The potential for E,A,S-mediated adversity is considered to have been sufficiently investigated. Overall, there is strong weight of evidence to indicate that <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ does not affect the estrogen, androgen, steroidogenesis modalities. - The potential for T-mediated adversity is considered to have not been sufficiently investigated. Overall, there is not sufficient weight of evidence to indicate that <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ affects the thyroid modality by a mode of action that is specific to the rat and, as such, it cannot be concluded there are no indications of endocrine adversity of relevance to humans. A MoA (mode of action) has been described in the ED assessment report regarding thyroid histopathological changes on rats. <p>Due to the age of some of the studies, they do not include all of the endocrine endpoints that would be required according to modern guidelines. But due to BPR article 90 restrictions, additional assays cannot be asked for.</p> <p>Therefore, regarding the endocrine disruption properties, no conclusion can be drawn as insufficient data is available for the assessment.</p>
Other effects	<p>In the historical data of MGK's employees, no serious adverse reaction to Pyrethrins have been reported. A few minor cases of dermatitis were noted, however, the implication of Pyrethrins is not confirmed. In the manufacturing plant of BRA over the last 25 years of manufacturing, no serious health problems have been recorded in workers either. Minor skin irritations have been observed following the harvesting operation however they have disappeared after washing with clean water and at worst within 1-2 days. Once the harvested crop material has been extracted, no cases of any health issues from handling the extracted and refined products have been noted. A few cases such as dermatitis, pruritus, nausea, a stinging sensation in the nasal and upper pharyngeal mucosa, moderate shortness of breath, cough productive of white phlegm without haemoptysis, fatigue, headache, dizziness and sensitisation were reported among the general population. However, people had recovered after few hours or in one case after 2 years.</p>

3.2 REFERENCE VALUES

Table 3.2 Reference values

Study	NOAEL/ LOAEL	Overall assessment factor	Value
AEL _{short-term}	Neurotoxicity 20 mg/kg bw/d total pyrethrins (31 mg/kg bw/d extract) based on observed tremors	100	0.20 mg/kg/d total pyrethrins (0.31 mg/kg bw/d extract)
AEL _{medium-term}	1-year study in dog 14 mg/kg bw/d total pyrethrins (22 mg/kg bw/d extract) based on hepatic damages	100	0.14 mg/kg/d total pyrethrins (0.22 mg/kg bw/d extract)
AEL _{long-term}	2-year study in rats 4.37 mg/kg bw/d total pyrethrins (6 mg/kg bw/d extract)	100	0.04 mg/kg/d total pyrethrins (0.06 mg/kg bw/d extract)
ADI	2-year study in rats 4.37 mg/kg bw/d total pyrethrins (6 mg/kg bw/d extract)	100	0.04 mg/kg/d total pyrethrins (0.06 mg/kg bw/d extract)
ARfD	Neurotoxicity study 20 mg/kg bw/d total pyrethrins (31 mg/kg bw/d extract) based on observed tremors.	100	0.20 mg/kg/d total pyrethrins (0.31 mg/kg bw/d extract)

4. SUMMARY OF THE ENVIRONMENTAL RISK ASSESSMENT

4.1 FATE AND BEHAVIOUR IN THE ENVIRONMENT

The fate and distribution in the environment was derived from studies on pyrethrin 1, since pyrethrin 1 represents the predominant analogue and a typical member (or paradigm) for the pyrethrum family. Therefore, it was regarded as feasible to make extrapolations from pyrethrin 1 to the active substance (*Chrysanthemum cinerariaefolium* extract from supercritical CO₂). Hence, it was also considered to be justified to model the fate of total pyrethrins in the environment based on characteristics of pyrethrin 1.

Fate and behavior in the environment based on the physico-chemical properties of the active:

Fate and behaviour in air

The data presented predicts that the active substance would degrade rapidly in air under daylight conditions. Moreover, considering the substance's Henry's Law Constant (7.83E-02 Pa/m³/mol at 25°C), vapour pressure (6.9E-05 Pa at 25°C) and octanol water partition coefficient (Log Kow: 5.59), none of the active will be lost to air according to the estimation of the fate of chemicals in a wastewater treatment plant as outlined in the TGD which is based on the SimpleTreat model 4.0. Therefore, emissions to air during the waste disposal stage need not be further considered. This data is important for the indoor-intended use products which contain the active.

Furthermore, this substance is unstable in the atmosphere to react with OH radicals and O₃ of troposphere. Its half-life in this compartment was calculated to be 1.28 h in the presence of OH radicals and 17.13 min in the presence of O₃. No major metabolites (at >10% AR) were detected in air.

Therefore, it is expected that the concentrations of the active substance in air will be negligible for all the uses presented in this dossier.

Fate and behaviour in aquatic compartment (including sediment)

Chrysanthemum cinerariaefolium extract from supercritical CO₂ (represented by pyrethrin 1) is characterized in the aquatic compartment by the following:

- Having a low water solubility, 0.23 mg/L at 20°C
- Being hydrolytically stable in water at 25°C, in the pH-range from 5 to 7, and unstable to pH 9 with a DT₅₀ value of 17 days; with chrysanthemic acid as the single major metabolite. The concentration of chrysanthemic acid (at pH 9) increased in proportion to the decrease of Pyrethrin 1, accounting for 61% of the applied radioactivity (AR) after 30 days (Selim, 1995).
- Not readily biodegradable under the test conditions assessed (CO₂ Evolution – Modified Sturm Test). According to the OECD 301B guideline, substances are readily biodegradable in the CO₂ evolution test if CO₂ production is equal to or greater than 60% of the theoretical value within 10 days of the level achieving 10%.
- Undergoing rapid primary degradation in aerobic natural waters. In the Hein et al. (2017) test, calculated SFO DT₅₀ values for pyrethrin 1 ranged from 6.7-10.7 days (at 20 ± 2°C). The main degradation product was pyrethrolone which reached a maximum of 9.5% AR after 21 days and then decreased to 2.8% AR at the last sampling interval. Several non-identified fractions were detected but these were minor and/or composed of several peaks. Mineralisation reached a maximum of 7%.
- Under the influence of natural sunlight Pyrethrin 1 undergoes degradative processes in an aqueous solution at pH 7. After the irradiation numerous products

were formed. However, no compound was identified. Its photolytic half-life in water was estimated to be 11.8 h.

- Pyrethrin 1 disappears rapidly when applied to aerobic water/sediment systems, taken from the natural environment, as a consequence of its low water solubility and its high adsorption coefficient, which leads to a fast movement into the sediment. In the Robinson and Wisocky (1994) test, degradation in water seems to proceed initially by a combination of hydrolysis and oxidation to form chrysanthemic acid and several low level degradates. A clear pattern of build-up and decline emerged for chrysanthemic acid, resulting in a maximum occurrence in the water/sediment system of 21.2% of applied radioactivity on day 21. The minor degradates were detected at various intervals, none exceeding 5% of the initial concentration of pyrethrin 1 at any point. Mineralisation was a minor degradation pathway (4% CO₂ at day 30).

In the Witte (2007) study, performed in aerobic water/sediment systems (sampled from a pond and a creek), pyrethrin 1 was observed to move rapidly from the water phase into the sediment phase combined with a steadily increasing mineralization to CO₂ (32.0 and 50.7% AR at the end of the test for pond and creek systems, respectively) and breakdown in both aquatic and sediment phases to the metabolite Chrysanthemic acid. A substantial portion of the applied radioactivity became bound to sediment in both test systems. The amounts of bound residues were 29.3 and 39.5 % AR at the end of the test for pond and creek systems, respectively. In the sediment phase, chrysanthemic acid moves back to the water phase as it has a much better solubility and a lower adsorption (compared to pyrethrin 1); hence it is mainly found in the water phase during the test. In the pond system, chrysanthemic acid was detected at a maximum of 48.8% AR (on day 14) and 26.6% AR (on day 2) in the water and sediment phases, respectively, before declining thereafter; and in the total pond water/sediment system, it was detected up to a maximum of 65.6% AR after 14 days. In the creek system, chrysanthemic acid was detected at a maximum of 56.2% AR (on day 1) and 16.6% AR (on day 14) in the water and sediment phases, respectively, before declining thereafter; and in the total water/sediment system, it was detected up to a maximum of 66.8 % AR after 14 days.

- Based on the Robinson and Wisocky (1994) and Witte (2007) tests performed in aerobic aquatic environments, a geometric mean DT₅₀ of 5.27 days was derived at 20°C (i.e. 10 days normalized to 12°C).

Fate and behaviour in soil

Pyrethrin 1 rapidly degrades in soil under aerobic conditions, with a geometric mean half-life of 3.3 days at 20°C. The following five laboratory soil degradation studies were performed to determine the rate of degradation of the substance.

In the Robinson (1994) study, degradation was assessed in a sandy loam soil; a half-life of 2.5 days at 20°C was determined. Degradation seems to proceed initially by a combination of hydrolysis and oxidation to form several low-level metabolites (none formed at >10% AR). Residues in soil are initially extracted, but extended degradation is accompanied by the formation of residues that are bound to soil humus fractions. Soil residues are ultimately bound to soil humus fractions and mineralized (converted to carbon dioxide).

Mineralization and formation of bound residues also occurred in the Hein (2017) study, after pyrethrin 1 was applied onto a loam soil. In this study, a half-life of 2.96 days at 20°C was obtained and no significant metabolites (at >5 %) were detected.

In the studies performed by Fifi (2015a, 2015b and 2015c), pyrethrin 1 exhibited low persistence in the loam, loamy sand and sandy loam soils tested. Half-lives of 4.69, 3.9

and 3 days were determined at 20°C.

Adsorption/desorption study characterizes Pyrethrin 1 as immobile compound in soil according to estimated Koc values in different kinds of soil. Thus it can be expected that pyrethrin 1, once incorporated into the soil, will bind tightly to the soil particles.

Fate and behavior in the environment based on the representative products' use:

The active substance is used as an insecticide in the following representative products: Product A and Product B. Product A is a non-ready to use (non-RTU) ultra-low volume (ULV) aerosol; designed to target flying insects. It is for use by professionals only, as a spot application in indoor areas that are continuously occupied such as hospital wards, residential nursing homes and prisons. Product B is an electric vaporizer mat that provides protection from mosquitoes for about 8 hours. The vaporizer mat is solely for use by non-professionals, in private housings, during night-time hours.

Releases to the environment, from the use of the products, may occur during the preparation (relevant to Product A only), application and cleaning steps. To assess the worst-case use of the products, wet-cleaning has been assessed only; therefore, the STP is considered as the main "receiving compartment". Subsequent receiving compartments in the environment are therefore outdoor air (atmosphere), surface water, sediment, agricultural soil and groundwater. Emissions to these environmental compartments result from the cumulative emission from the preparation, application and/or cleaning steps, carried out indoors, following treatment with the products.

Table 4.4 Summary table of compartments exposed and assessed

Summary table on compartments exposed and assessed		
Compartment	Exposed (Y/N)	Assessed (Y/N)
STP	Y	Y (quantitative)
Surface water/sediment	Y	Y (quantitative)
Soil	Y	Y (quantitative)
Groundwater	Y	Y (quantitative)
Air	Y	Y (qualitative)

As indicated in the previous section, chrysanthemic acid was detected as the only relevant metabolite (present at >10% AR) in the hydrolysis (Selim, 1995) and aerobic water/sediment tests performed (Robinson and Wisocky (1994); Witte (2007)). Since in the Witte (2007) study, the highest maximum formations were obtained, these were considered in the risk assessment as a worst-case approach and are detailed in the following table. In the other studies performed (in air and soil), no major metabolites were detected.

Table 4.5 Summary table on relevant metabolites/degradants

Summary table on relevant metabolites/ degradants		
Metabolite/ degradant/transformation or reaction product	Compartment	% Active Substance
Chrysanthemic acid	Pond water/sediment system	65.6
	Pond water system	48.8
	Pond sediment system	26.6
	Creek water/sediment system	66.8
	Creek water system	56.2
	Creek sediment system	16.6

Input parameters used for calculating the fate and distribution of the active substance in the environment are presented in the table below.

Table 4.6 Summary table of relevant physico-chemical and fate and behaviour parameter of the active substance

Summary table on relevant physico-chemical and fate and behaviour parameter of the active substance – Pyrethrin 1			
	Value	Unit	Remarks
Molecular weight	328.4	g/mol	-
Melting point	142.27	°C	-
Boiling point	400.8	°C	-
Vapour pressure (25°C)	6.9E-05	Pa	-
Water solubility (20°C)	0.23	mg/L	-
Log Octanol/water partition coefficient (Log Kow)	5.59	Log 10	Kow = 389045.14
Organic carbon/water partition coefficient (Koc)	34674	L/kg	Log Koc = 4.54
Henry's Law Constant (25°C)	7.83E-02	Pa/m ³ /mol	
Bioconcentration, aquatic	500	L/kg _{wwt}	Whole body BCF
Biodegradability	Not biodegradable		
DT ₅₀ for hydrolysis	115	d (at 25°C)	At pH 7
DT ₅₀ for photolysis in surface water	11.8	h	
DT ₅₀ for biodegradation in surface water	10.7	d (at 20°C)	Using SFO kinetics DT ₅₀ values ranging from 6.7-10.7 days were obtained (Hein <i>et al.</i> , 2017). The highest DT ₅₀ was used as a worst-case.
DT ₅₀ for biodegradation in sediment	10	d (at 12°C)	5.27 d at 20°C
DT ₅₀ for degradation in air	1.28	hr	
DT ₅₀ for biodegradation in soil	3.3	d (at 20°C)	Geomean value from five different soil studies/half-lives.

Effects assessment

Predicted no effect concentrations (PNEC), used in the environmental risk assessment of the parent compound, are listed in the table below.

Since chrysanthemic acid is several orders of magnitude less toxic than the parent substance (Mantilacci, 2015a), it can be assumed that the PNEC for the metabolite is covered by the PNEC of the parent substance (ENV 3, ECHA TAB 2018). In this case, the PNEC values for freshwater and sediment are the only relevant values for the assessment of chrysanthemic acid – since the metabolite was identified in these compartments only at >10% AR.

Table 4.7 Summary table of calculated PNEC values

Summary table of calculated PNEC values	
Compartment	PNEC
STP	2.3E-02 mg/L
Freshwater	8.6E-05 mg/L
Sediment	83.5 µg/kg _{dw} , equivalent to 1.82E-02 mg/kg _{wwt}
Soil	9E-03 mg/kg _{dw} , equivalent to 7.94E-03 mg/kg _{wwt}
Oral (secondary poisoning)	2.93 mg/kg diet

A. Assessment of intrinsic properties and effects of the active substance

A.1. General substance information

A.1.1. Identity of the substance

Table A.1 Summary table on substance identity

Summary table on substance identity	
Common name (ISO name, synonyms)	<i>Chrysanthemum cinerariaefolium</i> , extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO ₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i> , ext.) pyrethrin 1, pyrethrin 2, cinerin 1, cinerin 2, jasmolin 1, jasmolin 2
Chemical name (EC name, CA name, IUPAC name)	<i>Chrysanthemum cinerariaefolium</i> , extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO ₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i> , ext.) Pyrethrin 1: (Z)-(S)-2-methyl-4-oxo-3-(penta-2,4-dienyl) cyclopent-2-enyl (1R)-trans-2,2-dimethyl-3-(2-methylprop-1-enyl) cyclopropane carboxylate Cinerin 1: (Z)-(S)-3-(but-2-enyl)-2-methyl-4-oxocyclopent-2-enyl (1R)-trans-2,2-dimethyl-3-(2-methylprop-1-enyl) cyclopropane carboxylate Jasmolin 1: (Z)-(S)-2-methyl-4-oxo-3-(pent-2-enyl) cyclopent-2-enyl (1R)-trans-2,2-dimethyl-3-(2-methylprop-1-enyl) cyclopropane carboxylate Pyrethrin 2: (Z)-(S)-2-methyl-4-oxo-3-(penta-2,4-dienyl) cyclopent-2-enyl (E)-(1R)-trans-3-(2-methoxycarbonylprop-1-enyl)-2,2-dimethylcyclopropane carboxylate Cinerin 2: (Z)-(S)-3-(but-2-enyl)-2-methyl-4-oxocyclopent-2-enyl (E)-(1R)-trans-3-(2-methoxycarbonylprop-1-enyl)-2,2-dimethylcyclopropane carboxylate Jasmolin 2: (Z)-(S)-2-methyl-4-oxo-3-(pent-2-enyl)cyclopent-2-enyl (E)-(1R)-trans-3-(2-methoxycarbonylprop-1-enyl)-2,2-dimethylcyclopropane carboxylate

EC number	<p><i>Chrysanthemum cinerariaefolium</i>, extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i>, ext.): 289-699-3</p> <p><i>Chrysanthemum cinerariaefolium</i>: 289-699-3</p> <p>Pyrethrins: 232-319-8</p> <p>Pyrethrin 1: 204-455-8</p> <p>Pyrethrin 2: 204-462-6</p> <p>Cinerin 1: 246-948-0</p> <p>Cinerin 2: 204-454-2</p> <p>Jasmolin 1: not available</p> <p>Jasmolin 2: not available</p>
CAS number	<p><i>Chrysanthemum cinerariaefolium</i>, extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO₂ (Redefined from Pyrethrins and Pyrethroids and <i>Chrysanthemum cinerariaefolium</i>, ext.): 89997-63-7</p> <p><i>Chrysanthemum cinerariaefolium</i>: 89997-63-7</p> <p>Total Pyrethrins: 8003-34-7</p> <p>Pyrethrin 1: 121-21-1</p> <p>Pyrethrin 2: 121-29-9</p> <p>Cinerin 1: 246-948-0</p> <p>Cinerin 2: 121-20-0</p> <p>Jasmolin 1: 4466-14-2</p> <p>Jasmolin 2: 1172-63-0</p>
other CAS numbers (e.g. deleted, related, preferred, alternate)	<p>CIPAC: Pyrethrins: 32</p> <p>EU Index: Pyrethrins: 613-022-00-6</p>
Molecular formula	<p>pyrethrin 1: C₂₁H₂₈O₃</p> <p>cinerin 1: C₂₀H₂₈O₃</p> <p>jasmolin 1: C₂₁H₃₀O₃</p> <p>pyrethrin 2: C₂₂H₂₈O₅</p> <p>cinerin 2: C₂₁H₂₈O₅</p> <p>jasmolin 2: C₂₂H₃₀O₅</p>
Molecular weight or molecular weight range	<p>pyrethrin 1: 328.4 g/mol</p> <p>cinerin 1: 316.4 g/mol</p> <p>jasmolin 1: 330.4 g/mol</p>

pyrethrin 2: 372.4 g/mol
 cinerin 2: 360.4 g/mol
 jasmolin 2: 374.4 g/mol

Information on optical activity and typical ratio of (stereo) isomers (if applicable and appropriate) -

Description of the manufacturing process and identity of the source (for UVCB substances only) This information can be found in the Appendix VI: Confidential Information.

Degree of purity (%) 100% w/w


Conclusion on the identity of the active substance

Chrysanthemum cinerariaefolium extract from supercritical CO₂ (synonym: Pyrethrum Extract) is an extract of the flower heads of *Chrysanthemum cinerariaefolium*. It contains Pyrethrins, which may be divided into the two groups: Pyrethrins I (consisting of pyrethrin 1, cinerin 1, and jasmolin 1) and Pyrethrins II (consisting of pyrethrin 2, cinerin 2 and jasmolin 2).

The technical concentrate (TK) includes the presence of solvent to reduce the viscosity of the extract to make it easier to pour and mix during formulation, and also helps to keep the additive in solution (additive is added to avoid oxidation of the pyrethrins). eCA Spain, ECHA and MSs agreed that the solvent should not be part of the active substance, since the stability study shows that *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ remains stable without the solvent, showing only a slight decrease in stability, which is not enough to support the inclusion of the solvent in the composition of the active substance. This decision is based on the legal definition of "substance" established in article 3(2) of the BPR, which excludes any solvent that may be separated from the substance without affecting the stability or changing its composition. Therefore, the solvent should not be considered as part of the substance.


Chrysanthemum cinerariaefolium extract from supercritical CO₂, as pure active substance, has a minimum purity of 100% w/w. It is placed in the market as a solution, technical concentrate (c.a. 50% pyrethrins).

Table A.2 Structural formula


Structural formula

pyrethrin 1


pyrethrin 2




cinerin 1



cinerin 2



jasmolin 1



jasmolin 2

Table A.3 Origin of the natural active substance or precursor(s) of the active substance

Origin of the natural active substance or precursor(s) of the active substance

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is an extract of the flower heads of *Chrysanthemum cinerariaefolium*. It contains Pyrethrins, which may be divided into the two groups: Pyrethrins I (consisting of pyrethrin 1, cinerin 1, and jasmolin 1) and Pyrethrins II (consisting of pyrethrin 2, cinerin 2, and jasmolin 2).

A.1.2. Composition of the substance (reference specifications)

Table A.4 Constituents

Main constituent(s)					
Constituent (chemical name)	Typical concentration (%(w/w))	Concentration range (%(w/w))	Current CLH in Annex V I Table 3.1 (CLP)	Current self-classification and labelling (CLP)	Remarks / Discussion
<i>Chrysanthemum cinerariaefolium</i> , extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with supercritical CO ₂ (<i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂)	100% w/w	-	-	<p>Acute Tox., Category 4; H302: Harmful if swallowed</p> <p>Acute Tox., Category 4; H332: Harmful if inhaled</p> <p>Skin sensitisation, Category 1B; H317: May cause an allergic skin reaction</p> <p>Short-term (acute) aquatic hazard, Category 1; H400: Very toxic to aquatic life</p> <p>Long-term (chronic) aquatic hazard, Category 1; H410: Very toxic to aquatic life with long lasting effects</p>	Refer to the Appendix VI: Confidential Information for further details.

Table A.5 Impurities

This information can be found in the Appendix VI: Confidential Information.

Table A.6 Additives

The information can be found in the Appendix VI: Confidential Information.

Table A.7a Concentration of constituents in batches used for (eco)toxicity studies and proposed specification

All details are included in the relevant tables within Section A for this report.

Table A.8b Concentration of constituents in batches used for (eco)toxicity studies and proposed specification

All details are included in the relevant tables within Section A for this report.

A.1.3. Physical and chemical properties of the active substance

Table A.9 Physical and chemical properties of the active substance

Property	Result	Test method applied or description in case of deviation	Remarks / Discussion / Justification for waiving	References
Aggregate state at 20°C and 101.3 kPA	-	-	-	-
Physical state (appearance) at 20°C and 101.3 kPA	Liquid	US EPA 63-3	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins)	Sundquist D.L., 1995 (BRA, MGK and SCJ)
	Clear Liquid	EPA/OCSP 830.6303	Pure active substance (82.6% pyrethrins)	Comb, T., 2021 (BRA)
Colour at 20°C and 101.3 kPA	Clear, Dark Orange	US EPA 63-2	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins)	Sundquist D.L., 1995 (BRA, MGK and SCJ)
	Yellow (10Y 9/8 on Munsell colour system)	EPA/OCSP 830.6302	Pure active substance (82.6% pyrethrins)	Comb, T., 2021 (BRA)
Odour at 20°C and 101.3 kPA	Aromatic Solvent-like Odour	US EPA 63-4	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins)	Sundquist D.L., 1995 (BRA, MGK and SCJ)
	No discernible odour	EPA/OCSP 830.6304	Pure active substance (82.6% pyrethrins)	Comb, T., 2021 (BRA)

Melting / freezing point	Pyrethrin 1: 142.3°C Pyrethrin 2: 132.1°C Jasmolin 1: 143.1 °C Jasmolin 2: 133.3 °C Cinerin 1: 135.2 °C Cinerin 2: 124.2 °C	EPIWIN QSAR	Determined by structure analysis. As the active substance is a plant extract, <u>an experimental determination for the purified active substance is not possible.</u>	(BRA, MGK and SCJ)
Boiling point	Pyrethrin 1: 400.8°C Pyrethrin 2: 421.8°C Jasmolin 1: 401.7 °C Jasmolin 2: 422.7 °C Cinerin 1: 390.1 °C Cinerin 2: 411.1 °C	EPIWIN QSAR	Determined by structure analysis. As the active substance is a plant extract, <u>an experimental determination for the purified active substance is not possible.</u>	(BRA, MGK and SCJ)
	>210°C	-	Values derives from MSDS (BRA, MGK and SCJ). In addition, decomposition of the active substance as manufactured (ca 50% pyrethrins) starts at temperatures above 270°C.	-
Acidity	Not applicable.	-	Study not required. Not relevant, the active substance is not water-solvent based.	-
Relative density	0.952 g/mL at 20°C	US EPA 63-7	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins)	Sundquist D.L., 1995 (BRA, MGK and SCJ)
	D ²⁰ ₄ = 1.01	EC Method A.3, OECD Method 109 and EPA/OCSPP 830.7300	Pure active substance (82.6% pyrethrins)	Comb, T., 2021 (BRA)

Absorption spectra data (UV/VIS, IR, NMR) and a mass spectrum	<p>UV-Vis absorption</p> <p>The test substance is not a pure material; therefore, it is impossible to calculate the molar extinction coefficients for the absorbance maxima. The contribution of each component in the test substance to each absorbance maximum, which is located between 225 and 230 nm - could not be separated.</p>	US EPA OPPTS 830.705	Premium PYROCIDE 175 (20%), batch No. 4AE10421 (isopropanol solution)	Sinning D.J., 2002 (BRA, MGK and SCJ)
	<p>UV-Vis absorption</p> <p>Ethanol solvent:</p> <p>Pyrethrin 1: max 225 nm, Epsilon 36420</p> <p>Cinerin 1: max 226 nm; Epsilon 17720</p> <p>Pyrethrin 2: max 229 nm; Epsilon 45850</p> <p>Cinerin 2: 227-9 nm; Epsilon 28946</p> <p>n-hexane solvent:</p> <p>Pyrethrin 1: max 222 nm</p> <p>Cinerin 1: max 220 nm</p> <p>Pyrethrin 2: max 227-9 nm</p> <p>Cinerin 2: max 226-8 nm</p>	Not stated	Natural Pyrethrins purified using a Celite column; Purity: Not specified; Batch number: Not specified	Chang S.C., 1961 (BRA, MGK and SCJ)
	<p>UV-Vis absorption</p> <p>Pyrethrin 1: max 222.5 nm; Epsilon 38800</p>	Not stated	Main biologically active components pyrethrins separated from commercial Pyrethrum extracts; Purity: Not specified;	Sawicki R.M. & Thain E.M., 1961 (BRA, MGK and SCJ)

	<p>Cinerin 1: max 221 nm; Epsilon 21100</p> <p>Pyrethrin 2: max 228 nm; Epsilon 47500</p> <p>Cinerin 2: max 229 nm; Epsilon 27900</p>		<p>Batch number: Not specified (n-hexane solutions)</p>	
	<p>IR Absorbance</p> <p>Pyrethrin 1: 2929 cm⁻¹: C-H asym; 1735 cm⁻¹: C=O; 1655 cm⁻¹: C=C; 1148 cm⁻¹: C-O stretch; 910 cm⁻¹: C-H out of the plane</p> <p>Cinerin 1: 2929 cm⁻¹: C-H asym; 2880 cm⁻¹: C-H sym; 1735 cm⁻¹: C=O; 1657 cm⁻¹: C=C; 1148 cm⁻¹: C-O stretch</p> <p>Jasmolin 1: 2932 cm⁻¹: C-H asym; 2882 cm⁻¹: C-H sym; 1735 cm⁻¹: C=O; ~1650 cm⁻¹: C=C; 1148 cm⁻¹: C-O stretch</p> <p>Pyrethrin 2: 2980 cm⁻¹: C-H asym; 1735 cm⁻¹: C=O; 1655 cm⁻¹: C=C; 1327, 1220, 1150 and 1055: C-O stretch; 910 cm⁻¹: C-H out of the plane</p> <p>Cinerin 2: 2936 cm⁻¹: C-H asym; 2886 cm⁻¹: C-H sym; 1735 cm⁻¹: C=O; 1657 cm⁻¹: C=C; 1148 cm⁻¹: C-O stretch</p> <p>Jasmolin 2: 2976 cm⁻¹: C-H asym; 2889 cm⁻¹: C-H sym; 1735 cm⁻¹: C=O; ~1650 cm⁻¹: C=C; 1148 cm⁻¹: C-O stretch</p>	<p>Not stated</p>	<p>Natural pyrethrum extract (19.26%); Batch number: Not specified (isopropanol solutions)</p>	<p>Moorman R. & Nguyen K.T., 1997 (BRA, MGK and SCJ)</p>

	<p>IR Absorbance Pyrethrins I and II: 905 cm⁻¹: =CH₂ terminal, CH out of the plane; Absorption at 800-1800 cm⁻¹.</p>	<p>Not stated</p>	<p>Pyrethrum extracts 25% (Belgian Congo and Kenya); Batch number: Not specified Method used: Not stated</p>	<p>Sawicki R. M. & Thain E.M., 1961 (BRA, MGK and SCJ)</p>
	<p>IR Absorbance Pyrethrin 1: 1703 cm⁻¹: C=O 1282, 1227, 1198 and 1149 cm⁻¹: C-O stretch 909 cm⁻¹: CH out of the plane 855 cm⁻¹: CH out of the plane Cinerin 1: 1703 cm⁻¹: C=O 1282, 1227, 1198 and 1149 cm⁻¹: C-O stretch 855 cm⁻¹: CH out of the plane Pyrethrin 2: 1703 cm⁻¹: C=O 1605 cm⁻¹: C=C 1266, 1220, 1170,1149 cm⁻¹: C-O stretch 909 cm⁻¹: CH out of the plane 756 cm⁻¹: CH out of the plane Cinerin 2: 1703 cm⁻¹: C=O 1605 cm⁻¹: C=C 1266, 1220, 1170, 1149 cm⁻¹: C-O stretch 756 cm⁻¹: CH out of the plane</p>	<p>Not stated</p>	<p>Purified natural pyrethrins and reconstituted esters; Purity: Not specified; Batch number: Not specified</p>	<p>Elliott M., 1961 (BRA, MGK and SCJ)</p>
	<p>The spectra of the NMR analysis of Pyrethrin 1 and 2 indicate that the structures of purified Pyrethrins are consistent with the proposed</p>	<p>Not stated</p>	<p>Natural pyrethrins separated from World Standard Pyrethrum Extract (1974); Purity: Not specified; Batch number: Not specified</p>	<p>Dickinson C.M., 1982 (BRA, MGK and SCJ)</p>

	<p>structures.</p> <p>MS Pyrethrin 1: m/z 168, 162, 133, 123, 105, 91 Cinerin 1: m/z 168, 150, 123, 107, 105, 93, 91 Jasmolin 1: m/z 164, 162, 133, 123, 107, 105, 93, 91 Pyrethrin 2: m/z 167, 160, 145, 133, 119, 107, 105, 91 Cinerin 2: m/z 167, 149, 121, 107, 105, 93, 91 Jasmolin 2: m/z 167, 162, 133, 121, 107, 105, 93, 91</p>	Not stated	9.26% pure Natural pyrethrum extract; Batch number: Not specified	Moorman R. & Nguyen K.T., 1997 (BRA, MGK and SCJ)
	<p>MS Pyrethrin 1: Ester moiety m/z 329 [MH+] m/z 357 [M+C₂H₅]+ Acid moiety m/z 169 [RC(OH)₂]+ m/z 151 [RCO]+ m/z 123 [R]+ Alcohol moiety m/z 189 [R+ C₂H₄]+ m/z 161 [R]+ Cinerin 1: Ester moiety m/z 317 [MH]+</p>	Not stated	Pyrethrins from purified 72% Pyrethrum extract; Batch number: Not specified	Class, T.J. <i>et al.</i> , 1989 (BRA, MGK and SCJ)

<p>m/z 345 [M+C₂H₅]⁺ Acid moiety m/z 169 [RC(OH)₂]⁺ m/z 151 [RCO]⁺ m/z 123 [R]⁺ Alcohol moiety m/z 177 [R+ C₂H₄]⁺ m/z 149 [R]⁺ Jasmolin 1: Ester moiety m/z 331 [MH]⁺ m/z 359 [M+C₂H₅]⁺</p> <p>Acid moiety m/z 169 [RC(OH)₂]⁺ m/z 151 [RCO]⁺ m/z 123 [R]⁺ Alcohol m/z 191 [R+ C₂H₄]⁺ m/z 163 [R]⁺ Pyrethrin 2: Ester moiety m/z 373 ester [MH]⁺ m/z 401 [M+C₂H₅]⁺ m/z 341 [MH-CH₃OH]⁺ Acid moiety m/z 213 [RC(OH)₂]⁺ m/z 195 acid [RCO]⁺</p>			
--	--	--	--

<p>m/z 167 acid [R]+ Alcohol moiety m/z 189 [R+ C₂H₄]+ m/z 161 [R]+ Cinerin 2: Ester moiety m/z 361 [MH]+ m/z 389 [M+C₂H₅]+ m/z 329 [MH-CH₃OH]+ Acid moiety m/z 213 [RC(OH)₂]+ m/z 195 [RCO]+ m/z 167 [R]+</p> <p>Alcohol m/z 177 [R+ C₂H₄]+ m/z 149 [R]+ Jasmolin 2: Ester moiety m/z 375 [MH]+ m/z 403 [M+C₂H₅]+ m/z 343 [MH-CH₃OH]+ Acid moiety m/z 213 [RC(OH)₂]+ m/z 195 [RCO]+ m/z 167 [R]+ Alcohol moiety m/z 165 [R+ C₂H₄]+ m/z 163 [R]+</p>			
--	--	--	--

			No determination will be performed using purified active substance, as the data provided relating to the individual pyrethrins is not impacted by the presence or absence of other components.	(BRA)
Vapour pressure	Pyrethrins: 3.13E-05 Pa Pyrethrin 1: 1.88E-04 Pa Pyrethrin 2: 1.21E-05 Pa Cinerin 1: 6.93E-05 Pa Cinerin 2: 2.77E-05 Pa Jasmolin 1: 2.92E-05 Pa Jasmolin 2: 1.12E-05 Pa Temperature: 20°C	EpiSuite MPBPWIN, US EPA, 2008		O'Carroll N., 2008 (BRA, MGK and SCJ)
Henry's constant	law Pyrethrin 1: 0.268 Pa m ³ /mol Pyrethrin 2: 0.0626 Pa m ³ /mol Temperature: 20°C	Calculated		O'Carroll N., 2008 (BRA, MGK and SCJ)
				No determination will be performed using purified active substance, as no new vapour pressure or water solubility data are being generated.
Surface tension			The water solubility of the experimental and estimated values of the known constituents (>90%) indicate that a surface tension test is not required.	(BRA, MGK and SCJ)

	55.0 mN/m at 20 °C	EC Method A.5 and OECD Method 115 (harmonised ring method)	Pure active substance (82.6% pyrethrins) 90% saturated test item solution used	Comb, T., 2021 (BRA)
Water solubility at 20°C	Determined at 20°C Pyrethrin 1: 0.23 mg/L Pyrethrin 2: 0.072 mg/L Estimated for 25°C Jasmolin 1: 0.027 mg/L Cinerin 1: 0.085 mg/L Jasmolin 2: 0.094 mg/L Cinerin 2: 0.301 mg/L	US EPA D-63-8, OPPTS 830.7840	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins) No determination will be performed using test material without solvent, as the active substance is a plant extract. An experimental determination for the purified active substance is therefore not possible. Data for the individual pyrethrins has already been provided.	Sinning D.J., 2005 (BRA, MGK and SCJ) (BRA)
Partition coefficient (n-octanol/water) and its pH dependency	Log Pow (n-octanol/water): Pyrethrin 1: 5.59 Cinerin 1: 5.54 Jasmolin 1: 6.04 Pyrethrin 2: 4.32 Cinerin 2: 4.26 Jasmolin 2: 4.74	OECD 117 (HPLC method)	Reference standard (≥ 97%) Lots: XX8-82-P1 XX8-82-C1 XX8-82-J1 XX8-82-P2 XX8-82-C2 XX8-82-J2	Mori V., 2015 (KPIC) (BRA, MGK and SCJ)

	Pyrethrin 1: 5.9 Pyrethrin 2: 4.3 Cinerin 1: 5.6 Estimated at 25 °C Jasmolin 1: 6.42 Jasmolin 2: 5.47 Cinerin 2: 4.98	Not stated	Pyrethrum Extract (purity and lot No. not stated) Effect of pH is not relevant as the test substance is neither acidic nor basic.	Briggs G.G. <i>et al.</i> , 1982 (BRA, MGK and SCJ)
			No determination will be performed using purified active substance, as the active substance is a plant extract. <u>An experimental determination for the purified active substance is therefore not possible.</u> Data for the individual pyrethrins has already been provided.	(BRA)

Thermal stability and identity of breakdown products	Thermally stable at room temperature below 150°C. Oxidative decomposition underwent at temperatures above 270°C.	OECD Guideline 113 (Differential Scanning Calorimetry)	<p>Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins)</p> <p>The results from tests in air showed the initiation of predominant exothermic effects at temperatures above 270°C indicating significant decomposition of the test substance. The decomposition was subsequently confirmed by the presence of a black residue on completion of the tests.</p> <p>In comparison, the results from the tests in nitrogen showed only a slight increase in the baseline which was consistent with the heating of the apparatus characteristically observed for all samples using the specified apparatus. It should be noted that the scale of the thermal assessment of the nitrogen samples is much smaller than the traces presented for the air samples.</p> <p>The conclusion from these observations was that the sample predominantly underwent oxidative decomposition at temperatures above 270°C.</p>	Comb A.L., 2008 (BRA, MGK and SCJ)
--	--	--	---	------------------------------------

Reactivity towards container material	Diluted pyrethrum extract was demonstrated to react with metals iron, copper, zinc and lead. Tin did not react with the pyrethrum extract.	Not stated	Pyrethrum Extract (25%) refined Pale product in Shellsol T	Chiu F.T. & Wu N.C., 1974 (BRA, MGK and SCJ)
	Stable for 1 year at 37.8°C in F-Style Tin and FL-HDPE. Stable for 1 year at room temperature in amber glass.	US EPA 63-17	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins)	Sundquist D. L., 1996 (BRA, MGK and SCJ)
Dissociation constant	Not applicable.	-	Pyrethrins are not able to dissociate in water.	-
Granulometry	Not applicable.	-	The active substance is a liquid.	-
Viscosity	85.5 mPa·s at 23 °C	US EPA 63-18	(Active substance as manufactured, technical concentrate, TK) (ca 50% pyrethrins)	Sundquist D.L., 1995 (BRA, MGK and SCJ)
	650 mPa.s at 20°C (shear rates approx. 25 – 75 rpm) 76 mPa.s at 40°C (shear rates approx. 50 – 150 rpm)	OECD Method 114, EPA/OCSP 830.7100 and CIPAC MT 192 (rotational viscometer)	Pure active substance (82.6% pyrethrins) As there was no significant change in the viscosity readings with shear rate, it can be concluded that the test item is a Newtonian liquid	Comb, T., 2021 (BRA)
Solubility in organic solvents, including effect of temperature on solubility	Hexane: >65 g/L Methanol: 45 g/L Xylenes: >56 g/L Acetone: > 65 g/L Dichloromethane: >65 g/L	Not stated. Approximately 10 g of test substance was mixed with 100 mL of solvent. The solutions were allowed to stand at 20°C overnight. The	Active substance as manufactured (technical concentrate, TK) (ca 50% pyrethrins) Different values observed on the solubility in organic solvents may be due to the different methods	Faber H., 2004 (BRA, MGK and SCJ)

	Temperature: 20°C	resulting solution was filtered through 0.45-micro filter. A 10 mL alicuot was prepared for analysis by heating to remove the solvent by evaporation. The residual was re-dissolved in hexane. Pyrethrins were determined by HPLC.	used by the applicants.	
Stability in organic solvents used in biocidal products and identity of relevant degradation products	Stable in isopropanol, petroleum distillate and propylene glycol at 5% for 2 weeks at 54°C	CPT-2164	5.0% pure Pyrethrum Concentrate; Batch number: Not specified	Bergman J.T., 2007 (BRA, MGK and SCJ)

Conclusion on the physical, chemical and technical properties of the active substance

Physico-chemical properties of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ were initially determined on the whole extract (i.e. in the active substance as manufactured (technical concentrate), with the solvent present). The solvent reduces the viscosity of the extract to make it easier to pour and mix during formulation, and also helps to keep the stabilizer in solution (stabilizer is added to avoid oxidation of the pyrethrins). However, the solvent should not be part of the active substance, since the stability study shows that *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ remains stable without the solvent, showing only a slight decrease in stability, which is not enough to support the inclusion of the solvent in the composition of the active substance.

Therefore, some physico-chemical properties (appearance – physical state, colour and odour -, relative density, surface tension and viscosity) have been tested again using the pure extract, without the solvent. Other physico-chemical endpoints are already sufficiently addressed or are technically not possible to determine using the pure active substance.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ (pure active substance) is a yellow liquid with no discernible odour. Its relative density D_4^{20} is 1.01 and its surface tension is 55.0 mN/m at 20 °C. The viscosity is 650 mPa.s at 20°C and 76 mPa.s at 40°C and it can be considered a Newtonian liquid.

Physico-chemical properties are, in general, determined on the whole extract, and not on individual constituents.

A.1.4. Physical hazards and respective characteristics

Table A.10 Physical hazards and respective characteristics

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
Explosives	-	-	<p>The molecular structure of Pyrethrins indicates that the substance has little or no explosive properties.</p> <p>However, this waiver is not acceptable. Structural considerations are insufficient for UVCB substances to conclude on this physical hazard.</p>	(BRA, MGK and SCJ)

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
	UN Recommendations on the Transport of Dangerous Goods, 7 th Ed., 2019, Appendix 6	DSC Screening determination for heat of decomposition	<p>Pure active substance (82.6% pyrethrins)</p> <p>DSC Exotherms: Exotherm onset temp (°C) 1st: 132.09 2nd: 252.86 Heat of decomposition (J/g) 1st: 56.069 2nd: 128.27</p> <p><i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ is not a candidate for classification as a UN Class 1 explosive substance as the total heat of decomposition was <500 J/g</p>	Siusiene, E., 2022 (BRA)
Flammable gases	-	-	Not applicable	-
Flammable aerosols	-	-	Not applicable	-
Oxidising gases	-	-	Not applicable	-
Gases under pressure	-	-	Not applicable	-
Flammable liquids	ASTM D 56 - Non-equilibrium ISO 1523 - Equilibrium	-	<p>The flash point of the active substance is 72°C, measured with the non-equilibrium method (ASTM D56) and 71°C with the equilibrium method (ISO 1523).</p> <p>(Active substance as manufactured, technical concentrate, TK) (ca 50% pyrethrins)</p>	Bergman J.T., 2008 (BRA, MGK and SCJ)

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
	EU Method A.9 (Flash Point)	Pensky-Martens apparatus (closed cup)	Pure active substance (82.6% pyrethrins) Flash point = 130.5 °C <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ is not flammable.	Siusiene, E., 2022 (BRA)
Flammable solids	-	-	Not applicable	-
Self-reactive substances and mixtures	-	-	The study does not need to be conducted because there are no chemical groups present in the molecule which are associated with explosive or self-reactive properties and hence the classification procedure does not need to be applied. This waiver is not acceptable. Structural considerations are insufficient for UVCB substances to conclude on this physical hazard.	-

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
	UN Recommendations on the Transport of Dangerous Goods, 7 th Ed., 2019, Appendix 6	DSC Screening determination for heat of decomposition	<p>Pure active substance (82.6% pyrethrins)</p> <p>DSC Exotherms: Exotherm onset temp (°C) 1st: 132.09 2nd: 252.86 Heat of decomposition (J/g) 1st: 56.069 2nd: 128.27</p> <p><i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ is not a candidate for classification as a UN Class 4, Division 4.1 self-reactive substance as the total heat of decomposition was <300 J/g</p>	Siusiene, E., 2022 (BRA)
Pyrophoric liquids	-	-	The study does not need to be conducted because the substance is known to be stable in contact with air at room temperature for prolonged periods of time (days) and hence the classification procedure does not need to be applied.	-
Pyrophoric solids	-	-	Not applicable	-
Self-heating substances and mixtures	-	-	Not applicable as the substance is a liquid.	-

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
Substances and mixtures which in contact with water emit flammable gases	-	-	The study does not need to be conducted because the organic substance does not contain metals or metalloids and hence the classification procedure does not need to be applied.	-
	UN Test N.5	Spontaneous ignition with water/evolution of flammable gas > 1L/kg/hr	Pure active substance (82.6% pyrethrins) No gas generation or spontaneous ignitions observed. Classified as not UN Division 4.3	Siusiene, E., 2022 (BRA)
Oxidising liquids	-	-	The molecular structure of Pyrethrins indicates that there is little or no potential for this material as an oxidising or reducing agent. However, this waiver is not acceptable. Structural considerations are insufficient for UVCB substances to conclude on this physical hazard.	(BRA, MGK and SCJ)
	UN Test O.2	Mean pressure rise	Pure active substance (82.6% pyrethrins) The 1:1 mixture of the test item and cellulose was observed to have a mean pressure rise time greater than that of a 1:1 mixture of 65 % nitric acid and cellulose. The test item is therefore exempt from classification as an oxidising liquid of UN Class 5, Division 5.1	Siusiene, E., 2022 (BRA)

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
Oxidising solids	-	-	Not applicable	
Organic peroxides	-	-	This study does not need to be conducted because the substance does not fall under the definition of organic peroxides according to GHS and the relevant UN Manual of Tests and Criteria.	-
Corrosive to metals	-	-	The study does need to be conducted since based on the chemical evaluation none of the components contain chemical groups, which could initiate an irreversible electrochemical reaction with metals leading to significant damage or destruction. However, this waiver is not acceptable, as it is not based on CLP.	-
	UN Test C.1	Corrosion to Metals (mass loss & pitting using steel and aluminium coupons)	Pure active substance (82.6% pyrethrins) The percentage mass losses on steel and aluminium were found to be <13.5 % over 7 days and no pitting was observed. <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ is therefore exempt from classification as a corrosive substance of UN Class 8, Packing group III (according to the UN Transport of Dangerous Goods Recommendations)	Siusiene, E., 2022 (BRA)

Hazard class / characteristics	Guideline and Method	Parameter(s)	Results / Waiver	Reference
Auto-ignition temperature (liquids and gases)			The product is not expected to be flammable according to the flash point and the explosives and oxidising properties. This hazard class is not correctly addressed. A test on the purified active substance should be conducted.	(MGK, BRA and SCJ)
	EU Method A.15 (Auto-Ignition Temperature (Liquids and Gases))		Pure active substance (82.6% pyrethrins) The autoignition temperature of <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ has been determined to be 308°C.	Siusiene, E., 2022 (BRA)
Relative self-ignition temperature for solids	-	-	Not applicable	-
Dust explosion hazard	-	-	Not applicable	-

Conclusion on the physical hazards and respective characteristics of the active substance

Physical hazards of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ were initially determined on the whole extract (i.e. in the active substance as manufactured (technical concentrate), with the solvent present). However, the solvent is not part of the active substance composition. Furthermore, some physical hazards were waived based on the chemical structure of the active substance, but structural considerations in UVCB substances are not acceptable.

Therefore, physical hazard tests have been repeated on the purified active substance, without the solvent.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ (pure active substance) is not considered explosive nor self-reactive. It is not flammable nor oxidising. It does not emit flammable gases in contact with water. It is not corrosive to metals.

In conclusion, no physical hazard has been identified for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ (pure active substance).

A.1.5. Assessment of physical hazards according to the CLP criteria**A.1.5.1. Assessment of physical hazards**

Chrysanthemum cinerariaefolium extract from supercritical CO₂ (pure active substance) does not meet any physical hazard class described in BPR.

A.2. Assessment of effects on Human Health

A.2.1. Toxicokinetics

A2.1.1 Short summary and overall relevance of the provided toxicokinetic information

Toxicokinetics and distribution

¹⁴C-Pyrethrin was administered orally to male and female Sprague-Dawley rats. In a preliminary study time peak blood level and blood half-life were determined. In three definitive experiments, absorption, distribution, excretion and metabolism were examined (Selim, 1995). The definitive studies were conducted using three groups of 5 rats/sex/group as follows: a single low-dose (Group 1: 10 mg/kg bw), a single high-dose (Group 2: 50 mg/kg bw for females and 100 mg/kg for males) and a repeated low-dose (Group 3: non-radiolabelled Pyrethrin 1 at 10 mg/kg bw for 14 days prior to administration of a single oral low-dose of ¹⁴C-Pyrethrin 1 on day 15). Two additional groups of 5 rats/sex/group were dosed exclusively for the purposes of isolation and identification of metabolites. In this ¹⁴C-Pyrethrin 1 was administered orally at a single oral low-dose (Group 4: 10 mg/kg bw) and at a single oral high dose (Group 5: 50 mg/kg bw for females and 100 mg/kg for males).

All rats survived the studies and no signs of toxicity were observed. The preliminary study showed that less of the administered ¹⁴C-Pyrethrin 1 was systemically absorbed by the male rats than by the female rats with peaks in blood after 5-6 and 6-8 hours, respectively. Male rats excrete the ¹⁴C-Pyrethrin 1 derived radioactivity faster than females, since the elimination half-time in males and females was 5.3 and 6.7 hours, respectively.

In all groups of the definitive studies the majority of radioactivity was excreted in the urine and faeces during the first 72 hours following administration. The levels of radioactivity expired as ¹⁴CO₂ were very low. Regardless of dosage regimen, males excrete a majority of the administered radioactivity in the faeces via the enterohepatic circulation while females excrete approximately equal amounts in the urine and faeces. Repeated administration of Pyrethrin 1 increased the rate of elimination of the ¹⁴C-Pyrethrin 1-derived radioactivity from the body in both males and females thus suggesting that repeated dose of ¹⁴C-Pyrethrin 1 results in an induction of the liver microsomal enzyme system.

Analysis of tissues at 7 days showed that in all dosing regimens, concentrations of radioactivity more than twice that of whole blood were found in liver, ovaries, carcass and fat. The radioactive residues in the fat were consistently higher than the residues in all other tissues analysed. The elimination of substantial amounts of ¹⁴C-Pyrethrin 1-derived radioactivity in the faeces over and extended length of time indicated that enterohepatic circulation played a role in the elimination of the compound and/or its metabolites from the body.

Metabolism

In the metabolism study, two additional groups of male and female rats were orally dosed with ¹⁴C-Pyrethrin 1 at single low-dose and single high-dose for isolation and identification of metabolites in urine and faeces. This study demonstrated that ¹⁴C-Pyrethrin 1 is metabolized to six compounds, having been identified two major metabolites (Metabolite C and E) along with 4 minor metabolites (A, B, D, and F). The results indicated that in the rat, Pyrethrin 1 is metabolized by two major metabolic pathways. One pathway is by oxidation of the double bond on the cyclopentene or the cyclopropane side of the molecule to form a diol, in addition to oxidation of the methyl groups on the side chain of the cyclopropane ring to form a carboxylic acid. The second pathway is by hydrolysis of the ester bond to form the corresponding acid and alcohol.




Figure 1: Proposed metabolic pathway for pyrethrin 1 in the rat

Dermal absorption

Two studies were provided.

██████████ A single dose, open label study to investigate the absorption and excretion of orally administered or dermally applied [¹⁴C]-labelled pyrethrin I (PI) to healthy male volunteers.

The clinical proportion of the study was conducted in compliance with "Declaration of Helsinki" (as revised in Edinburgh, Scotland, October 2000); the rules of good clinical practice U.S. FDA (protection of human subjects, 21 CFR, Ch. 1, Part 500; Institutional review boards, 21 CFR, Ch. 1, part 56; Investigational New Drug Application, 21 CFR, Ch.1, Part 312) and Rules of Good Clinical Practice European Community (Good clinical practice for trials on medicinal products in the European Community, III/3976/88-EN-Final).

The analytical part of the study was conducted in compliance with the FDA Good Laboratory Practice Regulation (GLP) as set forth in Title 21 of the U.S. Code of federal regulations, Part 58, issued December 22, 1978 (and all applicable revisions) and the applicable regulations of the OECD Principles of Good Laboratory Practice (as revised in 1997), ENV/MC/CHEM (98) 17, OECD, Paris, 1998.

The study protocol was reviewed and approved by an Independent Ethics Committee and written informed consents were obtained by all eligible volunteers. The study is therefore valid.

The fogger being tested contained pyrethrin as the active substance and co-formulants, the main of which is added to pyrethrin products to inhibit insects' microsomal enzymes

that detoxify pyrethrins.

This study was conducted in 4 human male volunteers. The test material was applied to the volar aspect of the non-dominant forearm and the application site was covered with a non-occlusive aluminium dome, which allowed free movement of air but helped prevent loss of radioactivity due to physical contact with the application site, secured in place with an adhesive bandage. The test material remained in contact with the skin for approximately 8 hours. After exposure, the protective devices were removed, and the skin was wiped with cotton swabs dipped in 2% soapy water. The application site was dried with two cotton swabs, and then swabbed with two cotton swabs dipped in isopropyl alcohol (IPA), followed by four cotton swabs dipped in soapy water and four swabs dipped in IPA. One sixth of the dose site was also stripped with tape approximately 45 hours after the removal of test material. The purpose of the tape stripping was to determine the amount residual radioactivity associated with the surface layer of the skin. The skin site was also swabbed with cotton swabs dipped in IPA on Day 5. All urine and faeces excreted by the volunteers were collected for five days following dose administration. The protective enclosures, swabs, rinses, gauze, tape strips, urine and faeces were analysed for total radioactivity in order to determine mass balance and the percent of administered dose that was absorbed. In order to obtain information on the onset, rate and duration of absorption, venous blood samples were collected from the ipsilateral and contralateral veins during and after the 8-h exposure period; after separation, plasma samples were analysed for total radioactivity.

For the dermal dosing solution, MGK provided the testing facility with the blank of a typical fogger formulation containing all non-radiolabelled components but the active substance, similar to the commercial formulation. Radiolabelled pyrethrins I (PI) was added to the blank of the fogger formulation and mixed well to provide a homogeneously labelled formulation. The test substance formulation containing [¹⁴C]-PI was administered dermally at approximately 50 µL of [¹⁴C]-PI at a mean dose of 12.25 µg/cm² (the dose was administered over an area of 4x6 cm or 24 cm²). The concentration of PI in the fogger formulation (dosing formulation) was 0.3 mg per 50 mg (0.6%). Each 50 mg of fogger formulation contained approximately 50 µCi of radioactivity. Volunteer identification, body weight and dose information for volunteers is shown below.

Volunteer Identification	Volunteer Body weight (kg)	Total µCi Applied	Total µg PI Applied*	Total PI Applied (µg/kg)	Dose formulation Administered (mg)
01 ER	65.4	44.2	280	4.28	47.6
02 PR	81.7	44.6	282	3.45	48.0
03 CH	75.1	47.2	299	3.98	47.8
04 HB	91.3	49.6	314	3.44	50.3
Mean	78.4	46.4	294	3.79	48.43
± Standard deviation	10.91	2.51	15.97	0.41	1.26

* Calculated based on specific activity

In the dermally dosed subjects, no elevation in radioactivity was seen in the contralateral and ipsilateral plasma samples, reflecting the very slow dermal absorption of the radiolabelled PI. This was confirmed by the very low percent of dosed radioactivity (0.22%) that crossed the skin and was excreted in urine. Urinary excretion was rapid, with the highest percent of dosed radioactivity excreted between 12 and 24 hours. The overall mean recovery of radioactivity was 104.28%. The majority of the applied radioactivity was accounted for on the surface of the skin, mainly the swabs, skin rinses, protective enclosure and gauze with a mean of 104.28% of the applied radioactivity. The first four soapy water dermal swabs removed a mean of 88.03% of the applied radioactivity. The remaining dermal swabs removed a mean of 12.06% of the applied radioactivity. Tape stripping and swabbing of 1/6 of the dosed area removed a mean of 0.06% of applied radioactivity, indicating there is no accumulation of radioactivity in the skin.

Since a good material balance was achieved and only 0.22% of the applied radioactivity was recovered in the excreta, the dermal absorption of PI was determined to be 0.22%. A summary of the data is shown in the tables below.

Time-course of urinary excretion Percent of applied radioactivity						
Time (hours)	Subject Number				Mean	Standard deviation
	01	02	03	04		
Pre-dose	0.00	0.00	0.00	0.00	0.00	0.00
0-4	0.00	0.00	0.01	0.02	0.01	0.01
4-8	0.04	0.03	0.04	0.06	0.04	0.01
8-12	0.05	0.04	0.04	0.06	0.05	0.01
12-24	0.09	0.09	0.06	0.08	0.08	0.01
24-36	0.08	0.08	0.00	0.00	0.04	0.05
36-48	0.02	0.00	0.00	0.00	0.01	0.01
48-60	0.00	0.00	0.00	0.00	0.00	0.00
60-72	0.00	0.00	0.00	0.00	0.00	0.00
72-84	0.00	0.00	0.00	0.00	0.00	0.00
84-96	0.00	0.00	0.00	0.00	0.00	0.00
96-120	0.00	0.00	0.00	0.00	0.00	0.00
Subtotal	0.28	0.24	0.15	0.22	0.22	0.05

Percent of applied radioactivity excreted in urine following dermal application of [¹⁴ C]-pyrethrin						
	Subject Number				Mean	Standard deviation
	01	02	03	04		
Subtotal	0.28	0.24	0.15	0.22	0.22	0.05
Percent of applied radioactivity excreted in faeces following dermal application of [¹⁴ C]-pyrethrin						
	Subject Number				Mean	Standard deviation
	01	02	03	04		
Subtotal	0.00	0.00	0.00	0.00	0.00	0.00
Percent of applied dose recovered in tape-stripping						
	Subject Number				Mean	Standard deviation
	01	02	03	04		
Subtotal	0.04	0.1	0.05	0.05	0.06	0.03
Percent of applied radioactivity recovered from skin surface						
	Subject Number				Mean	Standard deviation
	01	02	03	04		
Duoderm	0.74	0.61	0.32	0.82	0.62	0.22
Aluminium dome	0.48	0.74	1.61	0.44	0.82	0.54
Dermal swabs	104.06	100.05	98.71	97.55	100.09	2.84
Skin rinse	1.88	3.54	1.47	1.88	2.19	0.92
Gauze	0.30	0.45	0.16	0.18	0.27	0.13
Tape strips	0.04	0.10	0.05	0.05	0.06	0.03
Subtotal	107.5	105.49	102.32	100.92	104.06	2.99

Total recovery of radioactivity following dermal application of [¹⁴ C]-pyrethrin						
	Subject Number				Mean	Standard deviation
	01	02	03	04		
Surface radioactivity	107.5	105.49	102.32	100.92	104.06	2.99
Urine	0.28	0.24	0.15	0.22	0.22	0.05
Faeces	0.00	0.00	0.00	0.00	0.00	0.00
Total	0.28	0.24	0.15	0.22	0.22	0.05

Human *in vivo* Percutaneous absorption of pyrethrin and piperonyl butoxide. Fd Chem. Toxic. Vol. 32, No 1, pp. 51-53.

This study was conducted in six male volunteers from whom informed consent was obtained. The [¹⁴C] formulation, similar to the commercial formulation, was spread on the ventral forearm and after 30 minutes the application site was washed with soap and water. Urine was collected on day-1 in divided interval (0-4, 4-8, 8-12, 12-25 hours) and on days 2-7 as total daily excretions. The commercial formulation used contained 0.3% pyrethrin. [¹⁴C]-pyrethrin (3.8 mCi/mmol) was applied at 5.25 µCi (487 µg) in a dosing volume of 200 µL spread over 88 cm² = 5.53 µg/cm². Percutaneous absorption was determined by ¹⁴C urinary excretion, and the following formula: dose absorbed = (total ¹⁴C urinary excretion of topical dose in man ÷ total ¹⁴C urinary excretion of parenteral dose in monkey) x100. In monkeys, 22.5% of pyrethrins were excreted in urine following a parenteral dose. The percutaneous absorption of pyrethrin from the ventral forearm was calculated to be about 2% (1.9% ± 1.2), as shown in the table below.

Subject	Percentage absorbed
1	1.4
2	1.6
3	2.0
4	0.6
5	1.6
6	4.1
Mean ± SD	1.9 ± 1.2

A2.1.2 Values and conclusions used for the risk assessment

Value(s) used in the Risk Assessment – Oral absorption	
Value(s)	100%
Justification for the selected value(s)	Default absorption value of 100% used. Fast absorption occurred with >90% systematically absorbed in 5 to 8 hours (██████████).

Value(s) used in the Risk Assessment – Dermal absorption	
Value(s) **	0.3%
Justification for the selected value(s)	Evaluation of the Human volunteer dermal absorption studies highlighted several deficiencies in the study conducted by ██████████. The clinical portion of the study was not conducted according to good clinical practice, and the analytical component of the study was not conducted in accordance with GLP standards. The exposure period of the study was just 30 minutes, and the study was not a mass-balance study. Instead, the dermal absorption values were determined by comparing urinary excretion values from the human volunteers with urinary excretion values obtained from monkeys parenterally administered pyrethrins (exact route not specified). No analytical data were reported. In contrast, the study conducted by ██████████ was

	<p>conducted according to good clinical practice and GLP standards. The study was a mass-balance study that reported good recovery values with low variability between volunteers. The volunteers were dermally exposed for eight hours under controlled conditions and the analytical data are presented.</p> <p>As [REDACTED] was conducted in vivo in human volunteers, there should be no need to adjust the obtained dermal absorption value of 0.22%. However, as it cannot be excluded that the tested formulation differs from the commercial product(s) containing pyrethrins, results are re-evaluated in line with the EFSA guidance on dermal absorption (EFSA Journal 2017;15(6):4873) here below.</p> <p>In [REDACTED] absorption of pyrethrins is complete i.e. more than 75% excretion in urine occurs within half of the study period. The low residue removed by tape stripping can, therefore, be excluded for deriving the dermal absorption value.</p> <p>The preferred approach to addressing variability between replicates is to add a multiple of the standard deviation to the mean value. Although variability among the four volunteers is low (<25%), considering that 4 valid replicates are available, the multiplication factor is 1.6.</p> <p>Therefore, the final dermal absorption value for pyrethrins from biocidal product(s) is estimated to be 0.3% (i.e. 0.22 + (0.05 x1.6) %).</p> <p>A dermal absorption value of 0.3% is considered appropriate for the purposes of human health risk assessment for biocidal products containing pyrethrins as the active substance.</p>
--	--

**The dermal absorption value is applicable for the active substance and might not be usable in product authorization.

Value(s) used in the Risk Assessment – Inhalatory absorption	
Value(s)	100%
Justification for the selected value(s)	Default worst-case value.

A.2.2. Acute toxicity / STOT SE

Oral Acute Toxicity

Independent from the ratio of Pyrethrins I and Pyrethrins II, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ exhibited low and moderate acute oral toxicity to male and female rats, respectively. Transient clinical signs of toxicity in surviving animals were hyperactivity, ruffled fur and tremors ([REDACTED]). (KPIC)

Dose [mg/kg] Total pyrethrins/ Extract	Number of dead / number of investigated	Time of death (range)	Observations
229	-	-	
355	-	-	
560	-	-	
710	-	-	
794	-	-	
891/1365	0/5	-	All animals appeared normal throughout the 14-day observation period.
1410/2160	0/5	-	In the beginning of the test period one animal exhibited dark nasal staining. All the others appeared normal until the end of the test phase.
2230/3417	2/5	day 1	In the beginning appearance of ruffled fur and tremors. After 1 day two animals were found dead, after the second day the remaining animals appeared normal.
2660/4075	4/5	day 1	Initially 2 of 5 animals exhibited tremors, after 1 day four animals were found dead.
3550/5439	5/5	day 1	After one day all animals found dead without any clinical signs after immediately treatment and 4 h later on.
LD₅₀ value	2140 mg/kg bw (1730 - 2640 mg/kg bw) total pyrethrins; 3269 mg/kg bw (2651 - 4045 mg/kg bw) extract		

Dose [mg/kg] Total pyrethrins/ Extract	Number of dead / number of investigated	Time of death (range)	Observations
229/351	0/5	-	All animals appeared normal throughout the 14-day observation period.
355/544	1/5	1 day	After 1 day one rat was found dead, whereas all the other animals appeared normal throughout the observation period.
560/858	1/5	1 day	After 4 h all animals exhibited tremors, one day later three of them appeared normal, one was found dead and one exhibited dark nasal and and ocular staining.
710/1088	1/5	4 h	After 4 h one animal was found dead, the other exhibited tremors. After one day all animals appeared normal throughout the test period.
794/1217	3/5	4 h-1 day	After 2 h three animals exhibited tremors, after 4 h one was found dead. Two other rats were found dead after 1 day, the rest appeared normal throughout the test period.
891/1365	5/5	4 h-1 day	After 4 h three of the test animals were already found dead, the rest have died one day after treatment.
1410	-	-	
2230	-	-	
2660	-	-	
3550	-	-	
LD₅₀ value	700 mg/kg bw (500 – 990 mg/kg bw) total pyrethrins; 1073 mg/kg bw (766 – 1517 mg/kg bw) extract		

Oral acute toxicity study was conducted in Sprague-Dawley rats. *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was dosed to 6 groups of males (5/group) at concentrations between 710 and 5000 mg/kg bw total pyrethrins (1088 and 7661 mg/kg bw extract) and 7 groups of females (5/group) at concentration between 316 and 2000 mg/kg bw total pyrethrins (484 and 3064 mg/kg bw extract). In males, no deaths were recorded at the 710 and 891 mg/kg bw total pyrethrins (1088 and 1365 mg/kg bw extract) dose levels. There were 2 deaths at the 2230 mg/kg bw total pyrethrins (3417 mg/kg bw extract) dose level, 3 deaths at the 2660 mg/kg bw total pyrethrins (4076 mg/kg bw extract) dose level, 4 deaths at the 3550 mg/kg bw total pyrethrins (5439 mg/kg bw extract) dose level and 5 deaths at the 5000 mg/kg bw total pyrethrins (7661 mg/kg bw extract) dose level. In females, no deaths were recorded at the 316, 500 and 794 mg/kg bw total pyrethrins (484, 766, and 1217 mg/kg bw extract) dose levels. There were 2 deaths at the 944 mg/kg bw total pyrethrins (1446 mg/kg bw extract) dose level, 3 deaths at the 1120 mg/kg bw total pyrethrins (1716 mg/kg bw extract) dose level and 5 deaths at the 1260 and 2000 mg/kg bw total pyrethrins (1931 and 3064 mg/kg extract) dose levels. Clinical signs were observed in male rats at a dose level greater than 710 mg/kg bw total pyrethrins (1088 mg/kg bw extract) (ruffling and tremors) and in female groups at a dose level greater than 316 mg/kg bw total pyrethrins (484 mg/kg bw extract) (tremors and hyperactivity). All surviving animals gained weight throughout the study. At necropsy, no internal abnormalities were observed at doses of 891 mg/kg bw total pyrethrins (1365 mg/kg bw extract) or less for males and 794 mg/kg bw total pyrethrins (1217 mg/kg bw extract) or less for females. At doses higher than 891/794 mg/kg bw total pyrethrins (1365/1217 mg/kg bw extract) male/female observations such as muzzle staining, coloured material in the lower gastrointestinal tract and haemorrhagic lungs were noted. The oral LD₅₀ of Pyrethrum extract was found to be 2370 (1680 - 3350) mg/kg bw total pyrethrins (3631 (2574 - 5133) mg/kg bw extract) for males and 1030 (860 - 1240) mg/kg bw total pyrethrins (1578 (1318 - 1900) mg/kg bw extract) for females. The no effect level for clinical signs was 710 mg/kg bw total pyrethrins (1088 mg/kg bw extract) in males and 316 mg/kg bw total pyrethrins (484 mg/kg bw extract) in females (██████████). (BRA, MGK and SCJ)

Mortality data					
Dose (mg/kg bw) Total pyrethrins/ Extract	Males Mortality	Time of death – days (no of animals)	Dose (mg/kg bw) Total pyrethrins/ Extract	Females Mortality	Time of death - days (no of animals)
710/1088	0/5	-	316/484	0/5	-
891/1365	0/5	-	500/766	0/5	-
2230/3417	2/5	1 (2)	794/1217	0/5	-
2660/4076	3/5	1 (3)	944/1446	2/5	1 (2)
3550/5439	4/5	1 (3), 2 (1)	1120/1716	3/5	1 (3)
5000/7661	5/5	1(5)	1260/1931	5/5	0 (1), 1 (4)
-	-	-	2000/3064	5/5	0 (1), 1 (4)
LD ₅₀ value	2370 mg/kg bw total pyrethrins 3631 mg/kg bw extract		1030 mg/kg bw total pyrethrins 1578 mg/kg bw extract		

Dermal Acute Toxicity

The acute dermal toxicity of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was investigated in rabbits. A single dose of active ingredient was applied to the skin of 5 New Zealand White rabbit/sex at a concentration of 2000 mg/kg bw total pyrethrins (3064 mg/kg bw extract) using an occlusive material. After a 24-hour dermal exposure period to *Chrysanthemum cinerariaefolium* extract from supercritical CO₂, male and female rabbits presented very slight to well defined erythema, slight oedema and a stained test site. All males appeared normal after 6 days and all females after 9 days. All animals gained weight during the study. No internal abnormalities were observed at gross necropsy in any animal. The percutaneous LD₅₀ of *Chrysanthemum cinerariaefolium* extract from

supercritical CO₂ was found to be in excess of 2000 mg/kg bw total pyrethrins (3064 mg/kg bw extract) (). (KPIC and BRA, MGK and SCJ)

Dose (mg/kg)	Males Mortality	Time of death – days (no of animals)	Dose (mg/kg)	Females Mortality	Time of death - days (no of animals)
2000 LD ₅₀ value	0/5 > 2000 mg/kg bw total pyrethrins >3064 mg/kg bw extract	-	2000 LD ₅₀ value	0/5 > 2000 mg/kg bw total pyrethrins >3064 mg/kg bw extract	-

Inhalation Acute Toxicity

In order to assess the acute inhalation toxicity, 4 groups of 5 Sprague-Dawley rats/sex/group were exposed to *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ for 4 hours at the following analytical concentrations: 0 (acetone control), 0.69, 2.1 and 4.6 mg/L total pyrethrum (0, 1.06, 3.2, and 7.1 mg/L extract). *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was administered as a liquid aerosol by the inhalation route, using acetone as a vehicle. In males, no deaths were recorded at 0, 0.69 and 2.1 mg/L total pyrethrins (0, 1.06, and 3.2 mg/L extract) dose levels. There were 2 deaths at 4.6 mg/L total pyrethrins (7.1 mg/L extract) dose level. In females, there were no deaths recorded at 0 and 0.69 mg/L total pyrethrins (0 and 1.06 mg/L extract) dose levels. 2 rats died at 2.1 mg/L total pyrethrins (3.2 mg/L extract) dose level and 4 rats died at 4.6 mg/L total pyrethrins (7.1 mg/L extract) dose level. During exposure the animals exhibited laboured breathing, excessive salivation, decreased activity and eye closure. Among the acetone-control group, similar effects were observed with an additional excessive lacrimation and nasal discharge. However, tremors were also noted during the higher-level exposures. These observations were noted immediately following exposure but after several days the signs decreased significantly. All groups of animals lost weight on the day following treatment. However, recovery of weight occurred over time and all surviving animals were in excess of weight at the end of the study. Reddening of the lung and turbinates and lung oedema were the major post-mortem findings which were considered to be related to exposure to *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. The LC₅₀ was calculated to be 3.4 mg/L total pyrethrins (5.2 mg/L extract) for the combined sexes (2.3-4.9) (). (KPIC and BRA, MGK and SCJ)

Group	Concentration (mg/L)		Mortality			LC ₅₀ (14d)
	Nominal	Analytical	Males	Females	Total	Dose (mg/L) Total pyrethrins/ Extract
I	14	4.6 ± 0.5 ¹⁾	3/5	4/5	7/10	3.4 (2.3-4.9) 5.2 (3.5-7.5) ³⁾
II	- ²⁾	-	0/5	0/5	0/10	
III	5.6	2.1 ± 0.4	0/5	2/5	2/10	
IV	0.92	0.69 ± 0.08	0/5	0/5	0/10	

¹⁾ mean ± standard deviation

²⁾ vehicle control

³⁾ 95 % confidence limits

A.2.2.1. Acute oral toxicity

Table A.11 Summary table of animal studies on acute oral toxicity

Summary table of animal studies on acute oral toxicity						
Method, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity) Vehicle, Dose levels, Type of administration (gavage, in diet, other)	Signs of toxicity (nature, onset, duration, severity, reversibility, include concentrations)	Value LD ₅₀	Remarks (e.g. major deviations)	Reference
US EPA 81-1, OECD 401 GLP Reliability 1 Key	Rat Sprague-Dawley male + female 5 per dose/sex	<p>Pyrethrum Extract (FEK-99; 57.574%)</p> <p>Male: 891, 1410, 2230, 1660, and 3550 mg/kg bw total pyrethrins; 1365, 2160, 3417, 4075, and 5439 mg/kg bw extract</p> <p>Female: 229, 355, 560, 710, 794, and 891 mg/kg bw total pyrethrins; 351, 544, 858, 1088, 1217, and 1365 mg/kg bw extract</p> <p>Single dose (gavage)</p>	Transient clinical signs of toxicity in surviving animals were hyperactivity, ruffled fur and tremors.	<p>Male: 2140 mg/kg bw (1730-2640 mg/kg bw) total pyrethrins; 3269 mg/kg bw (2651-4045 mg/kg bw) extract</p> <p>Female: 700 mg/kg bw (500-990 mg/kg bw) total pyrethrins; 1073 mg/kg bw (766-1517 mg/kg bw) extract</p>	-	(KPIC) IIIA6.1.1


<p>EPA OPP 81-1 and OECD Guideline 401 GLP Reliability 1 Key</p>	<p>Rat Sprague-Dawley Male/Female 5/sex/group</p>	<p>Pyrethrum extract (FEK-99; 57.03%) Males: 710, 891, 2230, 2660, 3550, and 5000 mg/kg total pyrethrins; 1088, 1365, 3417, 4076, 5439, and 7661 mg/kg bw extract Females: 316, 500, 794, 944, 1120, 1260, and 2000 mg/kg total pyrethrins; 484, 766, 1217, 1446, 1716, 1931, and 3064 mg/kg bw extract Single dose (gavage)</p>	<p>Males >710 mg/kg bw total pyrethrins (1088 mg/kg bw extract): rufflings and tremors Females >316 mg/kg bw total pyrethrins (484 mg/kg bw extract): tremors and hyperactivity. Females >794 and Males >891 mg/kg bw (1217 and 1365 mg/kg bw extract respectively): muzzle staining, coloured material in the lower gastrointestinal tract and haemorrhagic lungs</p>	<p>Males: 2370 (1680 - 3350) mg/kg bw total pyrethrins; 3631 (2574 - 5133) mg/kg bw extract Females: 1030 (860 - 1240) mg/kg bw total pyrethrins; 1578 (1318 - 1900) mg/kg bw extract</p>		<p> (BRA, MGK and SCJ) IIIA 6.1.1</p>
--	---	---	--	---	--	--

Table A.12 Summary table of human data on acute oral toxicity

No data are available.

Table A.13 Summary table of other studies relevant for acute oral toxicity

No data are available.

A2.2.1.1 Short summary and overall relevance of the provided information on acute oral toxicity

Chrysanthemum cinerariaefolium extract from supercritical CO₂ exhibited low and moderate acute oral toxicity to male and female rats, respectively.

A2.2.1.2 Comparison with the CLP criteria

Classified as Acute Tox. 4, (H302: Harmful if swallowed) because, according to the CLP regulation 3.1.2.1. (Table 3.1.1), 300 mg/kg bw < LD₅₀ ≤ 2000 mg/kg bw.

A2.2.1.3 Conclusion on classification and labelling for acute oral toxicity

Acute toxicity Category 4, H302: Harmful if swallowed. ATE oral = 700 mg/kg bw total pyrethrins; ATE oral = 1073 mg/kg bw extract.

A2.2.1.4 Conclusion on acute oral toxicity related to risk assessment

Value used in the Risk Assessment – Acute oral toxicity	
Value	700 mg/kg bw total pyrethrins; 1073 mg/kg bw extract. Acute Tox. 4, H302: Harmful if swallowed
Justification for the selected value	Lowest LD ₅₀ value

A.2.2.2. Acute dermal toxicity

Table A.14 Summary table of animal studies on acute dermal toxicity

Summary table of animal studies on acute dermal toxicity						
Method, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), Vehicle, Dose levels, Surface area	Signs of toxicity (nature, onset, duration, severity, reversibility, include concentrations)	Value LD ₅₀	Remarks (e.g. major deviations)	Reference
EPA OPP 81-2 and OECD Guideline 402 GLP Reliability 1 Key	Rabbit New Zealand White Male/Female 5/sex/group	Pyrethrum Extract (FEK-99; 57.467%) 2000 mg/kg bw total pyrethrins; 3064 mg/kg bw extract Occlusive 24 h	No systemic effects. Males and females: slight-well defined erythema, slight edema and stained test site. Recovering after several days.	>2000 mg/kg bw total pyrethrins; >3064 mg/kg bw extract	-	■■■■■■ IIIA6.1.2 (KPIC) (BRA, MGK and SCJ)

Table A.15 Summary table of human data on acute dermal toxicity

No data are available.

Table A.20 Summary table of other studies relevant for acute dermal toxicity

No data are available

A2.2.2.1 Short summary and overall relevance of the provided information on acute dermal toxicity

Dermal application of 2000 mg/kg bw total pyrethrins (3064 mg/kg bw extract) did not cause death or systemic clinical signs of toxicity.

A2.2.2.2 Comparison with the CLP criteria

Not classified because, according to the CLP regulation 3.1.2.1. (Table 3.1.1), LD₅₀ > 2000 mg/kg bw.

A2.2.2.3 Conclusion on classification and labelling for acute dermal toxicity

Substance does not meet the criteria to be classified in this hazard class.

A2.2.2.4 Conclusion on acute dermal toxicity related to risk assessment

Value used in the Risk Assessment – Acute dermal toxicity	
Value	>2000 mg/kg bw total pyrethrins; >3064 mg/kg bw extract
Justification for the selected value	Highest dose tested.

A.2.2.3. Acute inhalation toxicity

Table A.21 Summary table of animal studies on acute inhalation toxicity


Summary table of animal studies on acute inhalation toxicity							
Method, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), form (gas, vapour, dust, mist) and particle size (MMAD) Actual and nominal concentration, Type of administration (nose only / whole body/ head only)	Signs of toxicity (nature, onset, duration, severity, reversibility, include concentrations)	Value LC ₅₀	Remarks (e.g. major deviations)	Reference	
US EPA 81-3, OECD 403 GLP Reliability 1 Key	Rat Sprague-Dawley male + female 5 per dose/sex	Pyrethrum Extract (FEK-99; 57.574%) MMAD = 2.6 µm ± 2.2 µm 0, 0.69, 2.1, and 4.6 mg/L total pyrethrins; 0, 1.06, 3.2, and 7.1 mg/L extract analytical concentration 4 h – Whole body	Post-mortem findings were: reddening of the lung and turbinates and lung oedema.	LC ₅₀ males: 3.9 mg/L (2.1-7.2 mg/L with 95% confidence limit) total pyrethrins; 6.0 mg/L (3.2-11.0 mg/L with 95% confidence limit) extract LC ₅₀ females: 2.5 mg/L (1.5-4.3 mg/L with 95% confidence limit) total pyrethrins; 3.8 mg/L (2.3-6.6 mg/L with 95% confidence limit) extract LC ₅₀ both sexes: 3.4 mg/L (2.3-4.9 mg/L with 95% confidence limit) total pyrethrins; 5.2 mg/L (3.5-7.5 mg/L with 95% confidence limit) extract	-	 (KPIC) (BRA, MGK and SCJ) IIIA6.1.3	

Table A.16 Summary table of human data on acute inhalation toxicity
No data are available.

Table A.17 Summary table of other studies relevant for acute inhalation toxicity
No data are available.

A2.2.3.1 Short summary and overall relevance of the provided information on acute inhalation toxicity

Inhalation exposure to different concentrations of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ as an aerosol to rats resulted in a combined LC₅₀ (both sexes) of 3.4 mg/L total pyrethrins (5.2 mg/L extract).

A2.2.3.2 Comparison with the CLP criteria

Classified as Category 4: Acute Tox. 4 (H332: Harmful if inhaled) because, according to the CLP 3.1.2.1. (Table 3.1.1), 1.0 mg/L < LC₅₀ ≤ 5.0 mg/L according to the toxicity test result.

A2.2.3.3 Conclusion on classification and labelling for acute inhalation toxicity

Acute toxicity Category 4 H332: Harmful if inhaled. ATE inhalation = 2.5 mg/L total pyrethrins (dusts and mists). ATE inhalation = 3.8 mg/L extract (dusts and mists).

A2.2.3.4 Conclusion on acute inhalation toxicity related to risk assessment

Value used in the Risk Assessment – Acute inhalation toxicity	
Value	2.5 mg/L total pyrethrins. 3.8 mg/L extract. Acute Tox. 4, H332: Harmful if inhaled.
Justification for the selected value	Lowest LC ₅₀ value.

A.2.2.4. Specific target organ toxicity – single exposure Category 1 and 2 (STOT SE 1 and 2)

A2.2.4.1 Short summary and overall relevance of the provided information on STOT SE 1 and 2

Effects observed in oral acute toxicity study were in males, no deaths at the 710 and 891 mg/kg bw total pyrethrins (1088 and 1365 mg/kg bw extract) dose levels. There were 2 deaths at the 2230 mg/kg bw total pyrethrins (3417 mg/kg bw extract) dose level, 3 deaths at the 2660 mg/kg bw total pyrethrins (4076 mg/kg bw extract) dose level, 4 deaths at the 3550 mg/kg bw total pyrethrins (5439 mg/kg bw extract) dose level and 5 deaths at the 5000 mg/kg bw total pyrethrins (7661 mg/kg bw extract) dose level. In females, no deaths were recorded at the 316, 500 and 794 mg/kg bw total pyrethrins (484, 766 and 1217 mg/kg bw extract) dose levels. There were 2 deaths at the 944 mg/kg bw total pyrethrins (1446 mg/kg bw extract) dose level, 3 deaths at the 1120 mg/kg bw total pyrethrins (1716 mg/kg bw extract) dose level and 5 deaths at the 1260 and 2000 mg/kg bw total pyrethrins (1931 and 3064 mg/kg bw extract) dose levels. Clinical signs were observed in male rats at a dose level greater than 710 mg/kg bw total pyrethrins (1088 mg/kg bw extract) (ruffling and tremors) and in female groups at a dose level greater than 316 mg/kg bw total pyrethrins (484 mg/kg bw extract) (tremors and hyperactivity). All surviving animals gained weight throughout the study. At necropsy, no internal abnormalities were observed at doses of 891 mg/kg bw total pyrethrins (1365 mg/kg bw extract) or less for males and 794 mg/kg bw total pyrethrins (1217 mg/kg bw extract) or less for females. At doses higher than 891/794 mg/kg bw total pyrethrins (1365/1217 mg/kg bw extract) male/female observations such as muzzle staining, coloured material in the lower gastrointestinal tract and haemorrhagic lungs were noted. The oral LD₅₀ of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was found to be 2370 (1680 - 3350) mg/kg bw total pyrethrins (3631 (2574 - 5133) mg/kg bw

extract) for males and 1030 (860 - 1240) mg/kg bw total pyrethrins (1578 (1318 - 1900) mg/kg bw extract) for females. The no effect level for clinical signs was 710 mg/kg bw total pyrethrins (1088 mg/kg bw extract) in males and 316 mg/kg bw total pyrethrins (484 mg/kg bw extract) in females (██████████). (BRA, MGK and SCJ)

Effects observed in dermal acute toxicity study were very slight to well defined erythema, slight oedema and a stained test site in males and females. All males appeared normal after 6 days and all females after 9 days. All animals gained weight during the study. No internal abnormalities were observed at gross necropsy in any animal. The percutaneous LD₅₀ of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was found to be in excess of 2000 mg/kg bw total pyrethrins (3064 mg/kg bw extract) (██████████). (BRA, MGK and SCJ)

Effects observed in inhalation acute toxicity study were in males, no deaths at 0, 0.69 and 2.1 mg/L total pyrethrins (0, 1.06 and 3.2 mg/L extract) dose levels. There were 2 deaths at 4.6 mg/L (7.1 mg/L extract) dose level. In females, there were no deaths recorded at 0 and 0.69 mg/L total pyrethrins (0 and 1.06 mg/L extract) dose levels. 2 rats died at 2.1 mg/L total pyrethrins (3.2 mg/L extract) dose level and 4 rats died at 4.6 mg/L total pyrethrins (7.1 mg/L extract) dose level. During exposure the animals exhibited laboured breathing, excessive salivation, decreased activity and eye closure. Among the acetone-control group, similar effects were observed with an additional excessive lacrimation and nasal discharge. However, tremors were also noted during the higher-level exposures. These observations were noted immediately following exposure but after several days the signs decreased significantly. All groups of animals lost weight on the day following treatment. However, recovery of weight occurred over time and all surviving animals were in excess of weight at the end of the study. Reddening of the lung and turbinates and lung oedema were the major post-mortem findings which were considered to be related to exposure to *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. The LC₅₀ was calculated to be 3.4 mg/L total pyrethrins (5.2 mg/L extract) for the combined sexes (2.3-4.9 mg/L total pyrethrins (3.5-7.5 mg/L extract)) (██████████). (BRA, MGK and SCJ)

Regarding the effects observed in the oral, dermal and inhalation toxicity studies, all of them are not organ-specific. Indeed the specific findings are those typical of the route of exposure (effects in the gastrointestinal or respiratory tracts or in the skin) or related with systemic toxicity.

In a neurotoxicity study, groups of 15 male Sprague-Dawley rats received by gavage a 10% w/v solution of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ in corn oil at doses of 0, 40, 125 or 400 mg/kg bw total pyrethrins (0, 61, 192, and 613 mg/kg bw extract), and 15 females received a 5% w/v solution of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ in corn oil at doses of 0, 20, 63 or 200 mg/kg bw total pyrethrins (0, 31, 97, and 306 mg/kg bw extract).

Five males and two females at the high dose died on the day of treatment, and a variety of acute neurological signs were observed in the other animals at this dose, including tremors, urogenital area wetness, salivation, perinasal encrustation, exaggerated startle response, decreased grip strength, hind leg splay, and increased body temperature. Tremors were also observed in three females at the intermediate dose. Measurements of motor activity on the day of treatment indicated increased fine movement and decreased rearing and ambulation in animals of each sex at the high dose and decreased fine movement, rearing and ambulation in males as the intermediate dose. This is likely due to an effect of treatment and a predisposition for lower activity of this group if compared to the control group during pre-treatment evaluation.

In addition, slight, statistically non-significant decreases in body weight were seen in males at the high dose on days 7 and 14. There was no evidence of any gross, treatment-related lesion. The microscopic changes were limited mainly to sections of the sciatic nerve and its branches. The histomorphological changes within the peripheral nerve sections indicated the presence of scattered degenerating nerve fibres or myelin sheaths. These changes were seen in only a few animals, were graded as minimal, and were not dose-

related.

Males		Females	
Dosage (mg/kg total pyrethrins (mg/kg bw extract))	Number surviving /number initiated	Dosage (mg/kg total pyrethrins (mg/kg bw extract))	Number surviving /number initiated
0 (0)	15/15	0 (0)	15/15
40 (61)	15/15	20 (31)	15/15
125 (192)	15/15	63 (97)	15/15
400 (613)	10/15	200 (306)	13/15

Time point	0 mg/kg bw total pyrethrins 0 mg/kg bw extract	40 mg/kg bw total pyrethrins 61 mg/kg bw extract	125 mg/kg bw total pyrethrins 192 mg/kg bw extract	400 mg/kg bw total pyrethrins 613 mg/kg bw extract
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD
0 hour	200.47 ± 11.259	201.07 ± 12.411	199.26 ± 10.176	195.81 ± 11.173
3 hours	239.88 ± 15.132	239.99 ± 14.168	238.98 ± 11.009	230.40 ± 14.976
7 days	281.94 ± 18.795	282.53 ± 20.391	281.99 ± 16.150	269.27 ± 19.463
14 days	320.53 ± 23.108	319.69 ± 24.069	318.91 ± 20.345	304.24 ± 27.035

S.D. - Standard deviation
0 hour - Pretreatment

Time point	0 mg/kg bw total pyrethrins 0 mg/kg bw extract	20 mg/kg bw total pyrethrins 31 mg/kg bw extract	63 mg/kg bw total pyrethrins 97 mg/kg bw extract	200 mg/kg bw total pyrethrins 306 mg/kg bw extract
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD
0 hour	146.30 ± 8.146	144.67 ± 10.838	142.98 ± 8.952	146.52 ± 7.365
3 hours	162.73 ± 9.563	162.22 ± 11.541	158.7 ± 9.985	162.02 ± 7.818
7 days	182.91 ± 11.666	181.95 ± 13.632	177.99 ± 14.047	179.8 ± 8.081
14 days	198.57 ± 15.492	200.49 ± 17.886	193.51 ± 17.053	194.94 ± 10.208

S.D. - Standard deviation
0 hour - Pretreatment

FEEDING LEVEL (mg/kg bw total pyrethrins (mg/kg bw extract))	0 (0)	40 (61)	125 (192)	400 (613)	0 (0)	20 (31)	63 (97)	200 (306)
	MALES				FEMALES			
NUMBER IN GROUP	15	15	15	15	15	15	15	15
SCIATIC NERVE								
myelin degeneration/myelin sheath swelling	0	0	0	2	0	0	0	1
minimal focal	0	0	0	2	0	0	0	1
myelin/axon degeneration	0	2	1	1	0	0	0	4
minimal focal	0	2	1	1	-	-	-	-
minimal multifocal	0	1	1	1	0	0	0	3
moderate multifocal	0	1	0	0	0	0	0	1
NERVE PERONEAL								
myelin/axon degeneration	-	-	-	-	0	1	0	2

minimal multifocal	-	-	-	-	0	1	0	1
moderate multifocal	-	-	-	-	0	0	0	1
NERVE TIBIAL								
myelin/axon degeneration	-	-	-	-	0	0	0	2
minimal multifocal	-	-	-	-	0	0	0	1
moderate multifocal	-	-	-	-	0	0	0	1

The NOEL/NOAEL was 40 mg/kg bw total pyrethrins (61 mg/kg bw extract) for male rats and 20 mg/kg bw total pyrethrins (31 mg/kg bw extract) for female rats (██████████). (KPIC and BRA, MGK and SCJ)

Taking into account that the observed effects are not specific for any organ, active substance does not classify for this hazard class.

A2.2.4.2 Comparison with the CLP criteria

No data are available to indicate that the active substance should be classified for STOT SE 1 or 2. Observed effects are not organ-specific and do not compromise the normal function of any organ as stated in CLP 3.8.2.1.7.3.

A2.2.4.3 Conclusion on classification and labelling for STOT SE 1 and 2

Chrysanthemum cinerariaefolium extract from supercritical CO₂ does not meet the EU criteria to be classified as STOT SE 1 or 2.

A.2.2.5. Specific target organ toxicity – single exposure Category 3 (STOT SE 3)

A2.2.5.1 Short summary and overall relevance of the provided information on STOT SE 3

Not classified.

A2.2.5.2 Comparison with the CLP criteria

No data are available to indicate that the active substance should be classified for STOT SE 3, H335: May cause respiratory irritation. Acute inhalation toxicity studies in animals are not enough to support this hazard class regarding CLP 3.8.2.2.1.d).

No data are available to indicate that the active substance should be classified for STOT SE 3, H336: May cause drowsiness and dizziness. Observed effects in the acute neurotoxicity study are treatment-related but they are strongly related with the behaviour noted in the pre-treatment phase. In some cases there are not a dose-response relationship.

A2.2.5.3 Conclusion on classification and labelling for STOT SE 3

Chrysanthemum cinerariaefolium extract from supercritical CO₂ does not meet the EU criteria to be classified as STOT SE 3.

A2.2.5.4 Overall conclusion on acute toxicity related to risk assessment

Value used in the Risk Assessment – Acute systemic toxicity	
Value	Rat LD ₅₀ oral = 700 mg/kg bw total pyrethrins (1073 mg/kg bw extract) Rat LD ₅₀ dermal = > 2000 mg/kg bw total pyrethrins (3064 mg/kg bw extract) Rat LC ₅₀ inhalation = 2.5 mg/L total pyrethrins (3.8 mg/L extract)
Justification for the selected value	Lowest LD ₅₀ or LC ₅₀ values.
Proposed classification	Acute Tox. 4, H302: Harmful if swallowed

Acute Tox. 4, H332: Harmful if inhaled.

Value/conclusion used in the Risk Assessment – Acute local effects

Value/conclusion	Not applicable
------------------	----------------

Justification for the selected value/conclusion	-
---	---

A.2.3. Skin corrosion and irritation

An acute skin irritation study was conducted using six New Zealand White rabbits. 0.5 ml of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was applied to the back of the animals then the area was wrapped for 4 hours using a semi-occlusive dressing. The wrapping was removed at the end of the four-hour skin contact period and the residual extract was removed with deionized water. The treated areas were examined for signs of erythema and edema within 30-60 minutes after patch removal. Readings were also made after 24, 48 and 72 hours. *Chrysanthemum cinerariaefolium* extract from supercritical CO₂, when applied dermally as supplied, produced an average primary skin irritation score of 0.33 (24-72 hours) (██████████). (BRA, MGK and SCJ)

Score (average of 6 animals investigated)	Time	Erythema	Edema
Average score Draize scores (0 to maximum 4)	60 min	0.67	0
	24 h	0.5	0
	48 h	0.33	0
	72 h	0.17	0
Average score	24h, 48h, 72h	0.17	0
Reversibility: *		c	c
Average time for reversibility		72 h	0
* c : completely reversible n c : not completely reversible n : not reversible			

Table A.18 Summary table of *in vitro* studies on skin corrosion/irritation

No data are available.

Table A.19 Summary table of animal studies on skin corrosion/irritation

Summary table of animal studies on skin corrosion/irritation						
Method, Duration of study, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), Vehicle, Dose levels of exposure	Results Average score for erythema/eschar and oedema (24, 48, 72 h) per animal, observations and time point of onset, reversibility, other adverse local/systemic effects, histopathological findings		Remarks (e.g. major deviations)	Reference
US EPA 81-5, OECD 404 GLP Reliability 1 Key	Rabbit New Zealand White 6 rabbits/group (sex not reported)	Undiluted Pyrethrum Extract (FEK-99; 57.03%)	Erythema: 24 h: 0.5 48 h: 0.33 72 h: 0.17 Mean: 0.33	Edema: 24 h: 0 48 h: 0 72 h: 0 Mean: 0	-	[REDACTED] (KPIC) IIIA6.1.4/1 [REDACTED] (BRA, MGK and SCJ) IIIA6.1.4/1

Table A.26 Summary table of human data on skin corrosion/irritation

No data are available.

A2.3.1 Short summary and overall relevance of the provided information on skin corrosion/irritation

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is not irritating.

A2.3.2 Comparison with the CLP criteria

Chrysanthemum cinerariaefolium extract from supercritical CO₂ does not meet the EU criteria to be classified as skin irritant. According to CLP 3.2.2.1.2.1, average primary skin irritation score (0.17) < 2.3.

A2.3.3 Conclusion on classification and labelling for skin corrosion/irritation

Not classified.

A2.3.4 Overall conclusion on skin irritation and corrosivity related to risk assessment

Conclusion used in the Risk Assessment – Skin irritation and corrosivity	
Value/conclusion	Not irritating
Justification for the value/conclusion	<i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ , when applied dermally as supplied, produced an average primary skin irritation score of 0.33 (24-72 hours).
Proposed classification	Not classified

A.2.4. Serious eye damage and Eye irritation

Eye irritation from *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was investigated in a group of six New Zealand White rabbits. A volume of 0.1 ml of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was instilled into the conjunctival sac of one eye of the animals, the other untreated eye served as a control. The treated eyes were examined at 1, 24, 48, and 72 hours, and at 4 and 7 days following instillation of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. The test article produced conjunctival irritation in all rabbit eyes at 24- and 48-hour examination. No conjunctival irritation was observed in any of the test eyes by the 72-hour reading. No corneal opacity or iritis were noted during the observation period (██████████). (KPIC and BRA, MGK and SCJ)

	Cornea	Iris	Conjunctiva	
			redness	chemosis
Animals investigated	6	6	6	6
60 min	0	0	1	2
24 h	0	0	1	1
48 h	0	0	0.5	0.16
72 h	0	0	0	0
Average 24h, 48h, 72h	0	0	0.5	0.39
Area effected				
Maximum average score	0	0	1	2
Reversibility*	c	c	c	c
Average time for reversion	72 h	72 h	72 h	72 h

*c : completely reversible

Table A.27 Summary table of *in vitro* studies on serious eye damage and eye irritation

No data are available.

Table A.28 Summary table of animal studies on serious eye damage and eye irritation

Summary table of animal studies on serious eye damage and eye irritation								
Method, Duration of study, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), Vehicle, Dose levels Duration of exposure	Results Average score for corneal opacity, iritis, conjunctival redness and conjunctival oedema (24, 48, 72 h) per animal, observations and time point of onset, reversibility				Remarks (e.g. major deviations)	Reference
US EPA 81-4, OECD 405 GLP Reliability 1 Key	Rabbit New Zealand White 6 rabbits/group (sex not reported)	Undiluted Pyrethrum Extract (FEK-99; 57.03%) 7 d	Cornea 24 h: 0 48 h: 0 72 h: 0 Mean: 0	Iris 24 h: 0 48 h: 0 72 h: 0 Mean: 0	Redness Conjunctiva 24 h: 1 48 h: 0.5 72 h: 0 Mean: 0.5	Chemosis 24 h: 1 48 h: 0.2 72 h: 0 Mean: 0.4	-	[REDACTED] (KPIC) IIIA6.1.4/02 [REDACTED] (BRA, MGK and SCJ) IIIA6.1.4/2

Table A.29 Summary table of human data on serious eye damage and eye irritation

No data are available.

A2.4.1 Short summary and overall relevance of the provided information on serious eye damage/eye irritation

The test article produced conjunctival irritation in all rabbit eyes. No conjunctival irritation was observed in any of the test eyes by the 72-hour reading. No corneal opacity or iritis was noted during the observation period.

A2.4.2 Comparison with the CLP criteria

It does not meet the EU criteria to be classified as eye irritant. According to CLP 3.3.2.1.2. (Table 3.3.2), mean values were: corneal opacity (0) and iritis (0) < 1 and redness conjunctiva (0.5) and chemosis (0.39) < 2.

A2.4.3 Conclusion on classification and labelling for serious eye damage/eye irritation

Not classified.

A2.4.4 Overall conclusion on eye irritation and corrosivity related to risk assessment

Conclusion used in Risk Assessment – Eye irritation and corrosivity	
Value/conclusion	Not classified
Justification for the value/conclusion	No irritation noted.
Proposed classification	Not classified

A.2.5. Skin sensitisation

Table A.30 Summary table of studies on skin sensitisation

Summary table of <i>in vitro</i> studies on skin sensitisation						
Method, Guideline, GLP status, Reliability, Key/supportive study	Test substance (including purity), Vehicle, Doses	Relevant information about the study (e.g. organism (e.g. bacteria), cell type, strains)	Results	Remarks (e.g. major deviations)	Reference	
Similar to OECD TG 442D <i>In vitro</i> sensitisation assay GLP Reliability 1 Supportive	Pyrethrum Extract Pale (50% w/w), Lot# 2016-5-BB 50%, 25%, 10%, 5%, 2%, 1%, 0.4% and 0.2% v/v total pyrethrins in EtOH (25%, 12.5%, 5%, 2.5%, 1%, 0.5%, 0.2%, and 0.1%)	EpiDerm™ from MatTek (reconstructed human epidermis (RHE) of normal, human-derived epidermal keratinocytes (NHEK))	SI > 2		Troese M., 2017 (KPIC)	
			Sensitiser			
			Conc.			SI
			0.2%			0.8
			0.4%			2.2
			1%			1.9
			2%			2.8
			5%			2.8
10%	2.8					
25%	4.1					
50%	4.4					
Similar to OECD TG 442D <i>In vitro</i> sensitisation assay GLP Reliability 1 Supportive	Refined Pyrethrum Concentrate (53.72% w/w), Lot# 10209 50%, 25%, 10%, 5%, 2%, 1%, 0.4% and 0.2% v/v total pyrethrins in EtOH (25%, 12.5%, 5%, 2.5%, 1%, 0.5%, 0.2%, and 0.1%)	EpiDerm™ from MatTek (reconstructed human epidermis (RHE) of normal, human-derived epidermal keratinocytes (NHEK))	SI > 2		Troese M., 2017 (MGK)	
			Sensitiser			
			Conc.			SI
			0.2%			1.0
			0.4%			1.6
			1%			1.6
			2%			1.7
			5%			2.9
10%	7.3					
25%	21.2					
50%	25.3					

Similar to OECD TG 442D <i>In vitro</i> sensitisation assay GLP Reliability 1 Supportive	PY-T-50 Pale Refined Pyrethrins (Pyrethrin Extract (50%, Lot# 0116-501-6101 25%, 12.5%, 5%, 2.5%, 1%, 0.5%, 0.2% and 0.1% v/v total pyrethrins in EtOH	EpiDerm™ from MatTek (reconstructed human epidermis (RHE) of normal, human-derived epidermal keratinocytes (NHEK))	SI > 2 Sensitiser <table border="1"> <thead> <tr> <th>Conc.</th> <th>SI</th> </tr> </thead> <tbody> <tr> <td>0.1%</td> <td>1.2</td> </tr> <tr> <td>0.2%</td> <td>1.8</td> </tr> <tr> <td>0.5%</td> <td>3.0</td> </tr> <tr> <td>1%</td> <td>3.3</td> </tr> <tr> <td>2.5%</td> <td>4.9</td> </tr> <tr> <td>5%</td> <td>6.3</td> </tr> <tr> <td>12.5%</td> <td>17.2</td> </tr> <tr> <td>25%</td> <td>32.2</td> </tr> </tbody> </table>	Conc.	SI	0.1%	1.2	0.2%	1.8	0.5%	3.0	1%	3.3	2.5%	4.9	5%	6.3	12.5%	17.2	25%	32.2		Troese M., 2017 (BRA)
Conc.	SI																						
0.1%	1.2																						
0.2%	1.8																						
0.5%	3.0																						
1%	3.3																						
2.5%	4.9																						
5%	6.3																						
12.5%	17.2																						
25%	32.2																						
Summary table of animal studies on skin sensitisation																							
Method, Duration of study, Route of exposure (e.g. topical/intradermal, induction/challenge if relevant), Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	Results (e.g. EC3-value or amount of sensitised animals at induction dose)	Remarks (e.g. major deviations)	Reference																		
US EPA 81-6 (1984), OECD 406 (1981), Modified Buehler method GLP Reliability 2 Key	Guinea pig, Hartley, Male Test substance: 10 Naive control: 10 Positive control: 10	Pyrethrum Extract Pale (FEK-99, Purity 57.03%)	0/10 Not sensitising	A minimal of 20 animals have to be used in the treatment group.	██████████ (KPIC) IIIA6.1.5 ██████████ (BRA, MGK and SCJ) IIIA6.1.5																		
LLNA Method NIH No. 99-4494, 1999, EPA OPPTS 870.2600, Final Guideline (March 2003), OECD No. 429, revised July 2010. GLP Reliability 1 Key	Mice, CBA/J, female 5	Pyrethrum Extract Pale (50% w/w), Lot# 2016-5-BB Two positive control substances: α-hexylcinnamaldehyde in DMF	EC3 = 4.0% Sensitiser	-	██████████ (KPIC)																		

		(25% HCA) and 1-chloro-2,4-dinitrobenzene in DMF (0.25% DNCB)			
LLNA Method NIH No. 99-4494, 1999, EPA OPPTS 870.2600, Final Guideline (March 2003), OECD No. 429, revised July 2010. GLP Reliability 1 Key	Mice, female 5 CBA/J,	Refined Pyrethrum Concentrate (53.72% w/w), Lot# 10209 Two positive control substances: α-hexylcinnamaldehyde in DMF (25% HCA) and 1-chloro-2,4-dinitrobenzene in DMF (0.25% DNCB)	EC3 = 7.1% Sensitiser	-	██████████ (MGK)
LLNA Method NIH No. 99-4494, 1999, EPA OPPTS 870.2600, Final Guideline (March 2003), OECD No. 429, revised July 2010. GLP Reliability 1 Key	Mice, CBA/J, female 5	PY-T-50 Pale Refined Pyrethrins (Pyrethrin Extract (50%)), Lot# 0116-501-6101 Two positive control substances: α-hexylcinnamaldehyde in DMF (25% HCA)	EC3 = 6.2% Sensitiser	-	Xxxxxxxxxx X (xxxx) (BRA)

		and 1-chloro-2,4-dinitrobenzene in DMF (0.25% DNCB)			
--	--	---	--	--	--

Table A.31 Summary table of human data on skin sensitisation

No data are available.

Table A.20 Summary table of other studies relevant for skin sensitisation

No data are available.

A2.5.1 Short summary and overall relevance of the provided information on skin sensitisation

To assess the contact dermal sensitization potential of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂, a guinea pig dermal sensitisation study (modified Buehler method) was developed. In preliminary dose-range-finding studies, 10 animals were exposed to 4 different concentrations of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ at the highest non-irritating concentrations (as supplied (50%), and at 75% v/v (32.5%), 50% v/v (25%), and 25% v/v (12.5%) in corn oil). Based upon the results of the dose-range-finding studies, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was dosed as supplied for induction and challenge studies.

In the main study a group of 10 albino guinea pigs were clipped on the left side and a gauze patch loaded with *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was applied and an elastic bandage was wrapped around the trunk of the animal. After 6 hours, the patch was removed, and the site was cleaned with deionised water. Examinations were performed 60 to 75 minutes and at 24 hours after patch removal using the Draize method. Animals were rested for one day before a second induction application was applied to the same skin site. This procedure was repeated three times weekly for a total of nine applications. Then animals had a two-week rest period. At the end of this rest period, a challenge application was made during 6 hours at a challenge site (on the right side). A group of 10 naïve control were treated with *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ in the same manner. This group served as the control challenge group. Examinations were performed 24 hours and 48 hours. Some signs of erythema were noted in various animals at different time points after each induction phase. However, no such signs were apparent at the challenge phase. The positive control, 1-chloro-2,4-dinitrobenzene, appears to be a dermal sensitizer to guinea pigs. In conclusion, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ does not appear to be a dermal sensitizer to guinea pigs (Troese, 2017). (KPIC and BRA, MGK and SCJ)

Further studies

Test items were tested in an *in vitro* sensitisation assay using in ethanol (EtOH) as vehicle. A RHE of NHEK was tested for the skin sensitising properties. Due to this test focuses in the activation of keratinocytes is similar to the OECD TG 442D. However, molecular markers are different between OECD assay and applicants' assay: while OECD TG 442D measures the activation of Nrf2-ARE signalling pathway, the test performed with the RHE of NHEK measures IL-18 secretion. This cytokine is related with activation of keratinocytes in inflammatory responses and can be used to discriminate contact sensitizers from irritants and low molecular weight respiratory allergens. The positive control 1-chloro-2,4-dinitrobenzene (0.15% DNCB) induced the expected stimulation indices confirming the validity of the assay. Also, the negative (2% lactic acid), vehicle and undosed controls do not induce an increment in the stimulation index. Following 24-hour exposure to the test materials, the assay media was sampled for IL-18 analysis by ELISA kit. To calculate the IL-18 measurement (pg/ml) for each sample, the optical density of the stopped reaction was measured at a wavelength of 450 nm, subtracting out a background reading for all samples at 620 nm. A Stimulation Index (SI), i.e., the fold change in IL-18 of a test substance as compared to the vehicle control, was calculated. A substance with an SI less than 1.6 was considered a non-sensitizer; a substance with an SI greater than or equal to 2.0 was considered a sensitizer.

The viability of the tissues at 24 hours was determined using methyl thiazole tetrazolium (MTT) uptake and reduction. The absorbance of each sample was measured at 540 nm using a reference wavelength of 690 nm. The viability was then expressed as a percent of control values, corrected for direct MTT reduction.

Also, test items were tested *in vivo* in a dermal sensitisation study following OECD TG 429 using in *N,N*-dimethylformamide as vehicle. Five female CBA/J mice per dose level was

tested for the skin sensitizing properties by a local lymph node assay. The positive controls α -hexylcinnamaldehyde (HCA, 85%) and 1-chloro-2,4-dinitrobenzene (DNCB) induced the expected stimulation indices confirming the validity of the assay. The mice were given an intraperitoneal injection of the thymidine analog 5-bromo-2'-deoxy-uridine (BrdU) approximately five hours prior to euthanasia. The auricular lymph nodes were combined for each animal and single-cell suspensions were generated in RPMI-10 medium. An aliquot of each cell suspension was taken for immunophenotyping analysis; the remaining cell suspensions were fixed with 85% ethanol. The cell suspensions were used to determine BrdU incorporation into the lymphocyte and the total number of cells in the nodes, for each individual animal.

The percentages of B⁺ and T⁺ cells and of I-A^{K+} and I-A^{K+}CD69⁺ cells were determined by flow cytometry and the B:T ratios were calculated. Test groups that show an increase greater than 25% (1.25-fold) when compared to the vehicle control in these LNC surface markers and have an SI value of 3 or more are considered dermal sensitizers. The positive controls (0.25% DNCB and 25% HCA) were both found to be sensitizing.

There were no effects on body weights, clinical signs of toxicity or mortality. None of the test item treatments resulted in increases in ear thickness of 25% or more; therefore, the test items were not considered irritating. (KPIC and BRA, MGK and SCJ)

Pyrethrum Extract Pale (50% w/w) (KPIC)

Application of the test item at 0.4% (v/v) in EtOH (0.2%) onward resulted in SI values greater than 2, with the only exception of the 1.0% (v/v) (0.5%) (SI = 1.9 – probable sensitizer), existing a dose-response relationship.

Topical application of the above test item at 5% and 25% (v/v) (2.5% and 12.5%) in DMF resulted in SI values greater than or equal to 3. At a concentration of 10% (v/v) (5%) an SI of 2.9 was achieved. Treatment with the test item induced an increase of greater than 25% (1.25-fold) over the vehicle control in % B, B:T ratio, and % I-A^{K+}CD69⁺. Treatment with the test item at 25% (v/v) (12.5%) also induced an increase greater than 25% over the vehicle control in % I-A^{K+}. The test item was only of borderline activity. However, this test item is considered a sensitizing substance. The EC3 calculated for the test item is 4.0% (v/v) (2%).

Refined Pyrethrum Concentrate (53.72% w/w) (MGK)

Application of the test item at 5.0% (v/v) in EtOH (2.5%) onward resulted in SI values greater than 2, existing a dose-response relationship. Between 0.4% and 2.0% (v/v) (0.2% and 1%) concentrations a SI greater than 1.6 (probable sensitizer) was achieved.

Topical application of the above test item at 5%, 10% and 25% (v/v) in DMF (2.5%, 5%, 12.5%) resulted in SI values greater than 3 at 10% and 25% (5% and 12.5%). Treatment with the test item at 10% and 25% (v/v) (5% and 12.5%) induced an increase of greater than 25% (1.25-fold) over the vehicle control in % B, B:T ratio, and % I-A^{K+}CD69⁺. Treatment with the test item at 25% (v/v) (12.5%) also induced an increase greater than 25% over the vehicle control in % I-A^{K+}. Therefore, this test item is considered a sensitizing substance. The calculated EC3 of Refined Pyrethrum Concentrate (53.72% w/w) is 7.1% (v/v) (3.5%).

PY-T-50 Pale Refined Pyrethrins (BRA)

Application of the test item at 1.0% (v/v) in EtOH (0.5%) onward resulted in SI values greater than 2, existing a dose-response relationship. At 0.4% (v/v) (0.2%) a SI = 1.8 (probable sensitizer) was achieved.

Topical application of the above test item at 10% and 25% (v/v) (5% and 12.5% total pyrethrins respectively) in DMF resulted in SI values greater than 3. Treatment with the test item induced an increase of greater than 25% at 2.5% (v/v) (1.25% total pyrethrins) (in % B, B:T ratio, and % I-A^{K+}CD69⁺), 5% (v/v) (2.5% total pyrethrins) (in B:T ratio and % I-A^{K+}CD69⁺), and at 10% and 25% (v/v) (5% and 12.5% total pyrethrins respectively) (%B, B:T ratio, % I-A^{K+}, and % I-A^{K+}CD69⁺). Therefore, this test item is

considered a sensitizing substance. The calculated EC₃ of PY-T-50 Pale Refined Pyrethrins is 6.2% (v/v) (3.1% total pyrethrins). The SIs at 25% (v/v) (12.5% total pyrethrins) of PY-T-50 Pale Refined Pyrethrins and HCA (weak sensitiser) are 4.3 and 5.9, respectively. HCA is a typical Cat. 1B sensitiser as shown in the ANNEX 1 of OECD TG 429 'performance standards' where EC₃ of HCA ranges from 4.4% to 14.7%. Since SI of PY-T-50 Pale Refined Pyrethrins is similar to that of HCA, both compounds are considered to have comparable potency for skin sensitisation. PY-T-50 Pale Refined Pyrethrins also should be categorized as Cat. 1B.

Mean stimulating indices observed in LLNA with Pyrethrum extracts

Test concentration	5% (v/v)	10% (v/v)	25% (v/v)
	2.5%	5%	12.5%
Pyrethrum Extract Pale (50% w/w)	3.5	2.9	3.0
Refined Pyrethrum Concentrate (53.72% w/w)	2.3	4.0	8.1
PY-T-50 Pale Refined Pyrethrins	1.8	6.7	4.3

SI ≥ 3 indicates a sensitizing response

Vehicle Control – (DMF) SI = 1.0, Positive Control – (25% HCA) 5.9, Positive Control – (0.25% DNCB) 10.1

Under the experimental conditions of these *in vivo* studies, the test items are considered sensitizing. The calculated EC₃ were 4.0% 7.1%, and 6.2% (v/v) (12.5%, 3.5%, and 4.4% total pyrethrins) for Pyrethrum-Extract 50%, Refined Pyrethrum Concentrate, and PY-T-50 Pale Refined Pyrethrins respectively. However, for Pyrethrum Extract Pale 50% there was no dose response for the SI (██████████). (KPIC and BRA, MGK and SCJ)

Although the results differ between Buehler method and LLNAs, active substance will be classified as skin sensitizer for different reasons: Buehler method only measures the adverse outcome in a subjective way while LLNA measures a key event in an objective way; the Buehler test only has to test one concentration while LLNA tests three different concentrations (LLNA was performed using three sources also); finally, the positive result in the LLNAs is supported by the positive result in the three *in vitro* sensitization assays.

A2.5.2 Comparison with the CLP criteria

Classified as Skin Sens. 1B (H317: May cause an allergic skin reaction) because, according to the CLP 3.4.2.2.3.3. (Table 3.4.4.), EC₃ > 2 % in the three LLNA studies.

A2.5.3 Conclusion on classification and labelling for skin sensitisation

Skin sensitisation, Category 1B, H317: May cause an allergic skin reaction.

A2.5.4 Overall conclusion on skin sensitisation related to risk assessment

Conclusion used in Risk Assessment – Skin sensitisation	
Value/conclusion	Sensitising
Justification for the value/conclusion	Refer to discussion above.
Proposed classification	Skin sensitisation, Category 1B, H317: May cause an allergic skin reaction

A.2.6. Respiratory sensitisation

No data are available.

A2.6.1 Short summary and overall relevance of the provided information on respiratory sensitisation

No data are available to indicate that the active substance is a respiratory sensitiser.

A2.6.2 Comparison with the CLP criteria

It does not meet the CLP criteria to be classified as respiratory sensitiser.

A2.6.3 Conclusion on classification and labelling for respiratory sensitisation

Not classified.

A2.6.4 Overall conclusion on respiratory sensitisation related to risk assessment

Based on the available information *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ does not meet the EU criteria to be classified as respiratory sensitiser.

A.2.7. Repeated dose toxicity/STOT RE

A.2.7.1. Short term repeated dose toxicity

A2.7.1.1 Short-term oral toxicity

Oral repeated dose toxicity

Mouse

In a 2-week toxicity study on mice, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was offered in the diet at concentrations of 5000 and 7000 ppm equivalent to 816 and 1142 mg/kg bw/d total pyrethrins (1250 and 1750 mg/kg bw/d extract) to 50 male animals/treatment group. One animal of the 7000 ppm dosage group died at study day 2. There was no effect on body weight but food consumption was significantly reduced at both dosage levels. A statistically significant increase in absolute and relative liver weights was found at both dosage levels (██████████). (KPIC)

Pyrethrum conc. ppm/mg/kg bw/d extract	Clinical signs	Mean bw at day 14 g ± S.D.*	Mean food consumption				Liver weights	
			g/mouse/d		g/kg bw/d		Absolute g	Relative %
			week 1	week 2	week 1	week 2		
0/0	0/15	33 ± 2.5	5.5 ± 0.50*	5.5 ± 0.45	173.8 ± 11.7	163.6 ± 11.4	1.89	5.82
5000/1250	0/50	34 ± 2.5	5.4 ± 0.61	5.5 ± 0.54	164.7 ± 12.3	161.6 ± 10.2	2.82 ²⁾	8.50 ²⁾
7000/1750	0/49	34 ± 2.4	5.2 ± 0.41 ¹⁾	5.2 ± 0.38 ¹⁾	162.5 ± 14.3 ²⁾	155.5 ± 10.1 ¹⁾	2.94 ²⁾	8.91 ²⁾

* standard deviation

¹⁾ significantly different from the control group: p < 0.05

²⁾ significantly different from the control group: p < 0.01

Rat

Chrysanthemum cinerariaefolium extract from supercritical CO₂ was offered in three different diets (see above) at dosage levels of 0, 940, 2810, 5640, and 9400 ppm in each diet *ad libitum* for two weeks equivalent to 0, 69, 207, 416, and 672 mg/kg bw/d total pyrethrins (0, 106, 317, 637, and 1030 mg/kg bw/d extract). The groups consisted of 10 male rats each. No mortality and no dose related clinical signs were observed. Animals fed the Purina diet exhibited the largest effect on body weight gain at the 9400 ppm (1030 mg/kg bw/d extract) dosage level with a statistically significant decrease at week 2. The decrease in the other diets was not statistically significant. No clear differences between the three diets were noted. Food consumption depressions paralleled body weight depressions. On the basis of the data obtained, Purina Certified Rodent Chow[®]# 5002 was selected as the diet to be employed in future studies with *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. 2-week feeding of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ containing diets has comparatively low effects to the rat. The only difference to control animals was found in the highest concentration group (9400 ppm (1030 mg/kg bw/d extract)) which showed less body weight gain in the second week of

the test. However, the body weight at the end of week 2 was always higher than at the end of week one ([REDACTED]). (KPIC)

Parameter	Control 0 ppm			940 ppm 106 mg/kg bw/d extract			2180 ppm 317 mg/kg bw/d extract			5640 ppm 637 mg/kg bw/d extract			9400 ppm 1030 mg/kg bw/d extract			+/- dose- response		
	A*	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C
number of animals examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	-	-	-
Mortality	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
clinical signs	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
body weight ^a	140	139	134	144	146	151	121	133	134	118	132	123	114	121	123	-	-	-
observ. period 1 ^b	27.4	26	25.5	+	+0.4	+1.8	-	-	+0.7	-	-	-	-	-	-			
				0.1			2.4	1.1		4.3 ²	3.0 ²	1.9	6.6- ₂	3.9 ²	2.7 ¹			
oserv. period 2 ^b	25.7	24.5	25.0	+	+1.2	+1.7	-	-	+0.1	-	-	-	-	-	+1.0	-	-	-
				0.7			1.8	0.9		2.9 ¹	2.0	0.8	2.4¹	1.5				

* different diets containing pyrethrum in different concentrations

A: Purina Rodent Chow® #5002

B: Ziegler NIH-07 open formula diet

C: Agway Certified Prolab 3200

^a average changing of weight (**g**) from the beginning of the study to study termination


^b average changes in food consumption (**g / animal / day**) between two observation periods. difference to control group

¹ significantly different from the control group; p<0.05

² significantly different from the control group; p<0.01

Table A.34 Summary table of oral short-term animal studies (usually 28-day studies)

Summary table of oral short-term animal studies (usually 28-day studies)						
Method, Duration of study, Route of exposure (gavage, in diet, other) Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including also target organs)	Remarks (e.g. major deviations)	Reference
14 days Non-GLP No guideline Reliability 2 Key	Rat Charles River Male 50 per group	Pyrethrum Extract 2 (Lot No. FE K87, FNB 86-2-36A; Purity 54.6%) 0, 940, 2810, 5640, and 9400 ppm total pyrethrins in diet daily; equivalent to 0, 69, 207, 416, and 672 mg/kg bw/d total pyrethrins (0, 106, 317, 637, and 1030 mg/kg bw/d extract)	NOAEL: 5640 ppm (416 mg/kg bw/ d total pyrethrins (637 mg/kg bw/d extract)) LOAEL: 9400 ppm (672 mg/kg bw/ d total pyrethrins (1030 mg/kg bw/d extract))	940, 2810, 5640 ppm (106, 317, and 637 mg/kg bw/d extract): no effects 9400 ppm (1030 mg/kg bw/d extract): body weight gain ↓	-	██████████ (KPIC) IIIA6.3.1/01

<p>14 days GLP No guideline Reliability 2 Key</p>	<p>Mouse Charles River CD-1 Male 50 per group</p>	<p>Pyrethrum extract 2 (Batch # 011831-00 task force blend # FEK- 99Pale; Purity 57.57%) 0, 5000 and 7000 ppm in diet daily; equivalent to 0, 816, and 1142 mg/kg bw/d total pyrethrins (0, 1250, and 1750 mg/kg bw/d extract)</p>	<p>NOAEL: Not determined LOAEL: 5000 ppm (816 mg/kg bw/ d total pyrethrins (1250 mg/kg bw/d extract))</p>	<p>5000 ppm (1250 mg/kg bw/d extract): feed intake ↓, liver weight ↑ 7000 ppm (1750 mg/kg bw/d extract): one mortality, feed intake ↓, liver weight ↑</p>	<p>-</p>	<p> (KPIC) IIIA6.3.1/02</p>
---	---	---	---	---	----------	--

Value used in the Risk Assessment – Short-term oral toxicity	
Value/conclusion	AEL _{short-term} 0.2 mg/kg bw/d total pyrethrins (0.31 mg/kg bw/d extract).
Justification for the value/conclusion	Based on the neurotoxicity study in rats given single oral doses, acute neurological disorders and behavioural effects were noted, with a NOAEL of 20 mg/kg bw.

A2.7.1.2 Short-term dermal toxicity

Chrysanthemum cinerariaefolium extract from supercritical CO₂ was administered dermally to five male and five female New Zealand white rabbits in the form of a 25% (w/v) mixture in vegetable oil at doses of 0, 100, 300, or 1000 mg/kg bw once daily of total pyrethrins (0, 153, 460, and 1532 mg/kg bw/d extract), 5 days per week for 3 weeks. Animals in the vehicle control group were given vegetable oil on the same regimen and at the same volume as the group receiving the high dose. One rabbit at 1000 mg/kg bw/d total pyrethrins (1532 mg/kg bw/d extract) was sacrificed in extremis on day 10. Macroscopic examination of this animal did not reveal the cause of death. A low incidence of desquamation and/or red raised areas on the skin at the application site was observed in all groups, including the vehicle controls. Several animals in the treated groups showed very slight to well-defined erythema of the skin at the application site, but no clear pattern with regard to treatment was seen for any of these findings. Microscopic evaluation revealed no evidence of systemic toxicity. The microscopic lesions at the application site included acanthosis, haemorrhage, hyperkeratosis, and chronic inflammation, although haemorrhage was observed only in the group given the vehicle alone. Thus, all of the dermal reactions appeared to be due to the vegetable oil.

The NOAEL for systemic effects was 1000 mg/kg bw/d total pyrethrins (1532 mg/kg bw/d extract), the highest dose tested [REDACTED]. (KPIC and BRA, MGK and SCJ)

Table A.36 Summary table of dermal short-term animal studies (usually 28-day studies)


Summary table of dermal short-term animal studies (usually 28-day studies)						
Method, Duration of study, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels, Surface area, Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including target organs)	Remarks (e.g. major deviations)	Reference
EPA 82-2 and OECD 410 21 days GLP Reliability 1 Key	Rabbit New Zealand White rabbits Male/Female 5/sex/group	Pyrethrum extract (Lot FEK-99; Purity 57.574%) 0, 100, 300, and 1000 mg/kg bw/d total pyrethrins (0, 153, 460, and 1532 mg/kg bw/d extract) 5 days /week	NOAEL: 1000 mg/kg bw/d total pyrethrins (1532 mg/kg bw/d extract)	1000 mg/kg bw/d total pyrethrins (1532 mg/kg bw/d extract): 72% animals (and 10-20% of animals treated with lower doses) developed very slight to well defined erytema of the skin at the application syte by day 7. Very slight edema was observed in one female on day 11 and 14. These lessions were partially reversible.	-	 (KPIC) (BRA, MGK and SCJ) IIIA6.3.2

Table A.37 Summary table of human data on short-term dermal toxicity
No data are available.

A2.7.1.3 Short-term inhalation toxicity

No data are available.

A2.7.1.4 Overall conclusion on short-term repeated dose toxicity related risk assessment

Value used in the Risk Assessment – Short-term repeated dose systemic toxicity	
Value	AEL _{short-term} 0.2 mg/kg bw/d total pyrethrins (0.31 mg/kg bw/d extract).
Justification for the selected value	Based on the neurotoxicity study in rats given single oral doses, acute neurological disorders and behavioural effects were noted, with a NOAEL of 20 mg/kg bw total pyrethrins (31 mg/kg bw extract).
Proposed classification	Not classified

Value/conclusion used in the Risk Assessment – Short-term repeated dose local effects	
Value/conclusion	Not applicable
Justification for the selected value/conclusion	-
Proposed classification	-

A.2.7.2. Sub-chronic repeated dose toxicity

A3.7.2.1 Sub-chronic oral toxicity

Mouse

Groups of 15 male and 15 female Charles River CD-1 mice were offered diets containing 0, 300, 1000, 3000, 10000, and 30000 ppm total pyrethrins over a period of 90 days, equivalent to 0, 47, 160, 460, and 1600 mg/kg bw/d total pyrethrins (0, 72, 245, 705, and 2451 mg/kg bw/d extract) for males and 0, 56, 200, 580, and 1800 mg/kg bw/d total pyrethrins (0, 86, 306, 889, and 2758 mg/kg bw/d extract) for females.

All animals of the highest dosage group died or were sacrificed *in extremis* by study day 10. Four males and two females of the 10000 ppm (2451/2758 mg/kg bw/d extract) dosage group also died within the first two days, with clinical signs that included tremors, pale exposed skin, dilated pupils, altered activity, laboured breathing, cold to touch, moribundity, and hunched posture. No treatment related signs of toxicity were observed in the other test groups. The group mean body weights and food consumption were similar for all groups with surviving animals. The liver was the only target organ found. The absolute weight of the liver and the liver/body weight were statistically significantly increased in males and females at 3000 and 10000 ppm (705/889 and 2451/2758 mg/kg bw/d extract), whereas slight but significant increase of the relative liver weight was also found for males in the 300 and 1000 ppm (72 and 245 mg/kg bw/d extract) test groups. A treatment-related increase in the incidence and/or severity of congestion in the liver was observed in surviving male and female mice at 10000 ppm (2451/2758mg/kg bw/d extract), and an increased incidence but only mild severity was found in < 15% of investigated animals at 3000 ppm (705/889 mg/kg bw/d extract); at 1000 ppm (245/306 mg/kg bw/d extract), only 2 of 15 mice showed mild congestion of the liver on macroscopic observation. In general, macroscopic congestion was slightly more pronounced in male than in female. An increased incidence of hepatocellular

hypertrophy was present in surviving male and female mice at 3000 and 10000 ppm (705/889 and 2451/2758 mg/kg bw/d extract). The NOAEL was 300 ppm, equal to 47 mg/kg bw/d total pyrethrins (72 mg/kg bw/d extract) for males and 56 mg/kg bw/d total pyrethrins for females (86 mg/kg bw/d extract) (KPIC)

	Control		300 ppm pyrethrins 72/86 mg/kg bw/d extract		1000 ppm pyrethrins 245/306 mg/kg bw/d extract		3000 ppm pyrethrins 705/889 mg/kg bw/d extract		10000 ppm pyrethrins 2451/2758 mg/kg bw/d extract		30000 ppm pyrethrins	
	m ^a	f ^a	m ^a	f ^a	m ^a	f ^a	m ^a	f ^a	m ^a	f ^a	m ^a	f ^a
Number of animals examined	15	15	15	15	15	15	15	15	15	14	14	3
Mortality	0/15	0/15	0/15	0/15	0/15	0/15	0/15	0/15	0/15	1/15	1/15	12/15
Clinical signs*	0/15	0/15	0/15	0/15	0/15	0/15	15/15	15/15	15/15	15/15	15/15	15/15
Body weight (diff. control %)	588	276	594 (-1.0)	280 (+1.4)	569 (-3.2)	277 (+0.4)	548 (-6.8)	267 (-3.3)	523 (-11.1) ₁₎	233 (-15.6) ₁₎	460 (-21.8) ₁₎	234 (-15.2) ₂₎
Food consumption g/animal/day (diff. control %)	28.0	18.6	28.5 (+1.8)	18.2 (-2.2)	27.5 (-1.8)	18.0 (-3.2)	26.5 (-5.4)	17.3 (-7.0)	26.7 (-4.6)	15.6 (-16.1) ₁₎	24.3 (-15.0) ₂₎	15.9 (-14.9) ₂₎
Liver												
Organ weight g	25.06	11.12	27.21	11.6	25.53	11.91	28.8	14.0 ¹⁾	36.42 ¹⁾	17.97 ¹⁾	39.71 ¹⁾	23.11 ¹⁾
Gross pathology												
Enlargement	2/15	2/15	0/15	0/15	2/15	0/15	3/15	0/15	10/15	1/14	12/14	2/03
Congestion	2/15	0/15	0/15	0/15	2/15	0/15	3/15	0/15	10/15	1/14	12/14	2/03
Microscopic pathology	3/10	2/10	1/10	1/10	1/10	1/10	1/10	1/10	3/10	1/10	-	-
Kidney												
Organ weight g	4.59	2.4	4.75	2.33	4.59	2.36	4.98	2.4	5.53 ¹⁾	2.33	4.91	2.36

¹⁾ statistically significant different from the control group p < 0.01
²⁾ statistically significant different from the control group: p < 0.05
^a number of animals affected/total number of animals

Rat

Groups of 15 male and 15 female Charles River CD rats received diets containing 0, 300, 1000, 3000, 10000, and 20000 ppm total pyrethrins over a period of 90 days, equivalent to 0, 17, 57, 170, 590, and 1200 mg/kg bw/d total pyrethrins (0, 26, 87, 261, 904, and 1839 mg/kg bw/d extract) for males and 0, 22, 74, 220, 710, and 1400 mg/kg bw/d total pyrethrins (0, 34, 113, 337, 1088, and 2145 mg/kg bw/d extract) for females.

During the first 3 to 7 days one female at the 10000 ppm (1088 mg/kg bw/d extract) dosage level, and one male (904 mg/kg bw/d extract) plus 12 females of the 20000 ppm (2145 mg/kg bw/d extract) dosage level died. All other animals survived. Clinical signs of toxicity were only found at animals from the two highest dosage groups. Signs seen were decreased defecation, tremors, increased respiration rate, increased activity and occasionally convulsions. Most signs only occurred during the first two

weeks of the study. Statistically significant decreases in mean body weights were observed during most or all of the study in males and females at 10000 and 20000 ppm (904/1088 and 1839/2145 mg/kg bw/d extract), and statistically significant decreases in mean food consumption were seen during most or all of the study in females at 10000 ppm (1088 mg/kg bw/d extract) and males and females at 20000 ppm (1839/2145 mg/kg bw/d extract) when compared with the respective control groups. Statistically significantly decreased mean values for haematocrit and haemoglobin were found in males at 20000 ppm (1839 mg/kg bw/d extract) and for erythrocytes, haematocrit, and haemoglobin in females at 10000 and 20000 ppm (1088 and 2145 mg/kg bw/d extract). Females at 3000 ppm (337 mg/kg bw/d extract) also showed a slightly decreased mean haemoglobin value. The treatment-related macroscopic findings consisted of enlargement and congestion of the liver in both, males and females, but primarily in males, at 10000 and 20000 ppm (904/1088 and 1839/2145 mg/kg bw/d extract); however, the macroscopic observation could not be confirmed microscopically. The absolute liver weight, the liver/body weight ratio and the liver/brain weight ratio were all statistically significantly increased in males at 10000 and 20000 ppm (904 and 1839 mg/kg bw/d extract) and in females at 3000, 10000 and 20000 ppm (337, 1088, and 2145 mg/kg bw/d extract).

Statistically significant increases of absolute kidney weights were observed at males in the 10000 ppm (904 mg/kg bw/d extract) group only; relative kidney weights were increased at both sexes in the 3000, 10000 and 20000 ppm (261/337, 904/1088 and 1839/2145 mg/kg bw/d extract) dosage group. In microscopic investigations of the kidneys from males small focal or multifocal areas of tubular degeneration and regeneration in the renal cortex were observed. Because of the low incidence of such lesions it cannot be proved if these effects are related to the test substance at all.

The NOAEL was 1000 ppm, equal to 57 mg/kg bw/d total pyrethrins (87 mg/kg bw/d extract) for males and 74 mg/kg bw/d total pyrethrins (113 mg/kg bw/d extract) for females [REDACTED]. (KPIC and BRA, MGK and SCJ)

Changes in body weight are shown in the table below:

Week of study	Control (mg/kg bw/d extract)		300 ppm (mg/kg bw/d extract)		1000 ppm (mg/kg bw/d extract)		3000 ppm (mg/kg bw/d extract)		10000 ppm (mg/kg bw/d extract)		20000 ppm (mg/kg bw/d extract)	
	Male (0)	Female (0)	Male (26)	Female (34)	Male (87)	Female (113)	Male (261)	Female (337)	Male (904)	Female (1088)	Male (1839)	Female (2145)
-1	325 ± 12.2	172 ± 7.5	323 ± 13.5	171 ± 7.6	316 ± 13.9	172 ± 7.6	322 ± 13.7	173 ± 7.4	318 ± 14.2	169 ± 6.6	319 ± 14.1	170 ± 7.1
Day 0	340 ± 14.8	182 ± 9.2	338 ± 15.1	182 ± 8.9	332 ± 18.1	183 ± 7.8	335 ± 16.6	181 ± 9.0	331 ± 15.2	176 ± 8.0	330 ± 14.2	177 ± 9.5
1	379 ± 19.0	197 ± 11.6	377 ± 15.0	196 ± 9.8	366 ± 18.0	196 ± 11.0	368 ± 23.5	193 ± 10.7	349 ² ± 19.9	180 ² ± 9.9	296 ² ± 22.6	164 ² ± 13.1
2	403 ± 21.4	205 ± 11.5	404 ± 17.4	207 ± 11.6	390 ± 37.8	208 ± 10.9	386 ± 25.8	206 ± 9.9	372 ² ± 24.1	191 ² ± 8.1	329 ² ± 23.6	187 ¹ ± 11.0
3	435 ± 22.8	222 ± 15.3	438 ± 22.6	222 ± 14.3	421 ± 47.4	219 ± 11.9	419 ± 29.7	221 ± 9.7	400 ² ± 25.1	203 ² ± 10.7	357 ² ± 24.9	203 ± 2.9
4	457 ± 26.6	229 ± 17.0	461 ± 25.2	232 ± 15.7	445 ± 49.7	230 ± 12.6	440 ± 40.6	225 ± 9.1	421 ¹ ± 30.7	209 ² ± 9.2	382 ² ± 30.1	213 ² ± 3.5
5	478 ± 28.2	234 ± 18.6	486 ± 23.1	238 ± 17.0	469 ± 48.6	239 ± 14.2	462 ± 40.6	234 ± 12.4	442 ¹ ± 30.3	216 ² ± 10.6	401 ² ± 29.1	220 ± 3.0
6	501 ± 28.3	243 ± 20.3	510 ± 28.7	249 ± 19.7	489 ± 52.5	251 ± 26.4	478 ± 44.5	242 ± 10.7	459 ¹ ± 31.5	221 ² ± 9.2	416 ² ± 31.3	228 ± 9.3
7	525 ± 33.6	254 ± 21.9	533 ± 31.9	257 ± 21.1	514 ± 50.0	251 ± 17.4	497 ± 42.8	250 ± 10.4	480 ² ± 38.3	228 ² ± 10.2	431 ² ± 32.3	235 ¹ ± 7.2
8	533 ± 34.8	257 ± 23.0	541 ± 31.2	261 ± 22.0	519 ± 52.7	256 ± 15.8	502 ± 44.4	249 ± 12.0	485 ² ± 40.3	229 ² ± 8.6	438 ² ± 30.1	235 ¹ ± 8.7
9	549 ± 34.8	259 ± 23.7	556 ± 33.5	265 ± 20.8	535 ± 57.7	264 ± 17.1	516 ± 44.5	259 ± 12.8	499 ² ± 42.7	232 ² ± 11.4	447 ² ± 32.8	240 ± 7.8
10	561 ± 36.7	269 ± 25.3	571 ± 34.0	272 ± 23.6	547 ± 61.4	269 ± 16.5	530 ± 45.6	264 ± 13.0	508 ² ± 44.3	234 ² ± 10.1	458 ² ± 34.6	244 ¹ ± 9.7
11	577 ± 37.3	273 ± 25.2	584 ± 41.6	279 ± 24.2	560 ± 64.7	274 ± 17.5	548 ± 49.2	266 ± 12.8	526 ¹ ± 46.6	236 ² ± 10.5	468 ² ± 34.5	244 ¹ ± 12.1

ES *Chrysanthemum cinerariaefolium*, extract from open and mature flowers of *Tanacetum cinerariifolium* obtained with supercritical CO₂

12	583 ± 41.8	277 ± 29.0	589 ± 39.1	279 ± 24.2	567 ± 62.2	277 ± 19.4	548 ± 47.8	267 ± 15.1	529 ¹ ± 48.4	239 ² ± 10.6	468 ² ± 35.9	243 ² ± 12.7
13	588 ± 38.7	276 ± 26.9	594 ± 44.8	280 ± 23.0	569 ± 59.8	277 ± 21.7	548 ± 56.2	267 ± 18.1	523 ² ± 54.1	233 ² ± 12.2	460 ² ± 37.6	234 ¹ ± 15.0

¹ Significantly different from the control group; p<0.05

² Significantly different from the control group; p<0.01

Parameter	Assessment time	0 ppm (mg/kg bw/d extract)		300 ppm (mg/kg bw/d extract)		1000 ppm (mg/kg bw/d extract)		3000 ppm (mg/kg bw/d extract)		10000 ppm (mg/kg bw/d extract)		20000 ppm (mg/kg bw/d extract)	
		Male (0)	Female (0)	Male (26)	Female (34)	Male (87)	Female (113)	Male (261)	Female (337)	Male (904)	Female (1088)	Male (1839)	Female (2145)
Erythrocytes (x10 ⁶ /cmm)	Week 13	8.09±0.49	7.55±0.29	7.92±0.43	7.23±0.57	8.02±0.41	7.59±0.31	8.03±0.24	7.28±0.32	7.82±0.48	6.96 ² ±0.45	7.59±0.66	6.85 ¹ ±0.27
Haemoglobin (g/dL)	Week 13	17.2±0.62	16.6±0.52	16.8±0.89	16.0±1.17	17.1±0.82	16.5±0.66	17.0±0.56	15.5 ² ±0.82	16.4±0.91	14.7 ² ±1.44	15.5 ² ±1.35	13.3 ² ±0.59
Haematocrit (%)	Week 13	50.0±1.76	49.4±1.57	49.6±2.39	47.5±3.55	49.9±2.30	49.4±1.70	49.8±1.44	46.7±2.22	49.3±2.46	45.5 ² ±2.94	46.4 ¹ ±4.36	40.7 ² ±1.57
MCV (microns ³)	Week 13	62±3.2	66±1.1	63±3.1	66±1.8	62±2.3	65±1.1	62±1.7	64±1.8	63±0.9	65±2.0	61±2.3	59 ² ±1.2
MCH (pg)	Week 13	21.3±0.95	22.0±0.45	21.3±0.94	22.2±0.75	21.3±1.00	21.8±0.57	21.2±0.80	21.3 ² ±0.61	21.0±0.50	21.1±1.38	20.4±0.75	19.5 ² ±0.68
MCHC (g/dL)	Week 13	34.5±0.66	33.6±0.39	33.9±0.52	33.7±0.51	34.2±0.51	33.5±0.44	34.2±0.57	33.2 ¹ ±0.49	33.3 ² ±0.62	32.3 ¹ ±1.85	33.3 ² ±0.80	32.7±0.57

¹ Significantly different from the control group; p<0.05

² Significantly different from the control group; p<0.01

Parameter	Assessment time	0 ppm (mg/kg bw/d extract)		300 ppm (mg/kg bw/d extract)		1000 ppm (mg/kg bw/d extract)		3000 ppm (mg/kg bw/d extract)		10000 ppm (mg/kg bw/d extract)		20000 ppm (mg/kg bw/d extract)	
		Male (0)	Female (0)	Male (26)	Female (34)	Male (87)	Female (113)	Male (0)	Female (0)	Male (26)	Female (34)	Male (87)	Female (113)
Aspartate aminotransferase (IU/L)	Week 13	92±16.1	98±64.5	83±10.8	83±10.3	95±15.4	80±8.5	201±301.8	84±17.1	138±123.0	79±18.0	75 ¹ ±15.6	76±15.0
Alanine aminotransferase (IU/L)	Week 13	37±6.3	49±50.3	31 ¹ ±3.3	31±8.1	37±8.2	29±4.0	194±444.6	32±15.5	206±398.8	42±18.1	55±26.9	45±26.0
Glucose (mg/dL)	Week 13	113±21.0	107±6.9	105±13.4	100±10.3	102±8.3	93 ² ±10.1	99±19.3	93 ² ±9.7	106±15.8	97±9.2	108±11.9	105±7.5
Urea nitrogen (mg/dL)	Week 13	13.5±1.59	15.0±2.24	13.8±1.58	15.3±1.73	13.4±2.06	16.1±2.12	14.3±2.28	16.5±2.94	17.6 ² ±3.60	19.3±7.67	18.1 ² ±2.35	19.4±3.25
Creatinine (mg/dL)	Week 13	0.5±0.11	0.6±0.11	0.5±0.07	0.7±0.11	0.5±0.09	0.7±0.07	0.6±0.14	0.7±0.09	0.6±0.08	0.7 ¹ ±0.13	0.6±0.09	0.9 ² ±0.12
Total bilirubin (mg/dL)	Week 13	0.2±0.06	0.2±0.07	0.2±0.04	0.2±0.05	0.2±0.05	0.2±0.03	0.1 ² ±0.05	0.1 ² ±0.05	0.2±0.04	0.2 ¹ ±0.05	0.2±0.00	0.2±0.00
Albumin (g/dL)	Week 13	3.8±0.19	4.1±0.19	3.8±0.24	4.1±0.33	3.9±0.16	4.2±0.16	3.9±0.15	4.2±0.13	4.1 ² ±0.22	4.7 ² ±0.24	4.4 ² ±0.12	4.9 ² ±0.21
Globulin (g/dL)	Week 13	3.1±0.18	2.9±0.23	3.1±0.37	2.9±0.22	3.1±0.36	2.8±0.19	3.0±0.25	2.9±0.21	3.0±0.23	2.9±0.22	2.7 ² ±0.30	3.1±0.06
Total protein (g/dL)	Week 13	6.9±0.31	7.0±0.37	6.8±0.45	7.0±0.46	6.9±0.37	7.0±0.31	6.9±0.30	7.2±0.23	7.1±0.37	7.6 ² ±0.41	7.1±0.38	7.9 ² ±0.15

¹ Significantly different from the control group; p<0.05

² Significantly different from the control group; p<0.01

Parameter	0 ppm (mg/kg bw/d extract)		300 ppm (mg/kg bw/d extract)		1000 ppm (mg/kg bw/d extract)		3000 ppm (mg/kg bw/d extract)		10000 ppm (mg/kg bw/d extract)		20000 ppm (mg/kg bw/d extract)	
	Male (0)	Female (0)	Male (26)	Female (34)	Male (87)	Female (113)	Male (0)	Female (0)	Male (26)	Female (34)	Male (87)	Female (113)
Body weight (g)	586±40.2	274±27.9	594±43.3	278±22.1	564± 69.8	273±26.7	549± 57.3	266±16.1	523 ² ±56.3	236 ² ±10.4	460 ² ±38.6	238 ¹ ±16.9
Brain/body weight (% x 10)	3.67±0.378	6.82±0.77	3.57±0.32	6.81±0.65	3.79±0.47	6.99±0.76	3.95±0.51	7.03±0.52	4.06 ¹ ±0.34	7.76 ² ±0.45	4.54 ² ±0.42	7.97 ¹ ±0.38
Adrenal/body weight (% x 10 ³)	11.2±1.62	25.3±4.00	11.7±2.77	26.2±4.49	11.5±2.55	25.4±4.19	11.1±2.78	28.8±4.00	13.1±2.59	28.5±4.37	14.8 ² ±3.12	25.4±4.23
Kidney (g)	4.59±0.44	2.27±0.21	4.75±0.41	2.34±0.13	4.59±0.53	2.25±0.20	4.98±0.54	2.40±0.15	5.53 ² ±0.51	2.33±0.24	4.91±0.45	2.36±0.37
Kidney/body weight (% x 10)	7.82±0.5	8.32±0.74	8.01±0.56	8.46±0.66	8.19±0.76	8.25±0.63	9.10 ² ±0.83	9.04 ¹ ±0.39	10.6 ² ±0.70	9.88 ² ±0.88	10.7 ² ±0.82	9.85 ² ±0.84
Kidney/brain weight (% x 10 ⁻²)	2.15±0.24	1.23±0.13	2.26±0.20	1.25±0.10	2.18±0.29	1.19±0.11	2.33±0.31	1.29±0.09	2.62 ² ±0.19	1.28±0.13	2.37±0.22	1.24±0.17
Liver (g)	25.1±3.42	11.1±0.90	27.2±3.93	11.6±1.19	25.5±4.07	11.9±1.57	28.8±4.05	14.0 ² ±1.24	36.4 ² ±6.20	18.0 ² ±1.97	39.7 ² ±5.23	23.1 ¹ ±2.84
Liver/body weight (%)	4.26±0.39	4.08±0.33	4.59±0.64	4.18±0.29	4.54±0.54	4.36±0.43	5.25 ² ±0.50	5.27 ² ±0.49	6.94 ² ±0.80	7.61 ² ±0.66	8.63 ² ±0.80	9.67 ² ±0.51
Liver/brain weight (% x 10 ⁻²)	11.8±1.79	6.04±0.70	12.9±1.93	6.20±0.80	12.1±2.14	6.32±0.95	13.5±2.156	7.54 ² ±0.90	17.2 ² ±2.54	9.86 ² ±1.18	19.2 ² ±2.56	12.2 ² ±1.25
Testis/body weight (% x 10 ⁻²)	6.59±0.67	--	6.24±1.08	--	6.67±1.26	--	7.05±0.93	--	6.99±0.77	--	8.21 ² ±0.73	--

¹ Significantly different from the control group; p<0.05

² Significantly different from the control group; p<0.01

Dog

In a range-finding study 2 beagle dogs/treatment /sex were offered diets, containing 0, 600, 1000, 3000 and 6000 ppm total pyrethrins for a period of 8 weeks, equal to 0, 18, 30, 86, and 170 mg/kg bw/d total pyrethrins (0, 28, 46, 132, and 261 mg/kg bw/d extract) for males and 0, 19, 29, 94, and 200 mg/kg bw/d total pyrethrins (0, 29, 44, 144, and 306 mg/kg bw/d extract) for females.

One male and both females died or were killed *in extremis* at the 6000 ppm (261/306 mg/kg bw/d extract) dosage level. Treatment related clinical signs were observed in the 3000 and 6000 ppm (132/144 and 261/306 mg/kg bw/d extract) test groups and included inappetence, thin appearance, ataxia, trembling, oily coat, impaired limb function and shallow breathing. Both sexes at 6000 ppm (261/306 mg/kg bw/d extract) lost weight whereas the other test groups did not differ from controls. At the end of the study, in the 3000 ppm and 6000 ppm (132/144 and 261/306 mg/kg bw/d extract) test groups haematocrit, haemoglobin and erythrocyte values were slightly reduced, but there were no other treatment-related haematological findings. Dogs sacrificed *in extremis* (6000 ppm (261/306 mg/kg bw/d extract) group) showed in addition increased values of leukocytes and segmented neutrophils.



Slightly decreased glucose, calcium, phosphorus and cholesterol values were found at the end of the study in males at 6000 ppm (261 mg/kg bw/d extract), and males and females at 3000 ppm (132/144 mg/kg bw/d extract) had slightly decreased cholesterol concentrations. The aspartate and alanine aminotransferase activities of males at 6000 ppm (261 mg/kg bw/d extract) were slightly increased at the end of dosing, and the surviving male at this dose had a very high creatinine phosphokinase value. The absolute weights of the liver in both males and females at 1000 and 3000 ppm (46/44 and 132/144 mg/kg bw/d extract) were increased in a treatment-related fashion, and the absolute weight of the testis at these doses appeared to decrease in a similar manner. The NOEL was 600 ppm, equal to 18 mg/kg bw/d total pyrethrins (28 mg/kg bw/d extract) for males and 19 mg/kg bw/d total pyrethrins (29 mg/kg bw/d extract) for females ([REDACTED]). (KPIC and BRA, MGK and SCJ)


Parameter	Assessment time	0 ppm (mg/kg bw/d extract)		600 ppm (mg/kg bw/d extract)		1000 ppm (mg/kg bw/d extract)		3000 ppm (mg/kg bw/d extract)		6000 ppm (mg/kg bw/d extract)	
		Male (0)	Female (0)	Male (28)	Female (29)	Male (46)	Female (44)	Male (132)	Female (144)	Male (261)	Female (306)
Calcium (mg/dL)	Pretest	11.1±0.49	11.4±0.07	11.4±0.14	11.1±0.28	11.3±0.07	11.1±0.00	11.3±0.14	11.2±0.07	11.5±0.21	12.0±0.07
	Week 8	11.8±0.21	11.8±0.14	11.6±0.21	11.5±0.00	11.6±0.21	11.7±0.21	11.6±0.35	11.4±0.21	10.6±0.00	N/A
Phosphorus (mg/dL)	Pretest	7.2±0.78	6.7±0.21	7.7±0.42	6.7±0.07	8.0±1.20	6.3±0.07	7.0±0.42	6.7±0.78	7.5±0.71	6.9±0.78
	Week 8	5.7±0.00	5.4±0.49	5.6±0.14	5.5±0.21	5.8±0.85	5.2±0.07	5.6±0.00	5.0±0.49	5.1±0.49	N/A
Urea nitrogen (mg/dL)	Pretest	17.6±6.51	14.2±2.40	20.6±0.64	15.7±1.70	16.2±4.24	18.0±3.75	16.5±3.54	13.0±1.20	12.5±1.06	15.8±3.68
	Week 8	12.7±1.98	15.5±4.67	16.8±1.70	14.6±0.35	13.1±2.05	15.3±3.39	15.8±0.14	14.6±4.53	18.5±5.30	N/A
Aspartate aminotransferase (IU/L)	Pretest	20±0.7	20±1.4	22±1.4	23±2.1	22±1.4	18±2.1	19±1.4	19±0.0	19±2.1	21±2.8
	Week 8	19±0.0	20±0.7	19±0.7	18±0.0	18±0.0	16±0.0	18±0.7	19±2.8	38±31.8	N/A
Alanine aminotransferase (IU/L)	Pretest	24±3.5	21±1.4	26±2.8	26±1.4	18±2.8	21±3.5	21±3.5	22±4.9	24±0.7	26±4.2
	Week 8	29±2.8	26±0.7	33±2.8	34±2.1	23±1.4	35±2.8	34±0.7	36±2.8	53±28.3	N/A
Creatine phosphokinase (IU/L)	Pretest	88±3.5	95±7.8	118±12.7	112±2.8	106±0.0	84±21.9	95±17.0	100±5.7	96±0.7	100±5.7
	Week 8	103±19.1	72±0.0	95±14.1	87±7.1	99±32.5	61±6.4	81±2.8	80±4.2	426±543.8	N/A
Cholesterol (mg/dL)	Pretest	194±31.1	194±10.6	159±4.2	159±7.8	187±1.4	225±29.7	185±9.2	162±32.5	165±31.1	160±60.8
	Week 8	224±28.3	170±4.2	164±14.1	137±17.0	173±14.1	174±26.2	138±2.8	123±12.0	144±46.7	N/A
Glucose (mg/dL)	Pretest	110±0.7	98±5.7	110±0.0	106±4.9	103±0.7	99±4.9	103±2.8	101±4.9	106±4.2	102±1.4
	Week 8	113±2.8	96±2.1	109±9.2	98±4.2	105±1.4	95±4.2	99±2.8	96±1.4	89±2.1	N/A

Parameter	0 ppm (mg/kg bw/d extract)		600 ppm (mg/kg bw/d extract)		1000 ppm (mg/kg bw/d extract)		3000 ppm (mg/kg bw/d extract)		6000 ppm (mg/kg bw/d extract)	
	Male (0)	Female (0)	Male (28)	Female (29)	Male (46)	Male (0)	Female (0)	Male (28)	Female (29)	Male (46)
Liver (g)	334±2.91	276±7.60	321±15.0	274±52.1	339±30.4	317±67.3	350±54.0	345±2.46	427±0.0	N/A
Testis (g)	16.1±0.24	--	15.1±0.7	--	14.0±0.90	--	12.3±3.90	--	11.7±0.0	--

N/A: not available

Table A.38 Summary table of oral sub-chronic animal studies (usually 90-day studies)

Summary table of oral sub-chronic animal studies (usually 90-day studies)						
Method, Duration of study, Route of exposure (gavage, in diet, other), Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including also target organs)	Remarks (e.g. major deviations)	Reference
OECD TG 408 13 weeks GLP Reliability 1 Key	Rat Charles River CD® Male/Female 15/sex/group	Pyrethrum Extract (Batch # 011831-00 Task force blend # FEK-99; Purity 57.574%) Males: 0, 17, 57, 170, 587, and 1188 mg/kg/d total pyrethrins (0, 26, 87, 261, 904, and 1839 mg/kg bw/d extract) Females: 0, 22, 74, 220, 712, and 1440 mg/kg/d total pyrethrins (0, 34, 113, 337, 1088, and 2145 mg/kg bw/d extract) Daily	NOAEL: 1000 ppm, equal to 57 mg/kg bw/d total pyrethrins (87 mg/kg bw/d extract) for males and 74 mg/kg bw/d (113 mg/kg bw/d extract) for females. LOAEL: 3000 ppm, corresponding to 170 mg/kg bw/d total pyrethrins (261 mg/kg bw/d extract) (males) and 220 mg/kg bw/d total pyrethrins (337 mg/kg bw/d extract) (females).	300 ppm (26/34 mg/kg bw/d extract): No statistically significant effect 1000 ppm (87/113 mg/kg bw/d extract): No statistically significant effect 3000 ppm (261/337 mg/kg bw/d extract): ↓ food consumption, ↓ haemoglobin, ↑ kidney weight, ↑ liver weight in females 10000 ppm and 20000 ppm (904/1088 and 1839/2145 mg/kg bw/d extract): Mortality, ↓ defecation, ↑ respiration rate, tremors, ↑ activity, convulsions, ↓ body weights, ↓ food consumption, ↓ haematological parameters. Liver enlargement and congestion. ↑ Liver weight. ↑ Kidney weight and renal toxicit	-	 (BRA, MGK and SCJ) IIIA6.4.1/1  (KPIC) IIIA6.4.1

Summary table of oral sub-chronic animal studies (usually 90-day studies)						
Method, Duration of study, Route of exposure (gavage, in diet, other), Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including also target organs)	Remarks (e.g. major deviations)	Reference
OECD TG 408 90-day GLP Reliability 1 Key	Mouse, Charles River CD-1 Male/Female 15/sex/group	Pyrethrum Extract (Batch # 011831-00 Task force blend # FEK-99; Purity 57.574%) 300, 1000, 3000, 10000, 30000 ppm in diet daily <i>ad libitum</i> ; equivalent to 0, 47, 160, 460, and 1600 mg/kg bw/d total pyrethrins (0, 72, 245, 705, and 2451 mg/kg bw/d extract) for males and 0, 56, 200, 580, and 1800 mg/kg bw/d total pyrethrins (0, 86, 306, 889, and 2758 mg/kg bw/d extract) for females	NOAEL: 300 ppm, corresponding to 47 mg/kg bw/d total pyrethrins (72 mg/kg bw/d extract) in male mice and 56 mg/kg bw/d total pyrethrins (86 mg/kg bw/d extract) in female mice LOAEL: 1000 ppm, corresponding to 160 mg/kg bw/d total pyrethrins (245 mg/kg bw/d extract) in male mice and 200 mg/kg bw/d total pyrethrins (306 mg/kg bw/d extract) in female mice	1000 ppm and 3000 ppm (245/306 and 705/889 mg/kg bw/d extract): hepatotoxicity 10000 ppm (2451/2758 mg/kg bw/d extract): mortalities, hepatotoxicity 30000 ppm: all animals died	-	 (KPIC) IIIA6.4.1/0 2

Summary table of oral sub-chronic animal studies (usually 90-day studies)						
Method, Duration of study, Route of exposure (gavage, in diet, other), Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including also target organs)	Remarks (e.g. major deviations)	Reference
Compared to OECD TG 409 8 weeks GLP Reliability 1 Key	Dog Beagle dog 2 sex/dose	Pyrethrum Extract (Batch # 011831-00 Task force blend # FEK-99; Purity 57.574%) 600, 1000, 3000, 6000 ppm total pyrethrins in diet daily <i>ad libitum</i> ; Males: 0, 18, 30, 86, and 170 mg/kg bw/d total pyrethrins (0, 28, 46, 132, and 261 mg/kg bw/d extract) Females: 0, 19, 29, 94, and 200 mg/kg bw/d total pyrethrins (0, 29, 44, 144, and 306 mg/kg bw/d extract)	NOAEL: 600 ppm (18 mg/kg bw/d total pyrethrins (28 mg/kg bw/d extract) (males) and 19 mg/kg bw/d (29 mg/kg bw/d extract) (females)) LOAEL: 1000 ppm (30 mg/kg bw/d total pyrethrins (46 mg/kg bw/d extract) (males) and 29 mg/kg bw/day (44 mg/kg bw/d extract) (females))	1000 ppm (46/44 mg/kg bw/d extract): hepatotoxicity, testes weight ↑ 3000 ppm (132/144 mg/kg bw/d extract): clinical signs, haematologic changes, hepatotoxicity 6000 ppm (261/306 mg/kg bw/d extract): mortalities	-	[REDACTED] (KPIC) IIIA6.3.1/03 [REDACTED] (BRA, MGK and SCJ) IIIA6.4.1/2

Table A.39 Summary table of human data on sub-chronic oral toxicity
 No data are available.

A2.7.2.2 Sub-chronic dermal toxicity

No data are available.

A2.7.2.3 Sub-chronic inhalation toxicity

Groups of 15 Sprague Dawley rats of each sex were whole body exposed to analytical concentrations of 0, 11, 30, 100, and 356 mg a.i./m³ for 13 weeks, 6 hours/day, generally 5 days/week and a minimum of 65 exposures in total. Determinations of pesticide size distribution showed an overall mass median aerodynamic diameter of 2.7 µm.

Two animals of the highest concentration group died but only one is considered potentially exposure related.

Clinical signs were observed from 30 mg/m³ onwards and included secretory signs such as nasal discharge and dried material in the facial area in both sexes. In the highest exposure level animals laboured breathing, excess lacrimation, tremors, increased activity and matted coat were also observed. There were no ocular effects.

Body weight gains in the two highest concentration groups were lower than in controls (both sexes). Anaemia by decreased haemoglobin, haematocrit and erythrocyte values was observed from 30 mg/m³ onwards in males and at 356 mg/m³ in females, accompanied with increased leukocyte counts. Significant differences in clinical chemistry parameters were seen primarily in the highest test group, however, the differences were not consistent between sexes. Several significant increases in organ weights or ratios were observed but only the liver weight increases appear to be exposure related. Morphologic abnormalities were observed in the nasoturbinal tissues, nasopharynx, larynx and the lungs. Changes observed in these tissues in all exposure groups, including the air control group, were inflammation, oedema, haemorrhage, emphysema, macrophages, lymphoid cells, mineralisation, glandular dilation and/or goblet cell hyperplasia. These observations were generally graded in the range of minimal to moderate for animals in all exposure groups. The incidence of several other findings indicative of irritation was increased over control. These included squamous/squamoid metaplasia of the pseudostratified columnar or cuboidal epithelium in the larynx and ventral diverticulum. Also, keratinisation of the metaplastic epithelium was observed in the larynx in animals from all treated groups. Morphological abnormalities in the larynx, nasoturbines, nasopharynx and lungs observed by light microscopy were considered to be localized responses indicative of a treatment-related effect.

The NOAEL for systemic effects was 11 mg a.i./m³ air (██████████). (KPIC and BRA, MGK and SCJ)

Parameter	n = 15/sex	0 control	11 (mg/m ³)	30 (mg/m ³)	100 (mg/m ³)	356 (mg/m ³)
Leucocytes x 10 ³ /µl	M	12.10	11.74	12.25	10.54	12.8
	F	8.51	7.9	8.5	8.37	11.25 ¹⁾
Erythrocytes x 10 ⁶ /µl	M	8.96	9.04	8.61 ¹⁾	8.60 ¹⁾	8.40 ²⁾
	F	8.26	8.37	8.41	8.08	8.01
Hematocrit %	M	47.4	47.4	45.3 ¹⁾	45.8	44.7 ²⁾
	F	45.8	46.1	46.1	44.5	43.3 ²⁾

Hemoglobin g/dl	M	15.7	15.7	15.1	15.3	14.9
	F	15.4	15.6	15.5	15.1	14.6 ²⁾

¹⁾ significantly different from the control group: p < 0.05

²⁾ significantly different from the control group: p < 0.01

Parameter	n = 15/sex	0 control	11 (mg/m³)	30 (mg/m³)	100 (mg/m³)	356 (mg/m³)
AST (GOT) (IU/l)	M	60	63	62	65	61
	F	64	64	63	59	63
ALT (GPT) (IU/l)	M	29	33	29	28	27
	F	29	25	25	21 ²⁾	23 ¹⁾
Total protein (g/dL)	M	6.5	6.4	6.3	6.2	6.2 ¹⁾
	F	7.0	6.9	6.7	6.7	7.0
Globulin (g/dL)	M	2.5	2.5	2.4	2.3	2.2 ²⁾
	F	2.2	2.4	2.3	2.2	2.3
Albumin/Globulin ratio (mg/dL)	M	1.6	1.6	1.6	1.8	1.8 ¹⁾
	F	2.3	1.9 ¹⁾	1.9 ¹⁾	2.0	2.1
Glucose (mg/dL)	M	155	169	151	143	148
	F	147	146	157	133	114 ²⁾
Creatinine (mg/dL)	M	0.5	0.6	0.5	0.5	0.6
	F	0.6	0.6	0.6	0.6	0.7 ¹⁾

AST- Aspartate Amino Transferase (\triangle GOT = Glutamic Oxaloacetic Transaminase)

ALT - Alanine Amino Transferase (\triangle GPT = Glutamic Pyruvic Transaminase)

¹⁾ significantly different from the control group: p < 0.05

²⁾ significantly different from the control group: p < 0.01

Parameter	Air control		10 mg/m ³		30 mg/m ³		100 mg/m ³		350 mg/m ³	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
Terminal body weight (g)	550±40.8	345±31.6	548±52.3	327±23.0	551±52.7	335±22.1	524±31.9	319*±22.7	520±61.8	307**±22.0
Kidneys organ/body weight (x 1000)	7.47±0.51	7.26±0.52	7.43±0.95	7.70±0.69	7.57±1.10	7.51±0.54	7.77±0.54	7.98**±0.47	8.36*±0.74	8.29**±0.53
Liver weight (g)	14.8±1.57	9.23±0.81	15.0±1.68	8.98±0.82	14.8±1.68	8.98±0.87	14.3±1.49	9.14±0.82	16.4±2.02	11.0**±1.08
Liver organ/body weight (x 100)	2.69±0.15	2.68±0.19	2.73±0.17	2.74±0.19	2.69±0.18	2.68±0.26	2.73±0.19	2.86±0.22	3.16**±0.37	3.57**±0.28
Liver organ/brain (x 1)	6.64±0.70	4.48±0.45	6.81±0.77	4.39±0.44	6.72±0.78	4.25±0.49	6.44±0.84	4.39±0.41	7.41±1.09	5.27**±0.54
Lungs organ/body weight (x 1000)	3.43±0.36	4.14±0.46	3.48±0.54	4.45±0.50	3.33±0.31	4.54±0.46	3.53±0.32	4.55±0.48	3.85*±0.34	4.90**±0.46
Brain organ/body weight (x 1000)	4.07±0.27	6.02±0.59	4.04±0.36	6.28±0.46	4.03±0.33	6.34±0.44	4.27±0.34	6.54*±0.53	4.32±0.63	6.81**±0.52

*, ** Significantly different from control

Table A.40 Summary table of inhalatory sub-chronic animal studies (usually 90-day studies)

Summary table of inhalatory sub-chronic animal studies (usually 90-day studies)						
Method, Duration of study, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), form (gas, vapour, dust, mist) and particle size (MMAD), Actual and nominal concentration, Type of administration (nose only / whole body/ head only), Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including also target organs)	Remarks (e.g. major deviations)	Reference
EPA 82-4 comparable to OECD TG 413 13 weeks GLP Reliability 1 Key	Rat Charles River CD® Male/Female 15/sex/group	Pyrethrum extract (FEK-99; Purity 57.574%) Nominal: 0, 38, 68, 230, 827 mg/m ³ Analytical: 0, 11, 30, 100 and 356 mg/m ³ 5 days/week - 13 weeks MMAD = 2.7 ± 1.7 µm Whole body	NOAEL: 11 mg/m ³ LOAEL: 30 mg/m ³	11 mg/m ³ : No statistically significant effect 30 mg/m ³ : anaemia in males 100 mg/m ³ : anaemia, clinical signs (tremors), haematological changes 356 mg/m ³ : one mortality, ↑ secretory signs, ↓ body weights, nonregenerative anaemia, changes in clinical chemistry parameters, ↑ liver weights and microscopic changes in the nasopharyngeal tissues, larynx and the lungs.	-	(KPIC) (BRA, MGK and SCJ) IIIA6.4.3

Table A.41 Summary table of human data on sub-chronic inhalation toxicity
 No data are available.

A2.7.2.4 Overall conclusion on sub-chronic repeated dose toxicity related risk assessment

Value used in the Risk Assessment – Sub-chronic repeated dose systemic toxicity	
Value	AEL _{medium-term} = 0.14 mg/kg bw/d total pyrethrins (0.22 mg/kg bw/d extract)
Justification for the selected value	Based on a 1-year dog study where hepatic damage was observed at 14 mg/kg bw/d total pyrethrins (22 mg/kg bw/d extract). Since pyrethrins, and therefore <i>Chrysanthemum</i> extract, have a rodent-specific hepatotoxic mechanism, it is more appropriate to choose a study conducted without rodents. Sub-chronic study in dogs had only two animals per group since it was a range-finding study, so it is preferable to derive the AEL _{medium-term} the one-year study in dogs.
Proposed classification	Not classified.

Value/conclusion used in the Risk Assessment – Sub-chronic repeated dose local effects	
Value/conclusion	Not applicable
Justification for the selected value/conclusion	-
Proposed classification	Not classified.

A.2.7.3. Long-term repeated dose toxicity

A2.7.3.1 Long-term oral toxicity

Chrysanthemum cinerariaefolium extract from supercritical CO₂ was administered to 4 beagle dogs/sex/group for 52 weeks at the following concentrations of actual Pyrethrins: 0, 100, 500, and 2500 ppm equivalent to 0, 2.6, 13.7 and 66 mg/kg bw/d total pyrethrins (0, 3.94, 20.99, and 101.7 mg/kg bw/d extract) for males and 0, 2.8, 14.2 and 75 mg/kg bw/d total pyrethrins (0, 4.35, 21.8, and 114.3 mg/kg bw/d extract) for females. All animals survived to study termination and clinical signs were no remarkable at all dosage levels, although some animals showed some aversion to the 500 and 2500 ppm (20.99/21.8 and 101.7/114.3 mg/kg bw/d extract), dietary concentrations during the first two weeks of the study. No significant differences were observed in mean body weight values between treated and control groups, except for the males at the 100 ppm (3.94 mg/kg bw/d extract) dosage level which gained more weight than the control animals and females at the 2500 ppm (114.3 mg/kg bw/d extract) dose which were slightly less than controls. Some changes and lesions were found in all dose groups; however, these findings were not considered to be treatment related. Those findings are: reduced consumption relative to the controls during the first two weeks of the study; ocular effects; statistically significant differences in parameters such as chloride, cholesterol, urea nitrogen or globulin; macroscopic lesions such as dilatation of the lateral ventricles of the brain and discoloration and mottling of the lungs; microscopic non-neoplastic lesions such as multifocal calcification of the medulla of the kidneys, pituitary cysts, parafollicular cell hyperplasia of the thyroid, dilatation of the lateral ventricles of the cerebrum and multifocal chronic interstitial pneumonia.

Some effects noted at 2500 ppm (101.7/114.3 mg/kg bw/d extract) were considered to be treatment related. These findings were: significant increase in absolute and relative liver weights in males; increase in total leukocytes and segmented neutrophils in females; decrease in values of erythrocytes, haemoglobin and haematocrit in males, although they were not significantly different from the control; increased alanine aminotransferase in females.

The NOAEL for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ under the conditions of this study is 500 ppm equivalent to 13.7 mg/kg bw/d total pyrethrins (20.99

mg/kg bw/d extract) in males and 14.2 mg/kg bw/d total pyrethrins (21.8 mg/kg bw/d extract) in females [REDACTED]. (KPIC and BRA, MGK and SCJ)

Parameter	Assessment time	0 ppm (mg/kg bw/d)		100 ppm (mg/kg bw/d)		500 ppm (mg/kg bw/d)		2500 ppm (mg/kg bw/d)	
		Male (0)	Female (0)	Male (3.94)	Female (4.35)	Male (20.99)	Female (21.8)	Male (101.7)	Female (114.3)
Leukocytes (x 10 ³ mm ³)	Pretest	8.3 ± 1.77	9.3 ± 1.55	9.1 ± 2.13	8.3 ± 1.65	10.2 ± 1.87	8.4 ± 1.65	8.9 ± 1.93	9.3 ± 1.97
	6	7.7 ± 0.89	9.5 ± 1.91	9.0 ± 1.00	9.3 ± 3.62	10.6 ± 2.34	8.0 ± 1.18	10.0 ± 1.33	10.7 ± 3.73
	12	8.2 ± 0.87	8.6 ± 2.11	8.6 ± 1.92	10.4 ± 2.86	11.4* ± 1.74	9.4 ± 2.21	9.2 ± 1.14	17.7* ± 6.23
Segmented neutrophils (x 10 ³ mm ³)	Pretest	4.2 ± 0.71	4.7 ± 0.93	5.2 ± 1.80	4.2 ± 1.10	4.5 ± 0.56	4.4 ± 0.97	4.5 ± 1.34	5.0 ± 1.37
	6	4.9 ± 0.74	6.3 ± 1.69	6.1 ± 0.82	6.4 ± 3.43	7.3* ± 1.41	4.9 ± 0.45	6.9 ± 1.42	6.4 ± 2.87
	12	5.3 ± 1.94	4.8 ± 2.43	5.5 ± 1.00	6.0 ± 2.05	6.7 ± 0.36	6.2 ± 2.19	5.3 ± 0.33	12.9* ± 6.02

* Significantly different from the control group; p < 0.05

Parameter	Assessment time	0 ppm (mg/kg bw/d)		100 ppm (mg/kg bw/d)		500 ppm (mg/kg bw/d)		2500 ppm (mg/kg bw/d)	
		Male (0)	Female (0)	Male (3.94)	Female (4.35)	Male (0)	Female (0)	Male (3.94)	Female (4.35)
Chloride (mEq/l)	Pretest	112 ± 1.3	113 ± 3.8	112 ± 3.2	111 ± 1.0	110 ± 2.2	113 ± 0.6	110 ± 3.9	115 ± 2.4
	6	121 ± 1.3	121 ± 2.2	119 ± 2.2	120 ± 2.2	120 ± 3.4	118 ± 2.4	118 ± 2.4	122 ± 2.5
	12	124 ± 1.5	124 ± 5.4	124 ± 2.5	120 ± 3.1	120* ± 0.6	121 ± 1.0	123 ± 1.9	123 ± 1.4
Creatine phosphokinase (IU/l)	Pretest	143 ± 37.2	127 ± 46.2	83* ± 8.4	96 ± 20.5	124 ± 12.1	105 ± 30.5	119 ± 64.3	108 ± 46.6
	6	56 ± 16.7	59 ± 6.6	60 ± 6.0	59 ± 17.5	54 ± 15.8	57 ± 9.3	45 ± 5.7	71 ± 23.8
	12	54 ± 20.9	50 ± 10.8	65 ± 14.8	83 ± 44.3	43 ± 10.2	44 ± 4.2	53 ± 27.5	53 ± 11.3
Cholesterol (mg/dl)	Pretest	184 ± 13.0	175 ± 31.5	138 ± 47.5	166 ± 34.2	161 ± 29.2	163 ± 11.1	180 ± 20.3	149 ± 14.4
	6	165 ± 14.8	241 ± 74.3	155 ± 19.2	162 ± 40.6	155 ± 16.9	150* ± 22.0	140* ± 2.2	133* ± 23.9
	12	168 ± 14.2	224 ± 73.1	159 ± 28.8	172 ± 42.3	158 ± 25.5	148 ± 26.2	156 ± 16.2	151 ± 21.2
Alanine aminotransferase (IU/l)	Pretest	30 ± 3.0	29 ± 6.2	27 ± 4.6	28 ± 3.3	32 ± 0.5	28 ± 7.1	27 ± 5.8	27 ± 4.2
	6	32 ± 5.9	29 ± 6.0	29 ± 1.9	25 ± 0.8	34 ± 5.3	28 ± 4.5	40 ± 12.9	37* ± 4.1
	12	43 ± 13.5	28 ± 4.9	34 ± 3.8	26 ± 2.8	54 ± 19.3	29 ± 4.0	52 ± 19.3	39† ± 4.1
Urea nitrogen (mg/dl)	Pretest	15.5 ± 2.14	14.4 ± 2.84	13.3 ± 1.52	15.0 ± 2.95	14.3 ± 3.30	14.3 ± 2.40	16.8 ± 3.43	17.1 ± 3.79
	6	17.4 ± 2.91	17.5 ± 0.99	15.7 ± 0.94	15.1 ± 1.36	14.9 ± 3.60	15.9 ± 1.47	18.7 ± 4.60	18.6 ± 2.17
	12	20.5 ± 2.14	20.9 ± 2.23	16.9 ± 1.71	15.9† ± 1.49	16.7 ± 4.01	17.1* ± 1.69	20.8 ± 5.93	16.2* ± 2.33
Globulin (g/dl)	Pretest	1.6 ± 0.10	2.1 ± 0.10	1.9 ± 0.22	1.8 ± 0.35	1.7 ± 0.15	1.5* ± 0.22	1.6 ± 0.25	1.8 ± 0.26
	6	2.7 ± 0.38	2.4 ± 0.46	3.0 ± 0.51	2.4 ± 0.16	2.6 ± 0.47	2.0 ± 0.37	2.3 ± 0.43	2.3 ± 0.17
	12	3.1 ± 0.33	2.8 ± 0.44	3.3 ± 0.56	3.2 ± 0.34	2.9 ± 0.63	2.3 ± 0.39	2.7 ± 0.28	2.6 ± 0.22

* Significantly different from the control group; p < 0.05

† Significantly different from the control group; p < 0.01

ppm	mg/kg bw/d extract	sex (n = 4)	Body weight (kg)	Liver (g)	Kidney (g)	Pituitary (mg)	Testis (g)	Ovary (g)	Heart (g)
0	0	m	12.9	293.3	54.7	81	17.5		103.6
		f	10.1	238.0	41.3	66		0.93	80.7
100	3.94	m	13.5	306.2	56.7	76	17.7		103.6
	4.35	f	11.0	264.6	45.9	66		1.1	82.2
500	20.99	m	15.9	348.5	61.6	82	18.7		110.5
	21.8	f	10.7	263.0	41.6	78		1.05	79.7
2500	101.7	m	13.1	381.9¹⁾	61.2	67	18.0		105.6
	114.3	f	9.7	298.2	43.8	66		1.53	83.9

¹⁾ significantly different from the control group: $p < 0.05$

Table A.42 Summary table of oral long-term animal studies

Summary table of oral long-term animal studies						
Method, Duration of study, Route of exposure (gavage, in diet, other), Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	NOAEL, LOAEL	Results (all dose levels including severity and magnitude of all effects, including target organs)	Remarks (e.g. major deviations)	Reference
EPA 83- 1 comparable to OECD TG 452 52 weeks GLP Reliability 1 Key	Dog Pure bred beagle Male/Female 4/sex/group	Pyrethrum Extract (Batch # 011831-00 Task force blend # FEK-99; Purity 57.574%) Males: 0, 2.57, 13.7, and 66.4 mg/kg/d total pyrethrins (0, 3.94, 20.99, and 101.7 mg/kg bw/d extract) Females: 0, 2.84, 14.2, and 74.6 mg/kg/d total pyrethrins (0, 4.35, 21.8, and 114.3 mg/kg bw/d extract) Daily	NOAEL: 500 ppm, equal to 13.7 mg/kg bw/d total pyrethrins (20.99 mg/kg bw/d extract) for males and 14.2 mg/kg bw/d total pyrethrins (21.8 mg/kg bw/d extract) for females LOAEL: 2500 ppm, corresponding to 66.4 mg/ kg bw/d total pyrethrins (101.7 mg/kg bw/d extract) (males) and 74.6 mg/kg bw/d total pyrethrins (114.3 mg/kg bw/d extract) (females)	100 ppm (3.94/4.35 mg/kg bw/d extract): No treatment related effect 500 ppm (20.99/21.8 mg/kg bw/d extract): No treatment related effect 2500 ppm (101.7/114.3 mg/kg bw/d extract): Treatment related effects; ↑ leukocytes, segmented neutrophils and alanine aminotransferase in females; ↓ erythrocytes, haemoglobin and haematocrit: ↑ relative and absolute liver weights in males were recorded.	-	<p>██████████ (KPIC) IIIA6.4.1/03</p> <p>██████████ (BRA, MGK and SCJ) IIIA6.5</p>

Table A.43 Summary table of human data on long-term oral toxicity
 No data are available.

A2.7.2.2 Long-term dermal toxicity

No data are available.

A2.7.2.3 Long-term inhalation toxicity

No data are available.

A2.7.3.4 Overall conclusion on long-term repeated dose toxicity related risk assessment

Value used in the Risk Assessment – Long-term repeated dose systemic toxicity	
Value	AEI _{long-term} = 0.04 mg/kg bw/d total pyrethrins (0.06 mg/kg bw/d extract)
Justification for the selected value	Based on a 2-year rat (carcinogenicity) study.
Proposed classification	Not classified.

Value/conclusion used in the Risk Assessment Long-term repeated dose local effects	
Value/conclusion	Not applicable
Justification for the selected value/conclusion	-
Proposed classification	-

A.2.7.4. Specific target organ toxicity – repeated exposure (STOT RE)

A2.7.4.1 Short summary and overall relevance of the provided information on STOT RE

Effects in liver have been observed in one short-term (14 days) repeated dose study by oral exposure in mouse. These effects consisted of a statistically significant increase in absolute and relative weight at 5000 ppm (1250 mg/kg bw/d extract). In two short-term studies by oral exposure in rat and by dermal exposure in rabbit no effects were observed (see A.2.7.1).

Also, effects in liver have been observed in four sub-chronic studies: three studies by oral exposure in mouse, rat and dog and one study by inhalation exposure in rats. These effects consisted of a statistically significant increase in absolute and relative weight, congestion and hepatocellular hypertrophy (see A.2.7.2).

Route of exposure	Species	Liver weight		Congestion	Hepatocellular hypertrophy
		Absolute weight	Relative weight		
Oral	Rat	♂ 904 mg/kg bw/d extract ♀ 337 mg/kg bw/d extract	♂ 904 mg/kg bw/d extract ♀ 337 mg/kg bw/d extract	904/1088 mg/kg bw/d extract	-
	Mouse	♂ 705 mg/kg bw/d ♀ 889 mg/kg bw/d	♂ 705 mg/kg bw/d ♀ 889 mg/kg bw/d	♂ 2451 mg/kg bw/d ♀ 2758 mg/kg bw/d	♂ 705 mg/kg bw/d ♀ 889 mg/kg bw/d
	Dog	46/44 mg/kg bw/d extract	-	-	-
Inhalation	Rat	♂ - ♀ 350 mg/m ³	350 mg/m ³	-	-

In a long-term study by oral exposition in dog an increase in absolute liver weight was statistically significant in males at 101.7 mg/kg bw/d extract (see A.2.7.3).

In two carcinogenicity studies by oral exposure in rat and mouse effects in liver had been observed. In rats, these effects consisted in increased and accentuated lobulations of the liver in males at 1000 ppm, and in increased incidence of hepatic adenoma in females at 3000 ppm. In mice, from 2500 ppm onwards discoloured dark livers and increased absolute and relative liver weights were observed in males and females, and vacuolar fatty changes in the livers of males only (see A.2.9).

Accompanying these effects, an alteration in red cell parameters (hemoglobin, hematocrit and red blood cells) is observed. These effects are considered secondary to hepatotoxicity since they do not occur in the absence of hepatic impairment, except in the case of the inhalation study (where there is no strong dose relationship). Furthermore, they only occur in rats, these effects are not observed in dogs. See table below:

Study	Species	Exposure	Route	Sex	Dose	Effect (blood)	Effect (liver)
1988	Rat	3-month	Oral	♂	0 ppm	No effects	No effects
					300 ppm	No effects	No effects
					1000 ppm	No effects	No effects
					3000 ppm	No effects	↑ relative weight
					10000 ppm	No effects	↑ relative and absolute weights
					20000 ppm	↓ hemoglobin, ↓ hematocrit	↑ relative and absolute weights
				♀	0 ppm	No effects	No effects
					300 ppm	No effects	No effects
					1000 ppm	No effects	No effects
					3000 ppm	↓ hemoglobin	↑ relative and absolute weights
					10000 ppm	↓ hemoglobin, ↓ hematocrit, ↓ erythrocytes	↑ relative and absolute weights
					20000 ppm	↓ hemoglobin, ↓ hematocrit, ↓ erythrocytes	↑ relative and absolute weights
1992			Inhalation	♂	0 mg/m ³	No effects	No effects
					11 mg/m ³	No effects	No effects
					30 mg/m ³	↓ hemoglobin, ↓ hematocrit, ↓ erythrocytes	No effects
					100 mg/m ³	↓ erythrocytes	No effects
					356 mg/m ³	↓ hemoglobin, ↓ hematocrit, ↓ erythrocytes	↑ relative and absolute weights
				♀	0 mg/m ³	No effects	No effects
					11 mg/m ³	No effects	No effects
					30 mg/m ³	No effects	No effects
					100 mg/m ³	No effects	No effects
					356 mg/m ³	↓ hemoglobin, ↓ hematocrit	↑ relative and absolute weights
1988*	Dog	2-month	Oral	♂	0 ppm	No effects	No effects
					600 ppm	No effects	No effects
					1000 ppm	No effects	No effects
					3000 ppm	No effects	↑ relative and absolute weights
					6000 ppm	No effects	↑ relative and

							absolute weights
				♀	0 ppm	No effects	No effects
					600 ppm	No effects	No effects
					1000 ppm	No effects	↑ absolute weight
					3000 ppm	No effects	↑ relative and absolute weights
					6000 ppm	No effects	Both animals were sacrificed <i>in extremis</i>
		1-year		♂	0 ppm	No effects	No effects
					100 ppm	No effects	No effects
					500 ppm	No effects	No effects
					2500 ppm	No effects	↑ relative and absolute weights
				♀	0 ppm	No effects	No effects
					100 ppm	No effects	No effects
					500 ppm	No effects	No effects
					2500 ppm	No effects	No effects

*No statistical analysis was performed.

Pyrethrins did not show toxic effects in a two-generation study in rat and in two teratogenicity studies in rat and rabbit (see A.2.10).

A2.7.4.2 Comparison with the CLP criteria

No data are available to indicate that the active substance should be classified for STOT RE. Observed effects are not organ-specific and these are produced at doses higher than those indicated in CLP 3.9.2.9.7. (Table 3.9.3).

A2.7.4.3 Conclusion on classification and labelling for STOT RE

Effects observed in liver are species-specific adaptive responses as explained in A.2.9.1, so *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ does not meet the EU criteria to be classified as STOT RE.

A.2.8. Genotoxicity / Germ cell mutagenicity

A total of six studies was performed. No indication for mutagenic potential of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was observed. No indications for genotoxicity were detected. (KPIC)

A.2.8.1. In vitro

***In vitro* unscheduled DNA synthesis assay (rat primary hepatocytes)**

Chrysanthemum cinerariaefolium extract from supercritical CO₂ was tested in the unscheduled DNA synthesis assay in male Fischer rat primary hepatocytes with a confirmatory assay. The test article was evaluated at 0.03, 0.1, 0.3, 0.6 and 1.0 µl/ml in both the initial and the confirmatory assay. In a parallel cytotoxicity assay (LDH activity measure) the test article was tested at eight dose levels ranging from 0.003 to 3.0 µL/mL in the initial assay and at seven dose levels ranging from 0.01 to 3.0 µL/mL in the confirmatory assay. The released LDH values from the preliminary cytotoxicity assay indicated that the presence of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ may have interfered with the LDH assay. Therefore, in this and subsequent cytotoxicity assays, the amount of LDH released from the test article + 1% Triton condition was defined as 100% toxicity. Using this procedure, relative toxicities of 92, 84 and 35% were observed at concentrations of 10, 3.0 and 1.0 µL/mL, respectively. When hepatocyte cultures were examined microscopically, toxicity was also assessed at 10, 3.0, 1.0 and 0.3 µL/mL. At lower concentrations, normal cellular morphology was observed. The results of both the initial and confirmatory UDS assays indicate that none of the test article doses caused significant increase in the mean number of net nuclear grains counts when compared to the appropriate solvent control (DMBA), which induced a significant increase in the average net nuclear counts. Therefore, it is concluded that *Chrysanthemum*

cinerariaefolium extract from supercritical CO₂ has not shown any evidence of causing DNA damage in rat liver in this *in vitro* test system (Curren, 1989). (BRA, MGK and SCJ)

***In vitro* gene mutation study in bacteria (*Salmonella* sp.)**

Chrysanthemum cinerariaefolium extract from supercritical CO₂ was tested in the *Salmonella*/mammalian-microsome plate incorporation assay using the following bacterial strains: TA98, TA100, TA1535, TA1537 and TA1538. The assay was carried out in the presence and in the absence of an S9 activation system. *Escherichia coli* WP2 uvrA, *E. coli* WP2 uvrA (pKM101), or *S. typhimurium* TA102 were not included in the assay to detect cross-linking mutagens as the guideline recommends. The test article concentrations used in each study with and without metabolic activation were: 8.8 - 8772 µg/plate in the dose range finding study and 292 - 8772 µg/plate in the mutagenicity and confirmatory assay.

In the dose range-finding study no appreciable toxicity was observed up to 8772 µg per plate. In the mutagenicity assay no positive responses were observed with any of the tester strains in the presence of microsomal enzymes or with tester strains TA100, TA1535 or TA1538 in the absence of microsomal enzymes. Tester strains TA98 and TA1537 exhibited 2.0-fold non-dose responsive increases in the absence of microsomal enzymes. However, non-dose responsive increases are not evaluated as positive.

In the confirmatory assay no positive responses were observed with any of the tester strains either in the presence or the absence of microsomal enzymes. Dose-responsive increases observed with tester strain TA100 in the presence and absence of microsomal enzymes were noted but as they were at less than a 2-fold increase, they were not evaluated as positive. *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ did not cause a positive response with any of the tester strains either in the presence or absence of microsomal enzymes prepared from Aroclor induced rat liver (San & Springfield, 1989). (KPIC and BRA, MGK and SCJ)

***In vitro* cytogenicity study in mammalian cells (CHO-K1)**

An *in vitro* mammalian chromosome aberration test was carried out on Chinese hamster ovary (CHO-K1) cells with the test substance *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. The test article concentrations used in each study with and without metabolic activation were: 0.03 - 300 µg/ml in the preliminary cytotoxicity assay and 6.25 - 150 µg/ml in the chromosome aberration assay. In the cytotoxicity assay cell growth inhibitions relative to the solvent control were 82% and 100% at 100 and 300 µg/ml in the non-activated test system and 53% and 100% at 100 and 300 µg/ml in the S9-activated test system. The average generation time was delayed from 12.2 to 24 hours at the 100 µg/mL dose level in the non-activated study and from 12.5 to 24 hours at the 100 µg/ml dose level in the S9-activated study. At the 300 µg/ml dose level, complete inhibition of the cell cycle was observed in both the non-activated and the S9-activated test systems. Toxicity (cell growth inhibition) in the chromosome aberration test was between 65% and 100% at concentrations between 50 and 150 µL/mL. Under these concentrations the toxicity was found to be more than 50% cell growth inhibition. The specific data were as follows:

Without S9 mix:

72% cell growth inhibition at 85 µg/ml (12 hours)

78% cell growth inhibition at 100 µg/ml (24 hours)

88% cell growth inhibition at 50 µg/ml (48 hours)

With S9 mix:

65% cell growth inhibition at 70 µg/ml (12 hours)

66% cell growth inhibition at 100 µg/ml (24 hours)

100% cell growth inhibition at 85 µg/ml (48 hours)

No statistically significant increases in chromosome aberrations were observed in the non-activated or S9-activated test systems relative to the solvent control group when cells are harvested at 12, 24 and 48 hours after treatment initiation. Under the conditions of this assay, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was concluded to be negative in the chromosome aberration assay using CHO-K1 cells (Curry, 1996). (BRA, MGK and SCJ)

In an almost equal assay, with the only difference being a narrower concentration range (6.25 - 100 µg/mL), the same results were observed (Curry, 1996). (KPIC)

***In vitro* gene mutation assay in mammalian cells (L5178Y)**

Pyrethrin was examined for its potential to induce gene mutations at the Thymidine Kinase (TK)-locus of cultured mouse lymphoma L5178Y cells, in both the absence and the presence of a metabolic activation system (S9-mix). The mutant colonies at concentrations of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ were scored using the criteria of small and large colonies. Three assays were carried out: Test 1: TK assay with and without metabolic activation at 0.39 - 1200 µg/mL. Test 2: TK assay at 3.0 - 50 µg/mL without metabolic activation, and at 3.0 - 100 µg/mL with metabolic activation. Test 3: TK assay with metabolic activation at 13 - 72 µg/mL. Cytotoxicity was determined in each test measuring the relative total growth (RTG).

The highest concentration of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ evaluated was 50 µg/mL in the absence of S9-mix and 85 µg/mL with metabolic activation. *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was cytotoxic to L5178Y cells with and without S9 mix since the RTG was decreased in a dose-related manner above a concentration of 25 µg/mL. At the highest concentrations, the cytotoxicity test indicated a RTG of 6% at 50 µg/mL without metabolic activation and of 3% at 85 µg/mL in the presence of S9-mix. In the mutagenic assay without metabolic activation, the mutant frequency (MF) was not significantly increased at any dose level in either the first or the second assay. However, regarding the first assay, in the presence of S9-mix, the positive control (10 µg/mL MCA) did not comply with the criteria of validation. The positive control and negative control had a similar TK mutant frequency whereas the TK mutant frequency for the positive control should have been higher than 400 TFT-resistant mutants per 10⁶ clonable cells and should have been at least twice that of the corresponding negative control. Therefore, this assay was not considered to be valid. About the second assay, in the presence of S9-mix, the mutant frequency was significantly increased at concentration of 85 and 52 µg/mL of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂, and equivocal response was observed at concentration of 61 and 26 µg/mL. The mutant frequencies were increased by 421, 118, 94 and 59 per 10⁶ clonable cells. In general, the amount of small and large colonies was more or less equal. Since 90% cytotoxicity was observed at 85 µg/mL in the presence of S9-mix, the increase in MF may be considered as an artefact. Since 90% cytotoxicity was observed at 85 µg/mL in the presence of S9-mix, the increase in mutant frequency may be considered as an artefact. Hence, the second assay was also not considered to be valid. In the third assay no significant increase of the MF at any dose level was observed. Since no positive and equivocal responses were obtained in the third assay, the findings in the second assay were not of significance.

It was concluded that, under the conditions used in this study, the test substance *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is not mutagenic at the TK-locus of mouse lymphoma L5178Y cells (Steenwinkel, 2001). (BRA, MGK and SCJ)

In an almost equal assay, three tests were performed at the same concentrations with the only difference being the concentrations of the second test with metabolic activation which are the same as without metabolic activation. *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ did not induce a reproducible significant increase in mutant frequency, neither in the S9-activated nor in the nonactivated system. Also no dose related increase in mutant frequency was observed.

On the basis of this study it is concluded that the test substance *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is not mutagenic at the TK-locus of mouse lymphoma L5178Y cells (Steenwinkel, 2001). (KPIC)

Further Studies

New GLP-compliant studies on genotoxic activity are provided for two Pyrethrum extracts, Pyrethrum-Extract 50% and Pyroicide 50%. These include *in vitro* gene mutation studies in bacteria (OECD TG 471), *in vitro* cytogenicity studies in mammalian cells (OECD TG 487) and *in vitro* gene mutation studies in mammalian cells (OECD TG 490). These are briefly summarised below.

Microbial Mutagenicity studies (*Salmonella* sp.)

The mutagenic activity of Pyrethrum-Extract 50% and Pyroicide 50% was investigated in reverse gene mutation assays in bacteria according to the testing guideline OECD TG 471. Strains TA98, TA100, TA1535, and TA1537 of *S. typhimurium* and tester strain *E. coli* WP2 uvrA were exposed to the test items with and without metabolic activation in two independent experiments.

Pyrethrum-Extract 50% (KPIC) was tested with and without metabolic activation at 7 dose levels in experiment I (3.16 - 2500 µg/plate) and experiment II (2 - 2000 µg/plate). In this case, precipitation was noted at concentrations of 1000 µg/plate and above, therefore testing was done only up to 2500 µg/plate in the first and 2000 µg/plate in the second experiment. Pyroicide 50% (MGK) was tested at concentrations of 31.6 - 5000 µg/plate in both experiments with and without metabolic activation. PY-T-50 Pale Refined Pyrethrins (BRA) was tested with and without metabolic activation at 8 dose levels in experiment I (3.16 - 5000 µg/plate) and at 7 dose levels in experiment II (2.0 - 2000 µg/plate).

No mutagenic effect (no increase in revertant colony numbers as compared with control counts) was observed (MF ≤ 1.5%) for any of the test items tested up to the top concentration in any of the 5 test strains in two independent experiments without and with metabolic activation.

All test items (Pyrethrum-Extract 50%, Pyroicide 50%, and PY-T-50 Pale Refined Pyrethrins) were found not to be mutagenic when tested on *S. typhimurium* strains TA98, TA100, TA1535, and TA1537 and tester strain *E. coli* WP2 uvrA with or without S9-mix activation and can, therefore, be considered non-genotoxic in this bacterial reverse mutation assay. (KPIC and BRA, MGK and SCJ)

***In vitro* cytogenicity study in mammalian cells (V79)**

In order to investigate the cytogenetic potential in Chinese hamster V79 cells, *in vitro* micronucleus assays following the testing guideline OECD TG 487 were performed with Pyrethrum-Extract 50% and Pyroicide 50%. The selected concentrations were based on the results of a pre-experiment for cytotoxicity. Two independent experiments were carried out for each extract.

With the test item Pyrethrum-Extract 50% (KPIC), experiment I employed an exposure period of 4 h with and without metabolic activation at concentrations of 0.0010 - 0.0050 µL/mL without S9-mix and 0.0125 - 0.050 µL/mL with S9-mix. Experiment II was carried out without metabolic activation at 0.50 - 0.010 µL/mL.

In experiment I without S9-mix, no increase of the cytostasis above 30% was noted at 0.0010 µL/mL. At 0.0025 µL/mL a cytostasis of 31% and at 0.0050 µL/mL a cytostasis of 68% was noted. In experiment I with S9-mix, no increase of the cytostasis above 30% was noted up to 0.0250 µL/mL. At 0.050 µL/mL a cytostasis of 61% was noted. In experiment II without S9-mix, no increase of the cytostasis above 30% was noted up to 0.0050 µL/mL. At 0.075 µL/mL a cytostasis of 32% was noted and at 0.010 µL/mL a cytostasis of 49%. In some cases, although cytostasis > 30%, there are not statistically significant differences with respect to the control group, so under the conditions of this assay, Pyrethrum-Extract 50% was concluded to be negative in the cytogenicity study using V79 cells.

With the test item Pyrocide 50% (MGK), experiment I employed an exposure period of 4 h with and without metabolic activation at concentrations of 0.00010 - 0.010 µL/mL without S9-mix and 0.0125 - 0.10 µL/mL with S9-mix.

Experiment II was carried out without metabolic activation at 0.0005 - and 0.025 µL/mL and 0.010 - 0.10 µL/mL with S9-mix.

In experiment I without S9-mix, no increase of the cytostasis above 30% was noted at 0.0040 µL/mL. At 0.0050 µL/mL a cytostasis of 63% was noted. In experiment I with S9-mix, no increase of the cytostasis above 30% was noted up to 0.050 µL/mL. At 0.060 µL/mL a cytostasis of 51% was noted. In experiment II without S9-mix, no increase of the cytostasis above 30% was noted up to 0.0050 µL/mL. At 0.075 µL/mL a cytostasis of 50% was noted. In experiment I with S9-mix, no increase of the cytostasis above 30% was noted up to 0.040 µL/mL. At 0.065 µL/mL a cytostasis of 48% was noted and at 0.085 µL/mL a cytostasis of 54% was noted. In some cases, although cytostasis > 30%, there are not statistically significant differences with respect to the control group, so under the conditions of this assay, Pyrocide 50% was concluded to be negative in the cytogenicity study using V79 cells.

With the test item PY-T-50 Pale Refined Pyrethrins (BRA), experiment I employed an exposure period of 4 h with and without metabolic activation at concentrations of 0.0025 - 0.0060 µL/mL without S9-mix and 0.025 - 0.10 µL/mL with S9-mix. Experiment II was carried out without metabolic activation at 0.025 - 0.010 µL/mL.

In experiment I without S9-mix, no increase of the cytostasis above 30% was noted at 0.0025 µL/mL. At 0.0040 µL/mL a cytostasis of 44% and at 0.0060 µL/mL a cytostasis of 63% was noted. In experiment I with S9-mix, no increase of the cytostasis above 30% was noted up to 0.050 µL/mL. At 0.10 µL/mL a cytostasis of 51% was noted. In experiment II without S9-mix, no increase of the cytostasis above 30% was noted up to 0.0050 µL/mL. At 0.010 µL/mL a cytostasis of 54% was noted. In some cases, although cytostasis > 30%, there are not statistically significant differences with respect to the control group, so under the conditions of this assay, PY-T-50 Pale Refined Pyrethrins was concluded to be negative in the cytogenicity study using V79 cells.

***In vitro* gene mutation study in mammalian cells (L5178Y)**

The three test items, Pyrethrum-Extract 50%, Pyrocide 50%, and PY-T-50 Pale Refined Pyrethrins were examined for the potential to induce gene mutations at the TK-locus of cultured mouse lymphoma L5178Y cells in both the absence and presence of an S9 metabolic activation system in two independent experiments. The two test items exhibited considerable and similar cytotoxicity with EC₅₀ values in the range of 19 to 56 ng/mL.

Table A.44 Cytotoxicity of the three test items (EC₅₀) and genotoxicity in mouse lymphoma L5178Y cells

	Cytotoxicity EC ₅₀		Genotoxicity	
	without S9 (4h) [µg/mL]	with S9 (24h) [µg/mL]	without S9	with S9
Pyrethrum-Extract Pale 50%	0.056	0.056	negative	negative
Pyrocide 50%	0.021	0.037	negative	negative
PY-T-50 Pale Refined Pyrethrins	0.019	0.044	negative	negative

The test item Pyrethrum-Extract Pale 50% (KPIC) was tested in experiment I at concentrations of 0.0025 - 0.12 µL/mL without metabolic activation, and at concentrations of 0.0005 - 0.075 µL/mL with metabolic activation. Experiment II was carried out at concentrations of 0.005 - 0.03 µL/mL without metabolic activation, and at concentrations of 0.02 - 0.085 µL/mL with metabolic activation. In experiments I and II no biologically relevant increase in the number of mutants was found after treatment with Pyrethrum Extract Pale 50% with or without metabolic activation. The Global Evaluation Factor (GEF)

was not exceeded (< 126) by the induced mutant frequency at any concentration. Additionally, colony sizing showed no clastogenic effects induced by the test item.

The test item Pyrocide 50% (MGK) was tested in experiment I at concentrations of 0.01 – 0.0375 µL/mL without metabolic activation, and at concentrations of 0.02 - 0.08 µL/mL with metabolic activation. Experiment II was carried out at concentrations of 0.001 - 0.02 µL/mL without metabolic activation, and at concentrations of 0.015 - 0.085 µL/mL with metabolic activation. In experiments I and II no biologically relevant increase in the number of mutants was found after treatment with Pyrocide 50% with or without metabolic activation. The GEF was not exceeded (< 126) by the induced mutant frequency at any concentration. Additionally, colony sizing showed no clastogenic effects induced by the test item.

The test item PY-T-50 Pale Refined Pyrethrins (BRA) was tested in experiment I at concentrations of 0.0025 – 0.035 µL/mL without metabolic activation, and at concentrations of 0.005 - 0.08 µL/mL with metabolic activation. Experiment II was carried out at concentrations of 0.0025 - 0.024 µL/mL without metabolic activation, and at concentrations of 0.01 - 0.075 µL/mL with metabolic activation. In experiments I and II no biologically relevant increase in the number of mutants was found after treatment with PY-T-50 Pale Refined Pyrethrins with or without metabolic activation. The GEF was not exceeded (< 126) by the induced mutant frequency at any concentration. Additionally, colony sizing showed no clastogenic effects induced by the test item.

Under the conditions of these assays following OECD TG 490, Pyrethrum-Extract 50%, Pyrocide 50%, and PY-T-50 Pale Refined Pyrethrins tested up to cytotoxic concentrations in the absence and presence of metabolic activation, did neither induce mutations nor had any chromosomal aberration potential.

Table A.45 Summary table of *in vitro* genotoxicity studies

Summary table of <i>in vitro</i> genotoxicity studies					
Method, Guideline, GLP status, Reliability, Key/supportive study	Test substance (including purity), Vehicle, Doses	Relevant information about the study (e.g. organism (e.g. bacteria), cell type, strains)	Results (including cytotoxicity and +/-S9 mix)	Remarks (e.g. major deviations)	Reference
Bacterial assay gene mutation EPA F, 84-2 GLP Reliability 1 Key	Pyrethrum Extract (Blend FEK-99; Purity 57.574% for KPIC and 57.55% for BRA, MGK and SCJ) 292 585, 877, 2924, 5848, and 8772 µg/plate ± S9	<i>S. typhimurium</i> TA98, TA100, TA1535, TA1537 TA 1538	+S9: - -S9: -	Cytotoxicity not evaluated. Non-mutagenic to bacteria	San R.H.C. & Springfield K.A. (1989) (KPIC) (BRA, MGK and SCJ) IIIA 6.6.1
Clastogenicity mammalian cells Chromosomal aberrations EPA F, 84-2 OECD 473 GLP Reliability 1 Key	Pyrethrum Extract Blend FEK-99 Code 96AC14; Purity 55.98% (w/w) 0, 6.25, 12.5, 25, 40, 50, 55, 70, 85, 100 and 150 µg/mL (BRA, MGK and SCJ) 6.25 - 100 µg/mL (KPIC)	Chinese hamster ovary cells (CHO-K1)	+S9: - -S9: -	Cytotoxicity at 300 µg/mL S-9 activated, at 100 and 300 µg/mL in the non-activated system	Curry P.T. (1996) (KPIC) (BRA, MGK and SCJ) IIIA6.6.2
Clastogenicity mammalian cells Chromosomal aberrations EPA F, 84-2 GLP	Pyrethrum Extract (Blend FEK-99; Purity 57.55%) 0.02 - 0.32 µL/mL	Chinese hamster ovary cells (CHO-K1)	+S9: - -S9: -	Cytotoxicity at the high dose in both the non-activated and the S-9 activated	Putman & Morris (1989) A 6.6.2/02 (KPIC)

Reliability 1 Key				studies	
Mammalian cells gene mutation OECD 476 GLP Reliability 1 Key	Pyrethrum Extract (Blend FEK-99; Purity 57.03%) Preliminary cytotoxicity test: 0, 0.39, 0.78, 1.6, 3.1, 6.2, 12.5, 25, 50, 100, 200, 400, 800 and 1200 µg/mL Mutagenic assay: test 1 without metabolic activation (and with metabolic activation for KPIC): 0, 3.0, 4.3, 6.1, 8.8, 12.5, 17.9, 25.6, 31, 36.5, 40.5, 45 and 50 µg/mL with metabolic activation 0, 3.0, 4.3, 6.1, 8.8, 12.5, 17.9, 25.6, 36.5, 52.2, 61.4, 72, 85 and 100 µg/mL test 2 with metabolic activation: 0, 13, 26, 52, 61 and 72 µg/mL	TK-locus L5178Y cells	+S9: - -S9: -	Cytotoxicity at 26 µg/mL in non-activated systems 52 µg/mL in S9-activated systems	Steenwinkel M-J.S.T. (2001) (KPIC) IIIA6.6.3/01 Steenwinkel M-J.S.T. (2001) (BRA, MGK and SCJ) IIIA6.6.3
Unscheduled DNA synthesis assay EPA 84-2, OECD 482 GLP Reliability 1 Key	Pyrethrum extract (Batch Blend FEK-99; Purity Pyrethrins I = 37.67% (w/w), Pyrethrins II = 19.98% (w/w), total Pyrethrins = 57.55%) Parallel cytotoxicity assay: Eight dose levels ranging from 0.003 to 3.0 µL/mL. Unscheduled DNA synthesis assay: Six dose levels ranging from 0.03 to 1.0 µL/mL.	Primary male rat hepatocyte cell culture	+S9: - -S9: -	As hepatocytes are used, the metabolic activation system is being tested. No increase in the mean number of net nuclear grains count. No DNA damages	Curren R.D. (1989) (BRA, MGK and SCJ) IIIA6.6

<p><i>In vitro</i> gene mutation study in bacteria OECD 471 EU Method B.13/14 EPA OPPTS 870.5100 GLP Reliability 1 Key</p>	<p>Pyrethrum Extract 50% (Batch number: 2016-5-BB; Purity 50.24%) Pre-experiment: 3.16, 10.0, 31.6, 100, 316, 1000, 2500 and 5000 µg/plate Experiment I: 3.16, 10.0, 31.6, 100, 316, 1000, 2500 µg/plate Experiment II: 2.00, 6.32, 20.0, 63.2, 200, 632, 2000 µg/plate</p>	<p><i>S. typhimurium</i> TA1535, TA1537, TA98, TA100 and <i>E. coli</i> WP2 uvrA</p>	<p>+S9: - -S9: -</p>	<p>Pyrethrum Extract 50% did not cause gene mutations by base pair changes or frameshifts in the genome of the tester strains used.</p>	<p>Schreib G. (2016a) (KPIC)</p>
<p><i>In vitro</i> gene mutation study in bacteria OECD 471 EU Method B.13/14 EPA OPPTS 870.5100 GLP Reliability 1 Key</p>	<p>Pyrocide 50% (Batch No: 10209; Purity 53.72%) Pre-experiment: 3.16, 10.0, 31.6, 100, 316, 1000, 2500 and 5000 µg/plate Experiment I and experiment II (with and without metabolic activation): 31.6, 100, 316, 1000, 2500 and 5000 µg/plate</p>	<p><i>S. typhimurium</i> TA1535, TA1537, TA98, TA 100 and <i>E. coli</i> WP2 uvrA</p>	<p>+S9: - -S9: -</p>	<p>Pyrocide 50% did not cause gene mutations by base pair changes or frameshifts in the genome of the tester strains used.</p>	<p>Schreib G. (2016b) (MGK)</p>
<p><i>In vitro</i> gene mutation study in bacteria OECD 471 EC Method B.13/14 EPA OPPTS 870.5100 GLP Reliability 1 Key</p>	<p>PY-T-50 (Batch number: 0116-501-610; Purity 49.35%) Pre-experiment: 3.16, 10.0, 31.6, 100, 316, 100, 316, 1000, 2500, 5000 µg/plate Experiment I: 3.16, 10.0, 31.6, 100, 316, 1000, 2500 µg/plate Experiment II: 2.0, 6.32, 20.0, 63.2, 200, 632, 2000 µg/plate</p>	<p><i>S. typhimurium</i> TA1535, TA1537, TA98, TA100 and <i>E. coli</i> WP2 uvrA</p>	<p>+S9: - -S9: -</p>	<p>PY-T-50 did not cause gene mutations by base pair changes or frameshifts in the genome of the tester strains used.</p>	<p>Schreib G. (2016c) (BRA)</p>
<p><i>In vitro</i> cytogenicity study in mammalian cells GLP</p>	<p>Pyrethrum Extract 50% (Batch number: 2016-5-BB; Purity 50.24%)</p>	<p><i>In vitro</i> Mammalian Micronucleus Assay V79 Cells</p>	<p>Negative</p>	<p>Pyrethrum Extract 50% did not induce structural</p>	<p>Donath C. (2016a) (KPIC)</p>

Reliability 1 Key	<p>Pre-experiment without S9: 0.0078, 0.0156, 0.0313, and 0.0625 µL/mL</p> <p>Pre-experiment with S9: 0.0078, 0.0156, 0.0313, 0.0625, 0.125, 0.25, and 0.5 µL/mL</p> <p>Experiment I without S9: 0.0010, 0.0025, 0.0040, and 0.0050 µL/mL</p> <p>Experiment I with S9: 0.0125, 0.025, and 0.050 µL/mL</p> <p>Experiment II without S9: 0.0050, 0.0075, and 0.010 µL/mL</p>			and/or numerical chromosomal damage in Chinese Hamster V79 cells.	
<i>In vitro</i> cytogenicity study in mammalian cells GLP Reliability 1 Key	<p>Pyrocide 50% (Batch No: 10209; Purity 53.72%)</p> <p>Pre-experiment without metabolic activation: 0.0078, 0.0156, and 0.0313 µL/mL</p> <p>Pre-experiment with metabolic activation: 0.0078, 0.0156, 0.0313, 0.0625, and 0.125 µL/mL</p> <p>Experiment I without S9: 0.0025, 0.0040, and 0.0050 µL/mL</p> <p>Experiment I with S9: 0.025, 0.050, and 0.060 µL/mL</p> <p>Experiment II without S9: 0.0025, 0.0050, and 0.0075 µL/mL</p> <p>Experiment II with S9: 0.040, 0.065, and 0.085µL/mL</p>	<i>In vitro</i> Mammalian Micronucleus Assay V79 Cells	Negative	Pyrocide 50% did not induce structural and/or numerical chromosomal damage in Chinese hamster V79 cells.	Donath C. (2016b) (MGK)
<i>In vitro</i> cytogenicity study in mammalian cells GLP	PY-T-50 (Batch number: 0116-501-610; Purity 49.35%)	<i>In vitro</i> Mammalian Micronucleus Assay V79 Cells	Negative	PY-T-50 Pale Refined Pyrethrins did not induce	Donath C. (2016c) (BRA)

<p>Reliability 1 Key</p>	<p>Pre-experiment without metabolic activation: 0.0078, 0.0156, and 0.0313 µL/mL</p> <p>Pre-experiment with metabolic activation: 0.0078, 0.0156, 0.0313, 0.0625, and 0.125 µL/mL</p> <p>Experiment I without S9: 0.0025, 0.0040, and 0.0060 µL/mL</p> <p>Experiment I with S9: 0.025, 0.050, and 0.10 µL/mL</p> <p>Experiment II without S9: 0.0025, 0.0050, and 0.010µL/mL</p>			<p>structural and/or numerical chromosomal damage in Chinese Hamster V79 cells.</p>	
<p><i>In vitro</i> gene mutation study in mammalian cells GLP Reliability 1 Key</p>	<p>Pyrethrum Extract 50% (Batch number: 2016-5-BB; Purity 50.24%)</p> <p>Pre-experiment I: 0.025, 0.05, 0.075, 0.1, 0.25, 0.5, 0.75, 1, 1.5, and 2 µL/mL</p> <p>Pre-experiment II without metabolic activation: 0.005, 0.01, 0.02, 0.04, 0.06, 0.08, 0.1, and 0.12 µL/mL</p> <p>Experiment I: Without metabolic activation: 0.0025, 0.0050, 0.010, 0.025, 0.05, 0.075, 0.1, and 0.12µL/mL With metabolic activation: 0.0005, 0.001, 0.0025, 0.005, 0.01, 0.025, 0.05, and 0.075 µL/mL</p> <p>Experiment II: Without metabolic activation: 0.005, 0.01, 0.014, 0.018, 0.022, 0.024, 0.026, and 0.03 µL/mL</p>	<p>Gene mutations (TK-locus) mouse lymphoma L5178Y cells</p>	<p>Negative</p>	<p>Pyrethrum extract 50% is considered to be non-mutagenic in the <i>in vitro</i> mammalian cell gene mutation assay (thymidine Kinase locus) in mouse lymphoma L5178Y cells.</p>	<p>Wallner B. (2016a) (KPIC)</p>

	With metabolic activation: 0.010, 0.02, 0.04, 0.05, 0.06, 0.07, 0.075, 0.08, and 0.085 µL/mL				
<i>In vitro</i> gene mutation study in mammalian cells GLP Reliability 1 Key	<p>Pyroicide 50% (Batch No: 10209; Purity 53.72%)</p> <p>Pre-experiment I: 0.025, 0.05, 0.075, 0.1, 0.25, 0.5, 0.75, 1, 1.5, and 2 µL/mL</p> <p>Pre-experiment II without metabolic activation: 0.005, 0.01, 0.015, 0.02, 0.025, 0.03, 0.035, and 0.04 µL/mL</p> <p>Experiment I:</p> <p>Without metabolic activation: 0.010, 0.015, 0.020, 0.025, 0.030, 0.0325, 0.035, and 0.0375 µL/mL</p> <p>With metabolic activation: 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.075, and 0.08 µL/mL</p> <p>Experiment II:</p> <p>Without metabolic activation 0.001, 0.002, 0.006, 0.010, 0.014, 0.016, 0.018, and 0.020 µL/mL</p> <p>With metabolic activation 0.015, 0.025, 0.035, 0.045, 0.055, 0.065, 0.075, and 0.08 µL/mL</p>	Gene mutations (TK-locus) mouse lymphoma L5178Y cells	Negative	Pyroicide 50% is considered to be non-mutagenic in the <i>in vitro</i> mammalian cell gene mutation assay (thymidine Kinase locus) in mouse lymphoma L5178Y cells.	Wallner B. (2016b) (MGK)
<i>In vitro</i> gene mutation study in mammalian cells GLP Reliability 1 Key	<p>PY-T-50 (Batch number: 0116-501-610; Purity 49.35%)</p> <p>Pre-experiment I: 0.025, 0.05, 0.075, 0.1, 0.25, 0.5, 0.75, 1, 1.5, and 2 µL/mL</p> <p>Pre-experiment II without metabolic activation: 0.001,</p>	Gene mutations (TK-locus) mouse lymphoma L5178Y cells	Negative	PY-T-50 Pale Refined Pyrethrins is considered to be non-mutagenic in the <i>in vitro</i>	Wallner B. (2016c) (BRA)

	<p>0.0025, 0.005, 0.01, 0.02, 0.04, 0.06, and 0.08 µL/mL</p> <p>Experiment I:</p> <p>Without metabolic activation: 0.0025, 0.0050, 0.010, 0.020, 0.030, and 0.035 µL/mL</p> <p>With metabolic activation: 0.005, 0.01, 0.02, 0.04, 0.06, and 0.08 µL/mL</p> <p>Experiment II:</p> <p>Without metabolic activation 0.0025, 0.0050, 0.010, 0.016, 0.018, 0.020, 0.022, and 0.024 µL/mL</p> <p>With metabolic activation 0.010, 0.025, 0.050, 0.055, 0.060, 0.065, 0.070, and 0.075 µL/mL</p>			mammalian cell gene mutation assay (thymidine Kinase locus) in mouse lymphoma L5178Y cells.	
--	--	--	--	---	--

Conclusion used in Risk Assessment – Genotoxicity <i>in vitro</i>	
Conclusion	Not genotoxic
Justification for the conclusion	<p>No mutagenic activity was observed for Pyrethrum-Extract 50%, Pyroicide 50%, and PY-T-50 Pale Refined Pyrethrins in a reverse gene mutation assay in bacteria according to the testing guideline OECD TG 471.</p> <p>No biologically relevant increase of the micronucleus frequency was noted in an <i>in vitro</i> micronucleus test following OECD TG 487 in Chinese hamster V79 cells. Neither test item induced structural and/or numerical chromosomal damage. When examined for the potential to induce gene mutations at the TK-locus of cultured mouse lymphoma L5178Y cells in both the absence and presence of an S9 metabolic activation system according to OECD TG 490, there was no induction of mutations nor was any chromosomal aberration potential observed.</p> <p>The three test items, Pyrethrum-Extract 50%, Pyroicide 50%, and PY-T-50 Pale Refined Pyrethrins, are considered non-genotoxic based on these new experimental data. The same lack of genotoxic activity was reported for the blend FEK-99 in a battery of <i>in vitro</i> tests addressing the same endpoints.</p>

A.2.8.2. *In vivo*

In order to investigate the cytogenetic potential in CFLP mice, *in vivo* micronucleus assay was performed with a mix of 28.4% Pyrethrin I and 25.0% Pyrethrin II. The selected concentrations were based on the results of a dose-finding study. Mice were treated by gavage in two dosages separated by an interval of 24 hours at total dose levels of 0.25, 0.5 and 1.0 mL/kg bw. A negative control group was treated with the vehicle, corn oil, alone. Six hours after the last dose, animals were killed by cervical dislocation and bone marrow smears were analysed microscopically by counting micronuclei in 2000 polychromatic erythrocytes per animal.

There was no increase in the number of micronuclei in treated animals as compared to control group, all counts were within the laboratory standard range for negative controls. In conclusion, the test item was not genotoxic *in vivo*.

Table A6.6.4-1. Summary of micronucleus results in male and female mice¹⁾

Dose mL/kg bw	Micronuclei per 2000 polychromatic erythrocytes per animal	
	Mean	Range
0	2.3	0 – 5
0.25	1.9	0 – 4
0.5	2.2	0 – 6
1.0	1.6	0 – 4

¹⁾ Male and female combined values because there were no sex differences in micronucleus formation

Table A.46 Summary table of *in vivo* genotoxicity studies


Summary table of <i>in vivo</i> genotoxicity studies						
Method, duration of study, Guideline, GLP status, Reliability, Key/supportive study	Test substance (including purity), Vehicle, Doses	Relevant information about the study (e.g. species and strain, sex, no per group, route, frequency of application, sampling times, duration of exposure)	Main effects, Observations (specify regarding dose and sampling time)	Remarks (e.g. major deviations)	Reference	
Micronucleus test in rodents Compared to 92/69/EEC (B.12) Non GLP Reliability 2 Key	Pyrethrum Extract (Batch number Refined Pale Concentrate Bulk 99/Pale; Purity 28.4% Pyrethrin I, 25.0% Pyrethrin II) 0.25, 0.5, 1.0 mL/kg bw	CFLP mice (male and female) 5/sex/dose Two with 24 h interval 6 h after last dose Oral (gavage)	No increase in micronuclei	Non-genotoxic	 (KPIC) IIIA6.6.4	

Table A.47 Summary table of human data on genotoxicity

No data are available.

Conclusion used in Risk Assessment – Genotoxicity <i>in vivo</i>	
Conclusion	Not genotoxic
Justification for the conclusion	No effects.

A2.8.2.1 Short summary and overall relevance of the provided information on germ cell mutagenicity

A total of seven *in vitro* studies were performed. No indication for mutagenic potential of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ was observed. No indications for genotoxicity were detected.

In support of this conclusion, the *in vivo* study also shows no evidence of genotoxicity.

A2.8.2.2 Comparison with the CLP criteria

It does not meet the EU criteria to be classified as mutagenic. There are not positive results in tests carried out in mammals and in mutagenicity *in vitro* tests of chromosome aberrations and gene mutation in mammal cells or reverse mutation in bacteria (CLP 3.5.2.2. (Table 3.5.1) and CLP 3.5.2.3.(8.)).

A2.8.2.3 Conclusion on classification and labelling for germ cell mutagenicity

Not classified due to inconclusive data.

A2.8.2.4 Overall conclusion on genotoxicity related to risk assessment

Conclusion used in the Risk Assessment – Genotoxicity	
Conclusion	Not genotoxic
Justification for the conclusion	No effects were observed in any of the studies performed.
Proposed classification	Not classified

A.2.9. Carcinogenicity

In a 2-year dietary chronic toxicity and oncogenicity study on rats *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ corresponding to 3000 ppm total pyrethrins (199/265 mg/kg bw/d extract males/females) caused effects in both sexes: body weight gains were occasionally reduced and an increased incidence of hyperplasia and follicular cell adenomas at the thyroid was observed. In addition, males showed increased and accentuated lobulations of the liver, increased activity of serum transaminases and increased numbers of keratoacanthomas, while in females the incidence of hepatic adenoma was increased slightly.

At 1000 ppm total pyrethrins (66/86 mg/kg bw/d extract males/females) led to an accentuated lobulation of the liver and an increase in the incidence of hyperplasia and follicular cell adenomas of the thyroid, all in males only.

The NOAEL was 100 ppm actual Pyrethrins, equal to 4 mg/kg bw/d total pyrethrins (6 mg/kg bw/d extract) in males and 5 mg/kg bw/d total pyrethrins (8 mg/kg bw/d extract) in females (██████████). (KPIC) (BRA, MGK and SCJ)

		Males					Females				
		Control 1	Control 2	100 ppm	1000 ppm	3000 ppm	Control 1	Control 2	100 ppm	1000 ppm	3000 ppm
Body weight (g)	Mean	646	616	592	607	589	419	506*	478	439	408
	SD	108.5	112.9	103.4	122.7	107.5	94.1	120.6	107.2	116.3	84.8
	N	32	27	22	38	33	26	15	20	28	32
Brain (g)	Mean	2.26	2.23	2.25	2.26	589	2.06	2.01	2.07	2.05	2.05
	SD	0.159	0.132	0.145	0.138	107.5	0.079	0.121	0.133	0.141	0.110
	N	33	27	22	38	33	27	15	20	28	32
Brain/Body weight (%x10)	Mean	3.58	3.72	3.94	3.80	3.97	5.19	4.21	4.54	4.95	5.27**
	SD	0.575	0.600	0.825	0.738	0.856	1.343	1.136	1.104	1.158	1.234
	N	32	27	22	37	33	26	15	20	28	32
Liver weight (g)	Mean	24.31	22.01	24.72	24.353	24.86	16.99	19.69	18.25	17.75	18.69
	SD	4.697	4.490	5.450	4.463	5.534	3.005	4.155	4.582	4.881	3.485
	N	33	26	21	39	33	26	15	20	28	32
Liver/Body weight (%)	Mean	3.83	3.64	4.25**	4.08	4.25**	4.18	3.99	3.86	4.09	4.67**
	SD	0.751	0.709	1.065	0.854	0.718	1.056	0.746	0.668	0.768	0.817
	N	32	26	21	38	33	25	15	20	28	32
Liver/Brain weight (%x10 ⁻²)	Mean	10.79	9.93	11.03	10.90	11.03	8.25	9.76	8.87	8.65	9.12
	SD	2.115	2.039	2.591	1.961	2.481	1.405	1.880	2.371	2.363	1.760
	N	33	26	21	38	33	26	15	20	28	32

*Significantly different from Control group 1 (p < 0.05)

**Significantly different from Control group 2 (p < 0.05)

		Males										Females									
		Control 1		Control 2		100 ppm		1000 ppm		3000 ppm		Control 1		Control 2		100 ppm		1000 ppm		3000 ppm	
		DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS
N (examined)		27	33	33	27	38	22	21	39	27	33	33	27	45	15	40	20	32	28	28	32
LIVER																					
Accentuated lobulation	Mild	3.7	12.1	6.1	11.1	5.3	31.8		30.8	11.1	27.3		14.8	6.7	13.3		5.0	12.5	14.3	3.6	9.4
	Moderate	3.7		6.1		5.3		19.1		7.4		3.0		2.2		2.5		3.1		7.1	
	Severe			3.0				4.8								2.5					
Adhesions	Mild					5.3															
Congested	Mild																			3.6	
Cyst, clear/NOS	Mild						4.6			3.7											3.1
	Moderate											3.7		6.7							
Discolored, tan	Mild		21.2		7.4	5.3	4.6	4.8	2.6	3.7	9.1		3.7	6.7	13.3	5.0			3.6		3.1
	Moderate	3.7		3.0			4.6													3.6	
	Severe			6.1					2.6									3.1			
Enlarged	Mild	3.7																		3.6	
	Moderate																	3.1			
	Severe											3.0									
Foci/focus, red/dark red/black/dark	No grade													2.2							
	Trace			3.0				4.8						2.2							
	Mild	18.5	36.4	27.3	44.4	29.0	45.5	14.3	43.6	22.2	36.4	15.2	44.4	22.2	53.3	15.0	55.0	15.6	60.7	25.00	50.0
	Moderate	18.5		6.1				9.5	2.6					6.7		2.5		6.3		10.7	

Foci/focus, tan/yellow/white/NOS	No grade					2.6		4.8														
	Trace			3.0																		
	Mild	18.5	18.2	6.1	11.1	10.5	13.6	23.8	7.7	11.1	18.2	18.2	18.5	2.2	26.7	5.0	5.0	9.4	3.6	10.7	21.9	
	Moderate			3.0				4.8				6.1				5.0		3.1				
	Severe												2.2				3.1					
Foci, gray	Mild	3.7																				
Foci, clear	Mild				3.7					3.0												
Firm	Mild				3.7										2.5							
Friable/rupture	Mild								2.6		6.1											
	Moderate			6.1			4.6									5.0	3.1			3.6		
Granular	Mild			3.0			4.6															
	Moderate																	3.1				
Mass		7.4	3.0			2.6				3.7	3.0		3.7	2.2				3.1	7.1	3.6	3.1	
Mottled	Mild	7.4		3.0		10.5						3.0	4.4		7.5		3.1					
	Moderate			6.1		2.6			3.7			12.1			5.0		3.1					
	Severe	3.7				5.3							2.2		2.5							
Nodule									2.6			3.0									3.6	
Pale	Mild	11.1				2.6																
	Moderate					2.6			3.7						2.5						3.6	
	Severe					5.3																
Tan	Moderate			6.1																		
THYROID																						
Discolored, tan Enlarged	Mild						4.6															
	Trace			3.0																		
	Mild				7.4		4.6				3.0		3.7						3.6			9.4
	Moderate							4.8	2.6	3.7			3.7	2.2								

DOS - Deaths and unscheduled sacrifices

TS - Terminal sacrifice

NOS - Not otherwise specified

	Males										Females											
	Control 1		Control 2		100 ppm		1000 ppm		3000 ppm		Control 1		Control 2		100 ppm		1000 ppm		3000 ppm			
	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC		
N (examined)	27	33	33	27	38	22	21	39	27	33	33	27	45	15	40	20	32	28	28	28	32	
LIVER																						
Within normal limits	14.8	21.2	12.1	22.2	18.4	13.6	9.5	7.7	29.6	12.1	48.5	22.2	44.4	20.0	50.0	45.0	31.3	21.4	32.1	9.4		
Abscess															2.5							
Autolysis too severe for diagnosis													2.2									
Bile duct hyperplasia	Trace	18.5	36.4	21.2	18.5	26.3	4.6	9.5	48.7	7.4	33.3	12.1	14.8	4.4	6.7	7.5	15.0	15.6	25.0	17.9	28.1	
	Mild	22.2	15.2	21.2	33.3	13.2	50	23.8	20.5	3.7	27.3		14.8	8.9		7.5	5.0	12.5	28.6		21.9	
	Moderate			3.0	11.1	2.6				7.4			3.7			2.5					3.1	
Cholangioma												3.7		6.7							3.1	
Cholangiofibrosis	Trace									3.7												
	Mild	3.7									3.0											
	Moderate	3.7	3.0																			

Cyst	Mild						4.6	4.8														
Cytoplasmic alteration, basophilic	Trace		12.1		11.1	2.6			2.6	3.7	6.1		14.8		13.3	2.5	15.0	3.1	7.1		9.4	
	Mild		9.1	6.1	11.1	5.3	4.6		10.3	3.7	3.0	6.1	25.9	8.9	46.7	2.5	15.0		10.7		15.6	
	Moderate		3.0		7.4			4.8	7.7	3.7	3.0	3.0		2.2	13.3	2.5	5.0		7.1	3.6	9.4	
Cytoplasmic alteration, clear	Trace		12.1		18.5		9.1		2.6		6.1		3.7		20.0						3.1	
	Mild		6.1		3.7				10.3	3.7	21.2		3.7		6.7				10.7		6.3	
	Moderate				3.7		4.6					3.0	3.7						3.6			
Cytoplasmic alteration, eosinophilic	Mild										6.1						5.0					
	Moderate								5.1													
Hematopoiesis, extramedullary	Trace				3.7							3.0		2.2		7.5		3.1		10.7		
	Mild							4.8				3.0										
	Moderate												3.7	4.4								
Fibrosis	Trace									3.7			7.4									
	Mild					2.6								2.2		2.5						
Hepatocellular adenoma		11.1	9.1	3.0				9.5	2.6		9.1			2.2						3.6	7.1	9.4
Hepatocellular carcinoma		3.7								3.7			3.7									
Hemorrhage	Trace						4.6															
	Mild					2.6	4.6															
Hemolymphoreticular neoplasm		7.4		3.0		7.9		4.8	2.6	11.1		3.0		2.2		5.0		6.3		7.1		
Inflammation	Trace	3.7	9.1	3.0		2.6	4.6	4.8	2.6	7.4				6.7	2.5			3.1			6.3	
	Mild	3.7	6.1	6.1	3.7	5.3					3.0							3.1		3.6		
	Moderate									3.7												
Metastatic tumor																		3.1		3.6		
Spongiosis hepatis	Trace	3.7	21.2	9.1	14.8	5.3	27.3	9.5	12.8	11.1	30.3			2.2					7.1			
	Mild		6.1		11.1				7.7		18.1										3.6	
	Moderate		3.0						2.6	7.4												
Necrosis	Trace	3.7	3.0	6.1					7.7			3.0		2.2					3.6	3.6	3.1	
	Mild			3.0		13.2		14.3		3.7		3.0		4.4	6.7	10.0		3.1		3.6		
	Moderate	3.7		3.0										2.2						10.7		
	Severe	3.7		3.0		7.9										2.5		3.1		3.6		
Pigment, brown	Trace			3.0		2.6								6.7								
	Moderate												3.7									
Telangiectasis	Trace	11.1																			3.1	
	Mild	3.7	6.1						2.6			3.0		4.4						3.6	6.3	
	Moderate					2.6			2.6			3.0				5.0		3.1	3.6			
Vacuolar change	Trace				3.7		4.6															
Vacuolar change, fatty	Trace		3.0	3.0		10.5		14.3	2.6			3.0	7.4	8.9		2.5		9.4		3.6		
	Mild	22.2		15.2		15.8	4.6	9.5	2.6	11.1	12.1	3.7	8.9		2.5		9.4		3.6			
	Moderate			9.1	3.7	2.6		4.8		7.4			2.2		5.0							
	Severe							4.8					7.4	2.2				3.1				
Cytoplasmic alteration, vacuolated	Mild											3.7				10.0						
	Moderate											3.0	3.7								3.1	
	Severe														2.5							

THYROID																					
Within normal limits		44.4	51.5	48.5	33.3	68.4	68.2	76.2	53.9	55.6	54.6	72.7	44.4	60.0	53.3	77.5	40.0	56.3	89.3	60.7	46.9
Cyst, colloid	Trace		3.0																		
	Mild			6.1							3.0			2.2							
	Moderate																				3.1
Cyst	Mild																				3.1
Corpora amylacea				3.0																	
Follicular adenoma		3.7	3.0			7.9		9.5	7.7	11.1	6.1						10.0	9.4		3.6	12.5
Follicular carcinoma					3.7	2.6			5.1		6.1		3.7	4.4							3.1
Hyperplasia	Trace						4.6		5.1								5.0				
	Mild	3.7	3.0						7.7	11.1	12.1			2.2				3.1		3.6	12.5
	Moderate					2.6								2.2							
Inflammation	Trace			3.0																	3.6
Mineralization	Trace	3.7	9.1	3.0	3.7		4.6				9.1										3.1
	Mild										3.0					2.5					
Parafollicular cell adenoma		11.1	9.1	9.1	11.1	10.5	4.6	4.8	5.1		12.1		11.1	4.4		2.5	20.0	9.4	3.6	3.6	15.7
Parafollicular cell carcinoma		3.7	3.0		7.4				5.1				7.4			2.5					
Parafollicular cell hyperplasia	Trace	7.4		3.0					2.6			3.0	3.7	4.4	6.7	5.0	15.0	9.4		7.1	3.1
	Mild	11.1	6.1		14.8		4.6		2.6	3.7	3.0	6.1	7.4	2.2	6.7	2.5	5.0		3.6		9.4
	Moderate		3.0																		
Ultimobranchial cyst		22.2	27.3	33.3	33.3	10.5	13.6	9.5	12.8	22.2	12.1	24.2	29.6	28.9	33.3	7.5	10.0	18.8	7.1	21.4	6.3

DOS – Deaths and unscheduled sacrifices

SAC – Terminal sacrifice

In an 18-month dietary oncogenicity study on mice, from 2500 ppm total pyrethrins (530/633 mg/kg bw/d extract) onwards discoloured dark livers and increased absolute and relative liver weights were observed in males and females, and vacuolar fatty changes in the livers of males only. Lung carcinomas in males were not treatment related. For females, there was no evidence for carcinogenic response.

The NOEL/NOAEL was 100 ppm actual Pyrethrins, equal to 14/17 mg/kg bw/d total pyrethrins (22/26 mg/kg bw/d extract) for males/females [REDACTED]. (KPIC and BRA, MGK and SCJ)

		Males					Females				
		Control 1	Control 2	100 ppm	2500 ppm	5000 ppm	Control 1	Control 2	100 ppm	2500 ppm	5000 ppm
Body weight (g)	Mean	40	40	40	40	40	35	35	35	36	34
	SD	3.5	3.3	3.5	3.0	3.2	3.6	3.4	3.7	3.8	3.0
	N	42	48	44	40	44	45	41	49	44	42
Brain (g)	Mean	0.53	0.53	0.53	0.54	0.54	0.54	0.55	0.55	0.55	0.54
	SD	0.031	0.029	0.029	0.035	0.030	0.027	0.038	0.038	0.031	0.026
	N	42	48	44	40	44	45	41	49	44	42
Brain/Body weight (%x10)	Mean	13.3	13.4	13.2	13.7	13.6	15.9	15.9	15.8	15.6	15.8
	SD	1.20	1.27	1.13	1.02	1.13	1.79	1.93	1.74	1.70	1.41
	N	42	48	44	40	44	45	41	49	44	42
Liver weight (g)	Mean	2.29	2.33	2.38	2.89***	3.10***	2.17	2.18	1.99***	2.68***	2.90***
	SD	0.352	0.333	0.398	0.347	0.412	1.016	0.340	0.409	0.792	0.464
	N	39	44	44	40	38	45	41	49	44	42
Liver/Body weight (%)	Mean	5.77	5.81	5.92	7.30***	7.91***	6.20	6.29	5.68**	7.48***	8.46***
	SD	0.775	0.717	1.010	0.712	0.751	2.197	0.710	1.042	1.662	0.987
	N	39	44	44	40	38	45	41	49	44	42
Liver/Brain weight (%x10 ⁻²)	Mean	4.33	4.38	4.50	5.37***	5.79***	4.01	4.00	3.65****	4.88***	5.40***
	SD	0.683	0.678	0.753	0.654	0.673	1.862	0.678	0.761	1.482	0.849
	N	39	44	44	40	38	45	41	49	44	42

*Significantly different from Control group 1 (p < 0.01)

**Significantly different from Control group 2 (p < 0.01)

***Significantly different from Control group 1 (p < 0.05)

****Significantly different from Control group 2 (p < 0.05)

		Males										Females									
		Control 1		Control 2		100 ppm		2500 ppm		5000 ppm		Control 1		Control 2		100 ppm		2500 ppm		5000 ppm	
		DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS	DOS	TS
N (examined)		18	42	12	48	16	44	20	40	16	44	15	45	19	41	11	49	16	44	18	42
LIVER																					
Accentuated lobulation	Mild					6.3	2.3	5.0	7.5		9.1									5.6	2.4
	Adhesions	Mild	2.4																		
	Moderate												5.3								
Cyst, clear	Mild																			4.6	
Autolysis	Severe	5.6																			
	Mild				2.1				2.5	6.3	43.2								15.9	5.6	31.0

Discolored, dark/brown	Moderate																				5.6
Discolored, tan	Moderate	5.6					6.3														5.6
	Severe						6.3					6.7									
Enlarged	Mild						6.3							10.5						2.3	
	Moderate			8.3						6.3			2.2							2.3	
Foci/focus, red/black/dark	Mild	5.6					4.6						6.7							2.3	2.4
	Moderate							5.0													
Foci/focus, tan/white	Trace																	6.3			
	Mild	5.6	2.4				18.8	2.3	5.0			12.5	6.8		4.4		9.1			5.6	2.4
	Moderate													5.3							
	Severe																	6.3			
Friable/rupture	Moderate							5.0													5.6
Granular	Mild												2.2		2.4					2.3	
	Moderate																			2.3	
Mass		5.6	14.3	8.3	20.8	6.3	6.8	20.0	20.0	6.3	20.5			5.3	2.4	9.1		6.3			2.4
Mottled	Moderate			8.3																	
Nodule		5.6	2.4	8.3	6.3		2.3			2.5		4.6		2.2		9.1	2.0				
Pale	Severe															9.1					
THYROID																					
Enlarged	Mild														2.4				6.3		
	Moderate																				2.4

DOS - Deaths and unscheduled sacrifices

TS - Terminal sacrifice

	Males										Females												
	Control 1		Control 2		100 ppm		2500 ppm		5000 ppm		Control 1		Control 2		100 ppm		2500 ppm		5000 ppm				
	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC	DOS	SAC			
N (examined)	18	42	12	48	16*	44*	20*	40*	16	44	15	45	19	41	11*	49*	16**	44*	18	42			
LIVER																							
Within normal limits	50.0	50.0	75.0	45.8	31.3	40.9	40.0	22.5	37.5	34.1	53.3	35.6	47.4	24.4	36.4	40.8	37.5	36.4	27.8	47.6			
Abscess	Severe																						
Amyloidosis	Trace																						
	5.6		2.4		8.3		4.6		2.5		2.3		8.9		10.5		9.8		6.1				
	5.6		2.4		4.2		25.0		5.0		7.5		25.0		2.3		20.0		10.5		4.9		9.1
Autolysis too severe for diagnosis	Moderate																						
					6.3												9.1						
Cystic dilatation	Mild																						
	Moderate																						
	Severe																						
Cyst	Mild																						
					2.1																2.3		
	Moderate																						
Cytoplasmic alteration, basophilic	Trace																						
					2.1										2.4								
	Mild																						
								5.0		2.3		2.2		5.3									
Moderate																							
								2.5				6.7											

Cytoplasmic alteration, clear	Trace				2.1																
Cytoplasmic alteration, eosinophilic	Moderate				2.1				2.5							2.0			2.3		
Hematopoiesis, extramedullary	Trace											6.7			2.4	9.1				5.6	4.8
	Mild	11.1		8.3								6.7									
	Moderate						6.3					6.7		5.3		9.1					
Fibrosis	Mild													5.3							
	Moderate									6.3											
Hepatocellular adenoma		5.6	9.5	8.3	22.9	6.3	6.8	10.0	22.5		22.7		2.2							5.6	
Hemangiosarcoma					4.2		4.6	5.0	2.5		2.3						6.3				2.4
Hepatocellular carcinoma		5.6	7.1	8.3	4.2	6.3					6.3	4.6									2.4
Hemorrhage	Mild							5.0				6.7									
Hemolymphoreticular neoplasm		5.6											4.4	10.5		9.1	2.0	12.5	6.8		4.8
Inflammation	Trace		14.3		27.1		31.8	5.0	22.5		18.2	6.7	33.3	10.5	46.3		32.7		36.4		26.2
	Mild	5.6	27.8		2.1	6.3	11.4		5.0		2.3		13.3	10.5	14.6		14.3	6.3	4.6		2.4
	Moderate		2.4										6.7	5.3							
	Severe										6.3										
Infarct	Severe													5.3							
Lymphocytic infiltration	Trace		2.4				9.1	5.0	7.5	6.3	2.3				9.8	9.1	2.0	6.3	9.1	5.6	7.1
	Mild						2.3						2.2			4.1					
Plasma cell infiltration	Mild											2.2									
Mineralization	Trace				2.1																5.6
Metastatic tumor		5.6						5.0													
Necrosis	Trace	5.6	2.4		4.2					6.3						12.3			2.3		
	Mild		2.4			6.3	4.6						5.3	2.4			6.3	2.3	5.6		
	Moderate				2.1																
Pigment, brown	Trace		2.4		2.1		4.6		5.0		4.6	13.3	6.7	15.8	17.1		10.2		11.4		11.9
	Mild		4.8								2.3	6.7	6.7				6.3		2.3		9.5
	Moderate									6.3	2.3	2.2								5.6	2.4
Vacuolar change, fatty	Trace		2.4			6.3			10.0	6.3	13.6										
	Mild				2.1	6.3				7.5	13.6										
	Moderate									2.5	2.3										
Cytoplasmic alteration, vacuolated	Mild				2.1																
THYROID																					
Within normal limits		77.8	76.2	66.7	70.8	-	-	-	-	50.0	86.4	66.7	73.3	79.0	78.1	-	-		-	50.0	78.6
Cyst, colloid	Trace	11.1	7.1		12.5	-	-	-	-		4.6		6.7			-	-		-		2.4
	Mild		2.4		4.2	-	-	-	-		4.6	6.7	2.2			-	-		-		2.4
	Moderate					-	-	-	-				2.2			-	-		-		

Amyloidosis	Trace		2.4	8.3	4.2	-	-	-	-	12.5		6.7	2.2	5.3	7.3	-	-		-		2.4
	Mild	11.1	9.5	8.3	6.3	-	-	-	-			6.7	11.1	10.5	2.4	-	-		-	5.6	2.4
	Moderate			8.3	4.2	-	-	-	-	12.5	4.6	6.7	2.2		9.8	-	-	6.3	-	27.8	
	Severe		2.4	8.3		-	-	-	-	18.8		6.7		5.3		-	-		-	11.1	2.4
Hemolymphoreticular neoplasm						-	-	-	-					5.3		-	-		-		
Hyperplasia	Mild					-	-	-	-				2.2		2.4	-	-		-		2.4
Inflammation	Trace					-	-	-	-							-	-		-		4.8
	Mild					-	-	-	-							-	-		-		2.4
	Moderate					-	-	-	-	6.3						-	-		-		
Mineralization	Trace					-	-	-	-							-	-		-		5.6

DOS – Deaths and unscheduled sacrifices

SAC – Terminal sacrifice

*For thyroid, no animals were examined.

**For thyroid, only one animal was examined.

Table A.48 Summary table of carcinogenicity studies in animals

Summary table of carcinogenicity studies in animals						
Method, Duration of study, Route of exposure, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels, Duration of exposure	NOAEL, LOAEL	Results (Please indicate any results that might suggest carcinogenic effects, as well as other toxic effects, for all dose levels)	Remarks (e.g. major deviations)	Reference

<p>Oral 2-year Diet EPA 83-5, OECD 453 GLP Reliability 1 Key</p>	<p>Rat Charles River CD male+female 60/sex/dose</p>	<p>Pyrethrum extract (Blend FEK-99; Purity 57.574%) 0 ppm 100 ppm 1000 ppm 3000 ppm in diet daily <i>ad libitum</i>; equivalent to 0, 4, 43, and 130 mg/kg bw/d total pyrethrins (0, 6, 66, and 199 mg/kg bw/d extract) for males and 0, 5, 56 and 173 mg/kg bw/d total pyrethrins (0, 8, 86, and 265 mg/kg bw/d extract) for females</p>	<p>NOAEL: 100 ppm Males: 4 mg/kg bw/d total pyrethrins (6 mg/kg bw/d extract) Females: 5 mg/kg bw/d total pyrethrins (8 mg/kg bw/d extract) LOAEL: 1000 ppm Males: 43 mg/kg bw/d total pyrethrins (66 mg/kg bw/d extract) Females: 56 mg/kg bw/d total pyrethrins (86 mg/kg bw/d extract)</p>	<p>3000 ppm (199/265 mg/kg bw/d extract): hepatotoxicity, thyroidal tumours, liver adenoma (females) body weight ↓ 1000 ppm (66/86 mg/kg bw/d extract): hepatotoxicity (males)</p>	<p>-</p>	<p>██████████ (KPIC) IIIA6.7/01 ██████████ (BRA, MGK and SCJ) IIIA6.7/1</p>
--	---	---	---	--	----------	---

<p>Oral 18-months diet EPA 83-2, OECD 451 GLP Reliability 1 Key</p>	<p>Mouse Charles River CD male+female 60/sex/dose</p>	<p>Pyrethrum extract (Blend FEK-99; Purity 57.574%) 0 ppm 100 ppm 2500 ppm 5000 ppm in diet daily <i>ad libitum</i>; equivalent to 0, 14, 346, and 686 mg/kg bw/d total pyrethrins (0, 22, 530, and 1051 mg/kg bw/d extract) for males and 0, 17, 413, and 834 mg/kg bw/d total pyrethrins (0, 26, 633, and 1278 mg/kg bw/d extract) for females</p>	<p>NOAEL: 100 ppm males: 14 mg/kg bw/d total pyrethrins (22 mg/kg bw/d extract) Females: 17 mg/kg bw/d total pyrethrins (26 mg/kg bw/d extract) LOAEL: 2500 ppm Males: 346 mg/kg bw/d total pyrethrins (530 mg/kg bw/d extract) Females: 413 mg/kg bw/d total pyrethrins (633 mg/kg bw/d extract)</p>	<p>5000 ppm (1051/1278 mg/kg bw/d extract): hepatotoxicity 2500 ppm (530/633 mg/kg bw/d extract): hepatotoxicity</p>	<p>-</p>	<p>[REDACTED] (KPIC) IIIA6.7/02 [REDACTED] (BRA, MGK and SCJ) IIIA6.7/2</p>
---	---	--	---	--	----------	---

Table A.49 Summary table of human carcinogenicity data
No data are available.

Table A.50 Summary table of other relevant studies for carcinogenicity
No data are available.

A2.9.1 Short summary and overall relevance of the provided information on carcinogenicity

Rat

Groups of 60 Charles River CD rats of each sex received diets containing 0, 100, 1000, or 3000 ppm actual Pyrethrins for 104 weeks, equal to 0, 4, 43, and 130 mg/kg bw/d total pyrethrins (0, 6, 66, and 199 mg/kg bw/d extract) for males and 0, 5, 56, and 170 g/kg bw/d total pyrethrins (0, 8, 86, and 265 mg/kg bw/d extract) for females.

No mortalities and no clinical signs attributable to the administration of the test substance were found. Body weights were significantly depressed in both sexes in the 3000 ppm (199/265 mg/kg bw/d extract) treatment group during the first 78 weeks only, combined with a slight decrease in food consumption.

No test substance related organ weight changes and no haematological or urological changes were found. Statistically significant increases in the activity of serum transaminases were determined at most intervals of analysis at males of the 3000 ppm (199 mg/kg bw/d extract) dosage group.


Keratoacanthomas of the skin were increased at the highest dosage level in the male rats, however, due to the self-limiting nature of this lesion, it is unlikely that this finding has any true toxicological significance. Increased incidences of benign tumours of the thyroid were also observed and a statistically significantly higher incidence of hepatocellular adenomas was described in females at the high dose. Follicular adenomas and carcinomas were initially seen in the thyroid glands of rats at the high dose, which appeared to be related to treatment, but during a re-evaluation some of the carcinomas were reclassified as adenomas and some adenomas were reclassified as hyperplasia. After the re-evaluation, the incidence of hyperplasia was found to be enhanced in males and females, and the incidence of follicular adenomas was statistically significantly increased only in females at 3000 ppm (265 mg/kg bw/d extract). The NOEL/NOAEL was 100 ppm, equal to 4 mg/kg bw/d total pyrethrins (6 mg/kg bw/d extract) in males and 5 mg/kg bw/d total pyrethrins (8 mg/kg bw/d extract) in females.

To advance the understanding of the mechanisms by which Pyrethrins cause rodent tumours and to better understand the relevance of the observations in rats to humans, a mechanistic toxicity study in rats was carried out (■■■■■ (IIIA6.10/1); ■■■■■ (IIIA6.10/2)). The aim of both studies was to determine if the *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is a genotoxic carcinogen or the increase in the incidence of tumours follows a mechanism similar to that for phenobarbital as a consequence of microsomal enzyme induction. The first is focused on carcinogenic effects and the second on enzyme induction in the liver. The test substance (FEK-99) was the same as in the carcinogenicity study in rats (IIIA6.7/1) in the first study. The second study was carried out using liver samples from the previous study taken at different times. The following conclusions can be drawn:

- There is a consensus that pyrethrins are not genotoxic.
- Only benign tumours (adenoma) were observed – no carcinoma – in the 2-year oral toxicity bioassay in rats.
- There was no clear dose response, e.g. as a singularly phenomenon only in females maintained for two years on the (maximum tolerated) high dose (3000 ppm) a minimal increase in adenoma was noted.

- In male rats the incidences of adenoma were 5% (3 of 60 males) in both, the highest dose and the mid dose (3000 ppm and 1000 ppm (199 and 66 mg/kg bw/d extract)).
- Subchronic studies indicated that Pyrethrins induced signs of hepatotoxicity in male and female rats, this was confined to high dose levels (≥ 3000 ppm (199/265 mg/kg bw/d extract)).

Figure 2: Graphical presentation of the data obtained in the mechanistic study (14 days of treatment). Data are normalised for values of control animals. A clear parallelism is obvious for high dose Pyrethrins (hatched bars) and Phenobarbital (black bars) while low dose Pyrethrins (grey bars) parallel the untreated controls (white bars).



The mechanistic study was conducted with different concentrations of pyrethrins investigating several time points (7 d, 14 d, 42 d and 42 d treatment + 42 days recovery) including a negative control group and a phenobarbital-treated positive control group.

The parallelism of effects observed in the positive control (~1500 ppm phenobarbital) and the high dose Pyrethrins-treated groups is evident (Figure 2) and the conclusion may be drawn that both compounds induce tumours in the liver by a similar non-genotoxic mechanism at high dose levels. Notably, phenobarbital appears to be far more potent as compared to Pyrethrins, and indeed the effect of pyrethrins on the incidence of hepatic adenoma was only minimal.

Phenobarbital has been shown to induce liver enzymes and also hepatic cell proliferation via an activation of the constitutive androstane receptor (CAR), a nuclear receptor that forms heterodimers with the retinoid acid receptor to induce a number of proteins but not hepatic cytotoxicity. Similar mechanisms have been reported for synthetic analogs of Pyrethrins. Epidemiological studies showed no human cancer risk for Phenobarbital a widely used barbiturate and antiepileptic drug that is taken at considerable (pharmacologic) dose levels, therefore it is highly unlikely that exposure to Pyrethrins at levels below the ADI (0.04 mg/kg bw/d) will have an influence of human cancer risk.

Increased liver weights were observed at high dose levels in both male and female rats in the 2-year chronic bioassay and hepatotoxicity was also observed both in male and female rats in a subchronic study (90 days oral administration [redacted]). Increased liver weights were noted at the high dose level for male rats and for female rats both at 1000 ppm and 3000 ppm total pyrethrins (66/86 and 199/265 mg/kg bw/d extract) [redacted].

This is in concordance with the view, that hepatotoxicity was the mechanism in a 2-year feeding study with rats, where a marginal increase of benign tumours was noted in the liver of females.

It is a general consensus in the scientific community that benign liver tumours induced in

the rat that follow a non-genotoxic mechanism are highly specific to rodents and of no relevance for humans.

Thus, in the present case Pyrethrins with only a marginally tumorigenic activity in rodents act apparently by a mechanism similar to Phenobarbital, but the potency is about 5-10-fold lower with regard to biochemical effects. Phenobarbital has been shown to be non-carcinogenic to humans even when administered at pharmacologic levels.

The results show that Pyrethrins, in common with other non-genotoxic oncogens (phenobarbital) caused liver and thyroid gland tumours through a dose related proliferative response in the liver (replicative DNA synthesis and microsomal enzyme induction) and a secondary proliferative stimulation of thyroid follicular cells which is specific to rats. The lack of any effect of Pyrethrins at 100 ppm (6/8 mg/kg bw/d extract) confirms the threshold nature of this effect (██████████; Pfeil, 2003; ██████████). (BRA, MGK and SCJ)

Mouse

Groups of 60 Charles River CD-1 mice of each sex received diets containing 0, 100, 2500, and 5000 ppm total pyrethrins for 18 months, equal to doses of 0, 14, 350, and 690 mg/kg bw/d total pyrethrins (0, 22, 530, and 1051 mg/kg bw/d extract) for males and 0, 17, 400, and 830 mg/kg bw/d total pyrethrins (0, 26, 633, and 1278 mg/kg bw/d extract) for females.

One male and one female in the 5000 ppm (1051/1278 mg/kg bw/d extract) group were found dead during the first week of the study but no further treatment related deaths occurred. All 5000 ppm (1051/1278 mg/kg bw/d extract) animals exhibited increased activity when stimulated during the first week of study only. No dose related differences in body weight gain and food consumption were observed. There were no test substance related changes in the differential blood counts.

The only possibly test substance related changes at necropsy were:

- discoloured dark livers more common in males at 5000 ppm (1051 mg/kg bw/d extract) and in females at 2500 and 5000 ppm (633 and 1278 mg/kg bw/d extract)
- increased absolute and relative liver weights in both sexes at 2500 and 5000 ppm (530/633 and 1051/1278 mg/kg bw/d extract)
- vacuolar fatty changes in the livers of males at 2500 and 5000 ppm (530 and 1051 mg/kg bw/d extract))

The lung carcinomas in males were not treatment-related since the incidences at 2500 and 5000 ppm (530/633 and 1051/1278 mg/kg bw/ extract) were similar to those in the control groups. For females there was no evidence for carcinogenic response. The NOEL/NOAEL was 100 ppm total pyrethrins, equal to 14 mg/kg bw/d total pyrethrins (22 mg/kg bw/d extract) for males and 17 mg/kg bw/d total pyrethrins (26 mg/kg bw/d extract) for females ██████████. (KPIC and BRA, MGK and SCJ)

Table A.51 Compilation of some factors that may be taken into consideration in classification and labelling

No data are available.

A.2.9.2 Comparison with the CLP criteria

It does not meet the EU criteria to be classified as carcinogenic. There is a clear causal link between the *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ and the incidence of benign neoplasm in liver and thyroid established in rat and mouse. However, the mechanism has been characterized and can be concluded that there is no risk to human health (CLP 3.6.2.2.3.(b)) explained in the point A3.9.1 in this CLH report.

A.2.9.3 Conclusion on classification and labelling for carcinogenicity

Not classified.

A.2.9.4 Overall conclusion on carcinogenicity related to risk assessment

Conclusion used in Risk Assessment – Carcinogenicity	
Value/conclusion	Not carcinogenic.
Justification for the value/conclusion	From the testing detailed above there are no indications of carcinogenic potential.
Proposed classification	Not classified.

A.2.10. Reproductive toxicity

A.2.10.1. Sexual function and fertility

Table A.52 Summary table of animal studies on adverse effects on sexual function and fertility

Summary table of animal studies on adverse effects on sexual function and fertility								
Method, Duration of study, Route of exposure, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels Duration of exposure	NOAELs, LOAELs (e.g. maternal/parental toxicity, effects on sexual function and fertility)			Results (for all dose levels, specify critical effects on sexual function and fertility for parental animals (and offspring if relevant), report e.g. incidences and severity of the effects for all dose levels)	Remarks (e.g. major deviations)	Reference
Two generation Reproductive toxicity EPA Guideline 83-4 GLP Reliability 1 Key	Rat Charles River COBS® CD® Male and female 28 animals/sex/dose	Pyrethrum extract (Blend FEK-99; Purity 57.574%) 0, 12, 120 and 360 mg/kg bw/d total pyrethrins (0, 18, 184, and 552 mg/kg bw/d extract)	NOAEL Parental 12 mg/kg bw/d total pyrethrins (18 mg/kg bw/d extract) for toxicity, 360 mg/kg bw/d total pyrethrins (552 mg/kg bw/d extract) for reproductive parameters	NOAEL F1 12 mg/kg bw/d total pyrethrins (18 mg/kg bw/d extract) for toxicity, 360 mg/kg bw/d total pyrethrins (552 mg/kg bw/d extract) for reproductive parameters	NOAEL F2 12 mg/kg bw/d total pyrethrins (18 mg/kg bw/d extract) for toxicity, 360 mg/kg bw/d total pyrethrins (552 mg/kg bw/d extract) for reproductive parameters	360 and 120 mg/kg bw/d total pyrethrins (552 and 184 mg/kg bw/d extract): Parental: body weight↓; F1, F2: neonatal bodyweight↓ No effects on reproductive parameters	-	(KPIC) (BRA, MGK and SCJ) IIIA6.8.2

Table A.53 Summary table of human data on adverse effects on sexual function and fertility

No data are available.

Table A.54 Summary table of other relevant studies for sexual function and fertility

No data are available.

A2.10.1.1 Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility

Chrysanthemum cinerariaefolium extract from supercritical CO₂ was administered to 4 groups of 28 Charles River CD[®] rats sex/group at concentrations of 0, 12, 120 and 360 mg/kg bw/d total pyrethrins (0, 18, 184, and 552 mg/kg bw/d extract) over a two-generation period. Each group of 28 males and females formed the initial F0 parental generation. These animals were treated for a minimum of 77 days prior to the first of two matings. Weanlings from the F1b litters were randomly selected (28 rats/sex/group) to become parents of the F1 generation. These animals were treated for a minimum of 95 days prior to being mated twice to produce the F2a and F2b litters.

No treatment related effects were noted with respect to clinical signs, body weights or food consumption for the parental rats in the F0 generation. Body weights and food consumption for the parental rats in the F1 generation were significantly reduced at 360 mg/kg bw/d total pyrethrins (552 mg/kg bw/d extract) males/females, and slightly reduced at 120 mg/kg bw/d total pyrethrins (184 mg/kg bw/d extract) in males when compared with controls. These reductions were considered to be treatment related. Administration of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ at 360 mg/kg bw/d total pyrethrins (552 mg/kg bw/d extract) resulted in significantly reduced pup body weights for the male and female offspring for both matings of both generations. The mean body weight values for the pups at 120 mg/kg bw/d total pyrethrins (184 mg/kg bw/d extract) were also lower than controls during the F1b and F2a lactation periods. Reproductive performance and the other litter parameters assessed were not affected by ingestion of test diets at any level tested. Macroscopic and microscopic findings were considered to be spontaneous and/or incidental in nature and not related to administration of the test article.

The NOEL/NOAEL with respect to reproductive parameters is 360 mg/kg bw/d total pyrethrins (552 mg/kg bw/d extract) and with respect to parental and neonatal toxicity is 12 mg/kg bw/d total pyrethrins (18 mg/kg bw/d extract) (██████████). (KPIC and BRA, MGK and SCJ)

Weeks of Age	0 mg/kg bw/d extract (Control)		18 mg/kg bw/d extract		184 mg/kg bw/d extract		552 mg/kg bw/d extract	
	Male	Female	Male	Female	Male	Female	Male	Female
0	198 ± 12.3	145 ± 8.9	197 ± 12.7	146 ± 9.1	197 ± 12.8	142 ± 8.5	194 ± 12.0	148 ± 9.0
1	250 ± 13.3	170 ± 11.7	251 ± 16.0	171 ± 13.3	251 ± 14.9	167 ± 8.7	245 ± 14.7	171 ± 10.5
2	290 ± 15.9	186 ± 12.2	296 ± 18.3	189 ± 14.0	292 ± 18.3	180 ± 12.2	284 ± 16.5	187 ± 13.2
3	329 ± 19.1	207 ± 15.9	337 ± 21.3	209 ± 16.3	334 ± 21.5	202 ± 16.0	323 ± 20.5	205 ± 14.9
4	352 ± 23.0	217 ± 17.9	366 ± 24.4	220 ± 18.6	364 ± 26.0	215 ± 18.0	352 ± 23.1	215 ± 16.8
5	378 ± 26.7	228 ± 17.8	390 ± 33.5	234 ± 21.8	389 ± 29.3	225 ± 18.5	373 ± 27.5	227 ± 17.2
6	399 ± 29.3	235 ± 19.7	409 ± 33.3	239 ± 21.2	401 ± 36.2	233 ± 19.7	393 ± 28.4	237 ± 18.2
7	419 ± 31.4	246 ± 19.7	428 ± 5.6	247 ± 23.5	423 ± 36.0	240 ± 22.4	415 ± 31.3	245 ± 19.6
8	438 ± 34.4	252 ± 21.4	445 ± 37.4	255 ± 23.5	440 ± 38.2	246 ± 23.4	427 ± 31.7	251 ± 20.3
9	451 ± 35.8	259 ± 22.1	460 ± 40.3	261 ± 24.4	457 ± 39.7	250 ± 26.8	442 ± 33.5	257 ± 21.9
10	457 ± 38.5	260 ± 23.0	469 ± 41.9	263 ± 25.0	468 ± 40.4	254 ± 27.3	451 ± 34.4	260 ± 22.3
11	470 ± 38.6	269 ± 23.2	487 ± 45.1	271 ± 27.9	482 ± 43.8	264 ± 28.7	465 ± 36.4	265 ± 24.2
12a	472 ± 40.4	275 ± 35.9 (9)	487 ± 46.0	265 ± 36.1 (11)	483 ± 45.1	265 ± 29.9 (13)	465 ± 37.3	275 ± 24.4 (13)
13a	482 ± 42.5	278 (1)	495 ± 48.6	300 ± 74.2 (2)	490 ± 44.6	267 ± 13.3 (5)	474 ± 36.6	321 ± 10.7 (3)
14a	497 ± 44.3	-	508 ± 51.3	375 (1)	500 ± 46.4	310 (1)	485 ± 39.0	-
15a	502 ± 45.9	-	514 ± 51.4	360 (1)	505 ± 46.1	-	490 ± 40.5	-
16a	514 ± 45.3	285 ± 40.2 (4)	528 ± 53.3	361 (1)	518 ± 45.8	278 ± 33.1 (5)	505 ± 44.6	253 (1)
17a	518 ± 47.2	-	532 ± 54.5	-	529 ± 45.0	-	509 ± 45.2	-
18a	528 ± 46.9	294 ± 22.8 (15)	545 ± 61.2	298 ± 27.1 (15)	537 ± 48.8	297 ± 20.4 (14)	518 ± 43.5	287 ± 23.3 (14)
19a	534 ± 0.9	285 ± 26.2 (27)	551 ± 62.6	282 ± 28.9 (27)	538 ± 49.4	276 ± 30.1 (26)	523 ± 49.0	273 ± 25.1 (25)
20	541 ± 52.3	297 ± 26.5 (28)	559 ± 64.7	296 ± 8.5 (27)	549 ± 50.3	288 ± 29.7 (28)	533 ± 49.2	279 ± 24.5 (27)
21	548 ± 53.8	302 ± 25.5 (28)	568 ± 66.9	305 ± 32.6 (27)	557 ± 52.9	294 ± 30.4 (28)	537 ± 50.0	285 ± 24.0 (27)
22	557 ± 57.0	309 ± 27.1 (28)	580 ± 68.5	313 ± 36.1 (27)	568 ± 54.9	301 ± 33.0 (10)	546 ± 51.5	288 ± 29.0 (27)
23b	551 ± 56.3	323 ± 30.2 (13)	570 ± 67.0	329 ± 48.4 (10)	560 ± 55.1	310 ± 46.1 (10)	537 ± 53.0	294 ± 29.0 (15)
24b	557 ± 59.6	358 ± 45.8 (4)	577 ± 69.2	337 ± 66.1 (5)	567 ± 57.6	349 ± 27.8 (5)	542 ± 54.0	298 ± 8.9 (3)
25b	563 ± 57.9	352 ± 15.9 (3)	585 ± 72.9	364 ± 66.0 (4)	571 ± 59.3	392 ± 21.2 (2)	549 ± 54.9	303 ± - (1)
26b	562 ± 57.6	348 ± 6.4 (2)	585 ± 71.7	403 ± 52.0 (3)	569 ± 60.4	376 ± 16.3 (2)	544 ± 52.7	-
27b	574 ± 59.6	339 ± - (1)	597 ± 75.1	427 ± 4.2 (2)	581 ± 60.8	369 ± 14.8 (2)	558 ± 58.4	-
28	590 ± 61.6	350 ± - (1)	613 ± 77.6	446 ± - (1)	596 ± 63.6	378 ± 15.6 (2)	576 ± 60.3	-
29	592 ± 63.5	354 ± - (1)	615 ± 78.1	421 ± - (1)	601 ± 66	381 ± 14.8 (2)	580 ± 61.9	-

Note: Number of animals weighed was 28 unless otherwise stated in brackets of results column.

aF_{1a}= Gestation/lactation period

bF_{1b}= Gestation/lactation period

Weeks of study	0 mg/kg bw/d extract		18 mg/kg bw/d extract		184 mg/kg bw/d extract		552 mg/kg bw/d extract	
	Male	Female	Male	Female	Male	Female	Male	Female
5	152 ± 13	127 ± 10.3	154 ± 13.6	133 ± 7.7 *	145 ± 11.2	123 ± 13.6	135 ± 16.4**	121 ± 17.5
6	211 ± 16.8	157 ± 12.8	213 ± 16.2	163 ± 8.4 *	202 ± 14.9	153 ± 15.9	188 ± 22 **	154 ± 17.5
7	264 ± 20.4	177 ± 15.5	269 ± 20.3	186 ± 9.3 *	251 ± 20.0	173 ± 18.3	241 ± 23.5**	174 ± 18.0
8	319 ± 22	201 ± 18.1	324 ± 25.9	210 ± 12.4*	304 ± 21.3	191 ± 20.6	290 ± 25**	195 ± 21.7
9	364 ± 26.4	220 ± 19.0	368 ± 26.7 (27)	230 ± 14.3	349 ± 26.2	210 ± 25.8	330 ± 26.3**	210 ± 21.4
10	399 ± 28.9	232 ± 19.8	408 ± 29.9 (27)	244 ± 17.5	382 ± 30.0	224 ± 27.5	363 ± 30.4 **	221 ± 21.9
11	428 ± 31.1	243 ± 20.7	435 ± 34.3 (27)	258 ± 18.1 *	409 ± 32.1	233 ± 26.4	392 ± 30.7 **	231 ± 23.4
12	455 ± 33.1	254 ± 21.6	461 ± 38.5 (27)	268 ± 20.4	435 ± 36.3	243 ± 27.1	416 ± 30.9 **	240 ± 23.3
13	475 ± 37.7	262 ± 23.4	484 ± 42.6 (27)	278 ± 21.4 *	451 ± 39.1	247 ± 28.3	440 ± 37.3 **	247 ± 25.2
14	498 ± 38.3	271 ± 25.0	510 ± 44.4 (27)	285 ± 23.7	469 ± 41.9*	256 ± 28.2	458 ± 33.3 **	255 ± 27.5
15	417 ± 38.3	277 ± 25.7	528 ± 45.9 (27)	292 ± 23.8	487 ± 43.7*	256 ± 26.0 *	477 ± 33.5**	259 ± 28.6 *
16	526 ± 40.1	281 ± 25.8	536 ± 50.4 (27)	297 ± 26.2	500 ± 44.3	264 ± 29.7	487 ± 32.9 **	262 ± 27.9 *
17a	535 ± 44.2	285 ± 27.4	549 ± 55.5(27)	306 ± 28.1*	512 ± 48.7	272 ± 30.9 (25)	496 ± 38.5**	266 ± 29.3 *
18a	551 ± 45.5	290 ± 26.4	562 ± 55.2 (27)	309 ± 32.4 (24)	531 ± 54.6	278 ± 31.5 (25)	509 ± 39.2 **	270 ± 30.3 (25)
19a	559 ± 47	289 ± 42.8	570 ± 56.7 (27)	327 ± 30.3 (10)	532 ± 52.3	309 ± 25.5 (4)	516 ± 38 **	275 ± 33.6 (4)
20a	570 ± 45.3	-	576 ± 58.3 (27)	328 ± 31.2 ((6)	544 ± 56.3	306(1)	525 ± 39.9 **	262 (1)
21a	578 ± 48.8	-	588 ± 63.7 (27)	318 ± 53.6 (3)	553 ± 55.3 (27)	299 ± 10.6 (2)	536 ± 39.4 **	265 (2)
22a	586 ± 52.1	-	593 ± 64.0 (27)	325 ± 21.0 (3)	563 ± 58.3 (27)	299 ± 10.6 (2)	542 ± 39.0 **	302 ± 8.5 (7)
23a	597 ± 54.7	299 ± 29.7 (5)	612 ± 67.8 (27)	342 ± 41.0 (5)	573 ± 61.2 (27)	296 ± 35.0 (3)	550 ± 39.8 **	278 ± 31.7 (9)
24a	606 ± 54.8	306 ± 26.7 (5)	618 ± 66.6 (27)	349 ± 45.7 (8)	582 ± 61.8 (27)	310 ± 33.6 (5)	561 ± 43.7 **	283 ± 37.1(22)
25a	618 ± 58.9	311 ± 19.3(20)	631 ± 65.5 (27)	339 ± 44.3 *(17)	592 ± 63.4 (27)	298 ± 29.2 (20)	571 ± 43.2 **	289 ± 34.0*
26a	628 ± 58.2	304 ± 24.9 (27)	641 ± 68.3 (27)	331 ± 43.0 *(24)	598 ± 63.5 (27)	291 ± 29.1 (27)	575 ± 43.1**	284 ± 31.5 **
27a	636 ± 60.99	311 ± 26.7 (27)	652 ± 70.6 (27)	338 ± 40.6 **(27)	608 ± 66.0 (27)	301 ± 31.5	583 ± 47.8 **	289 ± 31.7 *(27)
28b	646 ± 62.1	316 ± 28.3 (27)	658 ± 68.5 (27)	345 ± 44.3 *	611 ± 68.6 (27)	309 ± 34.7(24)	591 ± 43.3 ** (26)	296 ± 38.4 (26)
29b	652 ± 62.8	315 ± 27.1 (26)	667 ± 66.7 (27)	349 ± 46.8 **(26)	622 ± 65.0 (27)	315 ± 35.1(26)	594 ± 46.2 **	301 ± 37.7(11)
30b	649 ± 63.1	336 ± 30.4(7)	666 ± 71.0 (25)	360 ± 37.2(12)	618 ± 65.5 (24)	334 ± 33.4 (7)	594 ± 48.6**	299 ± 43.4 (1)
31b	656 ± 64.7 (22)	400	668 ± 74.2 (19)	384 ± 39.5 (8)	628 ± 69.9 (24)	330 ± 42.7 (3)	598 ± 52.5 **	374
32b	664 ± 67.9	-	676 ± 75.5	426 ± 45.2 (3)	634 ± 72.6	338 ± 45.3 (2)	608 ± 49.7 ** (27)	-
33b	665 ± 67.8	-	678 ± 76.3	433 ± 0.7 (2)	628 ± 77.9	357 (1)	608 ± 53.8 ** (27)	-
34b	673 ± 68.6	-	688 ± 78.4	418 ± 0.0 (2)	644 ± 72.1	363 (1)	616 ± 53.7 ** (27)	-
35b	679 ± 71.9	-	695 ± 81.9	421 ± 4.2 (2)	650 ± 73.4	374 (1)	618 ± 52.6 ** (27)	-
36b	687 ± 72.1	-	698 ± 81.5	424 ± 10.6 (2)	655 ± 78.3	-	628 ± 51.8 ** (27)	-
37b	690 ± 74.0	-	703 ± 83.2	-	658 ± 81.1	-	634 ± 49.5 *(27)	-

38b	693±75.1	-	703 ± 81.2	-	661 ± 85	-	641 ± 53.4 *(23)	-
39b	692±76.3	-	698 ± 95.1	-	664 ± 86.3	-	629 ± 50.0 (16)	-

Note: Number of animals weighed was 28 unless otherwise stated in brackets of results column.

* =Significantly different from the control group; $p \leq 0.05$ ¹

**= Significantly different from the control group; $p \leq 0.01$

aF_{2a}= Gestation/lactation period

bF_{2b}= Gestation/lactation period

A2.10.1.2 Comparison with the CLP criteria

Chrysanthemum cinerariaefolium extract from supercritical CO₂ does not meet the EU criteria to be classified for sexual function and fertility. There are not findings related to the sexual function or the fertility (CLP 3.7.2.1.1. (Table 3.7.1 a))).

A2.10.1.3 Overall conclusion on sexual function and fertility related to risk assessment

Conclusion used in Risk Assessment – Effects on fertility	
Value/conclusion	No effects were observed.
Justification for the value/conclusion	Not classified

A.2.10.2. Developmental toxicity

Table A.55 Summary table of animal studies on adverse effects on development

Summary table of animal studies on adverse effects on development							
Method, Duration of exposure, Route of exposure, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels,	NOAELs, LOAELs (e.g. maternal, teratogenicity, embryotoxicity, offspring, parental, reproductive toxicity)	Results, maternal/parental (e.g. corrected body weight gain, for all dose levels)	Results, developmental (e.g. pup survival, structural abnormalities, altered growth, functional deficiencies, incidences and severity of the effects for all dose levels)	Remarks (e.g. major deviations)	Reference
Teratogenicity Test EPA 83-3 OECD 414 Day 6–15 post mating GLP Reliability 2 Key	Rat Charles River COBS® CD® rats Female 25/group	Pyrethrum extract (Blend FEK-99; Purity 57.574%) 0, 5, 25, and 75 mg/kg bw/d total pyrethrins (0, 8, 38, and 115 mg/kg bw/d extract)	NOAEL maternal toxicity >75 mg/kg bw/d total pyrethrins (115 mg/kg bw/d extract) NOAEL Teratogenicity Embryotoxicity 75 mg/kg bw/d total pyrethrins (115 mg/kg bw/d extract)	No treatment related effects	Critical effects Dams foetuses No treatment related effects	1. The test substance was administered daily from day 6 through 15 of gestation instead of from implantation to the day prior to scheduled caesarean section, as the guideline recommends. 2. Food consumption was not	<div style="background-color: black; color: white; padding: 2px;"> (KPIC) IIIA6.8.1/01 (BRA, MGK and SCJ) IIIA6.8.1/1 </div>

						recorded as the guideline recommends.	
<p>Teratogenicity Test EPA 83-3 OECD 414 Day 7–19 post mating GLP Reliability 2 Key</p>	<p>Rabbit New Zealand White SPF Female 16/group</p>	<p>Pyrethrum extract (Blend FEK-99; Purity 57.574%) 0, 25, 100 and 250 mg/kg/d total pyrethrins (0, 38, 153, and 383 mg/kg bw/d extract)</p>	<p>NOAEL maternal Toxicity: 25 mg/kg bw/d total pyrethrins (38 mg/kg bw/d extract) NOAEL Teratogenicity Embryotoxicity: 250 mg/kg bw/d (383 mg/kg bw/d extract)</p>	<p>Maternal (250 and 100 mg/kg bw/d total pyrethrins (383 and 153 mg/kg bw/d extract)) Body weight ↓ and excessive salivation and arched head</p>	<p>Critical effects Dams foetuses No effects on foetuses</p>	<p>1. The test substance was administered daily from day 7 through 19 of gestation instead of from implantation to the day prior to scheduled caesarean section, as the guideline recommends. 2. Individual body weights were recorded on gestation days 0, 7, 13, 20, 24 and 29 instead of at day 0, on the first day of dosing, at least every 3 days during the dosing period and on</p>	<p>(KPIC) IIA6.8.1/02 (BRA, MGK and SCJ) IIA6.8.1/2</p>

						the day of scheduled kill as the guideline recommends. 3. Food consumption was not recorded as the guideline recommends.	
--	--	--	--	--	--	---	--

Table A.56 Summary table of human data on adverse effects on development

No data are available.

Table A.57 Summary table of other relevant studies for developmental toxicity

No data are available.

A2.10.2.1 Short summary and overall relevance of the provided information on adverse effects on development

In addition to the Two-generation reproductive toxicity study (██████████), two teratogenicity studies have been carried out:

Rat

In a definitive study groups of 25 mated female Charles River rats were given *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ suspended in 0.5% methylcellulose orally by gavage at doses of 0, 5, 25, and 75 mg/kg bw/d total pyrethrins (0, 8, 38, and 115 mg/kg bw/d extract) on days 6-15 of gestation at a volume of 3 mL/kg. On day 20 of gestation, the foetuses were removed surgically for evaluation. One female in the 75 mg/kg/d total pyrethrins (115 mg/kg bw/d extract) dose group delivered its litter on gestation day 19, although this premature delivery was not considered to be treatment related.

No animals died or were killed *in extremis* during the study, and no treatment-related clinical signs were observed. Hair loss was the most frequent noted finding and was observed among all groups, including the control. The body-weight gains of the treated groups were comparable to those of the controls during treatment. No evidence of fetotoxicity was found, and morphological examination revealed no teratogenic effects at any dose tested. The following observations were noted among the treated groups but were not considered to be treatment related: folded retina in one foetus in the control, one in the 5 mg/kg/d (8 mg/kg bw/d extract) dose group, three in the 25 mg/kg/d total pyrethrins (38 mg/kg bw/d extract) dose group and two in the 75 mg/kg/d total pyrethrins (115 mg/kg bw/d extract) dose group; microphthalmia, anophthalmia and right-sided aortic arch occurred in single incidences in the control, 5 mg/kg/d total pyrethrins (8 mg/kg bw/d extract) dose group and 25 mg/kg/d total pyrethrins (38 mg/kg bw/d extract) dose group, respectively.

The NOAEL for maternal toxicity was 75 mg/kg bw/d total pyrethrins (115 mg/kg bw/d extract), and that for developmental toxicity was 75 mg/kg bw/d total pyrethrins (115 mg/kg bw/d extract), the highest dose tested ██████████. (KPIC and BRA, MGK and SCJ)

Rabbit

In a definite study groups of 16 inseminated female New Zealand white SPF rabbits were randomly assigned to receive Pyrethrins at doses of 0, 25, 100 and 250 mg/kg/d total pyrethrins (0, 38, 153, and 383 mg/kg bw/d extract) orally by gavage on days 7-19 of gestation at a volume of 3 mL/kg in methylcellulose. On day 29 of gestation, the foetuses were removed surgically for evaluation.

All animals survived to the end of treatment. One doe aborted in the high-dose group and whole litter resorption occurred for an additional high-dose doe, but it is not clear if these findings were related to treatment. Maternal weight loss or reduced body weight gain during the treatment period and excessive salivation and arched head post-dose was observed in few animals of the mid- and high-dose groups. There were no treatment-related effects on foetal development including teratogenicity. The NOEL/NOAEL for maternal toxicity was 25 mg/kg bw/d total pyrethrins (38 mg/kg bw/d extract) and that for developmental toxicity was 250 mg/kg bw/d total pyrethrins (383 mg/kg bw/d extract), the highest dose tested ██████████. (KPIC and BRA, MGK and SCJ)

	Total pyrethrins dosage level (mg/kg bw/d)											
	Extract (mg/kg bw/d)											
	0 (Control) 0			25 38			100 153			250 383		
	No.	%	± SD	No.	%	± SD	No.	%	± SD	No.	%	± SD
Animals on study	16	-	-	16	-	-	16	-	-	16	-	-
Animals that were gravid	16	-	-	15	-	-	15	-	-	16	-	-
Animals that died	0	-	-	0	-	-	0	-	-	0	-	-
Animals that aborted near term	0	-	-	0	-	-	0	-	-	1	-	-
Animals examined at Caesarean section	16	-	-	16	-	-	16	-	-	15	-	-
Nongravid	0	-	-	1	-	-	1	-	-	0	-	-
Gravid:	16	-	-	15	-	-	15	-	-	15	-	-
Does with resorption only:	0	-	-	0	-	-	0	-	-	1	-	-
Does with viable foetuses:	16	-	-	15	-	-	15	-	-	14	-	-
Viable foetuses/does:	7.6	-	3.22	6.9	-	2.80	6.5	-	2.67	6.7	-	3.20
Postimplantation loss/does:	0.4	-	0.73	1.0	-	1.81	1.1	-	1.16	0.7	-	1.18
Total implantations/does:	8.1	-	3.04	7.9	-	2.33	7.5	-	2.67	7.4	-	2.90
Corpora lutea/does:	11.3	-	3.87	11.5	-	2.68	10.9	-	2.17	11.0	-	3.76
Group mean preimplantation loss (%) ^a	-	28.3	-	-	32.3 ^d	-	-	30.7	-	-	31.2 ^c	-
Group mean postimplantation loss (%) ^b	-	5.4	-	-	12.7	-	-	14.2	-	-	8.2	-
Mean total body weight (grams):	39.0	-	5.56	41.2	-	7.24	42.6	-	6.42	41.8	-	8.31
Foetal sex distribution – male:	62	50.8	-	67	60.9	-	44	45.4	-	44	43.6	-
– female:	60	49.2	-	43	39.1	-	53	54.6	-	57	56.4	-

^a $\frac{\text{Total number of corpora lutea} - \text{Total number of implantations}}{\text{Total number of corpora lutea}} \times 100$

^b $\frac{\text{Total number of implantations} - \text{Total number of viable foetuses}}{\text{Total number of implantations}} \times 100$

^c Value does not include doe with regressing corpora lutea

^d Value does not include doe with number of corpora lutea less than number of implantations

Values of the treated groups did not differ significantly from those of the control group; $p > 0.05$

TABLE 1. Summary of Maternal Antemortem Observations

Observation	Pyrethrin Dosage Level (mg/kg/day)							
	0 (Control)		25		100		250	
	Number	%	Number	%	Number	%	Number	%
Number of animals observed	16	100	16	100	16	100	16	100
No visible abnormalities	11	69	12	75	13	81	7	44
Aborted near term							1	6
Hair loss	2	13	1	6			4	25
No stool							2	13
Small amount of stool	4	25	2	13	2	13	3	19
Soft stool					1	6		
Ocular discharge			1	6				
Material around nose							1	6
Incisors malaligned			1	6				
Incisor missing							1	6
Excessive salivation, post-dose					1	6	2	13
Head arched backward, post-dose					1	6	3	19
Labored breathing, post-dose							2	13

A2.10.2.2 Comparison with the CLP criteria

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is not classified for reproductive toxicity according to EU criteria due to there are not findings related to development (CLP 3.7.2.1.1. (Table 3.7.1 a))).

A2.10.2.3 Overall conclusion on effects on development related to risk assessment

Conclusion used in Risk Assessment – Effects on development	
Value/conclusion	No effects on development were observed.
Justification for the value/conclusion	Not classified

A.2.10.3. Effects on or via lactation

A2.10.3.1 Short summary and overall relevance of the provided information on adverse effects on or via lactation

An assay about the toxicokinetics and distribution of the *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ [REDACTED] shows that the a.i. accumulates in the fatty tissue. This propertie indicates the likelihood that the substance is present in potentially toxic levels in breast milk. However, in the Two-generation reproductive toxicity study [REDACTED] there are no effects on or via lactation.

A2.10.3.2 Comparison with the CLP criteria

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is not classified for reproductive toxicity according to EU criteria due to there are not findings related to effects on or via lactation, although substance is susceptible to being found in breast milk (CLP 3.7.2.1.1. (Table 3.7.1 b))).

A2.10.3.3 Overall conclusion on effects on development related to risk assessment

Conclusion used in Risk Assessment – Effects on or via lactation	
Value/conclusion	No effects on development were observed.
Justification for the value/conclusion	Not classified

A2.10.4 Conclusion on classification and labelling for reproductive toxicity

No effects were noted on reproduction. No classification required.

A2.10.5 Overall conclusion on reproductive toxicity related to risk assessment

No effects were noted on reproduction.

Conclusion used in the Risk Assessment – Reproductive toxicity	
Value	-
Justification for the selected value	No effects were noted on reproduction.
Proposed classification	Not classified.

A.2.11. Aspiration hazard

No data are available.

A2.11.1 Short summary and overall relevance of the provided information on aspiration hazard

Not classified.

A2.11.2 Comparison with the CLP criteria

It does not meet the EU criteria to be classified as aspiration hazard because, according to CLP 3.10.2. (Table 3.10.1), kinematic viscosity > 20.5 mm²/s.

A2.11.3 Conclusion on classification and labelling for aspiration hazard

Not classified.

A.2.12. Neurotoxicity

Table A.58 Summary table of animal studies on neurotoxicity

Summary table of animal studies on neurotoxicity						
Method, Duration of exposure, Route of exposure, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/ group	Test substance (including purity), Vehicle, Dose levels,	NOAEL, LOAEL	Results	Remarks (e.g. major deviations)	Reference
14 days EPA 81-8 GLP Single dose Reliability 1 Key	Rat Charles River CD 15 animal/sex group	Pyrethrum Extract (Batch Blend FEK-99/LS 92-37; Purity 57.467% (w/w)) Males: 0, 40, 125, and 400 mg/kg bw total pyrethrins; 0, 61, 192, and 613 mg/kg bw extract Females: 0, 20, 63, and 200 mg/kg bw total pyrethrins; 0, 31, 97, and 306 mg/kg bw extract Single dose	NOAEL: Males: 40 mg/kg bw total pyrethrins (61 mg/kg bw extract) Females: 20 mg/kg bw total pyrethrins (31 mg/kg bw extract) LOAEL: Males: 125 mg/kg bw total pyrethrins (192 mg/kg bw extract) Females: 63 mg/kg bw total pyrethrins (97 mg/kg bw extract)	400/200 mg/kg bw total pyrethrins (613/306 mg/kg bw extract) Males/females: Mortalities, acute neurological signs 125/63 mg/kg bw total pyrethrins (192/97 mg/kg bw extract) Males/females: Decreased motor activity in males Fine tremors in 3/15 females 40/20 mg/kg bw total pyrethrins (61/31 mg/kg bw extract) Males/females: No effects	-	██████████ (KPIC) IIIA6.9 ██████████ (BRA, MGK and SCJ) IIIA6.9

Table A.59 Summary table of human data on neurotoxicity
No data are available.

A2.12.1 Short summary and overall relevance of the provided information on neurotoxicity

In a neurotoxicity study, groups of 15 male Sprague-Dawley rats received by gavage a 10% w/v solution of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ in corn oil at doses of 0, 40, 125 and 400 mg /kg bw total pyrethrins (0, 61, 192, and 613 mg/kg bw extract), and 15 females received a 5% w/v solution of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ in corn oil at doses of 0, 20, 63 or 200 mg/kg bw total pyrethrins (0, 31, 97, and 306 mg/kg bw extract).

Five males and two females at the high dose died on the day of treatment, and a variety of acute neurological signs were observed in the other animals at this dose, including tremors, urogenital area wetness, salivation, perinasal encrustation, exaggerated startle response, decreased grip strength, hind leg splay, and increased body temperature. Tremors were also observed in three females at the intermediate dose. Measurements of motor activity on the day of treatment indicated increased fine movement and decreased rearing and ambulation in animals of each sex at the high dose and decreased fine movement, rearing and ambulation in males as the intermediate dose. This is likely due to an effect of treatment and a predisposition for lower activity of this group if compared to the control group during pre-treatment evaluation.

Time posttreat.	Males				Females			
	Pretreat.	3 h	7 d	14 d	Pretreat.	3 h	7 d	14 d
N	15	14	10	10	15	15-12 ¹	13	13
Cage posture								
Normal/awake	13	13	7	6	15	12	12	10
Normal/asleep	2	1	3	4	0	0	1	3
On side/prostrate	0	0	0	0	0	3	0	0
Cage clonic convulsions								
None	-	14	-	-	-	-	-	-
Explosive jumps	-	0	-	-	-	-	-	-
Cage fine tremors								
None	-	-	-	-	-	11	-	-
Whole body	-	-	-	-	-	4	-	-
Head	-	-	-	-	-	0	-	-
Cage coarse tremors								
None	-	1**	-	-	-	5**	-	-
Whole body	-	13	-	-	-	10	-	-
Cage palpebral closure								
Wide open	13	13	7	6	15	15	12	10
Half shut	0	0	0	0	0	0	0	0
Shut	2	1	3	4	0	0	1	3
Handling reactivity								
Reists	-	13	-	-	-	-	-	-
High resistance	-	1	-	-	-	-	-	-
Gait								
Normal	-	11	-	-	-	7**	-	-
Splayed	-	2	-	-	-	6	-	-
Hypotonic	-	1	-	-	-	0	-	-
Prostrate	-	0	-	-	-	1	-	-
Body position								
Normal	-	13	-	-	-	11	-	-
Hunched	-	1	-	-	-	0	-	-
On side	-	0	-	-	-	1	-	-
On stomach	-	0	-	-	-	2	-	-

Fine tremors								
None	-	11	-	-	-	7**	-	-
Whole body	-	3	-	-	-	7	-	-
Coarse tremors								
None	-	4**	-	-	-	7**	-	-
Whole body	-	10	-	-	-	7	-	-
Unusual behaviour								
None	-	-	-	-	-	13	-	-
Prostrate	-	-	-	-	-	1	-	-
Arousal								
Active/alert ²	-	13	8	7	15	8	11	12
Hyperactive	-	0	0	0	0	1	1	0
Inactive/alert	-	1	2	3	0	3	1	1
Inactive/Not alert	-	0	0	0	0	2	0	0
Palpebral closure								
Wide open	-	-	-	10	-	-	-	-
Slightly dropping	-	-	-	0	-	-	-	-
Defecation								
None	8	9	7	9	14	13	12	11
Normal	7	4	3	1	1	1	1	2
Soft	0	1	0	0	0	0	0	0
Urine								
None	6	4	7	6	3	10	9	7
Present	9	10	3	4	12	4	4	6
Rears (events)								
Mean	7.27	2.79	5.40	4.40	6.47	3.57	8.08	9.46
SD	2.71	2.16	4.03	2.95	3.54	4.89	5.95	5.58
Approach response								
Noticeable	14	13	-	-	-	9	-	-
None	0	0	-	-	-	1	-	-
Exaggerated	1	1	-	-	-	3	-	-
Startle response								
Noticeable	12	5**	-	-	15	6**	-	-
None	0	0	-	-	0	1	-	-
Exaggerated	3	9	-	-	0	6	-	-
Tail pinch response								
Noticeable	13	10	10	9	15	11	12	-
None	0	2	0	1	0	2	0	-
Exaggerated	2	2 ³	0	0	0	0	1	-
Pupil size								
Normal	-	13	10	10	-	12	8	7
Decreased	-	1	0	0	-	1	5	6
Visual placing								
Present	-	-	10	-	-	-	13	-
None	-	-	0	-	-	-	0	-
Muscle tone								
Normal	-	11	10	-	-	11	12	-
Decreased	-	3	0	-	-	2	1	-
Fur appearance								
Normal	-	12	-	-	-	-	-	-
Urine stains	-	2	-	-	-	-	-	-
Lacrimation								
None	-	-	-	-	-	12	-	-
Present	-	-	-	-	-	1	-	-
Salivation								

None	-	-	-	-	-	12	-	-
Excessive	-	-	-	-	-	1	-	-
Crust								
None	13	9*	10	-	-	8*	12	-
Eyes	2	0	0	-	-	1	0	-
Nose	0	5	0	-	-	4	1	-
Grip strength (fore) (kg)								
Mean	0.53	0.59	0.84	0.89	0.5	0.48**	0.77	0.75
SD	0.08	0.18	0.14	0.17	0.11	0.15	0.16	0.18
Grip strength (hind) (kg)								
Mean	0.44	0.46**	0.63	0.58	0.39	0.44	0.52	0.47
SD	0.12	0.13	0.11	0.07	0.08	0.09	0.09	0.11
Rectal temperature (°C)								
Mean	38.05	39.30**	38.17	37.95	38.06	39.68**	38.38	38.45
SD	0.05	0.57	0.57	0.47	0.46	0.43	8.08	0.63
Air righting								
Feet/Coordinated	-	-	-	10	-	-	-	-
Feet/Uncoordinated	-	-	-	0	-	-	-	-
Hind leg splay (cm)								
Mean	8.26	6.83*	8.77**	7.85	6.07	5.66	6.93	6.90
SD	1.17	1.28	1.11	1.44	0.85	1.14	1.28	1.23

*Significantly different from their respective control groups ($p < 0.05$).

**Significantly different from their respective control groups ($p < 0.01$).

¹One animal died following cageside observations and two animals were not fully evaluated in the open field or the manipulative portions of the FOB due to the deteriorating condition of the animals.

²Not significantly different from their respective control groups when evaluated as noticeable or none vs. exaggerated.

- The incidence for these endpoints was zero or the same for all doses.

In addition to the above observations, all animals were observed for the following endpoints: cage tonic convulsions, excessive vocalization, breathing pattern, clonic convulsions, tonic convulsions, palpebral closure, piloerection, exophthalmus, emaciation, and dehydration.

Time posttreat.	Males				Females			
	Pretreat.	4 h	7 d	14 d	Pretreat.	4 h	7 d	14 d
N	15	12	10	10	15	13	13	13
Fine movement								
Mean	294.40	657.60*	320.90	317.50	361.90	862.3*	456.70	473.80
S.D.	74.82	512.65	103.16	94.95	149.27	759.94	180.14	116.20
Ambulation								
Mean	198.90	157.90*	169.70	167.60	204.90	146.90*	268.20	272.20
S.D.	52.51	63.06	48.85	70.68	98.33	81.84	108.81	97.19
Rearing								
Mean	113.90	62.00*	128.70	136.30	137.50	66.00**	169.00	201.90
S.D.	31.42	52.55	35.20	38.90	59.98	48.11	64.52	77.28

*Significantly different from their respective control groups ($p < 0.05$).

**Significantly different from their respective control groups ($p < 0.01$).

In addition, slight, statistically non-significant decreases in body weight were seen in males at the high dose on days 7 and 14. There was no evidence of any gross, treatment-related lesion. The microscopic changes were limited mainly to sections of the sciatic nerve and its branches. The histomorphological changes within the peripheral nerve sections indicated the presence of scattered degenerating nerve fibres or myelin sheaths. These changes were seen in only a few animals, were graded as minimal, and were not dose-related.

	MALES	FEMALES
--	-------	---------

FEEDING LEVEL (mg/kg diet)	0	40	125	400	0	20	63	200
NUMBER IN GROUP	15	15	15	15	15	15	15	15
SCIATIC NERVE								
myelin degeneration/myelin sheath swelling	0	0	0	2	0	0	0	1
minimal focal	0	0	0	2	0	0	0	1
myelin/axon degeneration	0	2	1	1	0	0	0	4
minimal focal	0	2	1	1	-	-	-	-
minimal multifocal	0	1	1	1	0	0	0	3
moderate multifocal	0	1	0	0	0	0	0	1
NERVE PERONEAL								
myelin/axon degeneration	-	-	-	-	0	1	0	2
minimal multifocal	-	-	-	-	0	1	0	1
moderate multifocal	-	-	-	-	0	0	0	1
NERVE TIBIAL								
myelin/axon degeneration	-	-	-	-	0	0	0	2
minimal multifocal	-	-	-	-	0	0	0	1
moderate multifocal	-	-	-	-	0	0	0	1

The NOEL/NOAEL was 40 mg/kg bw total pyrethrins (61 mg/kg bw extract) for male rats and 20 mg/kg bw total pyrethrins (31 mg/kg bw extract) for female rats ([REDACTED]). (KPIC and BRA, MGK and SCJ)

A2.12.2 Comparison with the CLP criteria

Not classified.

A2.12.3 Conclusion on neurotoxicity related to risk assessment

Conclusion used in Risk Assessment – Neurotoxicity	
Value/conclusion	Not classified
Justification for the value/conclusion	Refer to Section A3.12.1.

A.2.13. Immunotoxicity

No data are available.

A.2.14. Endocrine disruption

A2.14.1 Short summary and overall relevance of the provided information on endocrine disruption

To evaluate a potential concern for endocrine disruption effects induced by *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ all available data were assessed for all levels of the OECD Conceptual Framework (Guidance for the identification of endocrine disruptors in the context of Regulations (EU) No 528/2012 and (EC) No 1107/2009).

A.2.14.1. Level 1

Existing data and non-test information, covered, physical and chemical properties, toxicological data from standardized or not-standardized tests, and read-across, chemical categories, QSARs and another *in silico* predictions, and ADME model predictions.

A few endocrine-related mechanistic studies were identified for pyrethrum flower extract

from the published literature. These include:

- Several *in vitro* study of hepatic enzyme induction in rat and human hepatocytes (Price *et al.*, 2007; Price *et al.*, 2008; Osimitz & Lake, 2009).
- An *in vitro* study of human androgen receptor (AR) binding and sex hormone binding globulin (SHBG) binding (Eil and Nisula, 1990).
- An *in vitro* study of antagonistic transactivation activity at the human AR (Vinggaard *et al.*, 2008).
- An *in vitro* study of agonist and antagonist activity at the human AR and the human estrogen receptors (ER)- α and - β (Kojima *et al.*, 2004).

This information explains the effects of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ in the thyroid gland as a secondary effect of the hepatic enzyme induction.

QSAR predictions using Derek Nexus software do not trigger any alert related with endocrine-disruptive properties.

A.2.14.2. Level 2

In vitro assays providing data about selected endocrine mechanism(s) and pathway(s).

ToxCast models and CERAPP do not give information about estrogen- and androgen-modalities. COMPARA displays information about pyrethrin 1, pyrethrin 2, cinerin 1, and cinerin 2. These compounds are inactive as agonist for androgen receptor and active as antagonist and binding (cinerin 2 was inactive for binding too). However, these results are not correlated with results in the studies of EDSP21 database, where those positive responses take place exceeding cytotoxic limits or with an efficacy < 50%.

Steroidogenesis-modality was negative in the EDSP21 database for the only one aromatase inhibition assay.

Thyroid-modality is more ambiguous due to the solvent used in the mix. The mechanism by which the *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ shows effects in the thyroid gland is well known, so those effects observed *in vitro* could be attributed to BHT.

A.2.14.3. Level 3

In vivo assays providing data about selected endocrine mechanism(s) and pathway(s).

No information on such *in vivo* assays is available for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Mammals	Information available in the current dossier
E, A modalities	
Uterotrophic bioassay in rodents (OECD TG 440)	N
Hershberger bioassay in rats (OECD TG 441)	N

A.2.14.4. Level 4

In vivo assays providing data on adverse effects on endocrine relevant endpoints.

Mammals	Information available in the current dossier
T-modality	
Repeated dose 28-day study (OECD TG 407)	N
Repeated dose 90-day study (OECD TG 408)	Y
Repeated dose 90-day study in non-rodents (OECD TG 409)	Y
Combined chronic toxicity and carcinogenicity studies (OECD TG 451-3)	Y

The thirteen-week dose range finding study in rats (OECD TG 408) has some organ weights (epididymis, prostate and seminal vesicle with coagulating glands and uterus) and

histopathologies (mammary gland, prostate and seminal vesicle with coagulating glands and uterus) missing. Although some key endpoints are not available in this study, those that are on it can complement other studies because the duration of this assessment is enough. None of the organ weight changes could be correlated with microscopic findings. The liver weight changes are possibly treatment related, since some of the livers were macroscopically enlarged. Treatment related macroscopic findings included enlargement and congestion of the liver in both sexes. This was more pronounced in the male than in the female rats. These macroscopic findings in the liver could not be confirmed microscopically. All other macroscopic findings were considered to be incidental. The only possible treatment related microscopic findings consisted of small focal or multifocal areas of tubular degeneration and regeneration in the renal cortex. However, due to the low incidence of this renal lesion, it was not possible to determine its relationship with the administration of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. A small number of other microscopic findings were observed in various organs at the different dosage levels, but they were considered to be incidental and not treatment related (██████████, 1988b).

The eight-week dose range finding toxicity study in dogs (OECD TG 409) has some organ weights (epididymis and uterus) and a histopathology (seminal vesicle) missed. This study complements the others using a different mammalian target. A small number of macroscopic lesions were observed in all treated groups, but were considered not to be related to the administration of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂. Also, the number of animals per group is too low and this does not let perform a statistical study (██████████, 1988a).

The one-year chronic toxicity study in dogs (OECD TG 452) has some organ weights (uterus, epididymis) and a histopathology (vagina) missed. However, these endpoints are present in other studies and the duration of this study lets see the evolution of the others endpoints clearly. About the histopathological changes in different organs related with EAS-modality the fact is the same as the other studies, the control group shows the same incidence as the treated groups, so an endocrine-related mechanism could be discarded (██████████, 1990a).

The two-year dietary toxicity and oncogenicity study in rats (OECD TG 453) has some organ weights (uterus and epididymis) missing. However, these endpoints are present in other studies and the duration of this study lets see the evolution of the others endpoints clearly. This study shows a great amount of histopathological changes and types of tumors that could suggest estrogen-, androgen- and steroidogenesis-related mechanisms. However these effects are present in the two control groups and the difference is not statistically significant, and they are also present in other organs that are not related with endocrine disruption. Due to at this widespread incidence, it seems that there is not an endocrine-disruption mechanism involved (██████████, 1990b).

The eighteen-month dietary oncogenicity study in mice (OECD TG 453) has some organ weights (uterus, ovary, epididymis and adrenal glands) missed. However, these endpoints are present in other studies and the duration of this study lets see the evolution of the others endpoints clearly. About the histopathologic changes in ovary, uterus, testes, epididymis and the other organs related with estrogen-, androgen- and steroidogenesis-mediated activity, the case is that those changes are more present in these organs, but the incidence is not very different in comparison with the two control groups. This fact and the absence of statistically significant changes or the lack of the weight of some organs make impossible to assert a mechanism related with endocrine disruption. Finally, the endpoints potentially sensitive to, but not diagnostic of, E, A, T, S have the same point. The histopathologic changes are present but are very similar in comparison with the two control groups. For these reasons, this study seems to indicate a non-related mechanism because the biological significance is relative (██████████, 1990c).

A.2.14.5. Level 5

In vivo assays providing more comprehensive data on adverse effects on endocrine

relevant endpoints over more extensive parts of the life cycle of the organism.

Mammals	Information available in the current dossier
EAST-modality	
Two-generation reproduction toxicity study (OECD TG 416)	Y

In the two-generation reproduction toxicity study (OECD TG 416), in general terms, there are not significant dose- or develop-related and histopathology-supported changes so this study does not show endocrine related effects or adversity in EAS-modality. Changes in adrenal glands weight could be a systemic toxicity related effect, but in any case there are not coherence between studies or between sexes. However, this study has some endpoints missing: change in AGD in pups, changes in estrus cyclicity in P and F1 generations, decreased age at VO and increased age at PPS in F1, changes in some organ weights (uterus, ovaries, testes (except for those of the males which did not sire a litter), epididymides, prostate, seminal vesicles (+ coagulating glands), thyroid and liver), some histopathologic changes (coagulating glands and thyroid) and sperm parameters (except for those of the males which did not sire a litter (numerical data not shown in the study)) in P and F1 generations. (██████████, 1989).

A2.12.2 Comparison with the CLP criteria

Not classified.

A2.12.3 Conclusion on endocrine disruption related to risk assessment

Conclusion used in Risk Assessment – Endocrine disruption	
Value/conclusion	Not classified
Justification for the value/conclusion	<p>Some concerns remain after the assessmet of all relevant available information of endocrine disrupting properties for <i>Chrysanthemum cinerariaefolium</i>, extract from open and mature flowers of <i>Tanacetum cinerariifolium</i> obtained with hydrocarbon solvents.</p> <p>EAS-modalities have been considered sufficiently investigated and no ED properties have been detected assessing all the gathered information.</p> <p>However, T-modality has not been considered sufficiently investigated and ED properties for this modality cannot be totally excluded. Since new studies cannot be requested due to this dossier is a backlog dossier a conclusion cannot be reached.</p> <p>The first draft CARs for this active substances (not redefined yet) were submitted to COM before September 2013. According to the agreement reached in the CA-meeting in March 2018 (CA-March18-Doc.7.3a-Final – EDs - Active substances under assessment) the applicant is not obliged to provide new studies, although they have the opportunity to do it. In consequence, the eCA does not need to reach a conclusion regarding this issue.</p> <p>For EAS-modalities, although several endpoints are missing in the two-generation reproduction toxicity study, overall, the available toxicological data on repeated dose toxicity studies does not indicate EAS adversity by the test substance, suggesting that the ED criteria for EAS modalities are not met.</p> <p>Literature, <i>in silico</i> and <i>in vitro</i> mechanistic ToxCast data were also collected and did not indicate EATS-mediated activity with the extract individual compounds.</p> <p>Therefore, and although the dataset seems not complete for EAS-</p>

	<p>modalities, in the absence of adversity and activity related to the endocrine system in the organs of both genders of different species and with different time of exposure duration (short-, long term and develop/reprotoxicity studies), no further studies (especially <i>in vivo</i>) should be required to conclude on EAS-modality.</p> <p>However, since a new OECD TG 416 or OECD TG 443 and a literature review will be requested in the renewal stage, the chosen scenario is 2a (ii).</p> <p>For T-modality, as a whole, data suggest that any thyroid effects of pyrethrum flower extract are species-specific (i.e., only seen in the rat) and occur with chronic exposure only and secondary to the induction of liver microsomal enzymes. However, the potential for T-mediated adversity is considered to have not been sufficiently investigated. Overall, there is not sufficient weight of evidence to indicate that pyrethrum flower extract affects the thyroid modality by a mode of action that is specific to the rat and, as such, it cannot be concluded there are no indications of endocrine adversity of relevance to humans.</p> <p>Since a new OECD TG 416 or OECD TG 443 and a literature review will be requested in the renewal stage, the chosen scenario is 2a (iii).</p> <p>Table 5: High level summary of the scenarios, including the next steps in the assessment; for a full description of the scenario, refer to Section 3.4.4</p> <table border="1"> <thead> <tr> <th>Adversity based on 'EATS-mediated' parameters</th> <th>Positive mechanistic OECD CF level 2/3 test</th> <th>Scenario</th> <th>Next step of the assessment</th> </tr> </thead> <tbody> <tr> <td>No (sufficiently investigated)</td> <td>Yes/No</td> <td>1a</td> <td>Conclude: ED criteria not met because there is no 'EATS-mediated' adversity</td> </tr> <tr> <td>Yes (sufficiently investigated)</td> <td>Yes/No</td> <td>1b</td> <td>Perform MoA analysis (postulate and document the MoA, see Section 3.5.1)</td> </tr> <tr> <td>No (not sufficiently investigated)</td> <td>Yes</td> <td>2a (i)</td> <td>Perform MoA analysis; additional information may be needed for the analysis</td> </tr> <tr> <td>No (not sufficiently investigated)</td> <td>No (sufficiently investigated)</td> <td>2a (ii)</td> <td>Conclude: ED criteria not met because no endocrine activity has been observed for the EATS modalities</td> </tr> <tr> <td>No (not sufficiently investigated)</td> <td>No (not sufficiently investigated)</td> <td>2a (iii)</td> <td>Generate missing level 2 and 3 information. Alternatively, generate missing 'EATS-mediated' parameters. Depending on the outcome of these tests move to the corresponding scenario</td> </tr> <tr> <td>Yes (not sufficiently investigated)</td> <td>Yes/No</td> <td>2b</td> <td>Perform MoA analysis (postulate and document the MoA, see Section 3.5.1)</td> </tr> </tbody> </table> <p>EATS: Estrogen, androgen, thyroid, steroidogenic; MoA: Mode of action.</p>	Adversity based on 'EATS-mediated' parameters	Positive mechanistic OECD CF level 2/3 test	Scenario	Next step of the assessment	No (sufficiently investigated)	Yes/No	1a	Conclude: ED criteria not met because there is no 'EATS-mediated' adversity	Yes (sufficiently investigated)	Yes/No	1b	Perform MoA analysis (postulate and document the MoA, see Section 3.5.1)	No (not sufficiently investigated)	Yes	2a (i)	Perform MoA analysis; additional information may be needed for the analysis	No (not sufficiently investigated)	No (sufficiently investigated)	2a (ii)	Conclude: ED criteria not met because no endocrine activity has been observed for the EATS modalities	No (not sufficiently investigated)	No (not sufficiently investigated)	2a (iii)	Generate missing level 2 and 3 information. Alternatively, generate missing 'EATS-mediated' parameters. Depending on the outcome of these tests move to the corresponding scenario	Yes (not sufficiently investigated)	Yes/No	2b	Perform MoA analysis (postulate and document the MoA, see Section 3.5.1)
Adversity based on 'EATS-mediated' parameters	Positive mechanistic OECD CF level 2/3 test	Scenario	Next step of the assessment																										
No (sufficiently investigated)	Yes/No	1a	Conclude: ED criteria not met because there is no 'EATS-mediated' adversity																										
Yes (sufficiently investigated)	Yes/No	1b	Perform MoA analysis (postulate and document the MoA, see Section 3.5.1)																										
No (not sufficiently investigated)	Yes	2a (i)	Perform MoA analysis; additional information may be needed for the analysis																										
No (not sufficiently investigated)	No (sufficiently investigated)	2a (ii)	Conclude: ED criteria not met because no endocrine activity has been observed for the EATS modalities																										
No (not sufficiently investigated)	No (not sufficiently investigated)	2a (iii)	Generate missing level 2 and 3 information. Alternatively, generate missing 'EATS-mediated' parameters. Depending on the outcome of these tests move to the corresponding scenario																										
Yes (not sufficiently investigated)	Yes/No	2b	Perform MoA analysis (postulate and document the MoA, see Section 3.5.1)																										

A.2.15. Further Human data

Medical surveillance on manufacturing plant personnel

58 people who had been employed in a Pyrethrum extract factory for 1 to 25 years were systematically examined on digestive, cardiovascular, respiratory, muscular and endocrine systems and on sense organs. Laboratory measurements were conducted with regard to haematology, blood electrolytes, blood sugar, serum transaminases, plasma testosterone and plasma thyroxine.

The individuals were found to be clinically healthy regardless of the degree of exposure to Pyrethrins. Minor pleural lesions of inflammatory type were found in people exposed to high concentrations of aerial Pyrethrum dust (daily concentrations of 6000 ppm of Pyrethrum powder, equivalent to 78 ppm Pyrethrum extracts). These lesions did not impair the health of the individuals and were probably a reaction to irritant dust. However, the

study is not acceptable and it is not possible to obtain conclusions because the original Document IV is not complete. Specifically all tables containing original data with current and previous ailments, results of the examination of the respiratory and cardiovascular systems and results of haematology and hormonal assays are not available (Gombe & Ogada, 1976). (KPIC)

Also, in a more recent update no health concerns affecting workers or other people in and around the factory were observed (Wangai, 2002). (KPIC)

The manufacturing plant of MGK has been extracting, refining and formulating with Pyrethrins for more than 70 years. According to historical data of MGK's employees, no serious adverse reaction to Pyrethrins have been reported. A few minor cases of dermatitis were noted, however, the implication of Pyrethrins is not confirmed.

The manufacturing plant of BRA in Tasmania has been harvesting and extracting Pyrethrins for many years. Over the last 25 years of manufacturing, no serious health problems have been recorded in workers. Minor skin irritations have been observed following the harvesting operation, however these effects are more likely to be due to the dustiness of the harvested crop. These skin irritations disappeared after washing with clean water and at worst within 1-2 days. Once the harvested crop material has been extracted, no cases of any health issues from handling the extracted and refined products have been noted. (BRA, MGK and SCJ)

Direct observations on clinical cases or poisoning incidents

Based on more than 80.000 calls, the Report to the American Association of Poison Control Centers concludes "that products containing Pyrethrins or pyrethroids can be used with the expectation of no undue risk" (Anonymous, 2001). (KPIC)

Occupational exposure in a Pyrethrum extract factory did not result in clinical symptoms even upon high exposure levels with the exception of minor pleural lesions of the inflammatory type that were attributed to the mechanical irritation of the high levels of dust and were not impairing the health of the individuals affected. (KPIC)

Pyrethrins have been on the market for many years and their applications/uses are numerous. Few isolated incidents have been reported in the literature. In any case, the data provided is not sufficient to establish a relationship between Pyrethrins and the effects and/or symptoms caused. Pyrethrins are not known to cause any serious poisoning incidents, the vast majority of exposures are relatively harmless and self-limited. The most common symptoms include: dermatitis with papules in moist areas, intense pruritus, nausea, a stinging sensation in the nasal and upper pharyngeal mucosa, moderate shortness of breath, a cough productive of white phlegm without haemoptysis, fatigue, headache, dizziness and sensitisation (Garcia-Bravo *et al.*, 1995; Paton & Walker, 1988; Wax & Hoffman, 1994). (BRA, MGK and SCJ)

Health records from industry or other sources

No health records have been reported from industry or any other sources. (BRA, MGK and SCJ)

Epidemiological studies on the general population

Analysis of exposures of insecticides containing Pyrethrins and/or Pyrethroids from 1994 to 1999 in the USA to humans was entered into the TESS© database. 81838 people were exposed by a variety of routes to consumer products. 41080 cases were reported to be related to products containing Pyrethrins or Pyrethroids.

One-third of the exposures were through ingestion, while inhalation, dermal and ocular exposures occurred in 27.8%, 26.2% and 10.7% of the cases, respectively. Exposure was unintended in 93.1% of cases, in or around the home in 93% and at work in 5% of cases. Nearly all cases (95%) involve an acute exposure (including continuous or repeated exposure lasting less than 8 hours) to Pyrethrin/Pyrethroid. The most frequently reported symptoms were ocular (22.8%), gastrointestinal (22.3%), dermal (21%), respiratory

(12.8%), miscellaneous symptoms (10.1%) and neurologic (9.7%). Children under 5 years were most likely to report ocular symptoms; children between the age of 5 and 9 were more likely to exhibit ocular and dermal symptoms. Adults were diagnosed with a variety of symptoms such as gastrointestinal (34.8%), dermal (33.3%), ocular (27.8%) and respiratory (25.1%). During the investigation, no deaths were registered as being due to Pyrethrins/Pyrethroids. The TESS report revealed a number of limitation of which the most significant is the non-distinction between Pyrethrins and Pyrethroids (Pegus, 2001). (BRA, MGK and SCJ)

Diagnosis of poisoning including specific signs of poisoning and clinical tests

Symptoms include dermatitis with papules in moist areas, intense pruritus, bullae, nausea, moderated shortness of breath and chest tightness, a cough productive of white phlegm, fatigue, headache and dizziness.

Contact dermatitis is the most common effect (HSDB database, 2001). It causes a mild erythematous vesicular dermatitis with papules in moist area and intense pruritus. In some cases, blisters appear. Oedema and skin cracking develop in severe cases. The clinical manifestations of inhalation exposure to Pyrethrins can be local or systemic. Localised reactions confined to the upper respiratory tract include rhinitis, sneezing, scratchy throat, oral mucosal oedema, and even laryngeal mucosal oedema. Localised reactions of the lower respiratory tract include coughing, shortness of breath and chest pain (Paton et al, 1988). An asthma like reaction occurs with acute exposures in sensitised patients. Allergic rhinitis and pneumonitis were also noted.

Pyrethrins also trigger conjunctival oedema and hyperaemia as if can be irritating to eyes and mucous membrane (HSDB database, 2001). (BRA, MGK and SCJ)

Sensitisation/allergenicity observations

Pyrethrum sensitivity has been reported in a number of cases via different forms. Contact dermatitis is the most common allergic reaction. In few cases, bullae appear. Oedema and cracking develop in severe cases. Local anaphylaxis was noted and described by dermatitis and sudden swelling of face.

Anaphylactic reactions are usually characterised by pallor, tachycardia, diaphoresis. A 43-year-old woman with a history of asthma and ragweed allergy, had an anaphylactic shock following the use of a shampoo containing Pyrethrins for the treatment of head lice (HSDB, 2001). About 50% of persons sensitive to ragweed exhibit cross-sensitivity to Pyrethrum. (BRA, MGK and SCJ)

Specific Treatment in Case of an Accident or Poisoning: First Aid Measure, antidotes and Medical Treatment, if Known

There is no specific antidote for Pyrethrin poisoning. Treatment is symptomatic and supportive. Pulmonary sequelae are treated symptomatically with airway maintenance, oxygen by mask at 10 to 15L/min, and ventilatory assistance as dictated by patient status. Circulatory support may require intravenous fluids and rarely, pressor agents. Pharmacologic treatment of bronchospasm and anaphylaxis uses the standard drugs and management protocols. Bronchospasms are treated with inhaled β_2 -agonist and oral or parenteral corticosteroids. To assist eye irritation, prescription of proparacaine hydrochloride is recommended.

Vitamin E topical application (Vitamin E oil i.e., dl- α -tocopheryl acetate) is highly effective in relieving paraesthesia. Antihistamines are effective in controlling most allergic reactions. Severe asthmatic reactions, particularly in predisposed persons, may require administration of inhaled β_2 -agonist and/or systemic corticosteroids. Inhalation exposure should be carefully avoided in the future. Anaphylaxis-type reactions may require subcutaneous epinephrine and respiratory support. Contact dermatitis may be treated by applications of topical corticosteroid preparations. (BRA, MGK and SCJ)

Prognosis following poisoning

Two cases found in the literature were described. After the use of a pesticides containing Pyrethrum extract, a 41-year-old farmer developed an erythematous papular lesion of the hands. The farmer gave up the use of both pesticides, and two years later remained free of dermatitis (Garcia-Bravo *et al.*, 1995). Following the application of a flea-killer containing Pyrethrum extract, a young man presented symptoms of shortness of breath, a cough productive of white phlegm and nausea. After 4 hours of emergency department treatment and observation, the man had recovered (Paton & Walker, 1988).

When an allergic reaction was established, future contact with the substance should be avoided. (BRA, MGK and SCJ)

Table A.60 Summary table of further human data

No data are available.

A.2.16. Other data

No data are available.

A.3. Environmental effects assessment

A.3.1. Fate and distribution in the environment

The *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ contains Pyrethrins among other constituents, which may be divided into the two groups Pyrethrins I (consisting of pyrethrin 1, cinerin 1, and jasmolin 1) and Pyrethrins II (consisting of pyrethrin 2, cinerin 2 and jasmolin 2). Among these compounds, Pyrethrin 1 was selected as a surrogate for all the pyrethrins to generate the environmental fate data since it is representative of all other components, it is the component with the highest concentration (53%) and because it is difficult to evaluate the environmental fate properties of a mixture. The ecotoxicological studies were performed with the whole extract and the results based on nominal or measured concentrations of total pyrethrins (the six components in Pyrethrin I and II).

A.3.1.1. Degradation

A3.1.1.1 Abiotic degradation

Hydrolysis

The results presented in Table A.61 show that the half-lives for Pyrethrin 1 at pH 5 and pH 7 were 687 and 527 days respectively (Selim S., 1995), only the half-life of 17 days, for the pH 9 buffer system, was considered to be meaningful because of the greatest amount of degradation observed over the 30 day test period. No degradation products > 10% of the applied dose were observed in the pH 5 and pH 7 samples. Therefore, ¹⁴C-Pyrethrin 1 was found to be stable in buffers at pH 5 and 7, over the 30-day study period. A single major degradation product (>10% dose) was observed in the pH 9 sample; it was designated as degradate A. Degradate A was identified as ¹⁴C-Chrysanthemic acid, produced by hydrolysis of the ester linkage of ¹⁴C-Pyrethrin 1. Its concentration at pH 9 increased in proportion to the decrease of Pyrethrin 1, accounting for 61% of the applied radioactivity (AR) after 30 days. The degradation reactions were given by pseudo-first-order kinetics.

Another study was presented (Perboni A., 2015) to assess abiotic hydrolytic transformations of the 6 Pyrethrum extract components (Pyrethrin 1, Cinerin 1, Jasmolin 1, Pyrethrin 2, Cinerin 2 and Jasmolin 2) in an aquatic system at pH values normally found in the environment (pH 4-9) under sterile conditions in the absence of light, according to OECD 111. At Tier 1, all six Pyrethrum extract components showed significant hydrolysis (i.e. 10% hydrolysis was observed after 5 days at 50 °C), apart from Pyrethrin 2 and Cinerin 2, both at pH 4. However, in order to describe hydrolysis of all components in a similar manner and facilitate the comparability of results, Tier 2 was performed with all six components. In the higher tier test, the hydrolysis of the test substances was monitored at pH 4.0 pH 7.0 and pH 9.0 and at 3 different temperatures: 15°C, 25°C and 45°C in the dark for a period of 30 days. Hydrolysis of Pyrethrum extract components was shown to be mainly driven by the pH, but also dependent on the temperature regime:

- At pH 4 significant but slow hydrolysis was only observed for some components.
- At pH 7 relevant hydrolysis was determined for all components, but generally only at elevated temperatures.
- At pH 9 significant hydrolysis was shown for all six components at all temperatures, resulting in fastest hydrolysis at 45 °C.

The DT50 values for hydrolysis was obtained considering FOCUS kinetics guidance, and SFO (single first order) turned out to be the kinetic model describing best the decrease of concentrations.

For risk assessment purposes in freshwater, the half-life of 115 d at pH 7 and 25°C for Pyrethrin 1 should be used.

For classification and labelling according to the CLP Guidance, data on hydrolysis e.g. OECD Test Guideline 111 might be considered for classification purposes only when the longest half-life t_{1/2} determined within the pH range 4-9 is shorter than 16 days. Based on the

above results, hydrolysis data for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ cannot be considered for classification purposes, since the longest half-life determined within the pH range 4-9 is longer than 16 days.

Table A.61 Summary table - Hydrolysis

Summary table - Hydrolysis								
Method, Guideline, GLP status, Reliability, Key/supportive study	pH	Temp. [°C]	Initial concentration, CO[mol/l]	TS	Half-life, DT ₅₀ [d]	Coefficient of correlation, r ²	Remarks	Reference
US-EPA Pesticide Assessment Guidelines, subdivision N, Series 161-1 GLP Reliability 2	5 7 9	25	14C-pyrethrin 1 (Batch number CFQ.7422; Purity 98.1%)		687 527 17	0.192 0.331 0.947	-	Selim S. (1995) IIIA-7.1.1.1.1 (BRA, MGK and SCJ) Doc III A7.1.1.1.1 (KPIC)
	7	12	pH 5: 0.31 to 0.38 mg ¹⁴ C-pyrethrin 1/L pH 7: 0.35 to 0.38 mg ¹⁴ C-pyrethrin 1/L pH 9: 0.35 to 0.38 mg ¹⁴ C-pyrethrin 1/L		1476			
OECD Guideline 111 (Hydrolysis as a Function of pH) GLP Reliability 1 Key	4, 7, 9	15°C; 25°C and 45°C	Pyrethrin 1 Lot/Batch: XX-82-P1 Purity:99.4%		DT ₅₀ ranged from 0.4 to 211.6 days DT ₅₀ for Pyrethrin 1 at 25°C is 115 d at pH 7. At 25°C and pH 9 values for DT ₅₀ ranged from 4.2 to 14.9 days.	-	DT ₅₀ values were calculated for Pyrethrum extract components showing significant hydrolysis. At pH 4 significant hydrolysis was only observed for Cinerin 2 at 45°C, for Jasmolin 1 at all temperatures and for Jasmolin 2 at 25°C and 45°C. At pH 7 significant hydrolysis was observed for all components at 45°C. At 25°C Pyrethrin 1, Cinerin 1, Jasmolin 1 and Jasmolin 2	Perboni, A. (2015) IUCLID 10.1.1.1.a (BRA, MGK, SCJ and KPIC)

		Pyrethrin 2 Lot/Batch: XX-82-P2 Purity: 99.3% Cinerin 1 Lot/Batch: XX-82-C1 Purity: 97.5% Cinerin 2 Lot/Batch: XX-82-C2 Purity: 99.3% Jasmolin 1 Lot/Batch: XX-82-J1 Purity: 99.3% Jasmolin 2 Lot/Batch: XX-82-J2 Purity: 96.6% 0.02 mg/L for each analyte			exhibited significant hydrolysis. At 15°C only Jasmolin 1 was not hydrolytically stable. At pH 9 all components were significantly hydrolysed, whereas at 25°C and 45°C fastest hydrolysis was measured for all components.	
--	--	--	--	--	--	--

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ for Pyrethrin 1 at 25°C is 115 d at pH 7.
Justification for the value/conclusion	Hydrolysis is not considered a relevant degradation pathway for Pyrethrins under environmental relevant conditions.

Phototransformation in water

A photolysis test with *Trans*-[cyclopropane-1-¹⁴C] Pyrethrin 1 was performed according to the US EPA Pesticide Assessment Guidelines, Subdivision N, 161-2 (comparable to the test OCDE guideline n° 316 "Phototransformation of chemicals in water - direct photolysis"). To determine the extent and rate of aqueous photolysis of ¹⁴C-Pyrethrin 1, test samples, containing nominal concentrations of Pyrethrin 1 ranging from 0.32 to 0.38 mg/L in an HEPES buffer at pH 7, were exposed to natural sunlight and maintained at a mean temperature of 25 ± 1°C for 72 hours. Confirmation of the identity and amount of ¹⁴C-Pyrethrin 1 was performed by HPLC with radioisotope detection. The isolated degradate E was analysed by GC/MS. Samples radioactivity was determined by LSC.

Results showed that the photolysis rate of Pyrethrin 1 and the degradate E (isomer of Pyrethrin 1), as principal degradation product (concentration greater than 10% of the initial measured dose -IMD), is consistent with first order kinetic with a half-life of 11.8 hours of sunlight and a coefficient of correlation of 0.99.

Table A.62 Summary table – Photolysis in water

Summary table – Photolysis in water									
Method, Guideline, status, Reliability, Key/supportive study	GLP	Initial molar TS concentration	Total recovery of test substance [% of appl. AS]	Photolysis rate constant (kcp)	Direct photolysis sunlight rate constant (kpE)	Reaction quantum yield (φcE)	Half-life (t _{1/2E})	Remarks	Reference
US-EPA Pesticide Assessment Guidelines, Subdivision N, Series 161-2 GLP Reliability 1 Key		0.32 mg/L to 0.38 mg pyrethrin 1 /L	Radiolabelled pyrethrin 1, product code CFQ.7422, Purity 98.1% Non-radiolabelled pyrethrin 1, product code NK9212, Purity 97% Buffer solution (pH 7): mean overall recovery = 97.4% (range: 91.3 to 102.5%)	Not stated	0.059 h ⁻¹	1.2 x 10 ⁻³ in water at pH 7	11.8 h (¹⁴ C-Pyrethrin 1 +E-isomer)		Selim, S. (1995) and Werle, H. (1991) Doc III IIIA-7.1.1.1.2 (KPIC, BRA, MGK and SCJ)

Conclusion

There are two phases in photodegradation pattern of ¹⁴C-Pyrethrin 1. ¹⁴C-Pyrethrin 1 is rapidly equilibrates with its 'E' isomer when they are exposed to natural sunlight in an aqueous solution buffered to a pH 7, followed by photolysis of both compounds forming numerous minor photolytic products. Therefore, ¹⁴C -Pyrethrin 1 is rapidly photolysed when is exposed to natural sunlight in a pH 7-buffered solution, with a half-life of 11.8 hours.

Value used in Risk Assessment	
Value/conclusion	Photolytic half-life in water estimated to be 11.8 h.
Justification for the value/conclusion	Refer to page above.

Estimated photo-oxidation in air

Following the physical and chemical properties and the structure of Pyrethrins it is assumed that degradation and persistence of the active substance mainly depends on reaction with hydroxyl-radicals and the average concentration of hydroxyl-radicals in air. Total OH rate constant was determined to be $281.1508 \times 10^{-12} \text{ cm}^3/\text{molec.} \cdot \text{sec.}$, mainly due to addition to olefinic bonds (96%) and hydrogen abstraction (4%). Other mechanisms do not contribute to hydroxyl radical estimations. The total rate of both, OH and ozone constant is very low. Half-life in the troposphere was calculated to be 27.391 min for overall OH rate constant and 29.562 min for ozone rate constant. Following the Atkinson calculation, the chemical half-life for Pyrethrum in the troposphere will be below 1 h. It is therefore concluded that Pyrethrins will not accumulate in air and will only be transported on very short distances.

The photochemical oxidative degradation half-life of Pyrethrin 1 in air was calculated according to the method developed by Atkinson², which is based on the structural activity relationship (QSAR's), by using the Atmospheric Oxidation Program v 1.91 (AOPWIN-software). These estimations were carried out with respect to the OH radical and ozone reactions, using a 12-hours-day with 300.95×10^{-12} and $96.32 \times 10^{-17} \text{ cm}^3/\text{molecule} \cdot \text{sec}$, respectively. The half-lives for the hydroxyl and ozone reactions in air are estimated to be 25.59 and 17.13 minutes, respectively.

A half-life of 76.8 minutes for Pyrethrin 1 in air has been estimated using a 24-hour-day and assuming an OH radical concentration of $5 \times 10^5 \text{ radicals} \cdot \text{cm}^{-3}$ according to AOPWIN version 1.91 and following recommendations of ECHA Guidance on Risk Assessment, Chapter 2.3.6.3³.

Table A.63 Summary table – Photo-oxidation in air

Summary table – Photo-oxidation in air					
Model	Light protection (yes/no)	Estimated daily (24h) OH concentration [OH/cm ³]	Overall OH rate constant [cm ³ /molecule/sec]	Half-life [min]	Reference
No Guideline available		1.5×10^6	300.95×10^{-12}	25.59	O'Carroll, N. (2005) IIIA-7.3.1 (BRA, MGK and SCJ)

² Atkinson, R. (1988). Estimation of Gas-Phase Hydroxyl Radical Rate Constants for Organics Chemicals. Environ. Toxicol. Chem. 7:435-442.

³ Guidance on BPR: Vol IV Environment Parts B+C, Version 2.0 October 2017

Estimation method by AOPWIN version 1.91		5 x 10 ⁵		76.8	IIIA-7.3.1 (BRA, MGK and SCJ)
Calculation			281.1508 x 10 ⁻¹²	<1 hour	Oellrich (2000), Doc III A7.3.1 (KPIC)
No Guideline available		7 x 10 ¹¹ (ozone concentration)	96.32 x 10 ⁻¹⁷ (Ozone reaction rate constant)	17.13	IIIA-7.3.1 (BRA, MGK and SCJ)

Value used in Risk Assessment	
Value/conclusion	Photo-oxidation in air DT ₅₀ = 76.8 min
Justification for the value/conclusion	Considering all the available information, Pyrethrin 1 is not expected to volatilise in air in significant quantities. The small amounts that may volatilise are rapidly degraded via reaction with OH radicals and ozone. Based on the short half-life of Pyrethrin 1 in the atmosphere, accumulation and contamination by wet or dry deposition in the atmosphere are not to be expected. Therefore, air will not be an environmental compartment of concern for Pyrethrin 1 used as insecticide for household.

Photolysis on soil

¹⁴C-pyrethrin 1 was found to rapidly photolyse when exposed to sunlight on the surface of soil to form numerous degradates including CO₂. No single degradate represented more than 10% of the initially applied radioactivity. Within 24 h, a mean of less than 17% of the initial concentration of ¹⁴C-pyrethrin 1 remained. Non-exposed control samples had more than 55% of the initial concentration of ¹⁴C-pyrethrin 1 remaining at the end of the 24 h period. The photolysis is consistent with first order kinetics with a half-life of 12.9 h. The ¹⁴C-pyrethrin 1 in the non-exposed test system also degraded during the 24 h test period, but at a slower rate than the exposed samples. The half-life of the non-exposed samples was determined to be 82.9 h. The results indicate that ¹⁴C-pyrethrin 1 in the exposed samples was degraded via both photolytic (fast) and non-photolytic (slower) pathways.

The Spain CA has adjusted values of calculated DT₅₀ at 25°C to the equivalent at 12°C using equation 28 in the ECHA Guidance⁴. Thus, the DT₅₀ of Pyrethrin 1 under the average EU outdoor temperature was calculated in 36.12 hours.

Table A.64 Summary table – Photolysis on soil

Summary table – Photolysis on soil					
Guideline /Test method	Initial molar TS concentration [mg.kg-1]	Temperature [°C]	Total recovery of test substance [% of appl.a.s.]	Half-life (t _{1/2E}) [h]	Reference
USEPA FIFRA N-161-3, 40 CFR Sec. 158.130 GLP Reliability	¹⁴ C-pyrethrin 1 (Batch number CFQ.7422; Purity 95.9%)	24 ± 2°C	Mean overall recovery = 96.6%	12.9 hours	Testman R. (1994), Doc III A7.2.2.4 (KPIC) IIIA-7.2.2.4

⁴ Guidance on BPR: Vol IV Environment Parts B+C, Version 2.0 October 2017

1 Key	9.3 mg pyrethrin 1/L			(BRA, MGK and SCJ)
----------	-------------------------	--	--	--------------------------

A3.1.1.2 Biotic degradation

A3.1.1.2.1 Biodegradability (ready)

One biodegradation study according to OECD guideline 301B and the US EPA Method 835.3110 (CO₂ evolution test) is available for the Refined Pyrethrum Extract. The test substance was added to two vessels containing mineral salts medium inoculated with activated sludge at a nominal test concentration of 10 mg Carbon/L. Sodium benzoate was used as reference substance. An additional mixture containing Sodium benzoate and Refined Pyrethrum Extract was established in order to assess the potential of the test substance for microbial inhibition.

The results in Table A.65 show that mean cumulative CO₂ production by mixtures containing Refined Pyrethrum Extract accounted for 46% by the end of the test on day 29. The reference substance, Sodium benzoate, was degraded by 66% after 7 days and 93% after 29 days in the absence of Refined Pyrethrum Extract, and by 69% after 7 days in its presence, which confirmed that Refined Pyrethrum Extract was not inhibitory to the activity of the microbial inoculum. (BRA, MGK and SCJ)

Results of the standard test indicate that Pyrethrum extract is neither readily nor inherently biodegradable. However, since Pyrethrum extract shows a low solubility in water, the lack of microbial degradation could be a consequence of the limited availability of Pyrethrum extract for micro-organisms.

In addition another test was performed according to OECD 301 (Koopmans 1995). Pyrethrum Extract was added to two vessels containing mineral salts medium inoculated with activated sludge at a nominal test concentration of 12 mg TOC/L. The relative degradation values calculated from the measurements performed during the test period revealed no significant (>10%) degradation of Pyrethrum Extract (see Table below).

Table A.65 Summary table - biodegradation studies (ready/inherent)

Summary table - biodegradation studies (ready/inherent)											
Method, Guideline, GLP status, Reliability, Key/supportive study	Test type	Test parameter	Inoculum			Additional substrate	Test substance concentration	Degradation		Remarks [positive control]	Reference
			Type	Concentration	Adaptation			Incubation period	Degree [%]		
OECD 301 B GLP Reliability 2 Key	Ready Biodegradability	CO ₂ evolution test	Activated sludge	Refined Pyrethrum Extract (Batch number FEK-99; purity 57.03%) 10 mL sludge/L mineral medium	No	No	10 mg/L	29 days	46	-	Barnes, S. (2002) IIIA-7.1.1.2.1 (MGK, BRA and SCJ)
OECD 301 B GLP Reliability 2 Key	Ready Biodegradability	CO ₂ evolution test	Activated sludge	Pyrethrum Extract (Batch 94/10.7; Purity 25.14% total Pyrethrins) 10 mL sludge/L mineral medium	No	No	15 mg Pyrethrum Extract /L	28 days	4.6%	-	Koopmans (1995), Doc III A7.1.1.2.1 (KPIC)

As the pass level (carbon dioxide production equal to or greater than 60% of the theoretical value) within the 28 days incubation period in this test on ready biodegradability is not fulfilled by Refined Pyrethrum Extract, this cannot be classified as readily biodegradable.

Value used in Risk Assessment	
Value/conclusion	Not readily biodegradable
Justification for the value/conclusion	See above

A3.1.1.3 Rate and route of degradation including identification of metabolites and degradation products

A3.1.1.3.1 Biological sewage treatment

Aerobic biodegradation

No data are available.

Anaerobic biodegradation

No data are available.

STP simulation test

No data are available.

A3.1.1.3.2 Biodegradation in freshwater

Aerobic aquatic degradation

The objective of the study from Hein and Moendel (2017) was to provide information on the degradation rate (DT 50, DT 90) and metabolism of Pyrethrin 1 in natural water. Additionally, information on the identity and quantity of transformation products in water including a mass balance was determined. Test systems containing natural water were treated using [cyclopentenone-2-¹⁴C] Pyrethrin 1, at two different nominal concentrations 10 µg ("low dose") Pyrethrin 1/L and 100 µg ("high dose") Pyrethrin 1/L. In the test, calculated SFO DT50 values for pyrethrin 1 ranged from 6.7-10.7 days (at 20 ± 2°C). The main degradation product was pyrethrolone which reached a maximum of 9.5% AR after 21 days and then decreased to 2.8% AR at the last sampling interval. Several non-identified fractions were detected but these were minor and/or composed of several peaks. Mineralisation reached a maximum of 7% by the end of the study.

Table A.66 Summary table – freshwater aerobic biodegradation

Summary table – freshwater aerobic biodegradation							
Method, Guideline, GLP status, Reliability, Key/supportive study	Test type	Exposure	Test substance concentration	Incubation period	Degradation (DT ₅₀)	Remarks	Reference
OECD Guideline for Testing of Chemicals, No 309 "Aerobic Mineralisation in Surface Water", Apr. 13, 2004 GLP Reliable 1	OECD	darkness under aerobic conditions in the laboratory at 20 ± 2°C	[cyclopentenone-2- ¹⁴ C] pyrethrin 1, (Batch CFQ42752; Purity >97.7%) 10 µg/L (i.e. "low dose") and 100 µg/L (i.e. "high dose")	62 days	Calculated SFO DT ₅₀ values ranged from 6.7-10.7 days	The major degradation product was pyrethrolone which reached a maximum of 9.5% AR after 21 days and then decreased to 2.8% AR at the last sampling interval.	Hein W. <i>et al.</i> , 2017, IUCLID 10.1.3.2 Doc IIIA / Section 7.1.2.1.1.1

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ water 20°C (d) 10.7
Justification for the value/conclusion	Using SFO kinetics DT ₅₀ values ranging from 6.7-10.7 days were obtained (Hein <i>et al.</i> , 2017). The highest DT ₅₀ was used as a worst-case.

Water/sediment degradation test

Pyrethrin 1 degraded at a very rapid rate when applied to an aerobic aquatic environment (1 ppm dosing) (Robinson and Wisocky, 1994). Degradation proceeded initially by oxidation to form chrysanthemic acid and a number of low level degradants. Residues in water and sediment were initially extracted but extended degradation was accompanied by the formation of residues that were bound to sediment humus fractions and appeared to be partially comprised of bound chrysanthemic acid. Mineralisation was an observed but minor degradation pathway (4% CO₂ at day 30). The principal extractives were apart from pyrethrin 1, chrysanthemic acid, the major metabolite, and three additional minor degradates. A clear pattern of build-up and decline emerged for chrysanthemic acid, resulting in a maximum occurrence in the water/sediment system of 21.2% of applied radioactivity on day 21. The minor degradates were detected at various intervals, none exceeding 5% of the initial concentration of pyrethrin 1 at any point. The half-life of the pyrethrin 1 in the water/sediment system tested was calculated to be 10.5 days following pseudo-first-order kinetics at 25 °C. This is equivalent to 29.7 days at 12 °C. Later the applicant recalculated endpoints based on FOCUS Guidance as shown below.

In this test, however, only one sediment was used for determination of degradation rates whereas Guidance indicates that at least two sediments and their associated water should be used for calculations. In addition, the test finished before guidance indicates, at day 30, when still 14.7 % of the substance was present.

In a second study conducted by Witte (2007), pyrethrin 1 was observed to degrade rapidly when applied to two separate water/sediment systems taken from the natural environment. This occurred via a rapid movement from the water phase into the sediment phase combined with a steadily increasing mineralization to CO₂ (30 – 51% at test end) and breakdown in both aquatic and sediment phases to the metabolite chrysanthemic acid (maximum 65.6 and 66.8%). A clear pattern of build-up and decline emerged for Chrysanthemic acid in the water and sediment phases of both systems. A substantial portion of the applied radioactivity (30 – 40% at test end) became bound to sediment in both test systems. Following first order kinetics, the half-lives for Pyrethrin 1 and chrysanthemic acid in the whole water/sediment system ranged from 1.6 to 2.4 days and 18 to 109 days, respectively.

The anaerobic metabolism of ¹⁴C-Pyrethrin was studied under laboratory conditions in a water sediment model system at an initial concentration of 10 mg/L at 25°C (Robinson and Wisocky 1995). With the extraction method used, 46.19% of the AR remained unextractable from sediment after 1 year. Three main extractable degradation products were found: Cyclopropane diacid (a maximum of 14.6% at 364 days), ¹⁴C-Chrysanthemic acid (a maximum of 12.6% at 31 days) and a reduced form of Pyrethrin 1 named Jasmolin 1 (a maximum of 10.0% at 180 days). Sediment fulvic fractions-bound residues were also found to contain ¹⁴C-Chrysanthemic acid as a terminal residue. All other degradation products were found at concentrations below or equal to 4% of the AR. In this system volatiles were formed in low amounts of 13.92% of the AR at day 365 (4% of the AR at day 30), so mineralization process was low. Therefore, mechanisms involved in degradation of ¹⁴C-Pyrethrin appear to be a combination of reductive and oxidative process.

¹⁴C-Pyrethrin dissipated under anaerobic aquatic sediment conditions with a calculated half-life of 86 days for the whole system (240.8 days reflecting the average EU outdoor temperature of 12°C).

Table A.67 Summary table – fresh water/sediment degradation

Summary table – fresh water/sediment degradation								
Method, Guideline, GLP status, Reliability, Key/supportive study	Exposure	Test system		Test substance concentration	Incubation period	Degradation (DT ₅₀)	Remarks	Reference
		Water	Sediment					
US EPA Subdiv. N, § 162-4 Chemistry: Environmental Fate GLP Reliability 2 Key	25 ±1°C	Pond	Pond in Lucama, North Carolina, USA	¹⁴ C-pyrethrin 1 (Batch number CFQ 7390; Purity 98.7% - 99.7%) ca. 10 ppm	30 days	10.5 days	pseudo-first-order kinetics	Robinson, R. A.; Wisocky, M.J. (1994), Doc III A7.1.2.2.2 (KPIC) (MGK, BRA and SCJ)
OECD 308 (April 2002), SETAC 1995 GLP Reliability 2 Key	20 ±2°C	Pond	Ensing, district of Enz, Germany Sandy silt	56.0 µg ¹⁴ C-pyrethrin 1 (56.6 µg unlabelled pyrethrin 1)	-	1.6 days whole system	-	Witte A. (2007), Doc III A7.1.2.2.2/02 (KPIC)
	20 ±2°C	Creek	Spiegelberg, district of Rems-Murr, Germany Sand	[cyclopropane-1- ¹⁴ C] pyrethrin 1 (Lot No. CFQ14811 Batch 1); Purity Radiochemical 98.4%) Pyrethrum Pale Extract (Batch number 2006/3-3/Pale; Purity Pyrethrins I: 30.65% w/w Pyrethrins II: 19.49% w/w Total Pyrethrins: 50.14% w/w)	-	2.4 days whole system	-	

Summary table – fresh water/sediment degradation								
Method, Guideline, GLP status, Reliability, Key/supportive study	Exposure	Test system		Test substance concentration	Incubation period	Degradation (DT ₅₀)	Remarks	Reference
		Water	Sediment					
US EPA Pesticide Assessment Guidelines, Subdivision N, Series 162 – 3 GLP Reliability 2	Anaerobic at 25°C in the dark Anaerobic at 12°C in the dark	24.28% (supernatant)	Sandy loam 1.3% Organic carbon pH=4.7 16.28% (extractable residues) 46.19 (PES)	<i>Trans</i> -[cyclopropane-1- ¹⁴ C] Pyrethrin 1 (Batch NB8309; > 97.99% (radiochemical purity)) 10 mg/L	364	86.1	-	Robinson, R.A. and Wisocky, M.J. (1995) IIIA-7.1.2.2.2 (BRA, MGK and SCJ)
1 Test according to OECD criteria								

Conclusion

Investigations of degradation and metabolism behaviour of ¹⁴C-Pyrethrin 1 in water-sediment systems were conducted under aerobic and anaerobic conditions in compliance with US EPA Pesticide Assessment Guidelines, Subdivision N, Series 162 – 4 and 3, respectively (comparable to OECD method 308). Pyrethrins 1 was observed to degrade fast under aerobic conditions, with DT₅₀ value of 10.5 days. Pyrethrin 1 was observed to degrade more slowly under anaerobic conditions, with DT₅₀ value of 86 days. A common metabolite was identified in both systems Chrysanthemic acid which can be partially bioavailable for aquatic and sediment organisms. Mineralization of Pyrethrin 1 reached a maximum of 50% in the study by Witte 2007. Considering the anaerobic degradation there is an important percentage of bound residues (non-extractable residues) in the sediment.

According to the CLP Guidance a substance is not rapidly degradable unless it is demonstrated to be primarily degraded biotically or abiotically e.g. via hydrolysis, in the aquatic environment with a half-life <16 days (corresponding to a degradation of > 70 % within 28 days), and it can be demonstrated that the degradation products do not fulfill the criteria for classification as hazardous to the aquatic environment. *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ does not meet these requirements.

The DT₅₀ values of the study by Robinson and Wisocky (1994) were recalculated based on the FOCUS guidance.

Please find below details of the DT_{50 system} values for pyrethrin 1 used for PEC calculations for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Water / sediment system	pH water phase	pH sed	t °C	DT50/DT90 system (d)	St. (r ²)	DT50/DT90 water (d)	St. (r ²)	DT50/DT90 sed (d)	St. (r ²)	Method of calculation
Sandy loam*	6.8	4.7	25 20	7.93/26.4 11.83/39.38	χ ² =7.3	n.c.	-	n.c.	-	SFO
Sandy silt**	8.34	7.2	20	5.26/17.55	0.947	1.3/4.3	0.986	n.c.	-	SFO
Sand**	8.24	7.4	20	2.36/7.85	0.870	0.7/2.3	0.867	n.c.	-	SFO
Geometric mean (DT _{50system} 20°C):				5.27						
DT50 12°C				11.2						

* = Robinson and Wisocky (1994)

** = Witte (2007)

n.c. = not calculated

SFO = Single First Order kinetic

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ for biodegradation in water/sediment 11.2 d (at 12°C) (Pyrethrin 1)
Justification for the value/conclusion	5.27 d at 20°C

A4.1.1.3.3 Biodegradation in seawater

Seawater degradation study

No data available.

Seawater/sediment degradation study

No data available.

A4.1.1.3.4 Higher tier degradation studies in water or sediment

No data available.

A4.1.1.3.5 Biodegradation during manure storage

No data available.

A4.1.1.3.6 Biotic degradation in soil

No data available.

A4.1.1.3.7 Laboratory soil degradation studies

Aerobic biodegradation

Pyrethrin 1 degraded in soil under aerobic conditions following treatment of a sandy loam soil at approximately 1 ppm (Robinson, 1994). Degradation occurred with the rapid loss of extractable soil residues, the formation of CO₂ (43.41% by day 181) and soil bound residues. Total pyrethrin 1 content in organic extracts declined from an average level of 85.42% of the applied radioactivity on day 0 to less than 1% by day 59 after application. Degradation proceeded initially by a combination of hydrolysis and oxidation to form a number of low level metabolites. Residues in soil were initially extracted, but extended degradation was accompanied by the formation of residues that were bound to soil humus fractions and ultimately mineralized, 43.41% CO₂ at the end of the test. In addition to the evolution to CO₂, four main metabolites (A to D) were found along with four metabolite regions where no single metabolite could be defined. None of the metabolites identified reached a concentration higher than 5%.

The objective of the study by Hein W. (2017) was to determine the degradation route and rate of [Cyclopentenone-2- ¹⁴C]Pyrethrin 1 under aerobic conditions in soil at 20 °C. The concentration of Pyrethrin 1 and possible metabolites was determined throughout the test duration only in soil (loam), including the formation of volatile products such as carbon dioxide generated by mineralization. Material balances were established at each sampling interval which showed the distribution of Pyrethrin 1 and any formed metabolites as a function of time. In the test, Pyrethrin 1 decreased continuously during the course of the study. Pyrethrin 1 was degraded mainly by mineralisation (51.07 % (mean) of the AR at day 120) and formation of bound residues (33.51 % (mean) of the AR at day 120). No other significant metabolites (>5 %) were detected.

For comparative purposes, additional studies (Fifi 2015 a, b and c) were conducted to determine the rate of degradation of the six components of Pyrethrum extract (Pyrethrin 1, Cinerin 1, Jasmolin 1, Pyrethrin 2, Cinerin 2 and Jasmolin 2) in 3 soils under aerobic conditions at 20°C in the dark. All six pyrethrin esters exhibit very low persistence with DT50 values ranging from 0.4 (pyrethrin 2) to 10.7 days (jasmolin 1), calculated according to FOCUS 2006 as far as possible. Based on these data it is concluded that pyrethrin 1 is representative of the pyrethrins.

The soil metabolites analysed were not found in relevant amounts throughout the incubation period. Soil concentrations of Chrysanthemic acid, Pyrethric acid, Pyrethrolone,

Cinrolone and Jasmolone were always below the LOQ. In addition Pyrethric acid, Cinrolone and Jasmolone did not exceed the LOD.

The half-life of pyrethrin 1 was calculated to be 3.3 days (pseudo-first-order kinetics). Four main metabolites (A to D) were designated. Metabolite D was identified as chrysanthemic acid. None of these metabolites amounted at any time to more than 10% of the initial pyrethrin 1 concentration.

Table A.68 Summary table – aerobic biodegradation in soil- laboratory study

Method, Guideline, GLP status, Reliability, Key/supportive study	Exposure	Test system				Test substance concentration	Incubation period	Degradation DT ₅₀	Remarks	Reference
		Soil origin	Soil type	pH	OC%					
US EPA Pesticide Assessment Guidelines, Subdivision N, Series 162 – 1 GLP Reliability 2 Key	Aerobic at 25 °C in the dark, 12 °C, 10 °C	Grand Forks County, North Dakota, USA	Sandy loam	6.4	2.2%	1 mg/kg (¹⁴ C-pyrethrin 1 (Batch number CFQ 7390; Purity 96.08% - 97.14% (radiochemical purity))	181 days	1.88 d at 25°C 2.47 d at 20°C 4.68 d at 12°C	The DT _{50soil} of 1.88 d (study value at 25°C, recalculated using SFO) normalized to 12°C is: 5.32 d. The DT _{50soil} of 2.47 d (study value of 1.88 d normalized to pF2 and 20°C) normalized to 12°C is: 4.68 d.	Robinson (1994), Doc III A7.2.1 (KPIC) IIIA-7.2.2 (BRA, MGK and SCJ)
OECD: Guideline 307; Aerobic and Anaerobic Transformation in Soil, April 24, 2002 GLP Reliability 1 Key	Aerobic at 25 °C in the dark, 19.4°C ± 0.7°C and a soil moisture content of about pF 2.5	Mußbach, Germany	Loam	7.28	1.73	0.1 mg/kg dwt [cyclopentenone-2- ¹⁴ C] pyrethrin 1, Batch CFQ42752; Radiochemical purity: >99%	120 days	DT ₅₀ value was 4.05 days using first order kinetics. The DT ₅₀ value was recalculated to 2.96 days at 20 °C and pF 2.	Pyrethrin 1 was degraded mainly by mineralisation (51.07% (mean) of the AR at day 120) and formation of bound residues (33.51% (mean) of the AR at day 120). No other significant metabolite (>5%) was	Hein W. (2017), IUCLID 10.2.1 Doc IIIA / Section 7.2.1

Method, Guideline, GLP status, Reliability, Key/supportive study	Exposure	Test system				Test substance concentration	Incubation period	Degradation DT ₅₀	Remarks	Reference
		Soil origin	Soil type	pH	OC%					
									detected.	
OECD: Guideline 307; Aerobic and Anaerobic Transformation in Soil, April 24, 2002 GLP Reliability 1 Key	aerobic conditions at 20°C and 45% MWHC in darkness	Loam Soil (batch F2.40815), Lufa-Speyer, Lufa code 2.4	Loam	7.30	2.21	0.1 mg/kg dwt pyrethrin extract containing six pyrethrin esters (pyrethrin 1, cinerin 1, jasmolin 1, pyrethrin 2, cinerin 2 and jasmolin 2)	120 days	DT ₅₀ values 2.1 and 9.0 days. The pseudo SFO DT ₅₀ value for pyrethrin 1 normalised to pF 2 was 4.69 days.	-	Fifi (2015a), IUCLID 10.2.1 Doc IIIA / Section 7.2.1
OECD: Guideline 307; Aerobic and Anaerobic Transformation in Soil, April 24, 2002 GLP Reliability 1 Key	aerobic conditions at 20°C and 45% MWHC in darkness	Loamy Sand Soil (batch F2.20815), Lufa-Speyer, Lufa code 2.2	Loamy Sand Soil	5.25	1.56	0.1 mg/kg dwt pyrethrin extract containing six pyrethrin esters (pyrethrin 1, cinerin 1, jasmolin 1, pyrethrin 2, cinerin 2 and jasmolin 2)	120 days	DT ₅₀ values 0.4 days and 5.4 days. The pseudo SFO DT ₅₀ value for pyrethrin 1 normalised pF2 was 3.9 days.	-	Fifi (2015b), IUCLID 10.2.1 Doc IIIA / Section 7.2.1
OECD: Guideline 307; Aerobic and Anaerobic Transformation in Soil, April 24, 2002 GLP Reliability 1 Key	aerobic conditions at 20°C and 45% MWHC in darkness	Sandy loam soil (batch F5M0815), Lufa-Speyer, Lufa code 5M	Sandy loam soil	7.40	1.00	0.1 mg/kg dwt pyrethrin extract containing six pyrethrin esters (pyrethrin 1, cinerin 1, jasmolin 1, pyrethrin 2, cinerin 2 and jasmolin 2)	120 days	DT ₅₀ values 0.5 days and 5.3 days. The pseudo SFO DT ₅₀ value for pyrethrin 1 normalised to 20°C and pF 2	-	Fifi (2015c), IUCLID 10.2.1 Doc IIIA / Section 7.2.1

Method, Guideline, GLP status, Reliability, Key/supportive study	Exposure	Test system				Test substance concentration	Incubation period	Degradation DT ₅₀	Remarks	Reference
		Soil origin	Soil type	pH	OC%					
								was 3.0 days.		

In summary, the rate of aerobic degradation of pyrethrin 1 was investigated in 5 soils at 20-25°C and various moisture contents (45% MWHC, 75% of pF 2.5, pF 2.5) in darkness. Degradation was rapid and results, including normalised values, are shown in the table below.

Table A.69: Laboratory kinetic and statistical analysis of pyrethrin 1

Study	System	Kinetic model	pH	Mo	Parameter (K, K ₁ , k ₂ , g, t _b , α, β)	χ ² , %-error	DT ₅₀ [days]	DT ₉₀ [days]	SFO DT ₅₀ 20°C, pF ₂
Robinson, 1994	North Dakota (sandy loam)	SFO	6.4	85.17	k = 0.3687	12.9	1.9	6.3	2.5
Hein, 2017	Mußbach (loam)	SFO	7.28	90.2	k = 0.171	6.89	4.1	13.5	2.96
Fifi, 2015a	Loam	FOMC	7.30	-	n/s	6.16	6.6*	21.9	4.69
Fifi, 2015b	Loamy Sand	FOMC	5.25	-	n/s	13.37	3.9*	12.8	3.9
Fifi, 2015c	Sandy loam	SFO	7.40	-	n/s	5.7	3.0	9.9	3.0
Geomean									3.3

* as SFO

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ for biodegradation in soil 3.3 d (at 20°C) (Pyrethrin 1)
Justification for the value/conclusion	Refer to the table above.

Anaerobic biodegradation

No data available.

A4.1.1.3.8 Higher tier degradation studies in soil

Field dissipation studies (field studies, two soil types)

Pyrethrum (measured as Pyrethrins 1) applied to bare ground soil in a single application of Pyrenone Crop Spray at the maximum labelled seasonal rate (0.52 kg/ha) exhibited a half-life of approximately 1-2 h. No residues of Pyrethrin 1 above the 0.1 ppm limit of quantitation were detectable in any soil core samples by 1 day after application and beyond. Storage stability data indicated that Pyrethrins 1 residues from all samples should have been stable until analysis. Transit stability data indicated stability of Pyrethrins I in frozen soil. Results from the field trial suggest that residues of Pyrethrum dissipated completely over a 2 day period.

Table A.70 Summary table – Field dissipation

Summary table – Field dissipation										
Method, Guideline, GLP status, Reliability, Key/supportive study	Site	Application rate (g AS/ha)	Surface	Soil type	Soil texture	Test duration	Degradation DT ₅₀ ^a	Degradation DT ₉₀ ^a	Remarks	Reference
USEPA Subdivision N, Section 164-1 GLP Reliability 2 Key	California/USA	0.52 kg a.i./ha	Bare ground soil	Sandy loam	Sandy loam	179 days	0.04	0.15	k1 (1/day) = 15.475	Hattermann D.R. (1992), Doc III A 7.2.2.2/01&02 (KPIC)
	Georgia/USA	Pyrenone Crop Spray (Batch number M60008		Sandy loam	Sandy loam		0.04	0.12	k1 (1/day) = 19.202	
	Michigan/USA	; contains 6% Pyrethrum and the synergist Piperonyl Butoxide. Pyrethrum again is composed of Pyrethrins I (pyrethrin 1, cinerin 1, jasmolin 1) and Pyrethrins II (pyrethrin 2, cinerin 2, jasmolin 2))		Sandy loam	Sandy loam		0.08	0.26	k1 (1/day) = 8.988	
^a degradation values presented as single first order values										

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ for biodegradation in soil 3.3 d (at 20°C) (Pyrethrin 1)
Justification for the value/conclusion	Refer to Section A4.1.1.3.7

A4.1.1.3.9 Short summary and overall relevance of the provided information on degradation and conclusion on rapid degradation

Hydrolysis

The half-lives for Pyrethrin 1 at pH 5 and pH 7 were 687 and 527 days respectively. At pH 9 the substance half life was 17 days. Temperature of the test was 25°C. No degradation products > 10% of the applied dose were observed in the pH 5 and pH 7 samples. A single major degradation product, identified as Chrysanthemic acid (>10% dose) was observed in the pH 9 sample. Its concentration at pH 9 increased in proportion to the decrease of Pyrethrin 1, accounting for 61% of the applied radioactivity (AR) after 30 days. The degradation reactions were given by pseudo-first-order kinetics.

In another study (Perboni A., 2015) abiotic hydrolytic transformations of the 6 Pyrethrum extract components (Pyrethrin 1, Cinerin 1, Jasmolin 1, Pyrethrin 2, Cinerin 2 and Jasmolin 2) in an aquatic system at pH values normally found in the environment (pH 4-9) under sterile conditions in the absence of light, according to OECD 111. In the higher tier test, the hydrolysis of the test substances was monitored at pH 4.0 pH 7.0 and pH 9.0 and at 3 different temperatures: 15°C, 25°C and 45°C in the dark for a period of 30 days. The half-life of 115 d at pH 7 and 25°C for Pyrethrin 1 was determined.

Biodegradability (ready)

Results of the standard tests with Pyrethrum Extract indicate that the active substance is not readily degradable. However, since Pyrethrum extract shows a low solubility in water, the lack of microbial degradation could be a consequence of the limited availability of Pyrethrum extract for micro-organisms.

Water and Water/sediment degradation test

In water, calculated SFO DT₅₀ values for pyrethrin 1 ranged from 6.7-10.7 days (at 20 ± 2°C). The main degradation product was pyrethrolone which reached a maximum of 9.5% AR after 21 days and then decreased to 2.8% AR at the last sampling interval. Several non-identified fractions were detected but these were minor and/or composed of several peaks. Mineralisation reached a maximum of 7% by the end of the study.

In aerobic water/sediment systems Pyrethrins 1 was observed to degrade under aerobic conditions, with DT₅₀ values ranging from 1.6 to 10.5 days. A substantial portion of the applied radioactivity became bound to sediment in the tests. A common metabolite was identified in both systems: Chrysanthemic acid, which can be partially bioavailable for aquatic and sediment organisms. Mineralization of Pyrethrin 1 reached a maximum of 50% under aerobic conditions.

Aerobic biodegradation in soil

In soil, the half-life of pyrethrin 1 was calculated to be 3.3 days at 20°C (pseudo-first-order kinetics, geomean of 5 studies). Four main metabolites (A to D) were designated. Metabolite D was identified as chrysanthemic acid. None of these metabolites amounted at any time to more than 10% of the initial pyrethrin 1 concentration. Maximum mineralization reached was 51.03 %.

Conclusion

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is a complex substance of natural origin. According to the Guidance on the Application of the CLP criteria (version 5, July 2017) a complex substance, such as UVCBs, should be regarded as not rapidly degradable if the constituents that are not rapidly degradable constitute a significant part of the substance, e.g. more than 20%, or for a hazardous constituent an even lower content.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ fate is represented by Pyrethrin 1. Hydrolysis data for this component yields half-lives > 16 days across different pHs at 25°C. According to the Guidance on the Application of the CLP criteria (version 5, July 2017), data on hydrolysis might be considered for classification purposes only when the longest half-life determined within the pH range 4-9 is shorter than 16 days. Thus, hydrolysis cannot be considered for classification purposes in this case.

In water tests Pyrethrin 1 was not ultimately degraded, not meeting the guideline requirement of a half-life <16 days (corresponding to a degradation of > 70 % within 28 days).

In water/sediment the substance primarily degraded with DT50 values ranging from 1.62 to 10.5 days at tests temperature and transformed into metabolites hazardous to the aquatic environment or non-identified metabolites. A part of the substance also bound to sediment.

Further, ready biodegradation available for Pyrethrum Extract showed that the substance is not ready biodegradable.

Based on the above *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is considered not rapidly degradable.

A.3.1.2. Distribution

A4.1.2.1 Adsorption onto/desorption from soils

The binding potential of radiolabelled pyrethrin 1 was studied in four soils using the batch equilibration method (Reynolds & Robinson, 1994). A 1:100 soil-to-solution ratio was used, and the equilibration time for both adsorption and desorption was three hours. At a rate of 0.75 ppm, 70.86%, 75.22%, 77.30% and 49.43% of total radioactivity adsorbed to solids of sandy loam, silty clay loam, silt loam and sand, respectively. The major radioactive product was the parent compound (pyrethrin 1). Low levels of the chrysanthemic acid, a hydrolysis product, were also detected in organic extracts from the adsorption phase, but none exceeded 5.16% of the applied radioactivity. The adsorption and desorption constants obtained in the study indicated that parent chemical was immobile in all of the examined soils and was not readily desorbed from the soil matrices. The mobility potential of pyrethrin 1 in the examined soils is classified as immobile. (KPIC)

The results from the study conducted by Mori, V. (2015) confirm that the Koc for Pyrethrin 1 is 34674 L/Kg and this value will be used in the environmental risk assessment.

Table A.71 Summary table – Adsorption/desorption

Summary table – Adsorption/desorption										
Method, GLP Reliability	Guideline, status,	Soil	Adsorbed AS [%]	K _a	K _{aoc}	K _d K _{doc} K _a /K _d	K _f	1/n	Remarks	Reference
USEPA Subdivision N, Section 163-1 GLP Reliability 2	N	Sandy loam, North Dakota	70.86	268	12472	2332 108679 0.1149	N/A	N/A	N/A	Reynolds & Robinson (1994), Doc III A 7.1.3/01 (KPIC) IIIA-7.2.3.1 (BRA, MGK and SCJ)
		Silty clay loam, Mahaska	75.22	310	16190	1151 60133 0.2693	N/A	N/A	N/A	
		Silt loam, Dundee	77.30	430	74175	2600 448257 0.1654	N/A	N/A	N/A	
		Sand, Wakulla	49.43	198	37847	965 184767 0.2052	N/A	N/A	N/A	
OECD Guideline 121 (Estimation of the Adsorption Coefficient (K _{oc}) on Soil and on Sewage Sludge using High Performance Liquid Chromatography (HPLC)) GLP Reliability 1 Key		N/A	N/A	N/A	Pyrethrin 1=34674 Cinerin 1=21136 Jasmolin 1=50119 Pyrethrin 2=7762 Cinerin 2=4732 Jasmolin 2=10471	N/A	N/A	N/A	Mori, V. (2015), IUCLID 10.1.2 Doc IIIA Section A7.2.3.1	

K_a = Adsorption coefficient

K_{aoc} = Adsorption coefficient based on organic carbon content

K_d = Desorption coefficient

K_{doc} = Desorption coefficient based on organic carbon content

K_a / K_d = Adsorption / Desorption distribution coefficient

Table A.72 Summary table – Adsorption/desorption metabolite/ degradant/ transformation- or reaction product*
No data available

Value used in Risk Assessment	
Value/conclusion	Organic carbon/water partition coefficient (Koc)= 34674 L/Kg Log Koc = 4.54
Justification for the value/conclusion	The fate and distribution in the environment was derived from studies on pyrethrin 1, since pyrethrin 1 represents the predominant analogue and a typical member (or paradigm) for the pyrethrum family. Pyrethrin 1 is considered to represent the worst case, as it has a higher Koc value than Pyrethrins 2 and, as the most sensitive compartment is sediment, the adopted approach to exposure assessment is consequently very conservative.

A4.1.2.2 Higher tier soil adsorption studies

No data available.

A4.1.2.3 Volatilisation

Regarding volatilisation, please see Part A, section 1.3 Physical and chemical properties of the active substance.

A.3.1.3. Bioaccumulation

Measured aquatic bioconcentration

A bioconcentration test with bluegill sunfish (*Lepomis macrochirus*) exposed to the initial measured concentration of 90.5 ng/L ¹⁴C-Pyrethrin 1 under flow-through system was performed. After an uptake phase of 28 days the fish were transferred to clean water for 15 days (deuration phase). After 28 days the bioconcentration factors were determined to be 127, 873 and 471 for edible tissue, non-edible tissue and whole body, respectively. All BCF values refer to the total amount of radioactivity (sum of radiolabelled parent, metabolites and mineralization products). Pyrethrin 1 and two major metabolites (Chrysanthemic acid and another metabolite which was not identified) were separated in the analysis of edible and non-edible (viscera) tissues.

In addition to the BCF steady state the applicant derived a BCF kinetics which resulted in a BCF = 500. This would be the preferred value for risk assessment as indicated in Guidance R7c and also because in the last consecutive three measure points, the test substance concentration in fish varies more than 20%.

The accumulation of Pyrethrin 1 was reversible as approximately half of the [¹⁴C] residues from both edible and non-edible tissues had deparated within 1 day of the initiation of the deuration period (half-life was determined to be 1 day). Therefore, the deuration rate was quite fast. (BRA, MGK and SCJ)

Table A.73 Summary table – Measured aquatic bioconcentration

Summary table – Measured aquatic bioconcentration										
Method, Guideline, GLP status, Reliability, Key/supportive study	Exposure	Log Kow of AS	Initial concentration of AS	Steady state BCF	Uptake rate constant (K1)	Depur. rate constant (K2)	Depur. time (DT ₅₀)	Metabolites	Remarks	Reference
OECD 305 US EPA 165-4 GLP Reliability 2 Key	Flow through 28 d	Pyrethrin 1: log Kow = 5.34 (20°C) Pyrethrin 2: log Kow = 3.79 (20°C)	90.5 [14C] Pyrethrin 1 and Pyrethrin 1 (Batch number Radiolabelled: CFQ6932, CFQ7390, CFQ7422; non-radiolabelled: NK9304) (Purity Radiolabelled: 97.0%, 99.6%, 98.1%; non-radiolabelled: 98.9%)	Edible 127X Non-edible 873X Whole Body 479X	Edible 183/1.39 Nonedible 1270/1.36 Whole Body 662/1.32	Edible 1.39 Nonedible 1.36 Whole Body 1.32	1 d	Chrysanthemic acid: %TRR ^a Hexane fraction: 5.4 Methanol fraction: 24.1%TRR ^b Hexane fraction: 1.2 Methanol fraction: 31.7 Metabolite #3: %TRR ^b Hexane fraction: 11.6 Methanol fraction: 4.7	-	█ (1994) A7.4.3.3.1/01 (KPIC); IIIA-7.4.2. (BRA, MGK and SCJ) and KPIC

Measured logPow

The partition coefficient n-Octanol-Water log Pow of Pyrethrum Extract components (Pyrethrin 1, Cinerin 1, Jasmolin 1, Pyrethrin 2, Jasmolin 2 and Cinerin 2) was determined by HPLC according to OECD117.

The following logpow values were obtained:

- Pyrethrin 1: 5.59
- Cinerin 1: 5.54
- Jasmolin 1: 6.04
- Pyrethrin 2: 4.32
- Cinerin 2: 4.26
- Jasmolin 2: 4.74

Pyrethrin 1, which was considered the reference component for fate data, has a log kow of 5.59 above the cut-off level for bioaccumulation of the CLP Guidance equal to 4

Conclusion

For bioaccumulation assessment BCF measured values are preferred over logkow values. A BCF value of 500 was calculated, indicating the potential for bioaccumulation of the substance. However, the reported BCF values refer to the total amount of radioactivity (sum of radiolabelled parent, metabolites and mineralization products) and may not reflect the real BCF value of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Further, a reliable logpow = 5.59 was provided which is above the cut-off value = 4 of the Guidance to determine if a substance is bioaccumulative or not. Based on this value the substance is considered potentially bioaccumulative.

A.3.1.4. Monitoring data

No data are available.

A.3.2. Effects on environmental organisms

A.3.2.1. Aquatic compartment

A3.2.1.1 Freshwater compartment

Acute/short-term toxicity (freshwater)

FISH

Pyrethrum extract (FEK-99) was tested on Rainbow trout (*Oncorhynchus mykiss*) in a flow-through system during 96 hours at five different nominal concentrations from 1.3 to 10 µg total Pyrethrins/L. Two replicates of ten fish per test concentration were exposed to Pyrethrum extract. This test (██████████, 1994a) was carried out according to the U.S. EPA Pesticide Assessment Guidelines E, Section 72-1 which is equivalent to the OECD guideline 203 "Fish, Acute Toxicity Test". The test conditions were within the range demanded by OECD 203. Analytical monitoring showed that the measured concentrations were not ≥ 80% of the nominal concentration, thus the mean measured concentrations of total Pyrethrins were used. The 96 hour LC50 was calculated to be 5.2 µg total Pyrethrins/L (with 95% confidence intervals of 3.1 to 5.7 µg total Pyrethrins/L), based on mean measured concentrations.

Pyrethrum extract (FEK-99) was tested on Bluegill Sunfish (*Lepomis Macrochirus*) system during 96 hours at five different nominal concentrations from 3.2 to 25 µg total

Pyrethrins/L (mean measured from 3.1 to 14) . Two replicates of ten fish per test concentration were exposed to Pyrethrum extract. This test (██████, 1994b) was carried out according to the U.S. EPA Pesticide Assessment Guidelines E, Section 72-1 which is equivalent to the OECD guideline 203 "Fish, Acute Toxicity Test". From the data collected during the exposition, the 96-hours LC50 was determined to be 10 µg total Pyrethrins/L (95% C.I. of 7.8 to 14 µg/L) based on the mean measured concentrations of total Pyrethrins.

Pyrethrum extract (FEK-99) was tested on Sheepshead minnow (*Cyprinodon variegatus*) in a flow-through system during 96 hours at five nominal concentrations from 10 to 78 µg total Pyrethrins/L. Two replicates of ten fish per test concentration were exposed to Pyrethrum extract. This test (██████, 1994c) was carried out according to the U.S. EPA Pesticide Assessment Guidelines E, Section 72-3. From the data collected during this study, the 96-hours LC50 was determined to be 16 µg total Pyrethrins/L (95% C.I. of 14 to 18 µg/L) based on the mean measured concentrations of total Pyrethrins. The abnormal/behavioural effects, mortality, surfacing, loss of equilibrium, fish on the bottom of the test chamber, laboured respiration, dark discoloration, vertical orientation and/or quiescence, were observed in the 18 µg/L test concentration during the study.

INVERTEBRATES

The acute toxicity of Pyrethrins (Pyrethrum extract (FEK-99) to aquatic invertebrates was tested (Putt A.E., 1994a) in *Daphnia magna* with five test concentrations (mean measured concentrations from 2.2 to 14 µg/L). The test was carried out according to the U.S. EPA Pesticide Assessment Guidelines E, Section 72-2 which is equivalent to the OECD guideline 202 "Daphnia sp., Acute Immobilisation Test". The test conditions were within the range demanded by OECD 202. The 48 h EC50 for Pyrethrins was determined to be 12 µg total Pyrethrins/L (10 – 13 µg total Pyrethrins/L). Analytical monitoring of the test substance showed that the measured concentrations were not ≥ 80% of the nominal concentration, thus the mean measured concentrations of total Pyrethrins were used.

In addition the applicants presented data on the toxicity of the substance metabolites (Mantilaci 2015a) and on the toxicity of Pyrethrum Extract (FEK-99) and the 6 Pyrethrin esters (Mantilacci, S. 2015b) to *Daphnia magna* following OECD Guideline 202 "Daphnia sp., Acute Immobilisation Test and Reproduction Test":

- In the first study, young daphnids (less than 24 hours) are exposed to five pyrethrin metabolites (Pyrethrolone, Cinerolone, Jasmolone, Pyrethric acid, Chrysanthemic acid) at a range of concentrations for a period of 48 hours under static conditions. The test fulfils OECD 202 validity criteria. However, the range of concentrations selected is not enough to reliably estimate an EC50. The highest inhibition, only 40%, at the highest concentration tested, 371.86 µg/L, occurred for Chrysanthemic acid. Thus, in most cases, no EC50 was determined in the study and the values were extrapolated outside the range of the tested concentrations. Nevertheless, the test shows that the substance metabolites are less toxic than parent and that monitoring the metabolites during other studies is not necessary. No other effects were observed on the exposed organisms.

- In the second study, the acute immobilisation tests were performed, under semi-static conditions, to assess the effects of Pyrethrum Extract (FEK-99) and the 6 Pyrethrin esters (Pyrethrin 1, Cinerin 1, Jasmolin 1, Pyrethrin 2, Cinerin 2, Jasmolin 2) on *Daphnia magna*. 20 daphnids less than 24 hours old, were exposed to determinate concentrations of each test item for 48 hours. Probit analysis was used to evaluate the dose-response function and the endpoints with 95% confidence limits.. The test fulfilled validity criteria. However, test concentrations do not allow to reliably estimating an EC50 for Cinerin 1 and 2, Jasmolin 1 and 2 and Pyrethrin 2. In the test, applied concentrations for Cinerin 1, Pyrethrin 2 and Cinerin 2 only reach a maximum of 45%, whereas for Jasmolin 1 and Jasmolin 2y, 55% inhibitions is reached. For FEK-99 the EC50 = 28.09 µg/L and for Pyrethrin 1 EC50 = 272.81 µg/L.

Despite their deficiencies the above two test showed that metabolites and pyrethrins esters

are less toxic than parent.

ALGAE

Unicellular fresh water green alga, *Desmodesmus subspicatus* was exposed under static conditions for 72 hours to six concentrations of Pyrethrum Extract Pale 50 %/L with three replicates of each concentration (Dengler D., 2000). Six replicates of a blank control were run in parallel. Test substance concentrations tested were presented in mg Pyrethrum Extract Pale 50 %/L (from 5.3 to 100). The mean value of the cell concentration was plotted versus time to produce growth curves for Pyrethrum Extract Pale 50 %/L and for the control. This resulted in a NOErC = 30.9 mg/L, an ErC50 = 65.1 mg/L and a EbC50 = 29.0 mg/L (based on nominal concentrations).

However, in the test, due to the physical-chemical properties of Pyrethrum Extract: low water solubility, tendency to bind on surfaces and volatilisation with steam; and because the application procedure was performed by application of the product in acetone to the vessels, volatilisation of the acetone, adding of test medium to the vessels and shaking on flat bed shaker, most of Pyrethrum was bound to the vessel surfaces. The total pyrethrum in the test system was above 74 to 91% of nominal at the beginning and decrease to 61 to 73% at the end of the test. Since the test substance disappears in the test RMS recalculated endpoints based on measured concentrations considering 61% of nominal as a worst case (concentrations were only measured for three concentrations groups and this value maximises concentration loss). This resulted in an EC50 = 39.8 mg/L and EC10 = 19.7mg/L. Transforming this value to total pyrethrins an EC50 = 19.mg/L and and EC10 = 9.85mg/L was obtained.

These values are higher than water solubility and the endpoints were considered as water solubility 0.23mg/L.

Table A.77 Summary table – acute/short-term aquatic toxicity

Summary table – acute/short-term aquatic toxicity										
Method, Guideline, GLP status, Reliability, Key/supportive study	Species	Endpoint / Type of test	Test material	Exposure		Results			Remarks	Reference
				Design	Duration	NOEC	LC/ EC ₁₀	LC/ EC ₅₀		
Fish										
EPA, Subdivision E, Series 72, § 72-1 GLP Reliability 2 Key	Rainbow trout (<i>Oncorhynchus mykiss</i>)	Mortality/ acute	Pyrethrum extract (FEK-99) (Batch R92-254; Purity 57.5%)	Flow-through	96 hours	3.1 µg total pyrethrins/L		5.2 µg total pyrethrins /L	5 concentrations tested, deaths in highest dose group	██████████ (1994a) A7.4.1.1/01 (BRA, MGK, SCJ) (KPIC)
EPA, Subdivision E, Series 72, § 72-1 GLP Reliability 2	Bluegill sunfish (<i>Lepomis macrochirus</i>)	Mortality/ acute	Pyrethrum extract (FEK-99) (Batch R92-254; Purity 57.5%)	Flow-through	96 hours	5.4 µg total pyrethrins /L		10 µg total pyrethrins /L	5 concentrations tested, deaths in the two highest dose groups	██████████ (1994b) A7.4.1.1/02 (KPIC)
EPA, Subdivision E, Series 72, § 72-3 GLP Reliability 2	Sheepshead minnow (<i>Cyprinodon variegatus</i>)	Mortality/ acute	Pyrethrum extract (FEK-99) (Batch R92-254; Purity 57.5%)	Flow-through	96 hours	7.4 µg total pyrethrins /L		16 µg total pyrethrins /L	5 concentrations tested, deaths in the four highest dose	██████████ (1994c) A7.4.1.1 / 03 (KPIC)

									groups	
Invertebrates										
U.S. EPA Pesticide Assessment Guidelines, Subdivision E, Section 72-2 GLP Reliability 2	<i>Daphnia magna</i>	Immobilization/ acute toxicity	Pyrethrum extract (FEK-99) (Lot. R92-254; 57.48% w/w total Pyrethrins)	Flow through	48h	3.7 µg total pyrethrins /L		12 µg total pyrethrins /L	5 concentrations tested, immobilization in the two highest dose groups Measured concentration < 80% of nominal	Putt, A.E. (1994a) IIIA-7.4.1.2 (BRA, MGK, SCJ, KPIC)
OECD Guideline No. 202 GLP Reliability 2	<i>Daphnia magna</i>	Immobilization/ acute toxicity	Pyrethrolone (98.0 mg/L); Cinerolone (99.9 mg/L); Jasmolone (99.5 mg/L); Pyrethric acid (100.0 mg/L) & Chrysanthemic acid (100.0 mg/L)	Static	48h	Pyrethrolone: ≥393.97 µg/L Cinerolone: ≥367.42 µg/L Jasmolone: ≥398.41 µg/L Pyrethric acid: ≥469.15 µg/L Chrysanthemic acid: 34.92 µg/L	Pyrethrolone: LOEC >393.97 µg/L EC ₁₀ = 189.78 µg/L Cinerolone: LOEC >367.42 µg/L EC ₁₀ = 277.80 µg/L Jasmolone: n.d Pyrethric acid: n.d	Pyrethrolone: n.d Cinerolone: n.d Jasmolone: n.d Pyrethric acid: n.d Chrysanthemic acid: 609.95 µg/L	The test fulfills validity criteria. However test concentrations do not allow a reliable estimation of the EC ₅₀ .	Mantilacci, S. (2015a) Doc IIIA / Section A7.4.1.2 (BRA, MGK, SCJ, KPIC)

							LOEC>469.15 µg/L EC ₁₀ = 87.34 µg/L Chrysanthe mic acid: LOEC = 76.83 µg/L EC ₁₀ = 85.84 µg/L			
OECD Guideline No. 202 GLP Reliability 2	<i>Daphnia magna</i>	Immobility/ acute toxicity	Pyrethrum Extract (FEK-99) (57.03% w/w); Pyrethrin 1 (29.76%); Cinerin 1 (5.55%); Jasmolin 1 (1.81%); Pyrethrin 2 (15.63%); Cinerin 2 (3.18%); Jasmolin 2 (1.10%)	semi-static	48h	Pyrethrum Extract (FEK-99): NOEC: 1.94 µg/L; total Pyrethrins NOEC: 0.89 µg/L Pyrethrin 1: NOEC: 14.09 µg/L Cinerin 1: NOEC: 43.39 µg/L Jasmolin 1: NOEC: 23.48 µg/L Pyrethrin 2: NOEC: 10.67 µg/L Cinerin 2: NOEC: 23.48 µg/L Jasmolin 2: NOEC: 51.65 µg/L	Pyrethrum Extract (FEK-99): LOEC: 4.27 µg/L; total Pyrethrins LOEC: 1.96 µg/L Pyrethrin 1: LOEC: 30.99 µg/L Cinerin 1: LOEC: 95.45 µg/L Jasmolin 1: LOEC: 51.65 µg/L Pyrethrin 2: LOEC: 23.48 µg/L Cinerin 2: LOEC: 51.65 µg/L Jasmolin 2: LOEC: 113.64 µg/L	Pyrethrum Extract (FEK-99) EC ₅₀ 28.09 µg/L; total Pyrethrins EC ₅₀ 12.92 µg/L Pyrethrin 1: EC ₅₀ 61.08 µg/L Cinerin 1: EC ₅₀ 272.81 µg/L Jasmolin 1: EC ₅₀ 226.67 µg/L Pyrethrin 2: EC ₅₀ 262.79 µg/L Cinerin 2: EC ₅₀ 358.95 µg/L Jasmolin 2: EC ₅₀ 216.16 µg/L	The test fulfils validity criteria. However, test concentrations only allow to reliably estimate the EC ₅₀ for FEK-99 and Pyrethrin 1.	Mantilacci, S. (2015b) Doc IIIA / Section A7.4.1.2
Algae (growth inhibition) ¹							NOErC	EbC50	E_rC₅₀	

OECD 201 EEC Directive C.3 (92/69/EC) GLP Reliability 1	<i>Desmodemus subspicatus</i>	Growth and biomass inhibition	Pyrethrum extract Pyrethrum Extract Pale 50% (Batch 99/11-5 B; Purity 50.17%)	Static	72 hours	0.23 mg/L	0.23 mg/L	0.23 mg/L	6 concentrations tested, significant inhibitory effects from 30.9 - 100 mg Pyrethrum Pale extract/L for biomass and growth rate	Dengler D. (2000) A7.4.1.3/01 (KPIC)
Other aquatic plants										
-	-	-	-	-	-	-	-	-	-	-
¹ calculated from growth rate, if not available please include the biomass value (NOEbC/EbCx) or the unspecified NOEC/ECx value										

Value used in Risk Assessment	
Value/conclusion	<i>Oncorhynchus mykiss</i> EC50 = 5.20 µg total pyrethrins /L = 7.97 µg a.s./L considering the whole extract as the a.s.
Justification for the value/conclusion	Based on acute values, fish (<i>Oncorhynchus mykiss</i>) is the most sensitive species tested. The resulting lowest EC50 is 5.20 µg total Pyrethrins/L.

There are some additional tests which were evaluated under PPP regulation for aquatic organisms. They are included in Appendix VII.

Chronic/long-term toxicity (freshwater)

FISH

A flow-through sub-acute toxicity test (██████████, 1994d) was performed with Fathead minnow (*Pimephales promelas*) during an early life stage exposure. The test was carried out according to the U.S. EPA Pesticide Assessment Guidelines E, Section 72-4 which is equivalent to the OECD guideline 210 "Fish, Early-life Stage Toxicity Test". The test conditions were within the range demanded by OECD 210. Analytical monitoring showed that the measured concentrations were not $\geq 80\%$ of the nominal concentration, thus the mean measured concentrations of total Pyrethrins were used. The NOEC and LOEC were determined to be 1.9 µg total Pyrethrins/l and 3.0 µg total Pyrethrins/l, respectively, based on the effects observed for percent embryo hatch and larval growth (total length and wet weight).

INVERTEBRATES

Toxicity test about effects on reproduction and growth rate was performed with *Daphnia magna* under flow-through conditions (Putt A.E. 1994b). The test was carried out according to the U.S. EPA Pesticide Assessment Guidelines E, Section 72-4 which is equivalent to the OECD guideline 211 "Daphnia magna reproduction test". The test conditions were within the range demanded by OECD 211. Six nominal concentrations were tested from 0 to 4.0 µg total Pyrethrins/L (nominal concentrations of 0, 0.25, 0.50, 1.0, 2.0 and 4.0 µg total Pyrethrins/L). The mean measured concentrations of total Pyrethrins were 0.20, 0.41, 0.86, 2.0 and 3.6 µg/L. The biological results are based on the mean measured concentrations during the test.

The following effects of Pyretrum Extract on daphnids was assessed:

- Immobility of adult daphnids
- Number of living young daphnids
- Number of dead young daphnids
- Appearance and behaviour of adult daphnids at test termination (Day 21)
- Growth (total body length and dry weight)

Survival:

Survival of adult daphnids was determined on test days 0, 1, 2, 4, 7, 9, 11, 14, 16, 18 and 21. Following 11 days of exposure, the mean percent survival of organisms exposed to all treatment levels was within the range of 80 – 100%. Control survival averaged 95%.

Reproduction:

Measurements of offspring production was made on days 0, 1, 2, 4, 7 and three times per week thereafter through study termination (day 21). At each observation interval, the offspring were removed, counted and discarded.

Mean reproduction of 26, 18, 27, 27 and 8 offspring per female was observed at these

respective concentrations. Control reproduction averaged 29 offspring per female. An adverse effect on reproduction was noted in the study, and was considered the most sensitive indicator of the toxicity of pyrethrum extract to *Daphnia magna*. Daphnids exposed to 0.2, 0.41, 0.86, 2.0, and 3.6 µg total Pyrethrins/L released an averaged of 197, 198, 211, 150 and 24 offspring per female, respectively. Statistical analysis determined a significant reduction in reproduction at the 2.0 and 3.6 µg total Pyrethrins/L concentrations when compared to the reproduction of pooled control daphnids (207 offspring per female). No significant reduction in reproduction was observed at ≤ 0.86 µg total Pyrethrins/L when compared to the pooled control.

Immobilization:

Throughout the exposure period, no young were observed to be immobilized in any test concentration or control.

Body length and weight:

At test termination, the length and weight of each surviving adult daphnia were measured. Mean total body length of daphnids exposed to 0.2, 0.41, 0.86, 2.0 µg total Pyrethrins/L ranged from 5.32 to 5.46 mm and was statistically reduced as compared to the mean total length of the pool control organisms (5.4 mm). The mean total body length among daphnids exposed to the highest treatment level tested (3.6 µg total Pyrethrins/L) was 4.85 mm and was statistically different from the pooled control. Similarly, mean organism dry weight for daphnids exposed to the four lowest concentrations ranged from 1.47 to 1.52 mg and was statistically comparable to mean weight of the pooled control organisms (1.62 mg). The mean dry weight of daphnids exposed to the highest concentration tested, 3.6 µg total Pyrethrins/L, was 1.09 mg and was statistically reduced as compared to the dry weight of the pooled control organisms. Therefore, the NOEC for growth was determined to be 2.0 µg total Pyrethrins/L.

Comment:

Since there was no statistically significant difference between the survival, reproduction and growth of the organisms in the control and solvent control, the data of survival, reproduction and growth from both control groups were pooled. The pooled control data was used during the comparisons to establish treatment level effects.

Conclusion:

Based on the data of the test, it can be concluded that reproduction was the most sensitive indicator observed during the study.

The EC₅₀ was determined to be 3.6 µg total Pyrethrins/l. The LOEC was determined to be 2.0 µg total Pyrethrins/l. The NOEC was determined to be 0.86 µg total Pyrethrins/L, which is the key value.

Table A.78 Summary table – chronic/long-term aquatic toxicity

Summary table – chronic/long-term aquatic toxicity								
Method, Guideline, GLP status, Reliability, Key/supportive study	Species	Endpoint/ Type of test	Test material	Exposure		Results LOEC/NOEC/EC ₁₀ [specify the value]	Remarks	Reference
				Design	Duration			
Fish								
US EPA 72-4 GLP Reliability 2	Fathead minnow (<i>Pimephales promelas</i>)	Embryo hatch, survival and growth of larvae	Pyrethrum extract (FEK-99) (Batch R92-254; Purity 57.5%)	Flow-through	35 days	NOEC 1.9 µg total Pyrethrins/L LOEC 3.0 µg total Pyrethrins/L	5 concentrations tested, deaths in all dose groups	█ (1994d) A7.4.3.2 (KPIC and BRA, MGK and SCJ)
Invertebrates								
US EPA 72-4/ GLP Reliability 2 Key	<i>Daphnia magna</i>	Reproduction /chronic	Pyrethrum extract (FEK-99) (Batch R92-254; Purity 57.5%)	Flow-through	21 days	NOEC 0.86 µg total Pyrethrins/L LOEC 2.0 µg total Pyrethrins/L EC ₅₀ (EC _x) 3.6 µg total Pyrethrins/L	5 concentrations tested, effects observed in the 2 highest concentrations	Putt A.E. (1994b) A7.4.3.4 (KPIC and BRA, MGK and SCJ)
Algae ¹								
OECD 201 EEC Directive C.3 (92/69/EC) GLP Reliability 2	<i>Desmodesmus subspicatus</i>	Growth and biomass inhibition	Pyrethrum extract Pyrethrum Extract Pale 50% (Batch 99/11-5 B; Purity 50.17%)	Static	72 h	NOEC 0.23 mg/L (solubility limit)	6 concentrations tested, significant inhibitory effects from 30.9 - 100 mg Pyrethrum Pale xtract/L for biomass and growth rate	Dengler D. (2000) A7.4.1.3 (KPIC)
other aquatic plants								
-	-	-	-	-	-	-	-	-
1 calculated from growth rate, if not available please include the biomass value (NOEbC/EbCx) or the unspecified NOEC/ECx value								

Value used in Risk Assessment	
Value/conclusion	Daphnia magna NOEC (21 days) = 0.86 µg total pyrethrins/L = 1.32 µg a.s./L, considering the whole extract as the a.s.
Justification for the value/conclusion	Daphnia is the most sensitive species tested. The chronic toxicity to Daphnia magna was determined in a 21-day reproduction study. The resulting NOEC, based on numbers of offspring per adult, was determined to be 0.86 µg total Pyrethrins/L.

A3.2.1.2 Sediment compartment

Acute/short-term toxicity (freshwater sediment)

Three acute immobilisation tests with *Chironomus riparius* were performed for the chemical similarity report, with pyrethrum extract 49.35%, pyroicide 50% and pyrethrum extract pale 50% (Dabrunz, A., 2017 a, b and c respectively). They all follow OECD guidance 235 "Chironomus sp., Acute Immobilisation Test". In all the three tests, *Chironomus riparius* larvae (20 in 4 replicates at each concentration) were exposed in 100 mL glass vessels to eight different nominal test concentrations from 0.241 to 60.0 µg total pyrethrins/L for 48 hours under static test conditions. Statistical evaluation was performed with a probit analysis using linear max. likelihood regression to obtain EC10, 20 and 50. The NOEC was established based on the highest test item concentration at which immobilisation was not higher than the acceptable control immobilisation (15 % immobilisation). The endpoints are summarized in table A.79, based on geomean measured concentration. This table shows the lowest endpoint in Dabrunz 2017a study.

In this specific test, observations on immobilization of the *Chironomus riparius* were made after 24 and 48 hours. The immobilised *Chironomus riparius* were counted and abnormal behaviour was noted at test start and every 24 hours thereafter. Water temperature, pH and dissolved oxygen were recorded throughout the exposure period. *Chironomus riparius* were not fed during the test period. Analytical determinations for total Pyrethrins concentration were made from samples taken from each replicate of each test item group at the start and end of the study. Mortality data as absolute numbers of immobile daphnids and as percent of exposed animals is shown below:

Test-Substance Concentration Total Pyrethrins (nominal) [µg/l]	Immobile <i>Chironomus riparius</i> (mean)			
	Number		Percentage	
	24 h	48 h	24 h	48 h
Control	0	0	0	0
Solvent control	0	0	0	0
0.241	1	1	5	5
0.529	1	1	5	5
1.16	4	4	20	20
2.56	6	7	30	35
5.63	7	10	35	50
12.4	7	19	35	95
27.3	13	20	65	100
60.0	17	20	85	100

An EC50 = 3.11 µg total pyrethrins/L based on measured concentration was obtained. Table A.79 Summary table – acute/short-term toxicity to sediment dwelling organisms

Summary table – acute/short-term toxicity to sediment dwelling organisms										
Method, Guideline, GLP status, Reliability, Key/supportive study	Species	Endpoint/ Type of test	Test material	Exposure		Results			Remarks	Reference
				Design	Duration	NOEC	LC/EC ₁₀	LC/EC ₅₀		
OECD 235; GLP GLP Reliability 2 Key	<i>Chironomus riparius</i>	Acute immobilisation	Pyrethrum Extract (49.35%)	Static	48 h	0.497 µg total pyrethrins/L	1.04 µg total pyrethrins/L	3.11 µg total pyrethrins/L	Measured concentrations	Dabrunz, A. (2017a) (BRA) Doc IIIA / Section A7.4.3.5.1
OECD 235; GLP GLP Reliability 2	<i>Chironomus riparius</i>	Acute immobilisation	Pyrocide® 50%	Static	48 h	2.18 µg total pyrethrins/L	4.79 µg total pyrethrins/L	5.25 µg total pyrethrins/L	Measured concentrations	Dabrunz, A. (2017b) (MGK) Doc IIIA / Section A7.4.3.5.1
OECD 235; GLP GLP Reliability 2	<i>Chironomus riparius</i>	Acute immobilisation	Pyrethrum Extract Pale 50%	Static	48 h	3.72 µg total pyrethrins/L	10.0 µg total pyrethrins/L	9.96 µg total pyrethrins/L	Measured concentrations	Dabrunz, A. (2017c) (KPIC) Doc IIIA / Section A7.4.3.5.1

Value used for classification purposes	
Value/conclusion	<i>Chironomus riparius</i> EC50 = 3.11 µg total pyrethrins /L = 4.76 µg a.s./L
Justification for the value/conclusion	Based on acute values, this is the most sensitive species tested for water/sediment system.

Chronic/long-term toxicity (freshwater sediment)

The effects of sediment-incorporated test substance on the survival and growth of *Chironomus riparius* was determined under static test conditions during a 28-day exposure period (Thomas, S.T. and Krueger H.O., 2009) in accordance with the OECD test Guideline 218. Larvae of *Chironomus riparius* were exposed in a static system to nominal concentrations of 31, 63, 125, 250, 500 and 1000 µg.kg⁻¹ of refined Pyrethrum extract FEK 99 (57.03 % w/w total Pyrethrins) in a water-sediment system (spiked sediment) at 20±2°C. Initial LC50, NOEC y LOEC values submitted by the applicant were based on nominal concentrations. However, analytical monitoring of the test substance, only measured at the lowest and highest test concentration (31 µg Refined Pyrethrum Extract/kg sed. dw and 1000 µg Refined Pyrethrum Extract/kg sed. dw) showed a significant decrease in the test substance concentration over the exposure period. For example, at highest test concentration after 7 days the 11.6% and after 28 days the 0.51% of nominal concentration was found in the sediment. For this reason RMS recalculated endpoints based on measured concentrations resulting in an EC50 >54.6 µg/kg, a LOEC 56.46 µg/kg and a NOEC 28.22 µg/kg.

The concentrations of Pyrethrins were not maintained over the test period, deviating from nominal concentration more than 20 per cent, so the results should be based on measured concentrations (mean recovery of 9.9%).

However, the test presents deviations that make the study not valid, but to be used as supporting information:

- A significant difference occurred only at the highest concentration in the case of the emergence ratio (49% emergence whereas in control it is 74%). It is very likely that this significant difference can be attributed to biological variability. The emergence ratios for concentrations > 1000 µg/kg might not be significantly different to the control. No LC50 could be determined either, due to lack of mortality at the highest test rate supporting the interpretation that the significant effect at the highest rate for the emergence ratio is an outlier. Furthermore there are no confidence intervals given for the emergence ratio.
- The variability of the mean emergence ratios between the different test concentrations indicates that no clear dose-response relationship can be shown. In some cases the lower test rates have even lower or equal emergence ratios compared with the higher test rates. Dunnett's test which may not be the appropriate test due to the high variability of the data (there was a statistically significant difference (p<0.05) from the negative control using Dunnett's test).
- The stock solutions appeared slightly cloudy and white with a foamy surface which might be related to residues of detergents used to clean the vessels. The cloudiness increased with increasing concentration. The appearance of the stock solutions might have had an impact on the emergence ratio of the test organisms (being rather a suspension, an homogeneous distribution of the test substance in the sediment is not guaranteed).

In another test, *Chironomus riparius* midge larvae were exposed under static conditions for 28 days to eight concentrations of Pyrethrum Pale Extract (Heintze, 2001) added to the water phase of the system, following OECD test guideline 219. The results were EC 50 for emergence inhibition of the midge, the emergence rate dependent and the development rate dependent NOECs. Nevertheless, this test is not considered for the risk assessment, just as supporting information, as it presents important deficiencies: at the end of the test – after 28 days – the % of total pyrethrins detected in the overlaying water was ~1 % of initial amount applied, showing a transfer of pyrethrins from water to

sediment phase and a subsequent loss from test system by either degradation or volatilization.

In the only reliable test, *Chironomus riparius* midge larvae were exposed under static conditions for 28 days to eight concentrations of Pyrethrum Pale Extract (Gonsior, 2009). Four replicates holding 25 midge larvae each were run. During the period of expected emergence (normally starting at day 10 and lasting until day 25) a daily check of emerged midges was performed. The sex and number of emerging adults were recorded daily. Only the number of fully emerged male and female midges was counted. Test substance concentrations were 1.2, 2.39, 4.78, 9.56, 19.1 and 38.2 mg Pyrethrum Pale Extract 50%/kg sediment dw, and were added to the sediment phase of the system, according to OECD guideline 218. Analytical verification of pyrethrins concentrations in the main test was performed at 1.2 mg Pyrethrum Extract Pale/kg sed. dw, 38.2 mg Pyrethrum Extract Pale/kg sed. dw and solvent control at distinct sampling dates (0, 7 and 28 days after introduction of larvae).

According to the applicant, the EC50 for emergence inhibition of the midge *C. riparius* was estimated to be 6.47 mg Pyrethrum Extract Pale/kg sediment dw (3.24 mg total pyrethrins/kg sed. dw, nominal concentration). The emergence rate dependent NOEC was 4.78 mg Pyrethrum Extract Pale/kg sed. dw equivalent to 2.39 mg totals pyrethrins/kg sed. dw (nominal concentration). The development rate dependent NOEC was also 4.78 mg Pyrethrum Extract Pale/kg equivalent to 2.39 mg total pyrethrins/kg sed. dw (nominal concentration).

eCA recalculated endpoints based on measured concentrations. To consider the decline in test substance concentration, the geometric mean of the measured concentrations for the time 0, day 7 and day 28 for the nominal concentration 0.166 mg Refined Pyrethrum Extract/per vessel calculated. This results in a mean measured concentration of 0.116 mg Refined Pyrethrum Extract/kg sed. Dw vessel of pyrethrum, corresponding to a recovery of 69.88 %. Applying this % recovery to the nominal EC50, NOEC and LOEC, the values of measured endpoints were estimated:

Effect data	28 days [mg Pyrethrum Extract Pale 50 % mg/kg dw nominal]	28 days [mg total pyrethrins mg/kg sediment dw nominal]	measured concentration total pyrethrins mg/kg sediment dw
EC50	6.47	3.24	2.26
NOEC	4.78	2,39	1.67*
LOEC	9.56	4.78	1.29

* The geomean measured concentration results to be 0.116 mg/vessel of pyrethrum, which corresponds to a recovery of 69.88%. Applying this recovery to the nominal NOEC of 4.780 mg/kg dwt of pyrethrum, results in a concentration of 1.67 mg/kg dwt of total pyrethrins.

Table A.80 Summary table – chronic/long-term toxicity to sediment dwelling organisms*

Summary table – chronic toxicity to sediment dwelling organisms									
Method, Guideline, GLP status, Reliability, Key/supportive study	Species	Endpoint/ Type of test	Test material	Exposure		Results LOEC/NOEC/ EC ₁₀ [specify the value]	Remarks	Reference	
				Design	Duration				
OECD 218 GLP Reliability 2 Key	<i>Chironomus riparius</i>	Developmental time/emergence ratio of midges	Pyrethrum Extract Pale 50% (Batch 2008/7-2; Purity 50.08%)	Static	28 days	NOEC = 4.78 mg/kg dw NOEC=1670 µg a.i./kg dwt total pyrethrins Normalised to 10% Organic carbon content: NOEC= 8350 µg a.i./kg dwt total pyrethrins (from 2%OC) Corrected to wet weight sediment: NOEC= 1815 µg a.i./kg wwt total pyrethrins LOEC = 9.56 [mg/kg dw]; EC ₅₀ = 6.47 [mg/kg dw]	6 concentrations tested; toxic effects observed in the three highest dose concentrations	Gonsior (2009) A7.4.3.5.1/02 (KPIC)	

OECD 218 GLP Reliability 3 Supporting	<i>Chironomus riparius</i>	Mortality, growth and behavioural signs	Refined Pyrethrum extract (FEK 99; 57.03% w/w total Pyrethrins)	Static system	28 days	EC ₅₀ >56.46 µg/kg LOEC 56.46 µg/kg NOEC 28.22 µg/kg	(nominal conc. recalculated to mean measured conc.)	Thomas, S.T. & Krueger, H.O. (2009) IIIA-7.4.3.5.1 (BRA, MGK and SCJ)
OECD 219/ GLP Reliability 3 Supporting	<i>Chironomus riparius</i>	Developmental time/emergence ratio of midges	Pyrethrum Extract Pale 50% (Batch 99/11-5 B; Purity 50.17%)	Static	28 days	NOEC=0.009 6 mg/L/ LOEC=0.018 6 mg/L/ EC ₅₀ =0.0515 [mg/L]	8 concentrations tested, toxic effects observed in the six highest dose concentrations	Heintze (2001) A7.4.3.5.1 (KPIC)

A3.2.1.3 Marine compartment

Acute/short-term toxicity (seawater)

No data available.

Chronic/long-term toxicity (seawater)

No data available.

A3.2.1.4 Sea sediment compartment

Acute/short-term toxicity (sea sediment)

No data available.

Chronic/long-term toxicity (sea sediment)

No data available.

A3.2.1.5 Higher tier studies on aquatic organisms

No data available.

A.3.3. Overall summary of acute and chronic aquatic toxicity data and Comparison with the CLP criteria

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is a complex substance of natural origin. Its main active components are Pyrethrin I: Pyrethrin 1 (min 418.9 g/kg), Cinerin 1 (min 46.0 g/kg), Jasmolin 1. (min 28.8 g/kg); Pyrethrin II: Pyrethrin 2 (min 285.5 g/kg), Cinerin 2 (min 41g/kg) and Jasmolin 2 (min 21.8g/kg). As a main component of the mixture, Pyrethrin 1 has been considered as a surrogate for many of the fate data. Further the substance contains other plant material (max 88.1 g/kg), BHT (max 69.4 g/kg), water (max 2.7 g/kg) and a solvent when commercialised. These other components are not relevant from an ecotoxicological point of view. The substance is stable without the solvent. Hence, here, a classification is provided with and without the solvent and worst case outcome is proposed for classification.

A.3.3.1. Short-term (acute) aquatic hazard

The hazard categories for acute aquatic toxicity and their related criteria are set out in Regulation (EC) No 1272/2008 on classification, labelling and packaging (CLP) of substances and mixtures, Annex I, Section 4.1.

In this case, Chrysantemun extract should be considered as a UVCB (plant extract), and hence, be classified based on test data of a mixture as a whole. Please see Appendix VI for further information on the active substance as a UVCB.

According to Annex I: 4.1.3.3.1 of EC 1272/2008, when the mixture as a whole has been tested to determine its aquatic toxicity, this information can be used for classifying the mixture.

There are adequate toxicity data for the a.s. (UVCB) for fish, invertebrates and algae. In addition, there is data for Chironomus based on OECD 235 which is a test done in water-only vessels and hence valid for classification. The use of Chironomus riparius values is further justified by the insecticidal mode of action of the substance. According to Table 4.1.0.(a):

- Being *Chironomus riparius* the most vulnerable species with a LC₅₀ = 0.00311 mg total pyrethrins/l which is equivalent to 0.0037 mg Chrysantemun extract CO₂ without solvent (pyrethrins are at a concentration of 82.39% in the composition of the plant extract considered as the mixture), in table 4.1.0 (a), if LC₅₀ < 1 mg/l, then category acute 1 applies.
- In Annex I, table 4.1.3, if 0.001 < LC₅₀ < 0.01, then a multiplying factor M = 100 applies.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is classified as Aquatic

Acute 1; H400, M=100.

A.3.3.2. Chronic/ long-term aquatic hazard (including information on bioaccumulation and degradation)

Long-term toxicity:

- There are adequate chronic toxicity data for the three trophic levels for the mixture as a whole (plant extract). In addition, there is also chronic data for *Chironomus riparius*, the most sensitive acute species.
- It can be concluded that the most sensitive aquatic organisms, on the basis of their chronic toxicity endpoints, is *Daphnia magna* (lowest NOEC = 0.00086 mg/l).

Bioaccumulation:

According to the CLP Guideline the criteria for determining if a substance is potentially bioaccumulative is the following:

Valid/high quality experimentally determined BCF value → YES:

→ BCF ≥ 500: *The substance meets the criterion*

→ BCF < 500: *The substance does not meet the criterion*

Valid/high quality experimentally determined BCF value → NO:

→ Valid/high quality experimentally determined log K_{ow} value → YES:

→ log K_{ow} ≥ 4: *The substance meets the criterion*

→ log K_{ow} < 4: *The substance does not meet the criterion*

Reliable BCF measured values are preferred over logk_{ow} values. A BCF value of 500 was calculated, indicating the potential for bioaccumulation of the substance. However, the reported BCF values refer to the total amount of radioactivity (sum of radiolabelled parent, metabolites and mineralization products) and may not reflect the real BCF value of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Further a reliable logp_{ow} = 5.59 was provided which is above the cut-off value = 4. Based on this value the substance is considered potentially bioaccumulative.

Degradation:

According to the guidance on the application of CLP criteria (2017), the substance is considered to be non-rapidly degradable unless at least one of the following is fulfilled:

- a. The substance is demonstrated to be readily biodegradable in a 28-day test for ready biodegradability. The pass level of the test (70 % DOC removal or 60 % theoretical oxygen demand) must be achieved within 10 days from the onset of biodegradation; or
- b. The substance is demonstrated to be ultimately degraded in a surface water simulation test with a half-life of <16 days (corresponding to a degradation of >70 % within 28 days); or
- c. The substance is demonstrated to be primarily degraded biotically or abiotically e.g. via hydrolysis, in the aquatic environment with a half-life <16 days (corresponding to a degradation of >70% within 28 days), and it can be demonstrated that the degradation products do not fulfill the criteria for classification as hazardous to the aquatic environment

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is a complex substance of natural origin. According to the Guidance on the Application of the CLP criteria (version 5, July 2017) a complex substance, such as UVCBs, should be regarded as not rapidly degradable if the constituents that are not rapidly degradable constitute a significant part of the substance, e.g. more than 20%, or for a hazardous constituent an even lower content.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ contains 43.9% of Pyrethrin 1. Hydrolysis data for this component yields half-lives > 16 days across different

pHs at 25°C. According to the Guidance on the Application of the CLP criteria (version 5, July 2017), data on hydrolysis might be considered for classification purposes only when the longest half-life determined within the pH range 4-9 is shorter than 16 days. Thus, hydrolysis cannot be considered for classification purposes in this case.

In water simulation tests Pyrethrin 1 was not ultimately degraded, not meeting the guideline requirement of with a half-life <<16 days (corresponding to a degradation of > 70 % within 28 days).

In water/sediment the substance primary degraded with DT50 values ranging from 1.6 to of 10.5 days and transformed into metabolites hazardous to the aquatic environment or in non-identified metabolites. Hence, it cannot be demonstrated that the degradation products do not fulfill the criteria for classification as hazardous to the aquatic environment

Further, results of the standard test indicate that Pyrethrum Extract is not readily biodegradable as CO₂ production accounted for 46% by the end of the test on day 29 (section A4.1.1.2.1 Biodegradability (ready/inherent)).

Based on the above *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is considered not rapidly degradable (NRD).

The hazard categories for chronic aquatic toxicity and their related criteria are set out in Regulation (EC) No 1272/2008 on classification, labelling and packaging (CLP) of substances and mixtures, Annex I, Section 4.1, Table 4.1.0.(b). According to table 4.1.0 (b)(i), for non-rapidly degradable substances, for which there are adequate chronic toxicity data available:

- For *Daphnia magna* there is the lowest NOEC = 0.00086 mg total pyrethrins/l, equivalent to 0.00104 mg *Chrysanthemum cinerariaefolium* extract from supercritical CO₂, without solvent. If NOEC < 0.1 mg/l, then category chronic 1 applies.
- In Annex I, table 4.1.3, if 0.001 < NOEC < 0.01, a multiplying factor M = 10 applies for NRD substances.

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is classified as Aquatic Chronic 1; H410; M=10 (when considering the substance as total pyrethrins the M factor would be 100 based on a NOEC = 0.00086mg/L).

In conclusion, Acute 1, M=100, Chronic 1, M= 10 is the proposed classification for *Chrysanthemum cinerariaefolium*, extract from open and mature flowers of *Tanacetum cinerariifolium* obtained with supercritical CO₂.

A.3.3.3. Conclusion on classification and labelling for environmental hazards and comparison with the CLP criteria

Chrysanthemum cinerariaefolium extract from supercritical CO₂ should be classified with respect to the environment as Aquatic Acute 1, H400, M = 100 and Aquatic Chronic 1, H410, M = 10.

A.4. Assessment of additional hazards

A.4.1. Hazardous to the ozone layer

A.4.1.1. Short summary and overall relevance of the provided information on ozone layer hazard

Following the physical and chemical properties and the structure of Pyrethrins it is assumed that degradation and persistence of the active substance mainly depends on reaction with hydroxyl-radicals and the average concentration of hydroxyl-radicals in air. Total OH rate constant was determined to be $281.1508 \times 10^{-12} \text{ cm}^3/\text{molec.} \cdot \text{sec.}$, mainly due to addition to olefinic bonds (96%) and hydrogen abstraction (4%). Other mechanisms do not contribute to hydroxyl radical estimations. The total rate of both, OH and ozone constant is very low. Half-life in the troposphere was calculated to be 27.391 min for overall OH rate constant and 29.562 min for ozone rate constant. Following the Atkinson calculation, the chemical half-life for Pyrethrum in the troposphere will be below 1 h. It is therefore concluded that Pyrethrins will not accumulate in air and will only be transported on very short distances.

The photochemical oxidative degradation half-life of Pyrethrin 1 in air was calculated according to the method developed by Atkinson, which is based on the structural activity relationship (QSAR's), by using the Atmospheric Oxidation Program v 1.91 (AOPWIN-software). These estimations were carried out with respect to the OH radical and ozone reactions, using a 12-hours-day with 300.95×10^{-12} and $96.32 \times 10^{-17} \text{ cm}^3/\text{molecule} \cdot \text{sec.}$, respectively. The half-lives for the hydroxyl and ozone reactions in air are estimated to be 25.59 and 17.13 minutes, respectively.

A half-life of 76.8 minutes for Pyrethrin 1 in air has been estimated using a 24-hour-day and assuming an OH radical concentration of $5 \times 10^5 \text{ radicals} \cdot \text{cm}^{-3}$ according to AOPWIN version 1.91 and following recommendations of ECHA Guidance on Risk Assessment, Chapter 2.3.6.3.

Stratospheric ozone depletion can be excluded due to the very short half-life in air (DT_{50} in air = 17.133 min), as a result of gas phase reactions with ozone (O₃).

A.4.1.2. Comparison with the CLP criteria

Not hazardous to the ozone layer.

Conclusion on classification and labelling for hazardous to the ozone layer.

No classification and labelling required.

A.5. Additional Labelling

No further labelling required.

A.6. Assessment of exclusion criteria, substitution criteria and POP

A.6.1. Exclusion criteria

A.6.1.1. Assessment of CMR properties

Criteria (BPR Article 5[1])	Assessment
Active substances which have been classified in accordance with Regulation (EC) No 1272/2008 as, or which meet the criteria to be classified as, carcinogen category 1A or 1B	Active substance is not classified and does not meet the criteria to be classified as Carc. Cat. 1A or 1B.
Active substances which have been classified in accordance with Regulation (EC) No 1272/2008 as, or which meet the criteria to be classified as, mutagen category 1A or 1B	Active substance is not classified and does not meet the criteria to be classified as Muta. Cat. 1A or 1B.
Active substances which have been classified in accordance with Regulation (EC) No 1272/2008 as, or which meet the criteria to be classified as, toxic for reproduction category 1A or 1B	Active substance is not classified and does not meet the criteria to be classified as Repr. Cat. 1A or 1B.
Conclusion on CMR properties	The exclusion criteria in BPR Article 5(1)a-c are not met.

A.6.1.2. Assessment of endocrine disrupting properties

Criteria (BPR Article 5)	Assessment
Active substances which, on the basis of the criteria specified pursuant to the first subparagraph of paragraph 3 are considered as having endocrine-disrupting properties that may cause adverse effects in humans and to the environment.	The assessment should be made in accordance with the scientific criteria set out in Commission Delegated Regulation (EU) 2017/2100 and Guidance for the identification of endocrine disruptors in the context of Regulations (EU) No 528/2012 and (EC) No 1107/2009.
Conclusion on ED properties:	ED properties have not been sufficiently investigated. It is not possible to ask for additional assays due to BPR article 90. Thus, eCA cannot conclude if the active substance meets or not ED criteria.

A.6.1.3. PBT Assessment (following Annex XIII to Regulation (EC) No 1907/2006)

Assessment of persistence

A persistence assessment based on the criteria for identification of persistent or very persistent substances based on REACH Annex XIII was performed. The assessment was performed for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Screening

Chrysanthemum cinerariaefolium extract from supercritical CO₂ is not readily biodegradable and, therefore, a full assessment of persistence is required.

The persistence assessment is summarised in the tables below.

P Criteria	Assessment
T1/2 > 60 days in seawater, or	-
T1/2 > 40 days in fresh- or estuarine water, or	Photolytic half-life in water estimated to be 11.8 h, therefore, it can be concluded that <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ does not fulfil the criterion for persistence.
T1/2 > 180 days in seawater sediment, or	-
T1/2 > 120 days in freshwater- or estuarine sediment, or	DT ₅₀ for biodegradation in sediment is 10 d (at 12°C) (Pyrethrin 1) and 5.27 d at 20°C, therefore, it can be concluded that <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ does not fulfil the criterion for persistence.
T1/2 ≤ 120 days in soil.	DT ₅₀ for biodegradation in soil 3.3 d (at 20°C) (Pyrethrin 1), therefore, it can be concluded that <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ does not fulfil the criterion for persistence.

vP Criteria	Assessment
T1/2 > 60 days in sea-, fresh- or estuarine water, or	The vP criterion criteria not fulfilled for <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ .
T1/2 > 180 days in seawater-, freshwater- or estuarine sediment, or	
T1/2 > 180 days in soil.	
Conclusion on P / vP properties	<p><i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ does not degrade in a ready biodegradability test and is not assumed to degrade in sewage treatment plants. Pyrethrins are hydrolytically stable to environmental relevant pH but the photolysis is rapid according to laboratory experiments in water that show a photolytic half-life of 11.8 hours.</p> <p>Apart from this, <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ has potential for rapid association to the sediment and consequent degradation according to the estimated half-life in water-sediment system (DT₅₀ = 5.27 days at 20°C). Pyrethrins are rapidly photolysed when they are exposed to natural sunlight on the surface of soil (DT₅₀ = 12.9 hours) and they are relatively quickly degraded in soil under aerobic conditions (DT₅₀ = 3.3 days). Thus, the criterion for persistence established by Table 11-2 in the PBT assessment guidance from ECHA (<i>Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT/vPvB assessment</i>, Version 3.0, June 2017⁵) is not fulfilled.</p>

Assessment of bioaccumulation

An assessment of bioaccumulation potential was performed based on the criteria for

⁵ https://echa.europa.eu/documents/10162/13632/information_requirements_r11_en.pdf

identification of bioaccumulative or very bioaccumulative substances based on REACH Annex XIII criteria. The assessment was performed for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Screening

A substance is considered to have the potential to fulfil the criterion of bioaccumulation when the log K_{ow} exceeds 4.5 (according to the PBT/vPvB Guidance Document⁶).

The log K_{ow} for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is 5.59 and hence further assessment of the bioaccumulation potential of *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is required.

The bioaccumulative assessment is summarised in the tables below.

B Criteria	Assessment
BCF > 2000	The BCF for fish is 500 which is well below the threshold of 2000.
vB Criteria	Assessment
BCF > 5000	-
Conclusion on B/vB properties	<i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ has a log K _{ow} equal to 5.59 which is higher than the limit value of 4.5 indicated in the ECHA Guidance. Therefore, this substance may be considered potentially bioaccumulating. Nevertheless, a bioconcentration study of Pyrethrins in fish was submitted by the applicant. The BCF whole body value for Pyrethrins estimated in this study was 500, in addition, a fast depuration rate was estimated (DT _{50, depuration} = 1 day). Based on this result, Pyrethrins do not fulfil the bioaccumulation criterion established in Table 11-2 in the PBT assessment guidance from ECHA (<i>Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT/vPvB assessment</i> , Version 3.0, June 2017 ⁷) is not fulfilled.

Assessment of toxicity

An assessment of toxicity was performed based on the criteria for identification of toxic substances based on REACH Annex XIII criteria. The assessment was performed for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂.

Screening

According to the most sensitive endpoint available for *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ from a 21 day *Daphnia magna* study (NOEC 0.86 µg ai/L) the toxic criterion is fulfilled according to the PBT/vPvB Guidance Document.

The toxicity assessment is summarised in the tables below.

T Criteria	Assessment
------------	------------

⁶ ECHA (2014); Guidance on Information Requirements and Chemical Safety Assessment; Chapter R.11: PBT/vPvB assessment; Version 2.0; November 2014

⁷ https://echa.europa.eu/documents/10162/13632/information_requirements_r11_en.pdf

NOEC/EC ₁₀ (long-term) < 0.01 mg/L for freshwater or seawater organisms, or	The NOEC for daphnia magna (21 days) is 0.86 µg ai/L which fulfils the criterion for toxicity.
substance meets the criteria for classification as carcinogenic (category 1A or 1B), germ cell mutagenic (category 1A or 1B), or toxic for reproduction (category 1A, 1B or 2) according to the CLP Regulation, or	<i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ is not classified as carcinogenic (category 1A or 1B), germ cell mutagenic (category 1A or 1B), or toxic for reproduction (category 1A, 1B or 2) according to the CLP Regulation.
there is other evidence of chronic toxicity, as identified by the substance meeting the criteria for classification: specific target organ toxicity after repeated exposure (STOT RE category 1 or 2) according to the CLP Regulation.	<i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ is not classified for specific target organ toxicity after repeated exposure (STOT RE category 1 or 2) according to the CLP Regulation.
Conclusion on T properties	The toxicity criterion is fulfilled by <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ as the chronic NOEC for aquatic organisms is lower than 0.01 mg/L [limit value established Table 11-2 in the PBT assessment guidance from ECHA (<i>Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT/vPvB assessment</i> , Version 3.0, June 2017 ⁸). On the other hand, toxicity criteria for mammals are not fulfilled by <i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO ₂ because of it is not classified as carcinogenic, mutagenic, or toxic for reproduction.

Summary and overall conclusions on PBT or vPvB properties

Overall conclusion:

Chrysanthemum cinerariaefolium extract from supercritical CO₂ cannot be considered as persistent or bioaccumulating. The only criterion fulfilled is Toxicity as it is very toxic to aquatic organisms. Consistently, *Chrysanthemum cinerariaefolium* extract from supercritical CO₂ is not PBT or vPvB.

A.6.2. Substitution criteria

Substitution criteria (BPR, Article 10)	Assessment
One of the exclusion criteria listed in Article 5(1) is met but AS may be approved in	Not met.

⁸ https://echa.europa.eu/documents/10162/13632/information_requirements_r11_en.pdf

accordance with Article 5(2)	
The criteria to be classified, in accordance with Regulation (EC) No 1272/2008, as a respiratory sensitiser are met	Not met.
The acceptable daily intake, acute reference dose or acceptable operator exposure level, as appropriate, is significantly lower than those of the majority of approved active substances for the same product-type and use scenario	Not met.
Two of the criteria for being PBT in accordance with Annex XIII to Regulation (EC) No 1907/2006 are met	Not met.
There are reasons for concern linked to the nature of the critical effects which, in combination with the use patterns, amount to use that could still cause concern, such as high potential of risk to groundwater, even with very restrictive risk management measures	Not met.
The AS contains a significant proportion of non-active isomers or impurities.	Not met.
Conclusion on substitution criteria	The substitution criteria in BPR Article 10(1)a-f are not met.

A.6.3. Assessment of long-range environmental transportation and Assessment of long-range environmental transportation and impact on environmental compartments

Assessment	
The active substance or a degradation product is a persistent organic pollutant (POP) listed in Annex I of EC 850/2004	The active substance is not listed in Annex I of EC 850/2004.
Assessment of long-range transport potential (LRTAP): Vapour pressure <1000 Pa and half-life in air > 2 days or Monitoring data in remote area showing that the substance is found in remote regions or Result of multimedia modelling	The vapour pressure of Pyrethrin 1, as a representative member of Pyrethrins, is 6.9E-05 Pa (25°C), its half-life in air is of 76.8 minutes (OH radicals) and 17.13 minutes (O ₃), indicating that the criterion for long-range transboundary atmospheric transport potential is not fulfilled.
The active substance or a degradation product is vP/vB or T?	The half-life of Pyrethrins in water is lower than two months, in sediment and soil the half-life is lower than 6 months. Therefore, IT cannot be considered as a persistent substance.

	<p>A log K_{ow} equal to 5.59 and a BCF_{whole body} of 500 was reported from a bioconcentration study of Pyrethrins in fish, where a fast depuration rate was observed (DT_{50, depuration} = 1 day). Thus, the criterion establish for bioaccumulating substances is not fulfil by Pyrethrins.</p> <p>Pyrethrins are very toxic to aquatic organisms (the most sensitive species was reported to be <i>D. magna</i>). Toxicity criteria for mammals are not fulfilled as it is not classified as carcinogenic, mutagenic, or toxic for reproduction.</p>
Conclusion on LRTAP/POP assessment	<p><i>Chrysanthemum cinerariaefolium</i> extract from supercritical CO₂ cannot be classified as a POP according to the Executive Body Decision 1998/2 on information to be submitted and the procedure for adding substances to annexes i, ii or iii to the protocol on persistent organic pollutants.</p>

B. Appendices

APPENDIX V: OVERALL REFERENCE LIST (INCLUDING DATA OWNER AND CONFIDENTIALITY CLAIM)

Section A

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
Anonymous	A3.2/02	1900	DETERMINATION OF VAPOUR PRESSURE IN REFINED PYRETHRUM EXTRACT Pyrethrum Board of Kenya, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	KPIC
Anonymous	A3.1.3/01	2000	SPECIFICATION PALE PYRETHRUM EXTRACT 50% Kenya Pyrethrum Information Centre, Oberalm, Austria Kenya Pyrethrum Information Centre Report-no. not applicable GLP: no Published: no	yes	KPIC
Anonymous	A6.12.2/01	2001	HUMAN EXPOSURES TO CONSUMER PRODUCTS CONTAINING PYRETHRINS AND PYRETHROIDS: REPORTS TO THE AMERICAN ASSOCIATION OF POISON CONTROL CENTERS 1994-1999 PEGUS Research, Inc., Salt Lake City, UT 84106 Pyrethrin Joint Venture Report-no. not stated GLP: no Published: no	yes	KPIC
Anonymous	A3.14/01	2006	MATERIAL SAFETY DATA SHEET - PYRETHRUM EXTRACT PALE 50% Kenya Pyrethrum Board, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not applicable GLP: no Published: no Submitted in: A3.3/01	yes	KPIC
Anonymous	A3.3/01	2006	MATERIAL SAFETY DATA SHEET - PYRETHRUM EXTRACT PALE 50% Kenya Pyrethrum Board, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not applicable GLP: no Published: no	yes	KPIC
Anonymous	A3.8/01	2006	MATERIAL SAFETY DATA SHEET -	yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			PYRETHRUM EXTRACT PALE 50% Kenya Pyrethrum Board, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not applicable GLP: no Published: no Submitted in: A3.3/01		
Anonymous	A6.1.4/02	1991	PRIMARY EYE IRRITATION - RABBITS Biosearch Incorporated, Philadelphia, PA 19134 Pyrethrin Joint Venture Report-no. 91-7316A GLP: yes Published: no	yes	PJV
Bienert, U. & Groer, M.	A7.5.1.2/01	1990	ACUTE TOXICITY (14 DAYS) OF PYRETHRUM EXTRACT TO EARTHWORMS EISENIA FOETIDA (SAVIGNY 1826) IN ARTIFICIAL SOIL IBACON, Rossdorf, Germany Kenya Pyrethrum Information Centre Report-no. 640021 GLP: yes Published: no	yes	KPIC
Bruske, L.J.	A3.4/04	1987	REPORT OF ANALYTICAL SERVICES, PYRETHRUM EXTRACT MASS SPECTROMETRY Shrader Lab. Inc., Vinewood, Detroit, Michigan, USA Kenya Pyrethrum Information Centre Report-no. 15335 GLP: no Published: no	yes	KPIC
Casida, J.E.	A3.4/03	1973	PYRETHRUM - THE NATURAL INSECTICIDE Academic Press, New York and London Report-no. not applicable GLP: no Published: yes	yes	-
Casida, J.E. , Quistad, G.B.	A6.12.3/01	1995	PYRETHRUM - A BENEFIT TO HUMAN WELFARE Oxford University Press, New York, Oxford Academic Press Report-no. not applicable GLP: no Published: yes	no	-
Casida, J.E., Quistad, G.B.	A5.4/01	1995	PYRETHRUM FLOWERS: PRODUCTION, CHEMISTRY, TOXICOLOGY AND USES - Oxford University Press, 1995, 217-233 Report-no. not applicable GLP: no	no	-

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			Published: yes		
Casida, J.E., Quistad, G.B.	A5.7/01	1995	PYRETHRUM FLOWERS: PRODUCTION, CHEMISTRY, TOXICOLOGY AND USES - Oxford University Press, 1995, 217-233 Report-no. not applicable GLP: no Published: yes Submitted in: A5.4/01	no	-
Comb, T.	A1.3	2021	PYRETHRINS: APPEARANCE AgroChemex Environmental Ltd, Aldhams Farm Research Station, Dead Lane, Manningtree, Essex, CO11 2NF United Kingdom Report ACE-21-459 GLP: yes Published: no	yes	MGK
Comb, T.	A1.3	2021	PYRETHRINS: RELATIVE DENSITY AgroChemex Environmental Ltd, Aldhams Farm Research Station, Dead Lane, Manningtree, Essex, CO11 2NF United Kingdom Report ACE-21-460 GLP: yes Published: no	yes	MGK
Comb, T.	A1.3	2021	PYRETHRINS: SURFACE TENSION AgroChemex Environmental Ltd, Aldhams Farm Research Station, Dead Lane, Manningtree, Essex, CO11 2NF United Kingdom Report ACE-21-461 GLP: yes Published: no	yes	MGK
Comb, T.	A1.3	2021	PYRETHRINS: VISCOSITY AgroChemex Environmental Ltd, Aldhams Farm Research Station, Dead Lane, Manningtree, Essex, CO11 2NF United Kingdom Report ACE-21-462 GLP: yes Published: no	yes	MGK
Curren, R.D.	A6.6	1989	Unscheduled DNA synthesis assay in rat primary hepatocytes with a confirmatory assay, Microbiological Associates, Inc.,	Yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			9900 Blackwell Road, Rockville, Maryland 20850, USA. report no. T8729.380009 GLP: Yes Published: No		
Curry, P.T.	A6.6.2/01	1996	CHROMOSOME ABERRATIONS IN CHINESE HAMSTER OVARY (CHO) CELLS Microbiol. Associates Inc., Rockville, Maryland, USA Pyrethrin Joint Venture Report-no. G96AC14.330001 GLP: yes Published: no	yes	PJV
Dabrunz, A.	A7.4.3.5.1	2017a	PY-T-50 Pale Refined Pyrethrins: Toxicity to the larvae of <i>Chironomus riparius</i> under Laboratory Conditions (Acute Immobilisation Test – Semi-Static). Rep. N° S16-07030 GLP: yes Published: no	Yes	BRA
Dabrunz, A.	A7.4.3.5.1	2017b	Pyrocide®50%: Toxicity to the larvae of <i>Chironomus riparius</i> under Laboratory Conditions (Acute Immobilisation Test – Semi-Static). Rep.N° S17-00132 GLP: yes Published: no	Yes	MGK
Dabrunz, A.	A7.4.3.5.1	2017c	Pyrethrum Extract Pale 50%: Toxicity to the larvae of <i>Chironomus riparius</i> under Laboratory Conditions (Acute Immobilisation Test – Semi-Static). Rep.N° S17-00133 GLP: yes Published: no	Yes	KPIC
Anonymous	A6.1.5	2017a	Local Lymph Node Assay in Mice (LLNA), MB Research Laboratories, 1765 Wentz Road, P.O. Box 178, Spinnerstown, PA 18968, US, Project No. MB 17-24906.26, Protocol No.5650A-08, 20 Apr 2018 GLP: yes Published: no	Yes	KPIC
Anonymous	A6.1.5	2017b	Local Lymph Node Assay in Mice (LLNA), MB Research Laboratories, 1765 Wentz Road, P.O. Box 178, Spinnerstown, PA 18968, US, Project No. MB 17-24904.26, Protocol No. 5650A-08, 20 Apr 2018 GLP: yes Published: no	Yes	MGK
Anonymous	A6.1.5	2017c	Local Lymph Node Assay in Mice (LLNA), MB Research Laboratories, 1765 Wentz Road, P.O. Box 178, Spinnerstown, PA 18968, US, Project No. MB 17-24905.26, Protocol No.	Yes	BRA

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			5650A-08, 20 Apr 2018 GLP: yes Published: no		
Dengler, D.	A7.4.1.3	2000	TESTING OF TOXIC EFFECTS OF PYRETHRUM EXTRACT PALE 50 % ON THE SINGLE CELL GREEN ALGA DESMODESMUS SUBSPICATUS ArGe GAB Biotech/IFU, Niefern- Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20001028/01-AAADs GLP: yes Published: no	yes	KPIC
Dengler, D.	A7.4.1.4	2003	ACUTE TOXICITY TESTING OF PYRETHRUM PALE EXTRACT ON ACTIVATED SLUDGE WITH THE RESPIRATION INHIBITION TEST ArGe GAB Biotech/IFU, Niefern- Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20031289/01-AAHT GLP: yes Published: no	yes	KPIC
Donath, C.	A6.6.2	2016a	<i>In Vitro</i> Mammalian Micronucleus Assay in Chinese Hamster V79 Cells with Pyrethrum Extreact 50%, Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, D-82152 Planegg/Munich, Germany, Study number 164165.GLP: yes Published: no	yes	KPIC
Donath, C.	A6.6.2	2016b	<i>In vitro</i> Mammalian Micronucleus Assay in Chinese Hamster V79 Cells with Pyroicide 50%, Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, D-82152 Planegg/Munich, Germany, study number 164338. GLP: yes Published: no	yes	MGK
Donath, C.	A6.6.2	2016c	<i>In vitro</i> Mammalian Micronucleus Assay in Chinese Hamster V79 Cells with PY-T-50 Pale Refined Pyrethrins, Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, D-82152 Planegg/Munich, Germany, study number 164099. GLP: yes Published: no	yes	BRA
Driz, M.T.	A4.2/01	1994	ANALYTICAL METHOD FOR THE DETERMINATION OF PYRETHRIN BY GAS CHROMATOGRAPHY AND PIPERONYL BUTOXIDE BY LIQUID CHROMATOGRAPHY IN SOIL, SEDIMENT AND WATER Pharmaco LSR Int. INC., East	yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			Millstone, NJ 08875 Pyrethrin Joint Venture Report-no. BD-047-92 GLP: yes Published: no		
A. P. Fifi	A7.2.2.1	2015a	Pyrethrum extract components – degradation study in loam soil according to OECD 307. BioTecnologie B. T. Srl, c/o Parco Tecnologico Agroalimentare dell’Umbria, Frazione Pantalla, 06059 Todi (PG), Italy GLP, unpublished	Yes	91/414 EU taskforce
A. P. Fifi	A7.2.2.1	2015b	Pyrethrum extract components – degradation study in sandy loam soil according to OECD 307. BioTecnologie B. T. Srl, c/o Parco Tecnologico Agroalimentare dell’Umbria, Frazione Pantalla, 06059 Todi (PG), Italy GLP, unpublished	Yes	91/414 EU taskforce
A. P. Fifi	A7.2.2.1	2015c	Pyrethrum extract components – degradation study in loamy sand soil according to OECD 307. BioTecnologie B. T. Srl, c/o Parco Tecnologico Agroalimentare dell’Umbria, Frazione Pantalla, 06059 Todi (PG), Italy GLP, unpublished	Yes	91/414 EU taskforce
Anonymous	A6.10/01	2002	DEFINITIVE MECHANISTIC TOXICITY STUDY IN RATS WITH PYRETHRINS Inveresk Research International, Tranent, Scotland Pyrethrin Joint Venture Report-no. 21029 GLP: yes Published: no	yes	PJV
Anonymous	A6.1.1/01	1991a	ACUTE ORAL TOXICITY, LD50 - RATS Biosearch Incorporated, Philadelphia, PA 19134 Pyrethrin Joint Venture Report-no. 91-7316A GLP: yes Published: no	yes	PJV
Anonymous	A6.1.2/01	1991b	ACUTE DERMAL TOXICITY, SINGLE LEVEL - RABBITS Biosearch Incorporated, Philadelphia, PA 19134 Pyrethrin Joint Venture Report-no. ITPY0027 GLP: yes Published: no	yes	PJV
Anonymous	A6.3.1/01	1987a	TWO WEEK DIETARY EXPLORATORY TOXICITY STUDY IN RATS International Research and Development Corp., Michigan, USA	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			Pyrethrin Joint Venture Report-no. 556-012 GLP: yes Published: no		
Anonymous	A6.3.1/02	1987b	2 WEEK-DIETARY TOXICITY STUDY IN MICE International Research and Development Corp., Michigan, USA Pyrethrin Joint Venture Report-no. 556-014 GLP: yes Published: no	yes	PJV
Anonymous	A6.4.1/01	1988a	EVALUATION OF PYRETHRUM EXTRACT IN A 13-WEEK DOSE RANGE FINDING STUDY IN RATS Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-010 GLP: yes Published: no	yes	PJV
Anonymous	A6.4.1/02	1988b	EVALUATION OF PYRETHRUM EXTRACT IN A 13-WEEK DOSE RANGE FINDING STUDY IN MICE Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-008 GLP: yes Published: no	yes	PJV
Anonymous	A6.3.1/03	1988c	EVALUATION OF PYRETHRUM EXTRACT IN AN EIGHT-WEEK DOSE RANGE FINDING TOXICITY STUDY IN DOGS Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-006 GLP: yes Published: no	yes	PJV
Anonymous	A6.4.1/03	1990a	EVALUATION OF PYRETHRUM EXTRACT IN A ONE YEAR CHRONIC TOXICITY STUDY IN DOGS Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-007 GLP: yes Published: no	yes	PJV
Anonymous	A6.7/01	1990b	EVALUATION OF PYRETHRUM EXTRACT IN TWO-YEAR DIETARY TOXICITY AND ONCOGENICITY STUDY IN RATS Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			Report-no. 556-011 GLP: yes Published: no		
Anonymous	A6.7/02	1990c	EVALUATION OF PYRETHRUM EXTRACT IN AN EIGHTEEN MONTH DIETARY ONCOGENICITY STUDY IN MICE Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-013 GLP: yes Published: no	yes	PJV
Anonymous	A6.3.2/01	1992	21-DAY REPEATED DOSE DERMAL TOXICITY STUDY WITH PYRETHRUM EXTRACT IN RABBITS Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-018 GLP: yes Published: no	yes	PJV
Anonymous	A6.12.1/01	1976	HEALTH OF MEN ON LONG-TERM EXPOSURE TO PYRETHRINS The Pyrethrum Marketing Board, Nakuru, Kenya, East Africa Pyrethrin Joint Venture Report-no. not stated GLP: no Published: no	yes	KPIC
Gonsior, G.	A 7.4.3.5.1/ 02	2009	ASSESSMENT OF SIDE EFFECTS OF PYRETHRUM PALE EXTRACT ON THE LARVAE OF THE MIDGE, CHIRONOMUS RIPARIUS WITH THE LABORATORY TEST METHOD eurofins-GAB GmbH, Niefern- Öschelbronn, Germany DKSH Switzerland Ltd. Report-no. S09-00063 GLP: yes Published: no	yes	DKSH
Hattermann, D.R.	A7.2.2.2/01	1992a	FIELD PHASE FOR PYRENONE CROP SPRAY (PYRETHRUM + PIPERONYL BUTOXIDE) FIELD DISSIPATION - TERRESTRIAL STUDY APPLIED TO BAREGROUND IN CALIFORNIA, GEORGIA, AND MICHIGAN - VOL.1 Bio/Dynamics, Inc., New Jersey, USA Pyrethrin Joint Venture Report-no. 91189 GLP: yes Published: no	yes	PJV
Hattermann, D.R.	A7.2.2.2/02	1992b	PYRETHRUM ANALYTICAL PHASE FOR PYRENONE CROP SPRAY (PYRETHRUM + PIPERONYL BUTOXIDE) FIELD DISSIPATION - TERRESTERIAL STUDY	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			APPLIED TO BAREGROUND IN CALIFORNIA, GEORGIA, AND MICHIGAN - VOL.2 Bio/Dynamics, Inc., New Jersey, USA Pyrethrin Joint Venture Report-no. 91189 GLP: yes Published: no		
W. Hein	A7.2.1		Degradation of [Cyclopentenone-2- ¹⁴ C]Pyrethrin I in Mußbach soil incubated under aerobic conditions at 20 °C in the dark. RLP AgroScience GmbH, Institut für Agrarökologie, Breitenweg 71, 67435 Neustadt, Germany GLP, unpublished	Yes	91/414 EU taskforce
W. Hein, M. Mömdel	A7.1.2.2	2017	[Cyclopentenone-2- ¹⁴ C]pyrethrin 1 "aerobic degradation in natural water". RLP AgroScience GmbH, Breitenweg 71, 67435 Neustadt, Germany GLP, unpublished	Yes	91/414 EU taskforce
Heintze, A.	A7.4.3.5.1	2001	ASSESSMENT OF SIDE EFFECTS OF PYRETHRUM EXTRACT PALE 50 % ON THE LARVAE OF THE MIDGE, CHIRONOMUS RIPARIUS WITH THE LABORATORY TEST METHOD ArGe GAB Biotech/IFU, Niefern-Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20001028/01-ASCr GLP: yes Published: no	yes	KPIC
Anonymous	A6.9/01	1993	ACUTE ORAL NEUROTOXICITY STUDY WITH PYRETHRUM EXTRACT IN RATS Bushy Run Research Center (BRRC) Pyrethrin Joint Venture Report-no. 92N1036 GLP: yes Published: no	yes	PJV
Anonymous	A6.1.3/01	1991	AN ACUTE INHALATION TOXICITY STUDY OF PYRETHRUM EXTRACT IN THE RAT Bio/Dynamics, Inc., New Jersey, USA Pyrethrin Joint Venture Report-no. 91-8331 GLP: yes Published: no	yes	PJV
Hoffmann, H.	A3.11/01	2000	PYRETHRUM PALE EXTRACT 99/11-5B PALE 50%, DETERMINATION OF PHYSICO-CHEMICAL PROPERTIES A.14 A.15 Aventis Research & Technologies, D-65926 Frankfurt a. M. Kenya Pyrethrum Information Centre Report-no. SI072-00	yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			GLP: yes Published: no		
Hoffmann, H.	A3.15/01	2000	PYRETHRUM PALE EXTRACT 99/11-5B PALE 50%, DETERMINATION OF PHYSICO-CHEMICAL PROPERTIES A.14 A.15 Aventis Research & Technologies, D- 65926 Frankfurt a. M. Kenya Pyrethrum Information Centre Report-no. SI072-00 GLP: yes Published: no Submitted in: A3.11/01	yes	KPIC
Anonymous	A6.6.4/01	1976	MICRONUCLEUS TEST ON PYRETHRUM EXTRACT Huntingdon Centre, England Pyrethrin Joint Venture Report-no. PYR 8/76707 GLP: no Published: no	yes	PJV
Koopmans, M.J.E.	A7.1.1.2.1	1995	DETERMINATION OF 'READY' BIODEGRADABILITY: CARBON DIOXIDE (CO ₂) EVOLUTION TEST (MODIFIED STURM TEST) WITH PYRETHRUM EXTRACT Notox B.V, 5231 DD 's- Hertogenbosch, The Netherlands Kenya Pyrethrum Information Centre Report-no. 141964 GLP: yes Published: no	yes	KPIC
Anonymous	A6.10/02	2002	AN INVESTIGATION OF SOME HEPATIC ENZYME ACTIVITIES IN LIVER SAMPLES DERIVED FROM INVERESK STUDY 455790: DEFINITIVE MECHANISTIC TOXICITY STUDY IN RATS WITH PYRETHRINS TNO BIBRA International Ltd. Surrey, UK Pyrethrin Joint Venture Report-no. 4024/2/2/2002 GLP: yes Published: no	yes	PJV
Leng, G., Gries, W., Selim, S.	A6.2/06	2006	BIOMARKER OF PYRETHRUM EXPOSURE not applicable Toxicology Lett, 162, 195-201 Report-no. GLP: no Published: yes	no	-
Lynn, S. P. & Hoxter, K.A.	A7.5.4.1	1991	AN ACUTE CONTACT TOXICITY STUDY WITH PYRETHRUM EXTRACT WITH THE HONEY BEE Wildlife International, Ltd., Easton, Maryland, USA Pyrethrin Joint Venture	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			Report-no. 326-104 GLP: yes Published: no		
Anonymous	A7.4.1.1/01	1994a	PYRETHRUM EXTRACT (FEK-99) - A FLOW-THROUGH ACUTE TOXICITY TEST WITH RAINBOW TROUT (ONCORHYNCHUS MYKISS) Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 93-11-5084 GLP: yes Published: no	yes	PJV
Anonymous	A7.4.1.1/02	1994b	PYRETHRUM EXTRACT (FEK-99) - A FLOW-THROUGH ACUTE TOXICITY TEST WITH BLUEGILL SUNFISH (LEPOMIS MACROCHIRUS) Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 93-8-4916 GLP: yes Published: no	yes	PJV
Anonymous	A7.4.1.1/03	1994c	PYRETHRUM EXTRACT (FEK-99) - A FLOW-THROUGH ACUTE TOXICITY TEST WITH SHEEPSHEAD MINNOW (CYPRINODON VARIEGATUS) Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 93-12-5081 GLP: yes Published: no	yes	PJV
Anonymous	A7.4.3.2/01	1994d	PYRETHRUM EXTRACT (FEK-99) - THE TOXICITY TO FATHEAD MINNOW (PIMEPHALES PROMELAS) DURING AN EARLY LIFE-STAGE EXPOSURE Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 93-10-4983 GLP: yes Published: no	yes	PJV
Maciver, D.R.	A5.3.1/05	1964	MOSQUITO COILS PART II. STUDIES ON THE ACTION OF MOSQUITO COIL SMOKE ON MOSQUITOES - Pyrethrum Post, Journal, 7 (3), 1964, 7-9 Report-no. not applicable GLP: no Published: yes	no	-
Mantilacci, S.	A7.4.1.2	2015a	5 pyrethrin metabolites - acute toxicity to daphnids (<i>daphnia magna</i>) under static conditions. rep.n°bt039/15.	Yes	91/414 EU taskforce

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			GLP: yes Published: No		
Mantilacci, S.	A7.4.1.2	2015b	Pyrethrum Extract (FEK-99) and 6 Pyrethrin esters - Acute toxicity to Daphnids (<i>Daphnia magna</i>) under semi-static conditions, Rep.N°BT038/15. GLP: yes Published: No	yes	91/414 EU taskforce
Mende, P.	A4.2/04	2006	VALIDATION OF AN ANALYTICAL METHOD FOR THE DETERMINATION OF PYRETHRUM IN WATER Eurofins-GAB GmbH, Niefern-Öschelbronn Kenya Pyrethrum Information Centre Report-no. 20051407/01-RVW GLP: yes Published: no	yes	KPIC
Mende, P.	A4.2/05	2006	VALIDATION OF AN ANALYTICAL METHOD FOR THE DETERMINATION OF PYRETHRUM IN SOIL Eurofins-GAB GmbH, Niefern-Öschelbronn Kenya Pyrethrum Information Centre Report-no. 20051407/01-RVS GLP: yes Published: no	yes	KPIC
Mori V.	A7.1.3	2015	Estimation of soil adsorption coefficient (K _{oc}) of Pyrethrum extract components by HPLC. Research Center Biospheres by Biotechnologie B.T., Parco Tecnologico Padano, Via A. Einstein – Loc. Cascina Codazza 26900 Lodi, Italy GLP, unpublished	yes	91/414 EU taskforce
Mori V.	A3.9	2015	Determination of the Partition Coefficient N-Octanol/Water (Log Pow) of Pyrethrum Extract Components By HPLC, Research Center Biospheres, Via A. Einstein-Loc. Cascina Codazza, 26900 LODI, Italy, Report CPU-004-15, 6 May 2015 GLP: yes Published: no	yes	91/414 EU taskforce
Morlock, G.	A4.2/03	2006	VALIDATION OF AN ANALYTICAL METHOD FOR THE DETERMINATION OF PYRETHRUM PALE EXTRACT IN AIR GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Kenya Pyrethrum Information Centre Report-no. 20051407/01-CMLU GLP: yes Published: no	yes	KPIC
Anonymous	A6.4.3/01	1992	A SUBCHRONIC (3-MONTH) INHALATION TOXICITY STUDY OF PYRETHRUM EXTRACT IN THE RAT VIA	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			WHOLE-BODY EXPOSURES Bio/Dynamics, Inc., New Jersey, USA Pyrethrin Joint Venture Report-no. 91-8335 GLP: yes Published: no		
Ochieng, C.D.	A3.9/01	1990	DETERMINATION OF PARTITION COEFFICIENT OF PYRETHRINS IN N- OCTANOL/WATER MIXTURE Pyrethrum Board of Kenya, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	KPIC
Oellrich, W.	A7.3.1/01	2000	PYRETHRUM EXTRACT, ESTIMATION OF THE PHOTOCHEMICAL OXIDATIVE DEGRADATION GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	KPIC
Oellrich, W.	A3.2.1	2001	PYRETHRINS, ANNEX II, POINT 2.3.2 HENRY'S LAW CONSTANT GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	KPIC
Oellrich, W.	A3.16/01	2002	PYRETHRINS, ANNEX II, POINT 2.15 OXIDIZING PROPERTIES GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. 104106-029001 GLP: no Published: no	yes	KPIC
Oellrich, W.	A3.1.1/01	2003	PYRETHRINS, ANNEX II POINT 2.1.1 MELTING POINT, POINT 2.1.2 BOILING POINT, POINT 2.3.1 VAPOUR PRESSURE DETERMINATION GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. 104108-001-01-rev01 GLP: no Published: no	yes	KPIC
Oellrich, W.	A3.1.2/01	2003	PYRETHRINS, ANNEX II POINT 2.1.1 MELTING POINT, POINT 2.1.2 BOILING POINT, POINT 2.3.1 VAPOUR PRESSURE DETERMINATION GAB Consulting GmbH, Lamstedt, Germany	yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			Kenya Pyrethrum Information Centre Report-no. 104108-001-01-rev01 GLP: no Published: no Submitted in: A3.1.1/01		
Oellrich, W.	A3.2/01	2003	PYRETHRINS, ANNEX II POINT 2.1.1 MELTING POINT, POINT 2.1.2 BOILING POINT, POINT 2.3.1 VAPOUR PRESSURE DETERMINATION GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. 104108-001-01-rev01 GLP: no Published: no Submitted in: A3.1.1/01	yes	KPIC
A. Perboni	A7.1.1.1.1.	2015	Pyrethrum extract components- hydrolysis study in water (according to OECD 111). Research Center Biospheres by Biotechnologie B.T., Parco Tecnologico Padano, Via A. Einstein – Loc. Cascina Codazza 26900 Lodi, Italy GLP, unpublished	Yes	91/414 EU taskforce
Pfersich, M.	A3.5/01	2001	SOLUBILITY OF PURIFIED ACTIVE SUBSTANCE IN WATER - Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	KPIC
Putman, D.L., Morris, M.J.	A6.6.2/02	1989	Chromosome aberrations in Chinese Hamster Ovary (CHO) cells Microbiol. Associates Inc., Rockville, Maryland, USA Pyrethrin Joint Venture Report-no. T8729.337 GLP, Unpublished	yes	PJV
Putt, A.E.	A7.4.1.2	1994a	PYRETHRUM EXTRACT (FEK-99) - ACUTE TOXICITY TO DAPHNIDS (DAPHNIA MAGNA) UNDER FLOW- THROUGH CONDITIONS Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 93-11-5082 GLP: yes Published: no	yes	PJV
Putt, A.E.	A7.4.3.4	1994b	PYRETHRUM EXTRACT (FEK-99) - THE CHRONIC TOXICITY TO DAPHNIA MAGNA UNDER FLOW-THROUGH CONDITIONS Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 94-1-5116	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			GLP: yes Published: no		
Reynolds, J.L., Robinson, R.A.	A7.1.3	1994	ADSORPTION AND DESORPTION OF [14C]PYRETHRIN 1 IN FOUR SOILS XenoBiotic Laboratories, Inc., Plainsboro, NJ 08536, USA Pyrethrin Joint Venture Report-no. RPT00156 GLP: yes Published: no	yes	PJV
Robinson, R. A.	A7.2.1/01	1994	AEROBIC SOIL METABOLISM OF [14C]PYRETHRIN 1 XenoBiotic Laboratories, Inc., Plainsboro, NJ 08536, USA Pyrethrin Joint Venture Report-no. RPT00204 GLP: yes Published: no	yes	PJV
Robinson, R. A.; Wisocky, M.J.	A7.1.2.2.2/0 1	1994	AEROBIC AQUATIC METABOLISM OF[14C]PYRETHRIN 1 XenoBiotic Laboratories, Inc., Plainsboro, NJ 08536, USA Pyrethrin Joint Venture Report-no. RPT00193 GLP: yes Published: no	yes	PJV
Anonymous	A6.1.4/01	1991a	PRIMARY SKIN IRRITATION - RABBITS Biosearch Incorporated, Philadelphia, PA 19134 Pyrethrin Joint Venture Report-no. 91-7316A GLP: yes Published: no	yes	PJV
Anonymous	A6.1.5/01	1991b	GUINEA PIG DERMAL SENSITIZATION - MODIFIED BÜHLER METHOD Biosearch Incorporated, Philadelphia, PA 19134 Pyrethrin Joint Venture Report-no. 91-7316A GLP: yes Published: no	yes	PJV
San, R.H.C., Springfield, K.A.	A6.6.1/01	1989	SALMONELLA/MAMMALIAN- MICROSOME PLATE INCORPORATION MUTAGENICITY ASSAY (AMES TEST) WITH A CONFIRMATORY ASSAY Microbiol. Associates Inc., Rockville, Maryland, USA Pyrethrin Joint Venture Report-no. T8729.501014 GLP: yes Published: no	yes	PJV
Anonymous	A6.8.1/01	1987a	EVALUATION OF PYRETHRUM EXTRACT IN A DEFINITIVE RAT TERATOLOGY STUDY Int. Res.and Develop.Corp. Mattawan,	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-002 GLP: yes Published: no		
Anonymous	A6.8.1/02	1987b	EVALUATION OF PYRETHRUM EXTRACT IN A DEFINITIVE RABBIT TERATOLOGY STUDY Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-004 GLP: yes Published: no	yes	PJV
Anonymous	A6.8.2/01	1989	TWO GENERATION REPRODUCTION STUDY IN RATS WITH PYRETHRUM EXTRACT Int. Res.and Develop.Corp. Mattawan, Michigan 49071, USA Pyrethrin Joint Venture Report-no. 556-005 GLP: yes Published: no	yes	PJV
Schmid, J.	A3.4/01	1990a	ANALYTICAL REPORT UV-VIS ABSORPTION SPECTRA OF PYRETRHUM EXTRACT ACCORDING TO OECD GUIDELINE 101 BioChem GmbH, Karlsruhe, Germany Kenya Pyrethrum Information Centre Report-no. GLP: no Published: no	yes	KPIC
Schmid, J.	A3.5/02	1990b	WATER SOLUBILITY OF PYRETHRUM EXTRACT ACCORDING TO CIPAC METHOD MT 157.1 BioChem GmbH, Karlsruhe, Germany Kenya Pyrethrum Information Centre Report-no. 947951 GLP: no Published: no	yes	KPIC
Schocken, M.J.	A7.4.3.3.1/0 1	1994	BIOCONCENTRATION STUDY WITH [14C] PYRETHRIN 1 IN BLUEGILL SUNFISH. Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 94-5-5258 GLP: yes Published: no	yes	PJV
Schreib, G.	A6.6.1	2016a	Reverse Mutation Assay using Bacteria (<i>Salmonella typhimurium</i> and <i>Escherichia coli</i>) with Pyrethrum- Extract 50%, Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, 82152 Planegg/Munich, Germany, Report	Yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			number 164164 GLP: yes Published: no		
Schreib, G.	A6.6.1	2016b	Reverse Mutation Assay using Bacteria (<i>Salmonella typhimurium</i> and <i>Escherichia coli</i>) with Pyrocide 50%, Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, 82152 Planegg/Munich, Germany, Report number 164337. GLP: yes Published: no	Yes	MGK
Schreib, G.	A6.6.1	2016c	Reverse Mutation Assay using Bacteria (<i>Salmonella typhimurium</i> and <i>Escherichia coli</i>) with PY-T-50 Pale Refined Pyrethrins, Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, 82152 Planegg/Munich, Germany, Report number 164098. GLP: yes Published: no	Yes	BRA
Schyler, K.C., Massing, R,A,	A5.3.1/04	1974	OBSERVATIONS ON GROUND ULV APPLICATIONS OF SYNERGISED PYRETHRINS ON NON-TARGET INSECTS AND MOSQUITOES IN CENTRE COUNTY, PENNSYLVANIA - Pyrethrum Post, Journal, 12 (4), 1974, 142-144 Report-no. not applicable GLP: no Published: yes	no	-
Anonymous	A6.2/03	2004	A Single Dose, Open Label Study to Investigate the Absorption and Excretion of Orally Administered or Dermally Applied (14C-Labeled Pyrethrin I (PI)) to Healthy Male Volunteers. Pyrethrin Joint Venture Consumer Specialty Product Associations, Inc. 900 17th Street, N.W. Washington, D.C. 20006 Study No.: Sel 0204 GLP Published: No	Yes	PJV
Selim, S.	A7.3.2/01	1993	LABORATORY VOLATILITY OF PYRETHRIN 1 FROM SOIL Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P0693011 GLP: yes Published: no	yes	PJV
Selim, S.	A7.1.1.1.1	1995a	HYDROLYSIS OF PYRETHRIN 1 AS A	yes	PJV

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			FUNCTION OF PH AT 25°C Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P1092011 GLP: yes Published: no		
Anonymous	A6.2/01	1995b	PHARMACOKINETICS AND METABOLISM OF PYRETHRIN 1 IN THE RAT Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P1092006 GLP: yes Published: no	yes	PJV
Selim, S.	A7.1.1.1.2/0 1	1995b	AQUEOUS PHOTOLYSIS OF PYRETHRIN 1 Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P1192006 GLP: yes Published: no	yes	PJV
Siusiene,E.	A1.4	2022	PHYSICO/CHEMICAL TESTING ON A TEST ITEM ON PYRETHRUM EXTRACT DEKRA UK Ltd, Phi House, Southampton Science Park, Southampton, SO16 7NS, United Kingdom Report GLP3016010232R1/2021 GLP: yes Published: no	yes	MGK
St.Laurent, J.P.	A4.2/02	1995	DEVELOPMENT AND VALIDATION OF AN ANALYTICAL METHODOLOGY FOR THE DETERMINATION OF TOTAL PYRETHRINS IN WATER, ACETONE AND FISH TISSUE Spingborn Lab. Inc., Wareham, Massachusetts 02571, USA Pyrethrin Joint Venture Report-no. 93-9-4922 GLP: yes Published: no	yes	PJV
Steenwinkel, M- J.S.T.	A6.6.3/01	2001	GENE MUTATION TEST AT THE TK- LOCUS OF L5178Y CELLS WITH PYRETHRIN TNO Nutrition and Food Res., Zeist, The Netherlands Pyrethrin Joint Venture Report-no. 2503/06 GLP: yes Published: no	yes	PJV
Straube, D.	A7.5.2.1	2016a	PY-T-50 Pale Refined Pyrethrins; Effects on Reproduction of the Collembola <i>Folsomia candida</i> in	yes	BRA

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
			Artificial Soil with 5% Peat: ibacon GmbH, Arheilger Weg 17, 64380 Rossdorf, Germany: unpublished report no., 114001016 (15 Sep 2016). GLP: yes		
Straube, D.	A7.5.2.1	2016b	PY-T-50 Pale Refined Pyrethrins; Effects on Reproduction of the Predatory Mite <i>Hypoaspis aculeifer</i> in Artificial Soil with 5% Peat: ibacon GmbH, Arheilger Weg 17, 64380 Rossdorf, Germany: unpublished report no., 114001089 (15 Sep 2016).	yes	BRA
Straube, D.	A7.5.2.1	2016c	PYROCID [®] 50%: Effects on Reproduction of the Collembola <i>Folsomia candida</i> in Artificial Soil with 5% Peat: ibacon GmbH, Arheilger Weg 17, 64380 Rossdorf, Germany: unpublished report no., 110901016 (12 Oct 2016). GLP: yes	yes	MGK
Straube, D.	A7.5.2.1	2016d	<i>PYROCID[®] 50%: Effects on Reproduction of the Predatory Mite Hypoaspis aculeifer in Artificial Soil with 5% Peat: ibacon GmbH, Arheilger Weg 17, 64380 Rossdorf, Germany: unpublished report no., 110901089 (12 Oct 2016).</i>	Yes	MGK
Straube, D.	A7.5.2.1	2016e	Pyrethrum Extract Pale 50%: Effects on Reproduction of the Collembola <i>Folsomia candida</i> in Artificial Soil with 5% Peat: ibacon GmbH, Arheilger Weg 17, 64380 Rossdorf, Germany: unpublished report no., 114011016 (15 Sep 2016).	Yes	KPIC
Straube, D.	A7.5.2.1	2016f	Pyrethrum Extract Pale 50%: Effects on Reproduction of the Predatory Mite <i>Hypoaspis aculeifer</i> in Artificial Soil with 5% Peat: ibacon GmbH, Arheilger Weg 17, 64380 Rossdorf, Germany: unpublished report no., 114011089 (16 Sep 2016).	yes	KPIC
Sum, K.S., Kimani, S.M., Kuria, J.N.	A5.3.1/03	2005	EMPIRICAL BIO-EFFICACY OF PYRETHRINS AND PYRETHROIDS IN AEROSOL FORMULATIONS FOR MOSQUITO CONTROL - Pyrethrum Post, Journal, 20 (04), 2005, 140-149 Report-no. not applicable GLP: no	no	-

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			Published: yes		
Testman, R.	A7.2.2.4/01	1994	SOIL SURFACE PHOTOLYSIS OF PYRETHRIN 1. Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P1192007 GLP: yes Published: no	yes	PJV
Tiemann, J.	A3.2/03	2004	PYRETHRINS, ANNEX II POINT 2.3.1 VAPOUR PRESSURE OF PURIFIED A.S., POINT 2.3.2 HENRY'S LAW CONSTANT GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	KPIC
Troese, M.	A6.1.5	2017a	<i>In vitro</i> Sensitization Assay (IVSA) Pyrethrum Extract Pale (50% w/w), MB Research Laboratories, 1765 Wentz Road, P.O.Box 178, Spinnerstown, PA 18968, USA, MB 17-24906.19 {Not signed and dated} GLP: yes Published: no	Yes	KPIC
Troese, M.	A6.1.5	2017b	<i>In vitro</i> Sensitization Assay (IVSA) Refined Pyrethrum Concentrate (53.72% w/w), MB Research Laboratories, 1765 Wentz Road, P.O.Box 178, Spinnerstown, PA 18968, USA, MB 17-24904.19 {Not signed and dated} GLP: yes Published: no	Yes	MGK
Troese, M.	A6.1.5	2017c	<i>In vitro</i> Sensitization Assay (IVSA) PY-T-50 Pale Refined Pyrethrins (49.36% w/w), MB Research Laboratories, 1765 Wentz Road, P.O.Box 178, Spinnerstown, PA 18968, USA, MB 17-24905.19 {Not signed and dated} GLP: yes Published: no	Yes	BRA
Wachter, S.	A7.5.1.1	2000	ASSESSMENT OF THE SIDE EFFECTS OF PYRETHRUM EXTRACT PALE 50 % ON THE ACTIVITY OF THE SOIL MICROFLORA ArGe GAB Biotech/IFU, Niefern- Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20001028/01-ABMF GLP: yes Published: no	yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
Wallner, B.	A6.6.1	2016a	<i>In vitro</i> Mammalian Cell Gene Mutation Assay (Thymidine Kinase Locus/TK ^{+/-}), Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, D-82152 Planegg/Munich, Germany, Report number 164166. GLP: yes Published: no	Yes	KPIC
Wallner, B.	A6.6.1	2016b	<i>In vitro</i> Mammalian Cell Gene Mutation Assay (thymidine Kinase Locus/TK ^{+/-}) in Mouse Lymphoma L5178Y Cells with Pyrocide 50% Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, D-82152 Planegg/Munich, Germany, Report number 164339 GLP: yes Published: no	Yes	MGK
Wallner, B.	A6.6.1	2016c	<i>In vitro</i> Mammalian Cell Gene Mutation Assay (thymidine Kinase Locus/TK ^{+/-}) in Mouse Lymphoma L5178Y Cells with PY-T-50 Pale Refined Pyrethrins Eurofins Biopharma Product Testing Munich GmbH, Behringstrasse 6/8, D-82152 Planegg/Munich, Germany, Report number 164100 GLP: yes Published: no	Yes	BRA
Walter, D.	A 3.1.3/02	2008a	RELATIVE DENSITY OF "PALE" PYRETHRUM EXTRACT 50 % eurofins-GAB GmbH, Niefern-Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20071557/01-PCRD GLP: yes Published: no	yes	KPIC
Walter, D.	A3.7/01	2000a	SOLUBILITY OF PYRETHRUM EXTRACT PALE 50% IN ORGANIC SOLVENTS ArGe GAB Biotech/IFU, Niefern-Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20001028/01-PSBO GLP: yes Published: no	yes	KPIC
Walter, D.	A3.12/01	2000b	FLASH POINT OF PYRETHRUM EXTRACT PALE 50 % ArGe GAB Biotech/IFU, Niefern-Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20001028/01-PCFB GLP: yes Published: no	yes	KPIC
Walter, D.	A3.13/01	2005	SURFACE TENSION OF PYRETHRUM EXTRACT PALE 50% GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn	yes	KPIC

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			Kenya Pyrethrum Information Centre Report-no. 20051087/01-PCST GLP: yes Published: no		
Walter, D.	A 3.14/02	2008b	VISCOSITY OF "PALE" PYRETHRUM EXTRACT 50 % eurofins-GAB GmbH, Niefern- Öschelbronn, Germany Kenya Pyrethrum Information Centre Report-no. 20071557/01-PCVC GLP: yes Published: no	yes	KPIC
Warui, C.	A5.3.1/01	1996a	TO COMPARE VARIOUS AEROSOL FORMULATIONS IN THE CONTROL OF HOUSEFLIES MUSCA DOMESTICA Kenya Pyrethrum Board, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. 896/1136 GLP: no Published: no	yes	KPIC
Warui, C.	A5.3.1/02	1996b	TO COMPARE VARIOUS AEROSOL FORMULATIONS IN THE CONTROL OF COCKROACHES Kenya Pyrethrum Board, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. 896/1136 GLP: no Published: no	yes	KPIC
Werle, H.	A7.1.1.1.2/0 2	1992	PHOTOLYSIS STUDY (QUANTUM YIELD) OF PYRETHRUM IN WATER BioChem GmbH, Karlsruhe, Germany Kenya Pyrethrum Information Centre Report-no. 915040180 GLP: no Published: no	yes	91/414 EU taskforce
Werle, H.	A3.4/02	1994	REPORT IR-SPECTRUM PYRETHRUM EXTRACT BioChem GmbH, Karlsruhe, Germany Kenya Pyrethrum Information Centre Report-no. 945040321 GLP: yes Published: no	yes	KPIC
Anonymous	A6.2/02	1994	HUMAN IN VIVO PERCUTANEOUS ABSORPTION OF PYRETHRIN AND PIPERONYL BUTOXIDE - Food Chem Toxicol, 32, 51 - 53 Report-no. Not applicable GLP: no Published: yes	no	-
Witte, A.	A7.1.2.2.2/0 2	2007	DEGRADATION AND METABOLISM OF PYRETHRIN 1 IN TWO WATER/SEDIMENT SYSTEMS UNDER AEROBIC CONDITIONS -	yes	91/414 EU taskforce

ES

Chrysanthemum cinerariaefolium, extract from open and mature flowers of
Tanacetum cinerariifolium obtained with supercritical CO₂

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protection claimed yes/no	Owner
			LABORATORY TEST eurofins-GAB GmbH, Niefern-Öschelbronn, Germany Pyrethrum Board of Kenya Report-no. 20061275/01-CUWS GLP: yes Published: no		

Section B Product A (AquaPy)

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
Balluff, M.	B7.8.6/01		AQUAPY: A GREENHOUSE TOXICITY STUDY TO DETERMINE THE EFFECTS OF A 30 G AI L-1 PIPERONYL BUTOXIDE EW FORMULATION ON THE VEGETATIVE VIGOUR OF SIX SPECIES OF PLANTS GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20054072/S2-FGVV GLP: no Published: no	yes	BES
Blagden, S.M.	B6.1.3/01	1994	AQUA PYBUTHRIN: ACUTE INHALATION TOXICITY STUDY FOUR HOUR EXPOSURE (NOSE ONLY) IN THE RAT Safepharm Laboratories Limited, Derby, UK Bayer ES Report-no. GR94-0003 GLP: yes Published: no	yes	BES
Bocksch, S.	B7.8.2/01	2005	ASSESSMENT OF SIDE EFFECTS OF AQUAPY® (PBO+PYR EW 135+30A G) TO THE HONEY BEE, APIS MELLIFERA L., IN THE LABORATORY GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-BLEU GLP: yes Published: no	yes	BES
Boucaud, B.	B6.6/01	2006	OCCUPATIONAL MEDICAL EXPERIENCES WITH PYRETHRE not applicable Bayer ES Report-no. not applicable GLP: no Published: no	yes	BES
Bowron, M.J.	B5.10.2/05	1993	AN EVALUATION OF AQUA PYBUTHRIN IN DRY DEPOSIT AND WET WALKOVER TESTS AGAINST BLATTELLA GERMANICA Roussel Uclaf Environmental Health, Berkhamstedt Bayer ES Report-no. REPE 93-C8 GLP: no Published: no	no	BES
Clouzeau, J.	B6.1.1/01	1993a	ACUTE ORAL TOXICITY IN RATS Centre International de Toxicologie, Evreux, France Bayer ES Report-no. 9965 TAR GLP: yes Published: no	yes	BES

Clouzeau, J.	B6.1.2/01	1993b	ACUTE DERMAL TOXICITY IN RATS Centre International de Toxicologie, Evreux, France Bayer ES Report-no. 9966 TAR GLP: yes Published: no	yes	BES
Clouzeau, J.	B6.2.1/01	1993c	ACUTE DERMAL IRRITATION IN RABBITS Centre International de Toxicologie, Evreux, France Bayer ES Report-no. 9967 TAL GLP: yes Published: no	yes	BES
Clouzeau, J.	B6.2.2/01	1993d	ACUTE EYE IRRITATION IN RABBITS Centre International de Toxicologie, Evreux, France Bayer ES Report-no. 9968 TAL GLP: yes Published: no	yes	BES
Clouzeau, J.	B6.3/01	1993e	SKIN SENSITIZATION TEST IN GUINEA- PIGS Centre International de Toxicologie, Evreux, France Bayer ES Report-no. 9969 TSG GLP: yes Published: no	yes	BES
Dengler, D.	B7.7.1.1/03	2005	TESTING OF TOXIC EFFECTS OF AQUAPY ON THE SINGLE CELL GREEN ALGA PSEUDOKIRCHNERIELA SUBCAPITATA (FORMERLY SELENASTRUM CAPROCORNUTUM GAB Biotechn. GmbH & IFU Umweltanalytik GmbH, Germany Bayer ES Report-no. 20051085/01-AAPs GLP: yes Published: no	yes	BES
Dobrat, W., Martijn, A.	B4.1/01	1998	PYRETHRUM + PIPERONYL BUTOXIDE + MGK 264 TECHNICAL CONCENTRATES 32+33+345/TK/(M)/- - CIPAC Collaborative International Pesticides Analytical Council, H 1998, 239-242 Report-no. not applicable GLP: no Published: yes	no	-
Kölzer, U.	B7.8.4/01	2005a	ACUTE TOXICITY OF AQUAPY ON EARTHWORMS, EISENIA FETIDA USING AN ARTIFICIAL SOIL TEST GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-NLEf GLP: yes Published: no	yes	BES

Kölzer, U.	B7.8.5/01	2005b	ASSESSMENT OF THE SIDE EFFECTS OF AQUAPY® ON THE ACTIVITY OF THE SOIL MICROFLORA GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-ABMF GLP: yes Published: no	yes	BES
Lucas, J.R., Bowron, M.J.	B5.10.2/01	1994a	AN EVALUATION OF THE BIOLOGICAL ACTIVITY OF AQUAPY (TF2578) WHEN APPLIED OUT OF DOORS AS A ULV SPACE SPRAY AGAINST CAGED MUSCA DOMESTICA AND CULEX QUINQUEFASCIATUS Roussel Uclaf Environmental Health, Berkhamstedt Bayer ES Report-no. GB94-0121 GLP: no Published: no	no	BES
Lucas, J.R., Bowron, M.J.	B5.10.2/03	1994b	AN EVALUATION OF THE BIOLOGICAL PERFORMANCE OF AQUAPY WHEN APPLIED IN A 42 M3 CHAMBER AS A MIST AGAINST HOUSEFLIES AND CLOTHES MOTHS Roussel Uclaf Environmental Health, Berkhamstedt Bayer ES Report-no. GB94-0019 GLP: no Published: no	no	BES
Lucas, J.R., Bowron, M.J.	B5.10.2/04	1994c	THERMAL FOGGING OF AQUAPY (TF2578) AGAINST CULEX QUINQUEFASCIATUS Roussel Uclaf Environmental Health, Berkhamstedt Bayer ES Report-no. GB94-0033 GLP: no Published: no	no	BES
Mooney, M.	B6.6/02	1996a	A FIELD EVALUATION OF OPERATOR EXPOSURE TO AQUAPY SPACE SPRAY IN A TOBACCO WAREHOUSE Hoechst Schering AgrEvo GmbH, Frankfurt am Main, Germany Bayer ES Report-no. GD96-0002 GLP: no Published: no	yes	BES
Mooney, M.	B6.6/03	1996b	A FIELD EVALUATION OF THE PHYSICAL CHARACTERISTICS OF AQUAPY SPACE SPRAY IN A TOBACCO WAREHOUSE Hoechst Schering AgrEvo GmbH, Frankfurt am Main, Germany Bayer ES Report-no. GD96-0001 GLP: no Published: no	yes	BES

Moretto, A.	B6.5/01	1995	PIPERONYL BUTOXIDE - A MONOGRAPH PREPARED BY THE JOINT FAO/WHO MEETING ON PESTICIDES RESIDUES Istituto di Medicina del Lavoro, Padua, Italy Report-no. not applicable GLP: no Published: yes	no	-
Ochieng, C.D.	A3.9/01	1990	DETERMINATION OF PARTITION COEFFICIENT OF PYRETHRINS IN N- OCTANOL/WATER MIXTURE Pyrethrum Board of Kenya, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	PBK
Oellrich, W.	A7.3.1/01	2000	PYRETHRUM EXTRACT, ESTIMATION OF THE PHOTOCHEMICAL OXIDATIVE DEGRADATION GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	PBK
Oellrich, W.	A3.2.1/01	2001	PYRETHRINS, ANNEX II, POINT 2.3.2 HENRY'S LAW CONSTANT GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	PBK
Oellrich, W.	A3.1.1/01	2003	PYRETHRINS, ANNEX II POINT 2.1.1 MELTING POINT, POINT 2.1.2 BOILING POINT, POINT 2.3.1 VAPOUR PRESSURE DETERMINATION GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. 104108-001-01-rev01 GLP: no Published: no	yes	PBK
Repetto-Larsay, M.	B6.3/02	2005	PBO+PYR EW 135+30A G(AQUAPY) (PIPERONYLBUTOXID 139.86 G/L, PYRETHRUM 33.07 G/L) EVALUATION OF THE POTENTIAL DERMAL SENSITIZATION IN THE LOCAL LYMPH NODE ASSAY IN THE MOUSE Bayer CropScience, Sophia Antipolis, France Bayer ES Report-no. SA 05101 GLP: yes Published: no	yes	BES
Reynolds, J.L., Robinson, R.A.	A7.1.3/01	1994	ADSORPTION AND DESORPTION OF [14C]PYRETHRIN 1 IN FOUR SOILS XenoBiotic Laboratories, Inc., Plainsboro, NJ 08536, USA Pyrethrin Joint Venture Report-no. RPT00156 GLP: yes Published: no	yes	PJV

Robinson, R. A.	A7.2.1/01	1994	AEROBIC SOIL METABOLISM OF [14C]PYRETHRIN 1 XenoBiotic Laboratories, Inc., Plainsboro, NJ 08536, USA Pyrethrin Joint Venture Report-no. RPT00204 GLP: yes Published: no	yes	PJV
Robinson, R. A.; Wisocky, M.J.	A7.1.2.2/01	1994	AEROBIC AQUATIC METABOLISM OF [14C]PYRETHRIN 1 XenoBiotic Laboratories, Inc., Plainsboro, NJ 08536, USA Pyrethrin Joint Venture Report-no. RPT00193 GLP: yes Published: no	yes	PJV
Schmid, J.	A3.5/02	1990	WATER SOLUBILITY OF PYRETHRUM EXTRACT ACCORDING TO CIPAC METHOD MT 157.1 BioChem GmbH, Karlsruhe, Germany Kenya Pyrethrum Information Centre Report-no. 947951 GLP: no Published: no	yes	PBK
Selim, S.	A7.1.1.1/01	1995a	HYDROLYSIS OF PYRETHRIN 1 AS A FUNCTION OF PH AT 25°C Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P1092011 GLP: yes Published: no	yes	PJV
Selim, S.	A7.1.1.1.2/01	1995b	AQUEOUS PHOTOLYSIS OF PYRETHRIN 1 Biological Test Center, Irvine, CA 92713-9791-USA Pyrethrin Joint Venture Report-no. P1192006 GLP: yes Published: no	yes	PJV
Stäbler, D.	B7.7.1.1/01	2005a	ACUTE TOXICITY TESTING OF AQUAPY IN RAINBOW TROUT (ONCORHYNCHUS MYKISS) (TELEOSTEI, SALMONIDAE) GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-AAOm GLP: yes Published: no	yes	BES
Stäbler, D.	B7.7.1.1/02	2005b	ASSESSMENT OF TOXIC EFFECTS OF AQUAPY ON DAPHNIA MAGNA USING THE 48 H ACUTE IMMOBILIZATION TEST GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-AADm GLP: yes Published: no	yes	BES

Tolosa, M.	B5.10.2/02	2006	FIELD CONDITIONS EVALUATION OF ADULTICIDE EFFICACY BY TERRESTRIAL APPLICATION OF THE AQUAPY PREPARATION (AQUEOUS EMULSION BASED ON 30 G PYRETHRINS AND 135 G PIPERONYL BUTOXIDE/L) ON MOSQUITO PESTS OCHLEROTATUS CASPIUS, OC. DEDRITUS, AEDES VEXANS (DIPTERA-CULICIDAE) EID Méditerranée Bayer ES Report-no. EID 05049 GLP: no Published: no	yes	BES
Warmers, C.	B7.8.3/01	2005a	AQUAPY ®: TOXICITY TO THE PREDATORY MITE, TYPHLODROMUS PYRI SCHEUTEN (ACARI, PHYTOSEIIDAE) IN THE LABORATORY (RATE RESPONSE TEST) GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-NLTp GLP: yes Published: no	yes	BES
Warmers, C.	B7.8.3/02	2005b	AQUAPY ®: ACUTE TOXICITY TO THE APHID PARASITOID. APHIDIUS RHOPALOSIPHI DE STEFANI PEREZ (HYMENOPTERA, BRACONIDAE) IN THE LABORATORY (RATE RESPONSE TEST) GAB Biotechn. GmbH & GAB Analytik GmbH, Niefern-Öschelbronn Bayer ES Report-no. 20051085/01-NLAp GLP: yes Published: no	yes	BES

Doc II-B (RAID / Baygon Pyrethrum Mat)

Author(s)	Section point/ reference number	Year	Title Testing Facility Owner / Source (where different from owner) Report No GLP or GEP status (where relevant) Published or not	Data protec- tion claimed yes/no	Owner
Anonymous	B5.10.2/02	2000	WORK REQUEST REPORT, STANDARD GLASS CHAMBER TEST WITH MOSQUITOES (CULEX QUINQUEFASCIATUS) - SC JOHNSON & SON, INC., RACINE REPORT-NO. W-145904 GLP: NO PUBLISHED: NO	yes	SCJ
Dobrat, W., Martijn, A.	B4.1/01	1998	PYRETHRUM + PIPERONYL BUTOXIDE + MGK 264 TECHNICAL CONCENTRATES 32+33+345/TK/(M)/- - CIPAC Collaborative International Pesticides Analytical Council, H 1998, 239-242 Report-no. not applicable GLP: no Published: yes	no	-
Eberhardt, R., Wieland, K.	B6.6/01	2006	DETERMINATION OF EVAPORATION- KINETICS OF A SHORT TERM VAPORIZER PYRETHRUM MATS FORMULA NO: 63136- 002 BioGenius GmbH, Monheim, Germany SC Johnson & Son, Inc., Racine Report-no. Mo 3133 GLP: yes Published: no	yes	SCJ
Görgülü, N.	B4.1/02	2006	VALIDATION OF METHOD MV002: DETERMINATION OF PYRETHRUM IN CELLULOSE MATS BioGenius GmbH, Monheim, Germany SC Johnson & Son, Inc., Racine Report-no. Mo2992 GLP: yes Published: no	yes	SCJ
Kristopeit, K.A., Sosa, A., Petersen, J.	B5.10.2/03	1995	WORK REQUEST REPORT NOT APPLICABLE SC JOHNSON & SON, INC., RACINE REPORT-NO. NOT APPLICABLE GLP: NO PUBLISHED: NO	yes	SCJ
Moretto, A.	B6.5/01	1995	PIPERONYL BUTOXIDE - A MONOGRAPH PREPARED BY THE JOINT FAO/WHO MEETING ON PESTICIDES RESIDUES Istituto di Medicina del Lavoro, Padua, Italy Report-no. not applicable GLP: no Published: yes	no	-
Ochieng, C.D.	A3.9/01	1990	DETERMINATION OF PARTITION COEFFICIENT OF PYRETHRINS IN N- OCTANOL/WATER MIXTURE Pyrethrum Board of Kenya, Nakuru, Kenya Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no	yes	PBK
Oellrich, W.	A3.2.1/01	2001	PYRETHRINS, ANNEX II, POINT 2.3.2	yes	PBK

			HENRY'S LAW CONSTANT GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. not stated GLP: no Published: no		
Oellrich, W.	A3.1.1/01	2003	PYRETHRINS, ANNEX II POINT 2.1.1 MELTING POINT, POINT 2.1.2 BOILING POINT, POINT 2.3.1 VAPOUR PRESSURE DETERMINATION GAB Consulting GmbH, Lamstedt, Germany Kenya Pyrethrum Information Centre Report-no. 104108-001-01-rev01 GLP: no Published: no	yes	PBK
Selim, S.	A6.2/01	1995b	PHARMACOKINETICS AND METABOLISM OF PYRETHRIN 1 IN THE RAT Biological Test Center, Irvine, CA 92713- 9791-USA Pyrethrin Joint Venture Report-no. P1092006 GLP: yes Published: no	yes	PJV
Verwey, R.E.	B5.10.2/01	2005	WORK REQUEST REPORT, STANDARD GLASS CHAMBER TEST WITH MOSQUITOES (CULEX QUINQUEFASCIATUS) - SC JOHNSON & SON, INC., RACINE REPORT-NO. W-211682 GLP: NO PUBLISHED: NO	yes	SCJ
Wester, R.C., Bucks D.A.W., Maibach H.I.	A6.2/02	1994	HUMAN IN VIVO PERCUTANEOUS ABSORPTION OF PYRETHRIN AND PIPERONYL BUTOXIDE - Food Chem Toxicol, 32, 51 - 53 Report-no. Not applicable GLP: no Published: yes	no	-

APPENDIX VII: STUDY SUMMARIES

1) CAR studies summary

a) Summary studies for toxicology used for classification:

Doc IIIA Sections 1-6 PYRETHRINS PT18 (BRA, MGK and SCJ)

Doc III-A Sections 1-6 Chrysanthemum PT18 (KPIC)

b) Summary studies for degradation and ecotoxicology used for classification:

Doc IIIA Sections 7-9 PYRETHRINS PT 18_BRA

Doc IIIA Sections 7-9 PYRETHRINS PT 18_KPIC

IIIA7.1.2.1.1 Hein_Moendel 2017

IIIA7.2.1 Fifi 2015 a, b and c

IIIA7.2.1 Hein 2017

IIIA7.2-Perboni

IIIA7.4.1.2 - 5 Pyrethrin metabolites_Daphnia_Mantilaci

IIIA7.4.1.2 - Pyrethrum Extract_6 Pyrethrin esters_Daphnia_Mantilacci

IIIA7.4.3.5.1 - Pyrethrum Extract Pale 50%_Chironomus

IIIA7.4.3.5.1 - Pyrethrum Extract Pale 50%_Chironomus

IIIA7.4.3.5.1 - Pyrethrum Extract Pale 50%_Chironomus

2) Studies included in the report submitted for the EU peer review of active substances used in plant protection products (RAR)

- a) All these studies have been already evaluated under BPR in the present CAR and corresponding docs IIIA above mentioned, hence there is no further evaluation under RAR presented in this appendix, except for the following studies, whose summaries are included in document "Pyrethrins_RAR_Ecotoxicology studies summary not in CAR", attached below:

Test organism	Test substance	Duration, conditions	Toxicological endpoint	Title	Dossier file number
Rat; Sprague-Dawley	¹⁴ C-Pyrethrin 1; 97.81%	Single oral dose by gavage; 3 days observation period	Multiple applied high doses of ¹⁴ C-pyrethrin 1 decrease the excretion in urine and possibly its metabolism rate to chrysanthemum dicarboxylic acid.	Determination of the metabolic profile of ¹⁴ C-Pyrethrin 1 in rats when administered in four different dosing formulations. GLP, Unpublished	CA 5.1.1/03 Limoges, J. 1994
Mouse or rat liver microsomes	Pyrethrin I & Pyrethrin II	1 h incubation	No metabolites were found of toxicological concern.	Pyrethroid metabolism; Microsomal oxidase metabolites of (s)-bioallethrin and the six natural pyrethrins. Not GLP, Unpublished	CA 5.1.2/01 Class, T.J., Ando, T., & Casida, J.E. 1989
Rat, dog and human hepatocytes	[cyclopentenone-2- ¹⁴ C]Pyrethrin 1; 97.9%	0, 30, 60, 120, 180 or 240 min. incubation	There is no strong evidence for any radiolabelled unique or disproportionate human metabolites.	[cyclopentenone-2- ¹⁴ C]Pyrethrin 1: Comparative <i>In Vitro</i> Metabolism Using Rat, Dog and Human Hepatocytes GLP, Unpublished	CA 5.1.2/02 Paul, D. 2020
Rat; Sprague-Dawley	Pyrethrum Extract; 55.99%	Acute oral gavage; 14 days observation period	LD50 : Males : 3.81 g/kg bw. Females : 1.21 g/kg bw.	Acute oral toxicity, LD50-rats (Pyrethrum Extract 55.99%) Not GLP, Unpublished	IIA 5.2.1/03 Costello, B.A. 1986
Rabbit; New Zealand White	Pyrethrum Extract	6 h/day; 5 days consecutively	Slight to moderate erythema, whereas the formation of edema generally does not appear.	Letter: An exploratory 5-day dermal irritation study in New Zealand white rabbits using pyrethrum extract. Not GLP, Unpublished	IIA 5.2.4/02 Schoenig, G.P. 1991
Rat; COBS® CD®	Pyrethrum Extract; 57.57%	10 days consecutively; once/day	Dose levels of 5, 25 and 75 mg/kg bw/day are considered suitable for use in the definitive teratology study	Pyrethrum Extract, Range-Finding Teratology Study in Rats. GLP, Unpublished	CA 5.6.2/01 Schardein, J.L. 1987a
Rabbit; New Zealand White SPF	Pyrethrum Extract; 57.57%	13 days consecutively; once/day	Dose levels of 25, 100 and 250 mg/kg bw/day are considered suitable for use in the definitive teratology study.	Pyrethrum extract Range-Finding Teratology Study in Rabbits. GLP, Unpublished	CA 5.6.2/02 Schardein, J.L. 1987b
Rat; Charles River CD® (Sprague-Dawley)	Pyrethrum Extract; 57.47%	Acute oral gavage; 24 h observation period	A dosing solution containing 10% total pyrethrins and dose levels of 0.04, 0.125, and 0.4 g total pyrethrins/kg bw were selected for males. For females in the definitive acute neurotoxicity study, a dosing solution containing 5% total pyrethrins and dose levels of 0.02, 0.063, and 0.2 g total pyrethrins/kg bw were selected. In addition, 3 to 5 h was determined as the time for the first post-exposure evaluation time.	Peroral (Gavage) Neurotoxicity Probe Study with Pyrethrum Extract in CD® Rats. GLP, Unpublished	CA 5.7.1/01 Hermansky, S. J. & Hurley, J. M. 1993a

<p>Rat; Charles River CrI:CD® (SD)IGS BR</p>	<p>Pyrethrins</p>	<p>Acute oral gavage, 24 h observation period; observed twice daily</p>	<p>Pyrethrins are considered to be the least potent type I pyrethroid (Type 1-T syndrome - non-α-cyano substituent in the 3-phenoxybenzyl alcohol moiety of the molecule) based on FOB data. Pyrethrins are of significantly lower neurotoxicity than all other pyrethroid molecules tested.</p>	<p>Comparative functional observational battery study of twelve commercial pyrethroid insecticides in male rats following acute oral exposure. GLP, Unpublished</p>	<p>CA 5.7.1/03 Weiner, M.L., Nemec, M., Sheets, L., Sargent, D. & Beckenridge, C. 2009</p>
--	-------------------	---	--	---	---

- b) All these studies have been already evaluated under BPR in the present CAR and corresponding docs IIIA above mentioned, hence there is no further evaluation under DAR presented in this appendix, except for the following studies, whose summaries are included in document "Pyrethrins_DAR_Ecotoxicology studies summary not in CAR", attached below:

Test organism	Test substance	Duration, conditions	Toxicological endpoint	Title	Dossier number	file
Fish, acute and chronic						
<i>Danio rerio</i>	Pyrethrum extract (FEK-99)	96h flow through	LC ₅₀ = 19.8 µg /L	Fish acute toxicity test over 96 h under flow through conditions (OECD TG 203, 1992) Acute toxicity of refined pyrethrum concentrate on the zebrafish (<i>Danio rerio</i>) GLP	KCA 8.2.1/04	Teigeler 2013b GAB-034/4-32/A
<i>Gasterosteus aculeatus</i>	Pyrethrum extract (FEK-99)	96h flow through	LC ₅₀ = 10.9 µg /L	Fish acute toxicity test over 96 h under flow through conditions (OECD TG 203, 1992) Acute toxicity of refined pyrethrum concentrate on the three-spined stickleback (<i>Gasterosteus aculeatus</i>)	KCA 8.2.1/05	Teigeler 2013a GAB-034/4-32/G
<i>Cyprinodon variegatus</i>	Pyrethrins TGAI	33 days, flow through	NOEC = 3.5 µg a.s./L	Pyrethrins TGAI - Early Life-Stage Toxicity Test with Sheepshead Minnow (<i>Cyprinodon variegatus</i>) Following OPPTS Draft guideline 850.1400	KCA 8.2.2.1/02	Lee, M.R. 2012
Invertebrates, acute and chronic						
<i>Mysidopsis bahia</i>	Pyrethrum extract (FEK-99)	96h flow through	LC ₅₀ = 1.4 µg /L (measured concentration)	Pyrethrum extract (FEK-99) – acute toxicity to mysid shrimp (<i>Mysidopsis bahia</i>) under flow-through conditions GLP, Unpublished	IIA, 8.3.1.3/01	(B) Machado M.W 1994
<i>Crassostrea virginica</i>	Pyrethrum extract (FEK-99)	96h flow through	EC ₅₀ = 87 µg /L (measured concentration)	Pyrethrum extract (FEK-99) – acute toxicity to eastern oyster (<i>Crassostrea virginica</i>) under flow-through conditions GLP, Unpublished	IIA, 8.3.1.4/01	(B) Dionne E 1994

Chrysanthemum cinerariaefolium, extract from open and mature flowers of *Tanacetum cinerariifolium* obtained with supercritical CO₂

<i>Americamysis bahia</i>	Pyrethrum Stewardship Blend	28 days, flow through	NOEC = 0.25 µg pyrethrins/L	Pyrethrum Stewardship Blend - Life Cycle Toxicity Test with Mysids (<i>Americamysis bahia</i>) Following Draft OPPTS Guideline 850.1350	KCA 8.2.5.2/01 Lee, M.R. 2013
Algae					
<i>Selenastrum capricornutum</i>	Pyrethrum extract	72h static	E _b L ₅₀ > 56,7 mg/L (measured concentration) E _r L ₅₀ > 1,95 mg/L (measured concentration)	Refined pyrethrum extract - Algal growth inhibition assay GLP, Published	IIA, 8.4/01 (B) Jenkins C.A. 2003
<i>Scenedesmus subspicatus</i>	Pyrethrum extract	72h static	EC ₅₀ > 2,32 mg a.s/L (nominal concentration)	Natural Pyrethrum: algal inhibition test GLP, Published	IIA, 8.4/02 (B) Mead C, McKenzie, J. 2003
			EC ₅₀ > 1,27 mg a.s/L (measured concentration)		
Water/sediment organisms					
<i>Hyalella azteca</i>	Pyrethrum Stewardship Blend	96h flow through	LC50 = 0.76 µg/L	Pyrethrum Stewardship Blend - Acute Toxicity to Freshwater Amphipods (<i>Hyalella azteca</i>) Under Flow-Through Conditions	KCA 8.2.8/01 Bradley, M.W., 2013

DAR - ANNEX B.6 Toxicology and metabolism

Active substance: Pyrethrins

- 1) *Determination of the metabolic profile of ¹⁴C-Pyrethrin 1 in rats when administered in four different dosing formulations; Limoges J., 1994; Cross reference IIA 5.1.1/03*

Guidelines:

US EPA 85-1

GLP:

Yes, conducted under GLP/Officially recognised testing facilities.

The study is acceptable as additional data

Executive Summary:

Absorption and excretion patterns and the metabolic profile of Pyrethrin 1 were investigated when administered orally to male rats in the form of four different dosing formulations. Eight male rats were distributed into four groups. Three groups were administered 100 mg ¹⁴C-pyrethrin 1/kg bw in corn oil, or DMSO (5 mL of dosing mixture/kg bw) by gavage. In the fourth group rats were administered orally 4 doses of 400 mg ¹⁴C-pyrethrin 1/kg bw in DMSO at 12-hour intervals.

The mean percent of administered radioactivity found in the urine from the rats in the various dosing regimens in which ¹⁴C-Pyrethrin was administered in corn oil or DMSO ranged from 22.80% to 33.80%. The corresponding value for the group of animals administered ¹⁴C-Pyrethrin 1 as a food slurry was 27.69%. The mean percent of administered radioactivity found in the faeces from the rats in the various dosing regimens which did not involve administration of ¹⁴C-Pyrethrin 1 as a food slurry ranged from 55.01% to 63.20%. The corresponding value for the rats administered ¹⁴C-Pyrethrin 1 as a slurry in food was 38.37%. Except for the group of rats administered ¹⁴C-Pyrethrin 1 as a slurry in food (group 2), total recovery of radioactivity ranged from 84.28% to 90.44%. Total recovery in group 2 was 66.06%.

HPLC analysis of the urine revealed that all groups, regardless of the dosing regimen or vehicle used to administer the dose, produced the same major metabolites. However, when comparing the percent of the total radioactivity represented by chrysanthemum dicarboxylic acid (CDCA) in the urine from each group, the values are 47.52, 42.17, 41.26, and 26.77% for the single dose corn oil, single dose food slurry, single dose DMSO and multiple dose DMSO groups, respectively.

The vehicle (corn oil, food slurry, or DMSO) had little or no influence on the percent dose excreted in the urine or faeces or the percent dose represented by CDCA in the urine. The results indicate that multiple applied high doses of ¹⁴C-Pyrethrin 1 decrease the excretion rate in urine and possibly its metabolism rate to chrysanthemum dicarboxylic acid as well.

Materials and methods:

A. MATERIALS

1. Test Material 1: ¹⁴C-Pyrethrin 1

Purity: 97.81%

Specific activity: 54 mCi/mmol

Specification code: CFQ.7422

Test Material 2: Non-radiolabelled pyrethrin 1

Purity: 98.9%

Specification code: NK9304

Vehicle and/or positive: Mazola corn oil, DMSO, and a food slurry

2. Test Animals

Species: Rat

Strain: Sprague-Dawley (CrI:CD VAF)

Age: 7 weeks

Sex: Males

Weight at dosing: 275-370 g

Source: Charles River Breeding Laboratories, Portage, Michigan.

Acclimation period: A minimum of 7 days

Diet: Purina Certified Rodent Chow #5002 ad libitum

Water: Tap water ad libitum

Housing: Individually housed in stainless steel suspended cages

Environmental Conditions

Temperature: 62°F -71°F

Humidity: 13-62%

Air changes: A minimum of 7 per hour

Photoperiod: 12-hour light/dark cycle

B. STUDY DESIGN AND METHODS

7. Dates of work: 05 November 1993 to 13 December 1994

8. Animal assignment and treatment

Eight male rats (2/group) were administered a single oral dose of 100 mg ¹⁴C-Pyrethrin 1/kg bw in Mazola corn oil, food slurry, or DMSO (5 mL of dosing mixture/kg bw) by gavage. In one group rats were administered 4 doses of 400 mg ¹⁴C-Pyrethrin 1 each/kg bw in DMSO at 12-hour intervals.

Group no.	Males/group	¹⁴ C-Pyrethrin 1 dosage level mg/kg bw	Dose formulation	Route of administration
1	2	100	Corn oil	Oral, single
2	2	100	Food slurry	Oral, single
3	2	100	DMSO	Oral, single
4	2	4 x 400	DMSO	Oral, multiple

9. Administration of the dosing solution

The dosing solutions were delivered by gavage with an intubation needle equipped with a disposable syringe. All doses for groups 1, 2, 3, and 4 were administered on a constant volume basis, i.e. 5 mL of dosing mixture per kg of body weight. The actual amount of compound administered to each rat was measured as a differential between the weight of the syringe before and after dosing.

10. Observations and sampling

Urine, faeces, and water rinses were collected at the following time intervals: 0-24, 24-48, and 48-72 hours after administration and analysed for total radioactivity to determine excretion patterns. The urine was analysed for Pyrethrin 1 and chrysanthemum dicarboxylic acid. Sample combustion was performed by a Harvey Sample Oxidizer, LSC by a Beckmann LS spectrometer and metabolite characterisation in urine by HPLC.

11. Statistics

The mean and standard deviation were used to characterize the data where appropriate.

Results and discussion:

1. Observations

All rats survived, and no signs of toxicity were observed in groups 1 to 3 (single doses of 100 mg/kg ¹⁴C-Pyrethrin 1). In the multiple dose group (4 x 400 mg/kg ¹⁴C-Pyrethrin 1) animals reacted with twitching (during the first 5 hours after the first treatment) and spasms (during the first two hours after the first treatment). Afterwards no clinical signs of toxicity were noted, also not after redosing at hours 12, 24, and 36.

Table 6.1.1/03-1: Mean body weights, compounds and radioactivity of males and females in a preliminary blood kinetics test

Group no.	¹⁴ C-Pyrethrin dosage level mg/kg bw	Dose formulation	Mean body weight (kg) ± SD	Mean administered dose ±SD	
				mg Pyrethrin/kg	Total µCi
1	100	Corn oil	0.302±0.004	94.28±0.13	14.6±0.1
2	100	Food slurry	0.345±0.011	93.38±1.97	12.9±0.1
3	100	DMSO	0.296±0.029	100.59±0.28	14.2±1.3
4	4 x 400 ¹	DMSO	0.350±0.029	1547.08±22.46	28.6±2.0

¹Total of four doses, administered at 12-hour intervals.

2. Excretion

The total recovery of radioactivity excreted in urine and faeces was generally ranged from 84 to 90%, however, in group 2 (100 mg/kg in food slurry) the recovery was less due to incomplete transfer of dose from the syringe to the animals caused by the nature of the food slurry. A higher proportion of radioactivity was excreted in the faeces (38 to 63% of applied radioactivity) than in urine (23 to 34% of applied radioactivity). After single dose applications the excretion rate of radioactivity was highest during the first 48 hours.

Table 6.1.1/03-2: A comparison of total recovery of radioactivity expressed as a cumulative percent of administered radioactivity excreted in urine and faeces from rats administered various dose formulations

Sample	Group 1 (Mean ± SD)	Group 2 (Mean ± SD)	Group 3 (Mean ± SD)	Group 4 (Mean ± SD)
Urine	29.28 ± 1.99	27.69 ± 2.67	33.80 ± 1.51	22.80 ± 0.25
Faeces	55.01 ± 0.50	38.37 ± 5.81	56.64 ± 1.78	63.20 ± 13.72
Total recovery	84.28 ± 2.49	66.06 ± 3.14	90.44 ± 3.29	86.00 ± 13.97

Analysis of urine using HPLC from groups 1, 2, and 3 showed that 1.8 to 13.47% of administered radioactivity were excreted as chrysanthemum dicarboxylic acid (CDCA). However, in the urine from group 4, CDCA represented 4.74% of administered radioactivity.

Table 6.1.1/03-3: Amount of CDCA in the urine from male rats in various dose formulations

Group no.	% of dosed radioactivity present in the urine sample	% of radioactivity in urine as CDCA	% of dosed radioactivity excreted in urine as CDCA
1	28.35	47.52	13.47
2	26.50	42.17	11.18
3	29.81	41.26	12.30
4	17.69	26.77	4.74

Conclusion:

It is concluded that the vehicle (corn oil, food slurry, or DMSO) has little or no influence on the percentage of dose excreted in the urine or faeces or the percentage of dose represented by CDCA in the urine. On the other hand, the results indicate that multiple applied high-doses of ¹⁴C-Pyrethrin 1 decrease the excretion in urine and possibly its metabolism rate to chrysanthemum dicarboxylic acid as well.

A. ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

This was a GLP compliant study. The study report is in broad compliance with OECD 417, although the guidelines are not mentioned in the study report.

Conclusion:

The vehicle had little or no influence on the percentage dose excreted in the urine or faeces or the percentage of dose represented by CDCA in the urine. The results indicate that a lower percentage of the total applied radioactivity was excreted in urine following multiple high doses of ¹⁴C-pyrethrin 1, and the proportion of radioactivity excreted in urine as chrysanthemum dicarboxylic acid was also lower, when compared to percentages of dosed radioactivity excreted in urine and as CDCA, following a single low dose.

B. ASSESSMENT AND CONCLUSION BY RMS

The study was evaluated in the original DAR (2007) and was considered acceptable as additional data.

It is concluded that the vehicle has little or no influence on the percentage of dose excreted in the urine or feces or the percentage of dose represented by CDCA in the urine. On the other hand, the results indicate that multiple applied high doses of ¹⁴C-pyrethrin 1 decrease the excretion in urine and possibly its metabolism rate to chrysanthemum dicarboxylic acid as well.

- 2) *Pyrethroid metabolism; Microsomal oxidase metabolites of (S)-bioallethrin and the six natural pyrethrins; Class T.J., Ando T., and Casida J.E., 1989; Cross reference IIA*
5.1.2/01

Guidelines:

Non-guideline study

GLP:

No, not conducted under GLP/Officially recognised testing facilities.

The study is acceptable

Executive Summary:

Six natural pyrethrins were isolated from purified extract and were analysed together with their synthetic analog (S)-bioallethrin to describe the production of possible metabolites. A complete metabolic profile was developed which shows similarities in the metabolization of natural chrysanthemates (cinerin 1, jasmolin 1, pyrethrin 1) and pyrethrates (cinerin 2, jasmolin 2, pyrethrin 2). The metabolism of the chrysanthemates proceed mainly through oxidative processes while the pyrethrates are metabolised through a combination of hydrolytic and oxidative processes. No metabolites of toxicological concern were found.

Materials and methods:

A. MATERIALS

1. Test Material 1: (S)-bioallethrin (synthetic analogue)

Test Material 2: Cinerin 1

Test Material 3: Jasmolin 1

Test Material 4: Pyrethrin 1

Test Material 5: Cinerin 2

Test Material 6: Jasmolin 2

Test Material 7: Pyrethrin 2

2. Mammalian metabolic system

System: Mouse or rat liver microsomes

3. Test animals

Species: Rat

Strain: Albino

Sex: Male

Weight at dosing: Not stated

Source: Not stated

Acclimation period: Not stated

Diet: Not stated

Water: Not stated

Housing: Not stated

Environmental Conditions

Temperature: Not stated

Humidity: Not stated

Air changes: Not stated

Photoperiod: Not stated

B. STUDY DESIGN AND METHODS

1. Dates of work: 19 November 1992 to 19 December 1995

2. Method

The pyrethrum extract constituents and (S)-bioallethrin were analysed by high resolution gas chromatography (HRGC) with an electron capture detector (ECD) of HRGC-chemical ionization (CI)-mass spectrometry (MS). These methods are applied here to the rethrin metabolites after appropriate derivations. This study compares the metabolic fate of (S)-bioallethrin and the six natural pyrethrins in mouse and rat liver microsomal oxidase systems and of (S)-bioallethrin in rats.

a) Formation and analysis of microsomal metabolites

The substrate (0.1 μmol) was incubated with mouse or rat liver microsomes (0.1, 0.3, 1.0, 3.0 mg protein) and NADPH (0 or 2.4 μmol) in phosphate buffer (0.1 M, pH 7.4, 2 mL) for 1 hour at 37 °C. Following extraction of the aqueous and organic phases, analysis was conducted using HRGC coupled to CI-MS. The amount of substrate recovered was determined by HRGC-ECD relative to the internal standard. Recoveries of the chrysanthemates were 100 \pm 10% at 0.3 mg microsomal protein in the absence of NADPH. The extent of metabolism of the chrysanthemates was calculated from the loss of substrate with NADPH.

The microsomal metabolism of the pyrethrates in the presence and absence of PSCP (phenyl saligenin cyclic phosphonate, a potent esterase inhibitor) was also investigated. The extent of metabolism was determined by substrate loss for incubations with microsomes compared with no microsomes. Another investigation compared the extent of metabolism of the (E)-8' vs. (Z)-8' isomers of cinerin 1, jasmolin 1 and pyrethrin 1 incubated as a mixture with bioallethrin each at 0.014 μmol per incubation with 0.3 and 1.0 mg mouse microsomal protein and NADPH.

b) Formation and analysis of urinary metabolites

Male albino rats (not specified) were treated with (S)-bioallethrin orally by stomach tube (250 mg/kg followed after 6 hours with 500 mg/kg) or intraperitoneally (12.5 mg/kg followed after 6 hours with 25 mg/kg) using DMSO as the vehicle. No signs of toxicity were observed. Urine was collected 0-6 hours after the second treatments were analysed for free and conjugated urinary metabolites.

3. Abbreviations for chemicals

The abbreviations ol, al and acid. refer to alcohols, aldehydes and carboxylic acids, respectively. Trimethylsilyl ethers are designated as TMS, ethyl esters as Et, epoxides as epoxy, diols from hydrolysis of epoxides as dihydrodiol, and their TMS derivatives as dihydro (TMS)₂. Designations such as 5/6-ol and 6'/10'/11'-ol indicates that the available information does not differentiate among the specified positions. "PI" and "PII" refer to rearranged products shown in Figure 6.1.2/01-1 and 5.1.2/01-2.

Results and discussion:

1. Microsomal metabolites

Metabolism of the chrysanthemates (S)-bioallethrin, cinerin 1, jasmolin 1, and pyrethrin 1 by NADPH-dependent oxidases of mouse liver microsomes yields 13-18 metabolites in each case oxidized at the methyl, methylene, and alkenyl substituents to form alcohols, aldehydes, carboxylic acids, epoxides and dihydrodiols. Rat microsomes are more specific than rat mouse microsomes in hydroxylating the (E)-methyl substituent of the 2-methylpropenyl moiety compared with other molecular sites.

Table 6.1.2/01-1: Extent of oxidative metabolism by mouse and rat liver microsomes for (S)-bioallethrin (A), the six natural pyrethrins and the (E)-8'-isomers of C1, J1 and P1

Designation	NADPH-dependent loss of substrate, % ^a , at indicated level of microsomal protein (mg)						
	Mouse				Rat		
	0.1	0.3	1.0	3.0	0.3	1.0	3.0
A	25	50 ^b	80	95	20	50	60
C1c(C2)d	15	45	90 (93) ^d	-	-	75	-
J1c(J2)d	10	35	80 (98) ^d	90	30	50	60
P1c(P2)d	10	40	80 (99) ^d	-	-	50	-

A (S)-bioallethrin

^a Mean of two experiments differing by <10%


^b Substrate loss (%) with primary metabolites used as substrates: A-7.8-epoxy [(7S)-isomer] 35, [(7R)-isomer] 10; A-7'-ol 35 with no diastereomer difference; A-10-ol 25; A-10-al 30.

^c Mean o experiments of pyrethrates with NADPH±PSCP, metabolism of C2, J2 and P2 in the absence of NADPH is 25, 55 and 55% respectively, with PSCP and 55, 65 and 75%, resectively, without PSCP.

C=cinerin, J=jasmolin, P=pyrethrin

Figure 6.1.2/01-1: Partial metabolic pathways for (S)-bioallethrin (A), cinerin 1 (C1), jasmolin 1 (J1), pyrethrin


1 (P1) in mouse and rat liver microsomal oxidase systems and of (S)-bioallethrin in rats *in vivo*. Additional metabolites not designated as structures arise from other combinations of modifications in the acid and alcohol moieties. Brackets designate compounds tentatively identified from SeO₂-oxidation and indicated but not established as metabolites



2. Urinary metabolites

Metabolites in the urine of allethrin-treated rats include compounds modified in both the 2-methylpropenyl and allyl moieties as free carboxylic acids and glucuronides. The pyrethrates cinerin 2, jasmolin 2, and pyrethrin 2 undergo microsomal hydrolysis of the methoxycarbonyl group and oxidation of the butenyl, pentenyl, and pentadienyl substituents to alcohols, epoxides, and dihydrodiols.

Figure 6.1.2/01-2: Partial metabolic pathways for cinerin 2 (CII), jasmolin 2I (JII) and pyrethrin 2I (PII) in mouse liver microsomal oxidase systems. Although not observed, 10', 11'-epoxy-PI is included as a likely intermediate



Conclusion:

A complete metabolic profile was developed which shows similarities in the metabolism of natural chrysanthemates and of pyrethrates. The metabolism of the chrysanthemates proceeds mainly through oxidative processes while the pyrethrates are metabolised through a combination of hydrolytic and oxidative processes. No metabolites were found of toxicological concern.

A. ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

The study report is in broad compliance with OECD 417, although the guidelines are not mentioned in the study report. The study provides reasonable information to predict the absorption, distribution, and excretion of the natural Pyrethrins.

Conclusion:

Agreement with the conclusions of the study authors.

B. ASSESSMENT AND CONCLUSION BY RMS

The study was evaluated in the original DAR (2007) and was considered acceptable.

A complete metabolic profile was developed which shows similarities in the metabolism of natural chrysanthemates and of pyrethrates. The metabolism of the chrysanthemates proceeds mainly through oxidative processes while the pyrethrates are metabolised through a combination of hydrolytic and oxidative processes. No metabolites were found of toxicological concern.

3) [cyclopentenone-2-¹⁴C]Pyrethrin 1: Comparative In Vitro Metabolism Using Rat, Dog and Human Hepatocytes; Paul D., 2020; Cross reference IIA 5.1.2/02

Guidelines:

No detailed test guidelines for the conduct of comparative *in vitro* metabolism studies are currently available. The data requirement is based on the Commission Regulation (EU) No 283/2013, 5.1.1, in accordance with Regulation (EC) No 1107/2009.

GLP:

Yes, UK Good Laboratory Practice Regulations (Statutory Instrument 1999 No. 3106, as amended by Statutory Instrument 2004 No. 994); OECD Principles of Good Laboratory Practice (as revised in 1997), ENV/MC/CHEM(98)17; EC Commission Directive 2004/10/EC

The study is acceptable

Executive Summary:

The purpose of this study is to compare the metabolism of [cyclopentenone-2-¹⁴C]Pyrethrin 1 also written as [¹⁴C]Pyrethrin 1 using rat, dog and human hepatocytes.

[¹⁴C]Pyrethrin 1 (10 µM) was incubated (at 37 ± 1 °C) with either rat, dog or human hepatocytes (0.5 x 10⁶ viable cells/mL for all species) for 0, 30, 60, 120, 180 or 240 minutes. Incubations in the absence of hepatocytes were also conducted with [¹⁴C]Pyrethrin 1 for 0 and 240 minutes, to check the stability under incubation conditions. Incubation samples were analysed by HPLC with on-line radioactive monitoring. The proportions of metabolites produced and parent [¹⁴C]Pyrethrin 1 were quantified.

A summary of the metabolites detected is presented in Summary Table 6.1.2/02-1.

Table 6.1.2/02-1 Metabolites detected following incubation of Pyrethrin 1 (10 µM) with Rat, Dog and Human Hepatocytes after 30 to 240 minutes

Metabolite fraction designation	R1	R2	R3	R4	R5	R6	R7	R8	R9	R10	R11	R12	R13	R14
Rat	+	++	+	-	++	+	++	+	+	++	+	-	+	+
Dog	+	+	-	-	++	+	++	+	+	++	+	++	+	+
Human	-	-	-	-	+	+	++	-	+	++	+	+	+	+
Metabolite fraction designation	R15	R16	R17	R18	R19	R20	R21	R22	R23	R24	R25	R26	R27	[¹⁴ C]Pyrethrin 1 metabolism (%)†
Rat	+	-	-	-	+	+	++	+	-	+	+	-	-	> 99
Dog	-	+	++	+	++	-	++	-	+	+	+	+	+	96.1
Human	-	-	+	-	++	-	++	+	+	-	+	+	-	84.6

+ Present at between ≥ 1% and < 5% sample radioactivity in species

++ Present at ≥ 5% sample radioactivity in species

- Not detected at < 1% or above sample radioactivity in species

† Consumption of parent at 240 minutes compared to time zero

* present in rat and dog at earlier timepoints.

Materials and test methods:

A. MATERIALS

1. Test substance [cyclopentenone-2-¹⁴C]Pyrethrin 1

2. Chemical name (IUPAC) [(1S)-2-methyl-4-oxo-3-[(2Z)-penta-2,4-dienyl]cyclopent-2-en-1-yl]

(1R,3R)-2,2-dimethyl-3-(2-methylprop-1-enyl)cyclopropane-1-carboxylate

3. CAS number 121-21-1
4. Molecular formula C₂₁H₂₈O₃
5. Molecular weight 328.45
6. Lot number LY13TX/NP/01
7. Specific activity 2.11 GBq/mmol
8. Radioactive concentration 0.345 MBq/mL
9. Radiochemical purity 97.9% (HPLC)
10. Physical form Solution in Acetonitrile
11. Storage conditions -10 °C to -30 °C, under nitrogen and protected from light

B. STUDY DESIGN AND METHODS

All cryopreserved hepatocytes were obtained from BioIVT (formerly Celsis IVT and Bioreclamation IVT) and delivered stored frozen in liquid nitrogen. Details of the hepatocytes used in this study are as follows:

Species	Strain	Gender	Batch number	Number of donors	Number of vials used
Rat	Sprague Dawley	Male	VRO	9	2
Dog	Beagle	Male	SDS	3	3
Human	Not applicable	Mixed	QGS	10	2

All animal hepatocytes were supplied as a pool of male donors. Each vial used contained at least 5 million cells. Human hepatocytes were supplied as a mixed-gender pool.

1. Incubation of [¹⁴C]Pyrethrin 1 with Rat, Dog, and Human Cryopreserved Hepatocytes

Incubations of [¹⁴C]Pyrethrin 1 were conducted with rat, dog and human cryopreserved hepatocytes as follows:

- Concentration: 10 µM
- Incubation times: 0, 30, 60, 120, 180 and 240 mins
- Incubation temperature: 37 ± 1 °C
- Number of replicates: Two
- Cell concentration: 0.5 × 10⁶ viable cells/mL incubation medium
- Volume of incubation medium: Either 5.0 mL (hepatocyte-containing samples or 2.0 mL (no hepatocytes samples)

The incubation components were mixed together in well-plates (6-well format) for each sample as follows:

- Supplemented Williams' Medium E containing 2.5 × 10⁶ viable cells in 4950 µL
- [¹⁴C]Pyrethrin 1 (50 µL of 1 mM solution in acetonitrile) for hepatocyte-containing samples.

Following the final addition of [¹⁴C]Pyrethrin 1, 0 minutes samples were terminated immediately. The well-plates were then placed on a tilting mini rocker-shaker in a temperature-controlled incubator (set at 37 °C) to commence the incubation.

The following control incubations were also conducted in parallel:

- Incubation of [¹⁴C]Pyrethrin 1 for 0 or 240 minutes in the absence of hepatocytes (duplicate incubations at 10 μM),
- Positive control samples incubating 7-ethoxy[3-¹⁴C]coumarin ([¹⁴C]7EC) at a concentration of 25 μM in single replicate for 0, 30, 60, 120, 180 and duplicate for 240 minutes.

At the end of requisite incubation period, an aliquot (0.5 mL) was removed from each incubate and transferred to an aliquot (0.5 mL) of chilled acetonitrile to stop the reaction. Each sample was stored on Ice and then treated using an ultrasonic bath for approximately 10 minutes to fully disrupt the cells. The samples were then processed as described in section

2. Analysis of [¹⁴C]Pyrethrin 1 and Metabolites

i. Preparation of Reference Standards

Parent non-radiolabelled Pyrethrin 1 which was previously supplied at a concentration of 0.1 mg/mL in cyclohexane was dried down and then reconstituted in acetonitrile (0.5 mL) to remove the storage solvent. Reference standard, non-radiolabelled pyrethrolone was prepared by weighing 4 μL (equivalent to 4.12 mg) and dissolving in 1 mL of DMSO to give a pyrethrolone concentration of 4 mg/mL. The pyrethrolone solution was then diluted by combining an aliquot (100 μL) of the solution with 900 μL of purified water to obtain a concentration of 0.4 mg/mL.

ii. Processing of [¹⁴C]Pyrethrin 1 Incubation Samples

Prior to analysis, 250 μL of each terminated incubation sample was centrifuged at 14,000 rpm (approximately 18,600 × g, 15 minutes, 5 °C ± 3°C) to sediment the cell debris. Duplicate 10 μL aliquots were removed from the supernatant for LSC and 5 mL Ultima Gold added. The resulting supernatant (200 μL) was diluted with Pyrethrin 1 standard (0.1 mg/mL) 2.5 μL, purified water (600 μL) and metabolite reference standard pyrethrolone (0.4 mg/mL) 2.5 μL in a glass HPLC vial. The mixture was vortexed thoroughly, prior to transfer duplicate 25 μL aliquots were removed for LSC with 5 mL Ultima Gold added. The HPLC vial was transferred to an HPLC autosampler for injection (500 μL) into the HPLC. Samples were stored at approximately -70 ± 10°C prior to and on completion of analysis.

3. Characterisation of Isolated Hepatocytes

i. Determination of Initial Cell Viability by the Trypan Blue Exclusion Test

A solution of 0.08% (w/v) trypan blue was prepared by diluting 0.4% (w/v) trypan blue 1:4 (v/v) with supplemented Williams' Medium E. For each species, an aliquot (25 μL) of the initial hepatocyte cell suspension was mixed with an aliquot (50 μL) of 0.08% (w/v) trypan blue solution and an aliquot (175 μL) of supplemented Williams' Medium E. Each mixture was then loaded into a C-Chip haemocytometer (chamber depth 0.1 mm) and the number of viable and non-viable hepatocytes was determined in at least two 1 mm² grid areas. The total cell count for each 1 mm² area was multiplied by an appropriate scaling factor to give the total number of cells per mL and the mean value was calculated.

ii. Incubation of 7-Ethoxy[3-¹⁴C]coumarin (7-EC) with Hepatocytes

In parallel incubations, positive control incubations were conducted with ([¹⁴C]7-EC) as substrate. Incubations comprised of supplemented Williams' Medium, to give a final concentration of 7-EC of 25 μM in total volumes of either 5 mL or 4 mL. The final dimethylformamide (DMF) concentration was not greater than 1% (v/v). The incubations were performed in well-plates (6-well format) on a tilting mini rocker-shaker in a temperature-controlled incubator (set at 37 °C). At the end of the requisite incubation period, an aliquot (1 mL) was removed from each incubate and transferred to an aliquot (1 mL) of chilled acetonitrile to stop the reaction. Each sample was stored on ice, then treated using an ultrasonic bath for approximately 10 minutes to fully disrupt the cells. All samples were then

stored at ca. -20 °C until sample processing. Following storage at ca. -20 °C, the samples were transferred to clean labelled microcentrifuge tubes and centrifuged at 18,000 x g, at ca 4 °C for 15 minutes to pellet the cell debris. The resulting supernatants from 7-EC samples were transferred to clean microcentrifuge tubes and concentrated to dryness under nitrogen gas. The dried residues were reconstituted in 40 mM ammonium formate (pH 5) and approximately half of each of the resulting supernatants was transferred to separate HPLC vials for direct injection into the HPLC system. A further portion of each solution (450 µL) was transferred to a clean micro-centrifuge tube along with a solution of β-glucuronidase enzyme (50 µL of a 40,000 units/mL solution, Type H1 from Helix pomatia also containing sulfatase activity). This mixture was incubated for 1 hour at 37 °C. Samples were then transferred to HPLC vials for direct injection into the HPLC system. Positive controls for β-glucuronidase and sulfatase enzyme activities were determined by the production of free phenolphthalein from phenolphthalein glucuronic acid and p-nitrocatechol from p-nitrocatechol sulfate, respectively, upon the addition of 1M sodium hydroxide after incubation (1 hour at 37 °C). Both the phenolphthalein glucuronic acid and p-nitrocatechol sulfate were prepared in 40 mM ammonium formate (pH 5).

Results and discussion:

For each chromatogram obtained from the analysis of samples from the incubations of [¹⁴C]Pyrethrin 1 with rat, dog and human hepatocytes by HPLC, regions of radioactivity were assigned as metabolite fraction identities of R1 to R27 (each representing either a separated radioactive component or components where complete resolution could not be attained) and parent test item.

Not all metabolite fractions were present in every sample and some were only present in trace quantities. Regions of radioactivity that contained ≥ 5% of sample radioactivity were considered major (Whalley *et al.*, 2017), whilst those that contained 1 to 4.9% of sample radioactivity were considered minor. Assigned regions of radioactivity that contained < 1% of sample radioactivity were considered below the limit of quantification.

1. Pyrethrin 1 Stability

The mean proportions of [¹⁴C]Pyrethrin 1 (10 µM) remaining in the absence of hepatocytes were 97.3 and 86.7% following 0 and 240 minutes incubation, respectively. No notable breakdown products were noted over this period.

2. Rat Hepatocytes

The mean proportions of [¹⁴C]Pyrethrin 1 present in samples incubated with rat hepatocytes were 96.1, 28.8, 9.2 and 1.1% following 0, 30, 60 and 120 minutes incubation, respectively and BLQ thereafter. The mean extents of metabolism of [¹⁴C] Pyrethrin 1 expressed relative to time zero were 67.3, 86.9, 95, > 99 and > 99% after 30, 60, 120, 180 and 240 minutes incubation, respectively. All metabolites R1 to R27 inclusive measured BLQ at time zero. In these samples up to 19 metabolites (R1-R3, R5-R11, R13-R15, R19-R22, R24 and R25) were detected above the limit of quantification at various time points during the incubation.

Metabolites R2, R5, R7, R10 and R21 were formed as major metabolites (≥ 5% of sample radioactivity) by rat hepatocytes and accounted for a mean sample radioactivity respectively of 8.5, 23.0, 10.5, 10.1 and 12.8% after 240 minutes incubation.

The other prominent metabolite fractions (minor metabolites) detected following incubation with rat hepatocytes, R1, R3, R6, R8, R9, R11, R13, R14 and R20 accounted for a maximum mean of 4.9% of the total sample radioactivity following 240 minutes incubation. The mean proportions of the metabolites detected above the limit of quantification generally increased with incubation time.

3. Dog Hepatocytes

The mean proportions of [¹⁴C]Pyrethrin 1 present in samples incubated with dog hepatocytes were 97.9, 55.2, 31.1, 12.3, 5.0 and 1.8% following 0, 30, 60, 120, 180 and 240 minutes incubation, respectively. The mean extents of metabolism of [¹⁴C]Pyrethrin 1 expressed relative to time zero were 42.7, 66.8, 85.6, 92.9 and 96.1% after 30, 60, 120, 180 and 240 minutes incubation, respectively. All metabolites R1 to R27 inclusive for both replicate samples measured at BLQ at time zero. In these samples up to 22 metabolites (R1, R2, R5-R14, R16-R19, R21, R23-R27) were detected above the limit of quantification at various time points during the incubation. Metabolites R5, R7, R10, R12, R17, R19 and R21 were formed as major metabolites ($\geq 5\%$ of sample radioactivity) by dog hepatocytes and accounted for a mean sample radioactivity respectively of 8.8, 8.4, 8.5, 5.3, 7.9, 15.8 and 5.4% after 240 minutes incubation. The mean proportions of the metabolites detected above the limit of quantification generally increased with incubation time.

4. Human Hepatocytes

The mean proportions of [¹⁴C]Pyrethrin 1 present in samples incubated with human hepatocytes were 97.9, 77.3, 57.5, 28.4, 23.4 and 13.3% following 0, 30, 60, 120, 180 and 240 minutes incubation, respectively. The mean extents of metabolism of [¹⁴C]Pyrethrin 1 expressed relative to time zero were 20.6, 40.4, 69.5, 74.5 and 84.6% after 30, 60, 120, 180 and 240 minutes incubation, respectively. All metabolites R1 to R27 inclusive for both replicate samples measured at BLQ at time zero. In these samples up to 16 metabolites (R5-R7, R9-R14, R17, R19, R21-R23, R25 and R26) were detected above the limit of quantification at various time points during the incubation. Metabolites R7, R10, R19 and R21 were formed as major metabolites ($\geq 5\%$ of sample radioactivity) by human hepatocytes and accounted for a mean sample radioactivity respectively of 7.0, 5.7, 12.8 and 20.8% after 240 minutes incubation. The proportions of the metabolites detected above the limit of quantification generally increased with incubation time.

5. Comparison of Rat, Dog and Human Metabolism

Following the incubation of [¹⁴C]Pyrethrin 1 with rat, dog and human hepatocytes over 240 minutes, the most prominent human metabolites were R7, R10, R19 and R21 (major). R10 seemed to corresponded to the reference standard Pyrethrolone chromatographically which is a major metabolite of [¹⁴C]Pyrethrin 1. These metabolites were also observed in rat or dog hepatocytes. Metabolite R5 was a prominent major metabolite in the rat and dog, respectively, but a minor metabolite in the human. To a lesser degree, R8 was a minor metabolite in the rat and dog, respectively, and below the limit of quantification in human. Metabolite R2 was also a prominent major metabolite in the rat, a minor metabolite in the dog and below the limit of quantification in the human. Metabolite R19 was a prominent major metabolite in the dog and human, but present in the rat as a minor metabolite following 30 and 60 minutes incubation, respectively. R6, R9, R11, R13 and R14 were minor metabolites in all species. Taken together, there was no evidence for any radiolabelled unique human metabolites because metabolites identified following incubation with human hepatocytes were also detected in at least one animal species. In all species, the Phase I metabolite 7-hydroxycoumarin (7-HC) formed was subsequently conjugated to varying extents. The changes in the levels of the Phase I metabolite 7-HC following deconjugation indicated that all hepatocytes were metabolically viable and were capable of integrated Phase I/II metabolism under the incubation conditions used on this study. Therefore, the results generated for the incubation of these hepatocytes with [¹⁴C]Pyrethrin 1 are considered to be valid.

Conclusion:

Extensive metabolism of [¹⁴C]Pyrethrin 1-10 μM was observed across all species investigated with up to 27 metabolites (designated R1-R27) detected along with parent [¹⁴C]Pyrethrin 1. [¹⁴C]Pyrethrin 1 was metabolised to a slightly lesser extent in human hepatocytes than rat and dog hepatocytes in terms of the number of metabolites generated.

Following the incubation of [¹⁴C]Pyrethrin 1 with rat, dog and human hepatocytes over 240 minutes, the most prominent metabolites in human hepatocytes were R7, R10 and R21 (major, >5%). Based on the UV chromatogram R10 corresponds to the reference standard Pyrethrolone which is a major metabolite of [¹⁴C]Pyrethrin 1.

Taken together, there is no strong evidence for any radiolabelled unique or disproportionate human metabolites.

A. ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

Unaudited interim report for the comparative in vitro metabolism study using rat, dog and human hepatocytes, with [cyclopentenone-2-¹⁴C]Pyrethrin 1. The final report is to be issued in April 2020.

Conclusion:

Taken together, there is no strong evidence for any radiolabelled unique or disproportionate human metabolites.

B. Assessment and conclusion by RMS

The study was submitted for the renewal procedure and was considered acceptable.

There is no strong evidence for any radiolabelled unique or disproportionate human metabolites.

4) (A) *Acute oral toxicity, LD50 – RATS (Pyrethrum Extract 55.99 %); Costello B.A., 1986;*

Cross reference IIA 5.2.1/03

Guidelines:

Similar to 40 CFR, Sect. 163.81-1, Fed. Reg. , August 22, 1978; modified in accordance with revised EPA Pesticide Assessment Guidelines, Nov. 1982.

GLP:

No

GLP was not compulsory at the time when the study was performed (1986)

The study is acceptable

Executive Summary:

A total of 50 (25 males and 25 females) albino rats, weighing 140-244 g, were administered a single dose of Pyrethrum Extract (55.99 % purity) by gavage. Following administration, the animals were allowed food and water ad libitum for the 14 day observation period. Animals were observed frequently on the day of dosing, twice per day on weekdays and once per day on weekends and holidays for signs of toxicity and mortality. Individual weights were recorded on the day of dosing, weekly thereafter and prior to sacrifice. After euthanasia, gross necropsies were performed on all animals.

Oral LD ₅₀	males	= 3.81 g/kg bw
	femal	= 1.21 g/kg bw

Mortality started at dosages of 2510 mg/kg (females) and 1000 mg/kg (males). Females were more susceptible to Pyrethrum Extract than males. The main signs for intoxication were increased responsiveness to external stimuli, tremors, salivation and ruffled appearance. Gross pathology showed congested lungs (at males from 3980 mg/kg onwards, at females from 1260 mg/kg onwards).

Material and methods:

A. MATERIALS

1. Test Material: Pyrethrum Extract
Description: Yellow liquid
Lot/Batch #: FNB 86-2-18A
Purity: 55.99 % Pyrethrins
CAS #: As given in section 2
Stability of test compound: Not determined
2. Vehicle and/or positive control: No vehicle
3. Test animals
Species: Rat
Strain: Outbred Sprague-Dawley
Age: Young adult
Weight at dosing: 140-244 g
Source: Buckshire Corp. Perkasie, PA 18944
Acclimation period: At least 5 days
Diet: Wayne Rodent-Blox 8604, *ad libitum*
Water: Tap water, *ad libitum*
Housing: Animals were housed in groups of 3-5 animals by sex in labeled stainless steel suspended cages.

Environmental conditions

- | | |
|--------------|---|
| Temperature: | 21.1°C – 26.7°C |
| Humidity: | Relative humidity in %: 55 ± 25 |
| Air changes: | Not recorded |
| Photoperiod: | Alternating 12-hour light and dark cycles, artificial fluorescent light |

B. STUDY DESIGN AND METHODS:

1. In life dates: 09 June to 25 June 1986
2. Animal assignment and treatment

A total of 50 (25 males and 25 females) albino rats were administered a single dose of Pyrethrum Extract by gavage. Following administration, the animals were allowed food and water *ad libitum* for the 14 day observation period. Animals were observed frequently on the day of dosing, twice per day on weekdays and once per day on weekends and holidays for signs of toxicity and mortality. Individual weights were recorded on the day of dosing, weekly thereafter and prior to sacrifice. After euthanasia, gross necropsies were performed on all animals.

Table B.6. 14: Doses, mortality / animals treated

Dose (mg/kg bw)	Males	Females	Combined
630	-*	1/5	1/5
1000	0/5	1/5	1/10
1260	-	4/5	4/5
1580	0/5	3/5	3/10
2510	1/5	4/5	5/10
3980	2/5	-	2/5
6310	5/5	-	5/5

* not tested

3. Statistics

The data did not warrant statistical analysis.

Findings:

A. MORTALITY

Details are provided in Table B.6. 14. No mortalities occurred at 1580 mg / kg for male rats.

Oral LD₅₀ for males = 3.81 g/kg bw

for females = 1.28 g/kg bw

B. CLINICAL OBSERVATIONS

Females were more susceptible to Pyrethrum Extract (55,99 %) than males. The main signs for intoxication were increased responsiveness to external stimuli, tremors, salivation and ruffled appearance.

C. BODY WEIGHT

All animals had gained weight 7 and 14 days following dosis.

D. NECROPSY

Gross pathology showed congested lungs (at males from 3980 mg/kg onwards, at females from 1260 mg/kg onwards)

E. DEFICIENCIES

The study was not conducted under GLP, since GLP was not compulsory in 1986, when the study was performed. It was conducted in accordance with generally accepted scientific principles. A NOAL could not be derived from this study.

Conlcusions:

The oral LD₅₀ of the test compound was determined to be 3.92 g/kg for males and 1.28 g/kg for females. In accordance with the provisions of Council Directive 67/548/EEC, classification is not required. (Costello, B.A. 1986)

- 5) (A) Letter: An exploratory 5-day dermal irritation study in New Zealand white rabbits using Pyrethrum Extract; Schoenig G.P., 1991; Cross reference IIA 5.2.4/02

Guidelines:

None

GLP:

No

The study is not acceptable due to absent information about many parameters as purity of the test article, batch number, temperature, humidity and others.

Executive Summary:

Three groups of two rabbits (one male and one female) were used to assess the dermal irritation of Pyrethrum Extract upon repeated application at concentrations of 25%, 50% and 75% in corn oil. The test solution (4.0 mL/kg) was applied once daily for five consecutive days to the shaven intact skin on the back of each rabbit. After a skin contact period of 6 hours per day, the test article was wiped with tepid tap water and dried with disposable terrycloth (paper) towelling.

Animals were observed twice daily for mortality and overt toxicity and once daily for dermal observations using the Draize method. No mortalities and no test substance related clinical signs of systemic intoxication or influences on body weights were observed. Daily dermal application of 4 mL of 25, 50 or 75 % Pyrethrum Extract in corn oil for five continuous days caused time- and dosage-dependant formation of slight to moderate erythema to the rabbit.

Material and methods:

A. MATERIALS

1. Test Material:	Pyrethrum extract
Description:	Not stated
Lot/Batch #:	Not stated
Purity:	Not stated
CAS #:	As given in section 2
Stability of test compound:	Not determined
2. Vehicle and/or positive control:	corn oil
3. Test animals	
Species:	Rabbit
Strain:	New Zealand White
Age:	Young adult,
Weight at dosing:	Males: 2868 g-3119 g Females: 2565 g-3425 g
Source:	[REDACTED]
	Not recorded
Acclimation period:	Not recorded
Diet:	Not recorded
Water:	Animals were individually housed in labeled cages
Housing:	with perforated floors
Environmental conditions:	
Temperature:	No specification
Humidity:	Not reported
Air changes:	Not reported
Photoperiod:	Not reported

B. STUDY DESIGN AND METHODS

1. In life dates: 18 June to 23 June 1991

2. Animal assignment and treatment

Three groups of 1 male and 1 female rabbit each were used to assess the dermal irritation of the test substance at dosage concentrations of 25 %, 50 % and 75 % in corn oil (w/v). The test substance was applied once daily for five consecutive days to the shaven intact skin on the back of each rabbit, at a constant dose volume of 4.0 mL/kg. The

test substance was covered with a porous gauze dressing fixed with non-irritating tape throughout a 6-h daily exposure period, afterwards the test site was washed with tepid water and dried with disposable terrycloth (paper) towelling.

Animals were observed twice per day for mortality and overt toxicity and daily for irritation effects using the Draize method. Individual body weights were recorded at initiation and at study termination (after 5 days).

Findings:

No mortalities and no test substance related clinical signs of systemic intoxication. No test substance related influence on body weights.

The initiation scores are summarized in Table B.6. 18.

During the first 2 days no erythema were observed except one very slight erythema in the highest concentration group. From day 3 onwards a time and dose dependant increase becomes obvious, from no erythema (grade 0) at one male of the lowest concentration group on day 3 to moderate erythema (grade 3) at four animals of the higher concentration groups on day 5.

No edema were observed at the animals of all test substance concentrations during the first 4 days. One male of the highest concentration group showed very slight edema formation (grade 1) on days 5 and 6.

Daily dermal application of 4 mL 25, 50 or 75 % Pyrethrum Extract in corn oil for five continuous days causes time- and dosage-dependant formation of slight to moderate erythema to the rabbit, whereas the formation of edema generally does not appear (except very slight edema at one male during the end of the test period).

Table B.6. 18: Individual and mean skin irritation scores according to the Draize scheme

Test Substance Conc. %	Erythema and Eschar Formation						Edema Formation					
	25 %		50 %		75 %		25 %		50 %		75 %	
days	M	F	M	F	M	F	M	F	M	F	M	F
1	0	0	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	1	0	0	0	0	0	0
3	0	1	1	1	2	2	0	0	0	0	0	0
4	1	2	2	2	2	3	0	0	0	0	1	0
5	2	2	3	3	3	3	0	0	0	0	1	0
6	2	2	3	2	3	3						
average score, Draize scores (0 to maximum 4)	0.83	1.2	1.5	1.3	1.67	2.0	0	0	0	0	0	0.3

Conclusions:

Pyrethrum Extract was non-irritant to rabbit skin. On the basis of this study Pyrethrins do not warrant classification as being irritating to the skin. (Schoening, P 1991)

6) *Pyrethrum Extract, Range-Finding Teratology Study in Rats; Schardein J.L., 1979a;*

Cross reference IIA 5.6.2/01

Guidelines:

None

GLP:

EPA GLP practice standards May 2, 1984

The study is acceptable as supportive information

Executive Summary:

Mated Charles River COBS® CD® female rats, assigned to one control and five treatment groups of 5 animals each, were used in this range-finding study to determine dose levels of pyrethrins for a definitive teratology study. Dose levels of 37.5, 75, 150, 300 and 600 mg/kg bw/day in terms of actual pyrethrin content were administered orally by gavage as a single daily dose on days 6 through 15 of gestation at a volume of 3 mL/kg. The control group received the vehicle only, 0.5% methylcellulose, on a comparable regime. Uterine examinations were performed on all surviving females on gestation day 20.

Treatment-related maternal toxicity, in terms of mortality and convulsions and/or tremors, occurred at 150, 300 and 600 mg/kg bw/day. There was no treatment-related maternal mortality at 75 mg/kg bw/day, although tremors were observed in this group. No treatment-related clinical signs were observed at 37.5 mg/kg bw/day. No dose-related differences from controls were noted in the mean number of viable foetuses, the mean postimplantation loss and the mean number of implantations and corpora lutea of the dams treated with 37.5, 75 and 150 mg/kg/day.

Based on these results, dose levels of 5, 25 and 75 mg/kg bw/day were selected for use in a definitive teratology study.

Materials and methods:

A. MATERIALS

1. Test Material: Pyrethrum Extract Task Force Blend

Description: Amber liquid

Lot/Batch number: #FEK-99, 08/05/86

Purity: 57.574%

CAS#: Not reported

Stability of test compound: 24-hour stability of the test suspension was established

2. Vehicle: 0.5% methylcellulose

3. Test Animals

Species: Rat

Strain: COBS® CD®

Age/weight at receipt: 70-day old, weight at mating 220 to 225 g

Source: Charles River Laboratories, Inc., Portage, Michigan, USA.

Housing: Individually in suspended wire-mesh cages

Acclimatisation period: 12 days

Diet: Purina® Certified Rodent Chow® #5002

Water: Tap Water *ad libitum*

Environmental conditions: Temperature: 22 to 44°C

Humidity: 38 to 92%

12 hours of fluorescent light and 12 hours of darkness provided per 24 hours

B. STUDY DESIGN

1. In-life dates: 3 to 26 September 1986

2. Mating procedure

At the end of the acclimation period, all animals were weighed and subjected to a detailed physical examination. At this time, animals considered suitable for study were cohabited with stock males used exclusively for this purpose. One female and one male rat of the same strain and source were placed together for mating. The occurrence of copulation was determined by daily inspection for a copulatory plug. The day evidence of mating was detected was designated day 0 of gestation and the female was returned to an individual cage, assigned a permanent animal number and properly identified by ear tag.

3. Animal assignment

Mated females were consecutively assigned in a block design to one control and five treatment groups consisting of 5 rats each by the following procedures. The order in which the mated females were assigned corresponded to the day of the copulatory plug was observed and the order in which the animal appeared on the breeding record. The first mated female on the breeding record was assigned to the first group, the second mated female assigned to the second group; all remaining animals were assigned in this manner until the required number of mated females had been placed into each group.

Table 6.6.2/01-1 Number of animals and treatment groups

Group no.	Females/group	Treatment	Dose level (mg/kg bw/day)	Dose concentration (mg/mL)	Dose volume (mL/kg bw)
1	5	Vehicle	0	0	3
2	5	Actual pyrethrins	37.5	12.5	3
3	5		75	25	3
4	5		150	50	3
5	5		300	100	3
6	5		600	200	3

4. Dosage preparation and analysis

The appropriate amounts of pyrethrum extract for each group was suspended in the vehicle, 0.5% methylcellulose, using a tissue homogenizer. A correction factor of 1.7369 was used in test article calculations to adjust for the purity of the test article. The required amount of additional vehicle was then added to this suspension and the resulting mixture was shaken by hand. The test article was prepared daily (with two exceptions) at concentrations to permit the administration of dose levels of 37.5, 75, 150, 300 and 600 mg/kg bw/day at a dose volume of 3 mL/kg. The suspensions administered on occasions designated for sample collection were prepared the day before to allow time for analysis prior to administration. The 300 and 600 mg/kg bw/day dosing solutions were no longer prepared after the surviving females in these groups were humanely killed due to the occurrence of excessive toxicity at these dose levels.

Replicate samples were collected on approximately the first and last days of administration from the top, middle and bottom of the dosing suspensions and analysed for homogeneity and concentration of the active ingredient (pyrethrin) of the test article. An additional sample of each dosing suspension was collected on these same days and stored frozen for possible future analysis.

5. Dose administration

The test article was administered as a single daily dose in vehicle 0.5% methylcellulose, on days 6 through 15 of gestation. The prepared test article suspensions were stirred using a magnetic stir bar and stir plate during administration. Animals were dosed by oral gavage using a 1 cc glass syringe and 16-gauge, 7.6 cm long stainless-steel dosing needle. A constant dose volume of 3 mL/kg body weight was used, adjusted to the most recent body weight. The control group received the vehicle only on a comparable regimen.

C. METHODS

1. Maternal observations

Throughout the study, the animals were observed twice daily for mortality and overt changes in appearance and behaviour. The presence and duration of clinical signs of toxicity were recorded once daily on days 6 through 20 of gestation. Females not surviving to scheduled sacrifice were necropsied in an attempt to determine the cause of death.

Due to the occurrence of excessive toxicity in the dosed females at 300 and 600 mg/kg bw/day, all surviving animals at these doses were humanely killed on gestation days 5 or 6. No necropsy examinations were conducted on these animals.

2. Body weight

Individual body weights were recorded on gestation days 0, 6, 9, 12, 16 and 20.

3. *Post Mortem* Investigations

On gestation day 20 all females were sacrificed by carbon dioxide inhalation. Immediately following sacrifice, the uterus and ovaries were exposed by an abdominal incision and the number and location of the viable and non-viable fetuses, early and late resorptions and the total number of implantation and corpora lutea were recorded. The abdominal and thoracic cavities and organs of the dams were examined for grossly evident morphological changes and the carcasses discarded. Uteri from females that appeared non-gravid were opened and placed in 10% ammonium sulphide solution for detections of implantations.

4. Statistical analyses

Statistical analysis of data was not conducted.

Results and discussion:

Concentration analysis results

Analytical results of the periodic test article: 0.5% methylcellulose suspensions, approximately the first and last days of administration, contained mean concentrations ranging from 91 to 102% of target levels.

Homogeneity and stability results

Stability analysis showed that pyrethrum extract was stable for up to 24 hours in 0.5% methylcellulose suspensions under normal laboratory conditions. Homogeneity evaluation of pyrethrin in 0.5% methylcellulose showed mean homogeneity assay values for all groups analysed ranging from 91 to 102% of target concentration. These results indicate that a homogeneous test suspension was produced.

A. OBSERVATIONS

1. Mortality and maternal clinical signs of toxicity

Two, three and two females died post-dose on gestation day 6 or 7 in the 150, 300 and 600 mg/kg/day dose groups, respectively. Convulsions, and tremors in some instances, were observed post-dose on the day of death for these females. As a result of the toxicity observed in these groups, the surviving females in the 300 and 600 mg/kg bw/day groups were humanely sacrificed.

Tremors and/or convulsions were observed post-dose for two dams at 75 mg/kg bw/day, and in two rats which survived at 150 mg/kg bw/day. Wet red material around the nose and eyes

was also noted postdose for one of these females at 150 mg/kg bw/day. There were no treatment-related differences in the appearance or behaviour of the rats at 37.5 mg/kg bw/day.

Table 6.6.2/01-2 Summary of clinical observations

Observation	Pyrethrin Dose level (mg/kg bw/day)					
	0	37.5	75	150	300	600
Number of animals observed	5	5	5	5	4 ^a	3 ^b
No visible abnormalities	4	3	3	3	3	2
Died				2	3	2
Killed (due to excessive toxicity at this dose)					2 ^a	3 ^b
Hair loss	1	2	2		1	1
Tremors (post-dose)			2	3	1	
Convulsions (post-dose)				4	3	3
Material around nose and eyes, wet, red (post-dose)				1		

^a One female sacrificed gestation day 5, prior to initiation of antemortem observations

^b Two females sacrificed gestation day 5, prior to initiation of antemortem observations

B. BODY WEIGHT AND BODY WEIGHT GAIN

As a result of mortality and resultant early termination, the body weight gain of the 300 and 600 mg/kg bw/day dams could not be assessed during treatment. There were no treatment-related differences in body weight of the dams at 37.5, 75 and 150 mg/kg bw/day, during the treatment period (gestation days 6 through 15) or during the entire gestation period (gestation days 0 to 20).

Table 6.6.2.1-3 Intergroup comparison of group mean bodyweight gain (g)

Day	Pyrethrins (mg/kg bw/day) ^{a, b}					
	0	37.5	75	150	300	600
0 to 6	34	36	39	37	40	29
6 to 9	6	7	4	7	--	--
9 to 12	14	16	17	13	--	--
12 to 16	22	24	21	22	--	--
16 to 20	62	52	55	57	--	--
6 to 15 ^c (treatment period)	42	48	41	42	--	--
0 to 20	138	137	135	133	--	--

^a Non-gravid animals were not included in calculation of means

^b Values represent the mean of the individual changes in maternal body weight for these intervals

^c Gestation day 16 values were utilised to reflect the entire 10-day treatment period

-- Not applicable

C. FOOD CONSUMPTION

Food consumption was not recorded.

D. SACRIFICE AND PATHOLOGY

1. Gross pathology

No gross lesions were present at gross necropsy for the animals that died. No gross lesions were observed for the control and treated animals at necropsy.

2. Caesarean section data

Gestation day 20 uterine observation data was not available for the females in the 300 and 600 mg/kg/day dosage groups, since these groups were terminated early in the gestation period.

No dose-related differences were noted in the mean number of viable foetuses, the mean postimplantation loss and the mean number of implantations and corpora lutea of dams at 37.5, 75 and 150 mg/kg bw/day, in comparison to the control values.

Table 6.6.2/01-4 Caesarean section observations for all pregnant females

Observation	Pyrethrins (mg/kg bw/day)					
	0	37.5	75	150	300	600
Animals Assigned (Mated)	5	5	5	5	5	5
Animals that were gravid	5	5	4	5	1	0
Animal that died	0	0	0	2	3	2
Nongravid	--	--	--	0	2	2
Gravid	--	--	--	2	1	0
Humanely killed (not examined)	--	--	--	--	2	3
Animals examined at uterine examination	5	5	5	3	0	0
Nongravid	0	0	1	0	--	--
Gravid	5	5	4	3	--	--
Dams with resorptions only	0	0	0	0	--	--
Dams with viable foetuses	5	5	4	3	--	--
Viable foetuses/dam	14.4	12.6	13.0	14.0	--	--
Post-implantation loss/dam	0.4	2.2	2.0	0.3	--	--
Total implantations /dam	14.8	14.8	15.0	14.3	--	--
Corpora lutea/dam	15.8	16.0	16.0	15.0	--	--
Group mean preimplantation loss (%) ^a	6.3	7.5	6.3	4.4	--	--
Group mean post-implantation loss (%) ^b	2.7	14.9	13.3	2.3	--	--

^a Total number of corpora lutea- Total number of implantations x 100

Total Number of corpora lutea

^b Total number of implantations- Total number of viable foetuses x 100

Total Number of implantations

-- Not applicable

Conclusion:

Treatment-related maternal toxicity, in terms of convulsions and/or tremors and mortality, occurred at 150, 300 and 600 mg/kg bw/day. No excessive dose-related maternal mortality was evident at 75 mg/kg bw/day, although tremors were observed in this group. No treatment related clinical signs were observed at 37.5 mg/kg bw/day.

Based on these results, dose levels of 5, 25 and 75 mg/kg bw/day were selected for use in the definitive teratology study.

A. ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

This non-guideline range-finding study is supportive information.

Conclusion:

Provides justification for the dose levels selected in the definitive study.

B. ASSESSMENT AND CONCLUSION BY RMS

This non-guideline range-finding study in rat is considered appropriate for the intended aim of determine a dose levels of pyrethrins for a definitive teratology study.

On the basis of the results from this study, dose levels of 5, 25 and 75 mg/kg bw/day are considered suitable for use in the definitive teratology study.

Please note that for this study analytical methods for current standards are o longer considered acceptable (please refers to Vol 3 CA B.5). Anyway this study is considered fit for pourpose, pending on conclusion on storage stability issues.

- 7) *Pyrethrum extract Range-Finding Teratology Study in Rabbits; Schardein J.L., 1979b;*
Cross reference IIA 5.6.2/02

Guidelines:

None

GLP:

EPA GLP practice standards May 2, 1984

The study is acceptable as supportive information

Executive Summary:

Inseminated female New Zealand White SPF rabbits randomly assigned to one control and five treatment groups of 5 animals each were used in this range-finding study to determine dose levels of pyrethrins for a teratology study. Dose levels of 37.7, 75, 150, 300 and 600 mg/kg bw/day in terms of actual pyrethrin content were administered orally by gavage as a single daily dose on days 7 through 19 of gestation at a volume of 3 mL/kg. The control group received the vehicle only, 0.5% methylcellulose, on a comparable regimen. Uterine examinations were performed on all surviving females on gestation day 29. Treatment-related maternal toxicity in terms of mortality, tremors/convulsions and weight loss and foetotoxicity in terms of high post-implantation loss was observed at 600 mg/kg bw/day. Maternal toxicity in terms of weight loss during the treatment period and tremors were evident at 300 mg/kg bw/day. No clear treatment-related effects were observed at 37.5, 75 or 150 mg/kg bw/day. Based on these results, dose levels of 25, 100 and 250 mg/kg bw/day were selected for the definitive teratology study.

Materials and methods:

A. MATERIALS

1. Test Material: Pyrethrum Extract Task Force Blend

Description: Amber liquid

Lot/Batch number: #FEK-99, 08/05/86

Purity: 57.574%

CAS#: Not reported

Stability of test compound: 24-hour stability of the test suspension was established

2. Vehicle: 0.5% methylcellulose

3. Test Animals

Species: Rabbit

Strain: New Zealand White SPF

Age/weight at receipt: 4-month old/3160 to 4132 g on gestation day 0

Source: Hazleton Research Animals, Denver, Pennsylvania, USA.

Housing: Individually in wire cages with Deotized animal cage board waste pan litters

Acclimatisation period: 48 days

Diet: Purina® Certified Rabbit Chow® #5322 *ad libitum*

Water: Tap Water *ad libitum*

Environmental conditions: Temperature: 23 to 26°C

Humidity: 36 to 78%

12 hours of fluorescent light and 12 hours of darkness provided per 24 hours

B. STUDY DESIGN

1. In-life dates: 2 September to 1 October 1986.

2. Mating procedure

The females were approximately 5 ½ months old at the time of insemination and weighed between 3160 and 4132 g on gestation day 0. Approximately three weeks prior to insemination, the does were superovulated by an injection of 50 U.S.P. units of human chorionic gonadotropin via a marginal ear vein. Semen was collected from five proven male rabbits of the same breed and source. Semen was collected using an artificial vagina and the gelatinous plug was removed from the ejaculate. The semen was immediately evaluated for motility and was used for insemination only if the motility was 60% or greater, as assessed subjectively. The ejaculate was diluted with 3.0 mL of 0.9% sodium chloride for injection, U.S.P., at 33-37°C and 0.3 mL of this dilute semen was introduced into the anterior vagina of the female using an insemination pipette. Immediately after insemination, ovulation was induced by an injection of 100 U.S.P. units of human chorionic gonadotropin into the marginal ear vein. Semen from one male was used to inseminate an equal number of females in each group. Insemination procedures were performed on one day. The day of insemination was designated as day 0 of gestation.

3. Animal assignment

At the end of the acclimation period, all animals were weighed and subjected to a detailed physical examination. Animals considered suitable for study were randomly assigned to one control group and five treatment groups of 5 rabbits each using a weight stratified randomisation procedure. Bartlett's test for homogeneity of variance was applied to the groups; the groups were judged to be homogeneous.

Table 6.6.2/02-1 Number of animals and treatment groups

Group no.	Females/group	Treatment	Dose level (mg/kg bw/day)	Dose concentration (mg/mL)	Dose volumen (mL/kg bw)
1	5	Vehicle	0	0	3
2	5	Actual pyrethrins	37.5	12.5	3
3	5		75	25	3
4	5		150	50	3
5	5		300	100	3
6	5		600	200	3

4. Dosage preparation and analysis

The appropriate amount of pyrethrum extract for each group was suspended in the vehicle (0.5% methylcellulose) using a tissue homogeniser. A correction factor of 1.7369 was used in test article preparation calculations to adjust for the purity of the test article (57.574%). The required amount of additional vehicle was then added to this suspension and the resulting mixture was shaken by hand. The test article was prepared daily (with two exceptions) at concentrations to permit the administration of dose levels of 37.5, 75, 150, 300 and 600 mg/kg bw/day at a dose volume of 3 mL/kg. The suspensions administered on gestation days 7 and 15 were designated to be analysed for pyrethrin concentration and were prepared the day before to allow time for analysis prior to administration.

Stability analysis

Prior to initiation of the test article administration period, the 24-hour stability of pyrethrins in suspensions was assessed. In addition, a sample of each dosing solution was collected and frozen for possible future analysis.

Homogeneity and pyrethrin concentration analysis

Replicate samples were collected from preparations administered on gestation days 7 and 15, from the top, middle and bottom of the dosing suspensions and analysed for the homogeneity and concentration. An additional sample of each dosing suspension was collected at the same time and frozen for possible future analysis.

5. Dose administration

The dosing preparations were administered as a single daily dose on days 7 through 19 of gestation. The dosing preparation suspensions were stirred using a magnetic stir bar and stir plate during administration. The dosing preparations were administered by intragastric intubation using 35 cc disposable plastic (first two days of administration) or glass (remainder of administration period) syringes and 12-gauge, 15 cm long curved stainless-steel dosing needles. The control group received the vehicle only on a comparable regimen at a volume of 3 mL/kg. Individual doses were determined from the most recently recorded individual body weights.

C. METHODS

1. Maternal observations

Throughout the study, the animals were observed twice daily for mortality and overt changes in appearance and behaviour. The presence and duration of clinical signs of toxicity were recorded once daily on days 7 through 29 of gestation, although detailed observations recorded on days 20 through 29 were not required by protocol. Females not surviving to the scheduled sacrifice were necropsied in an attempt to determine the cause of death. Any female showing signs of abortion or premature delivery was sacrificed and necropsied on the day such evidence was observed and the aborted tissue was examined and discarded.

2. Body weight

Individual maternal body weights were recorded on gestation days 0, 7, 13, 20, 24 and 29.

3. *Post Mortem* Investigations

On gestation day 29, all surviving females were sacrificed by an injection of sodium pentobarbital via a marginal ear vein. Immediately following sacrifice, the uterus and ovaries

were exposed by an abdominal incision and the number and location of the viable and non-viable fetuses, early and late resorptions and the number of total implantations and corpora lutea were recorded. The abdominal and thoracic cavities and organs of the dams were examined for grossly evident morphological changes and the carcasses discarded. Maternal tissues were preserved in neutral buffered 10% formalin for possible histopathological examination as deemed necessary by the gross findings. Uteri from females that appeared non-gravid were opened and placed in 10% ammonium sulphide solution for detections of implantations.

Results and discussion:

Test article stability in 0.5% methylcellulose

Stability analysis showed that pyrethrum extract was stable for up to 24 hours in 0.5% methylcellulose suspensions under normal laboratory conditions.

Concentration analysis results

Test suspensions prepared for administration on gestation days 8 and 19 at 25, 100 and 250 mg/kg bw/day contained 90 to 104% of the target concentrations (means of assay values).

Homogeneity and stability results

Pyrethrin concentrations found in samples taken from the top, middle and bottom of the 25, 100 and 250 mg/kg bw/day dosing suspensions prepared for administration on gestation days 8 and 19 were within $\pm 5\%$ of the mean values. The dosing suspensions were homogeneous. Stability was also confirmed, as average levels at 24 hours ranged from 94 to 100% of the average zero hours concentrations.

E. OBSERVATIONS

1. Mortality

One doe died at 37.5 mg/kg bw/day and two does died at 600 mg/kg bw/day.

2. Maternal clinical signs of toxicity

No visible abnormalities were noted prior to death on gestation day 20 for the doe at 37.5 mg/kg bw/day. Findings for the does that dies on gestation days 18 and 20 at 600 mg/kg bw/day included tremors, convulsions, laboured breathing, hunched posture, excessive salivation and/or a wet, yellow-stained area around the mouth and nose. With the exception of the last finding, these pharmacotoxic signs generally occurred post-dose.

One doe at 37.5 mg/kg bw/day and one doe at 300 mg/kg bw/day aborted on days 22 and 25 of gestation, respectively. One doe at 600 mg/kg bw/day delivered on gestation day 29. Antemortem findings noted for these animals included emaciation (600 mg/kg bw/day), tremors (300 and 600 mg/kg bw/day), nasal discharge post-dose (300 mg/kg bw/day) and dark brown liquid in the cage pan (37.5 mg/kg bw/day).

Tremors, and convulsions in some cases, occurred in all the females at 600 mg/kg bw/day. Tremors also occurred in one female at 300 mg/kg bw/day. There were no treatment-related differences in appearance and behaviour of does at 37.5, 75 and 150 mg/kg bw/day.

F. BODY WEIGHT AND BODY WEIGHT GAIN

Dose-related mean body weight losses occurred for does at 300 and 600 mg/kg bw/day during the treatment period (gestation days 7 through 19). Similarly weight loss was noted for does at 600 mg/kg bw/day over the entire gestation period. Reduced weight gain relative to control was observed in dams at 300 mg/kg bw/day over the entire gestation period.

There were no consistent treatment-related differences in body weight gain in does at 37.5, 75 and 150 mg/kg bw/day, in comparison to control.

Table 6.6.2/02-2 Intergroup comparison of group mean bodyweight gain (g)

Day	Pyrethrins (mg/kg bw/day) ^a					
	0	37.5	75	150	300	600
0 to 7	157	157	172	206	203	141
7 to 13	47	105	-24	31	-94	-299
13 to 20	114	117	20	105	-121	-325
20 to 24	-21	-89	47	0	26	115
24 to 29	-39	-185	10	22	41	60
7 to 19 ^b (treatment period)	161	219	-4	136	-215	-514
0 to 29	258	266	226	365	160	-56

^a Values represent the mean of the individual changes in maternal body weight for these intervals

^b Gestation day 20 values were utilised to reflect the entire 10-day treatment period

G. FOOD CONSUMPTION

Food consumption was not recorded.

H. SACRIFICE AND PATHOLOGY

1. Gross pathology

There were no noteworthy maternal necropsy findings. There were no apparent treatment-related gross pathological changes in animals sacrificed at the end of the study, and for those does that aborted or those that died.

2. Caesarean section data

There was an increase in mean post-implantation loss and a resultant decrease in the mean number of viable foetuses at 600 mg/kg bw/day, relative to control. Slight increases in post-implantation loss were also recorded at 150 and 300 mg/kg bw/day; however, it was unclear if the magnitude of increase was of biological significance. No treatment-related differences in the values for the uterine examination parameters were noted at 37.5 and 75 mg/kg bw/day.

Table 6.6.2/02-3 Caesarean section observations for all pregnant females

Observation	Pyrethrins (mg/kg bw/day)					
	0	37.5	75	150	300	600
Animals Assigned (Mated)	5	5	5	5	5	5
Animals that were gravid	5	5	5	5	5	4
Animals that died	0	1	0	0	0	2
Nongravid	-	0	-	-	-	0
Gravid	-	1	-	-	-	2
Animal that aborted/delivered	0	1	0	0	1	1
Animals examined at uterine examination	5	3	5	5	4	2
Nongravid	0	0	0	0	0	1
Gravid	5	3	5	5	4	1

Does with rerptions only	0	0	0	0	0	0
Does with viable foetuses	5	3	5	5	4	1
Viable foetuses/doe	6.8	8.0	8.6	7.0	7.0	5.0
Post-implantation loss/doe	0.2	0.0	0.2	1.4	1.0	4.0
Total implantations/doe	7.0	8.0	8.8	8.4	8.0	9.0
Corpora lutea/doe	12.0	12.3	11.2	12.4	11.5	15.0
Group mean pre-implantation loss (%) ^a	41.7	35.1	21.4	32.3	30.4	40.0
Group mean post-implantation loss (%) ^b	2.9	0.0	2.3	16.7	12.5	44.4

^a Total number of corpora lutea- Total number of implantations x100

Total Number of corpora lutea

^b Total number of implantations- Total number of viable foetuses x100

Total Number of implantations

- Nota applicable

Conclusion:

Treatment-related maternal toxicity in terms of mortality, tremors/convulsions and weight loss and foetotoxicity in terms of high post-implantation loss were observed at 600 mg/kg bw/day. Maternal toxicity in terms of weight loss during the treatment period and tremors was evident at 300 mg/kg bw/day. No clear treatment-related effects were observed at 37.5, 75 or 150 mg/kg bw/day.

Based on these results, dose levels of 25, 100 and 250 mg/kg bw/day were considered appropriate for the definitive teratology study in rabbits.

A. ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

This non-guideline range-finding study is supportive information.

Conclusion:

Provides justification for the dose levels selected in the definitive study.

B. ASSESSMENT AND CONCLUSION BY RMS

This non-guideline range-finding study in rabbit is considered appropriate for the intended aim of determine a dose levels of pyrethrins for a definitive teratology study in rabbits.

On the basis of the results from this study, dose levels of 25, 100 and 250 mg/kg bw/day are considered suitable for use in the definitive teratology study.

Please note that for this study analytical methods for current standards are no longer considered acceptable (please refers to Vol 3 CA B.5). Anyway this study is considered fit for purpose, pending on conclusion on storage stability issues.

- 8) *Peroral (Gavage) Neurotoxicity Probe Study with Pyrethrum Extract in CD® Rats;*
Hermansky S.J. & Hurley J.M., 1993a; Cross reference IIA 5.7.1/01

Guidelines:

None

GLP:

Yes, conducted under GLP/Officially recognised testing facilities

The study is acceptable as supportive information

Executive Summary:

The probe study was divided into 2 phases. In Phase I, 1 male and 1 female rat/dose, were dosed by oral gavage, over a broad range of doses to generate preliminary information with which to select doses for more extensive testing in Phase II. In Phase I, clinical signs of toxicity were monitored hourly for several hours after dosing. In Phase II, groups of 2-4 rats/sex/group were employed and clinical signs of toxicity were recorded and specific evaluations for tremors, arousal state, and gait were made at least hourly for several hours and 24 hours after dosing.

Phase I

Males were treated with a solution of pyrethrum extract in corn oil, prepared to contain 25% total pyrethrins, at dose levels of 0.2, 0.4, 0.8, 1.4, and 2.5 g total pyrethrins/kg body weight. Animals treated with 1.4 and 2.5 g/kg, died within 5.5 hours of dosing. Tremors were observed at all dose levels within 2 hours after dosing. Tremors persisted through the 10-hour observation period for the male at 0.8 g/kg and through the 4-hour observation for the male at 0.4 g/kg. Tremors for the male at 0.2 g/kg were not observed after the 2-hour observation. Other observations recorded prior to death for the 1.4 and 2.5 g/kg animals included lying on stomach, labored respiration, salivation, urine stains, and prostration.

A dosing solution containing 10% total pyrethrins was selected for females in Phase I based on the results obtained for males during Phase I and LD50 information supplied by the Sponsor indicating that females were more sensitive to the test substance than males. Females were treated at dose levels of 0.05, 0.1, 0.2, 0.4, and 0.8 g total pyrethrins/kg body weight. Dose levels of 0.2, 0.4, and 0.8 g/kg resulted in death within 5 hours of dosing, in addition, tremors were recorded at all dose levels within 2 hours of dosing. Tremors persisted through the 3-hour observation period for the female at 0.1 g/kg. Tremors for the female at 0.05 g/kg were not observed after the 2-hour observation. Other observations recorded prior to death for the 0.4 and 0.8 g/kg animals were like those observed for males.

Phase II

Males were administered a solution of pyrethrum extract in corn oil at a concentration of 10% total pyrethrins at dose levels of 0.0, 0.04, 0.1, 0.2, and 0.4 g total pyrethrins/kg body weight. Animals in the control group were administered corn oil at a volume equal to the volume of test solution administered to the highest dose group. No mortality was observed. Tremors were observed in males dosed at 0.4 g/kg from the 2- through the 5-hour observation periods. No tremors were recorded for males in any other dose group at any observation period or in any animals at 24 hours after dosing. Salivation was observed for 1 animal in the 0.2 g/kg group. A possible decrease in the level of arousal compared to control was observed in the 3 highest dose groups at the 5- and 24-hour observation periods. The time to peak effect in the 0.4 g/kg group was estimated to be between 3 and 5 hours after administration.

In Phase II, females initially were treated with a solution of pyrethrum extract in corn oil at a concentration of 2.5% total pyrethrins at dose levels of 0.0, 0.025, 0.05, 0.1, and 0.15 g total pyrethrins/kg body weight. Animals in the control group were administered corn oil at a volume equal to the volume of test solution administered to the highest dose group. No females died at these dose levels. Clinical signs were limited to the 0.15 g/kg group and included fine tremors for 1 female 6 and 7 hours after treatment and altered gait for another

female at the 2-hour through 24-hour observation periods. Because of the limited number of clinical signs observed in these females, 12 additional females (2/dose level) were treated with a solution of pyrethrum extract in corn oil at a concentration of 5% total pyrethrins at dose levels of 0.0, 0.025, 0.05, 0.1, 0.15, and 0.2 g total pyrethrins/kg body weight. Animals in the control group were administered corn oil at a volume equal to the volume of test solution administered to the highest dose group. Tremors were observed for both female animals dosed at 0.2 g/kg and 1 of 2 females in each of the 0.15 and 0.1 g/kg groups. These tremors were generally described as fine and were observed for all 3 of these groups during the 3 and 4-hour observation periods and for the highest dose group during the 2, 5, and 6-hour observation periods. One animal from the 0.2 g/kg group was also described as hyperactive and as having perinasal encrustation. The time to peak effect was estimated to be between 3 and 5 hours after dose administration. Based upon the results of this study, dose levels of 0.04, 0.125, and 0.4 g total pyrethrins/kg body weight were selected for males in the definitive acute neurotoxicity study. These doses were to be administered as a solution of pyrethrum extract prepared in corn oil to contain 10% total pyrethrins. For females, dose levels of 0.02, 0.063, and 0.2 g total pyrethrins/kg body weight were selected for use in the definitive acute neurotoxicity study. These doses were to be administered as a solution of pyrethrum extract prepared in corn oil to contain 5% total pyrethrins. Based upon the time course of effects observed in this study, 3 to 5 hours was selected as the time for the first post-exposure evaluation in the definitive acute neurotoxicity study.

Material and methods:

A. MATERIALS

1. Test Material: Pyrethrum Extract

Description: Dark viscous liquid

Lot/Batch: LS92-37, Task Force Blend FEK-99

Purity: 57.467% (w/w)

2. Vehicle: Mazola corn oil

3. Test Animals

Species: Rat

Strain: Charles River CD[®] (Sprague-Dawley)

Age: 35 days

Sex: Male and female

Weight at dosing:

Source: Charles River Laboratories (Portage, MI)

Acclimation period: Approximately 15 days

Diet: Ground, certified Rodent chow[®] #5002 (Purina Mills, Inc.) *ad libitum*

Water: Tap water *ad libitum*

Housing: Individually housed

Environmental Conditions: Temperature: 66-77 °F
Humidity: 40-70%

Photoperiod: 12-hour light/dark cycle

B. STUDY DESIGN

1. In-life dates: March 1992 to April 1992

2. Animal assignment and treatment

Forty male and 40 female CD rats were assigned to the study. In phase I, 1 male and 1 female rat was assigned to each treatment group. For the first part of phase II, groups of 4 rats/sex/group were assigned to treatment and control groups using a weight stratified computerized randomization procedure. The animals selected for use in phase I of the study were randomly selected from the remaining animals after this randomisation procedure. For the second part of phase II, female animals not selected for either Phase I or phase II were assigned to 5 dose groups so that each group contained an animal with a high body weight and an animal with a low body weight. Two female control animals from the first part of phase II were selected for use as a control group in the second part of phase II.

In phase I, males were treated with a 25% total pyrethrins solution in corn oil, at dose levels of 0.2, 0.4, 0.8, 1.4, and 2.5 g total pyrethrins/kg bw and females were treated with corn oil solution that contained 10% total pyrethrins at dose levels between 0.05, 0.1, 0.2, 0.4 and 0.8 g total pyrethrins/ kg bw.

In phase II, a dosing solution containing 10% total pyrethrins in corn oil and dose levels of 0.0, 0.04, 0.1, 0.2, and 0.4 g total pyrethrins/kg bw were selected for males. Females initially were treated with a solution of 2.5% pyrethrins in corn oil and doses 0.0, 0.025, 0.05, 0.1, and 0.15 g total pyrethrins/kg bw. Due to the limited number of clinical signs observed in these females, 12 additional females (2/dose level) were treated 5% total pyrethrins in corn oil at dose levels of 0.0, 0.025, 0.05, 0.1, 0.15, and 0.2 g. Animals of the control group were treated with corn oil at a volume equivalent to that administered to the rats in the high dose treatment groups.

Table 6.7.1/01-1 Study Design

Study phase	Dose preparation concentration (% total pyrethrins in corn oil)		Dose/animal (g/kg bw)		Animals assigned	
	Male	Female	Male	Female	Male	Female
I	25%	10%	0.2	0.05	1	1
			0.4	0.1	1	1
			0.8	0.2	1	1
			1.4	0.4	1	1
			2.5	0.8	1	1
II	10%	2.5%	0	0	4	4
			0.04	0.025	4	4
			0.10	0.05	4	4
			0.20	0.10	4	4
			0.40	0.15	4	4
	--	5%	--	0	--	2*
			--	0.025	--	2
			--	0.05	--	2
			--	0.10	--	2
			--	0.15	--	2
			--	0.20	--	2
			--	--	--	--

* Two female control animals from the first part of phase II were selected for use as a control group in the second part of phase II.

3. Dose preparation

At least 2 hours prior to preparation of dosing solutions, the test substance was removed from the refrigerator and equilibrated to room temperature and mixed vigorously by manual inversion for at least 2 minutes. Due to light degradation of the test substance, all solutions of pyrethrum extract were prepared in a photographic dark room in flasks covered with black

electrical tape. The dosing solutions were prepared to contain 5% and 10% total pyrethrins by diluting the appropriate amount of test substance in corn oil. Each dosing solution was mixed for a minimum of 15 minutes on a magnetic stir plate. Each dose solution was prepared once, stored refrigerated, and used during a single day of dosing.

4. Statistics

The data for quantitative continuous variables were intercompared for the 3 treatment groups and the control group by use of Levene's test for equality of variances, ANOVA, and t-tests. Incidence data were compared using the Fisher's Exact Test. Incidence data for select FOB endpoints with ordered severity scores were analysed for group differences using Gamma, Kendall's Tau-B, Stuart's Tau-C, and Somers' D measures of association. A nested analysis of motor activity data was performed using repeated measures analysis of variance with dose as the grouping factor and test period and test session time as within subject factors. The epsilon-adjustment procedure (Greenhouse-Geisser correction) was used in the repeated measures analysis of motor activity data. All statistical analysis, except neuropathology frequency comparisons, were performed using BMDP statistical software. For all statistical tests the probability value of <0.05 (two-tailed) was used as the critical level of significance.

C. METHODS

1. Observations

All animals assigned to the study were observed for mortality twice each day, 7 days a week. Clinical signs of overt toxicity were documented once each day. Observations for clinical signs of toxicity or changes in behaviour were made approximately every hour, for several hours following dosing, during all phases of the study. Animals were observed, specifically, for signs of gait alteration, level of arousal, and tremors during phase II of the study.

2. Body weight

Body weights were recorded in the morning prior to dosing.

Results and discussion:

A. OBSERVATIONS

1. Clinical observations

In phase I, tremors were observed in all males within 2 hours of treatment. The tremors were described as severe at various time periods for the animals dosed at the 0.8, 1.4, and 2.5 g/kg dose levels. Tremors were downgraded to slight 9 and 10 hours after dosing for the animals treated with 0.8 g/kg. Additional observations recorded prior to death included lying on the stomach, labored respiration, salivation, urine stains, and prostration. Tremors were not observed after the 2-hour observation for the male dosed at 0.2 g/kg or after the 4-hour observation for the male dosed at 0.4 g/kg.

Tremors were observed in all female animals within 2 hours of treatment. Tremors were not noted in the female dosed at 0.05 g/kg after the 2-hour observation or in the female dosed at 0.1 g/kg following the 3-hour observation. The tremors for the animals dosed at 0.2, 0.4, and 0.8 g/kg were described as coarse and considered to be severe at several observation periods. The female dosed at 0.8 g/kg was observed lying on her stomach and salivating at the 3-hour observation and prostrate with fine tremors, salivation, and diarrhea at the 4-hour observation.

In phase II, effects for the males were largely limited to the high dose group (0.4 g/kg). Fine tremors were observed in 2 of 4 male animals dosed at 0.4 g/kg at the 2-hour observation period. At the 3- and 4-hour observations, all males in this group had tremors (3 of 4 animals had fine tremors and the remaining animal had coarse tremors). Five hours after treatment, fine tremors were observed in 3 of these 4 animals. No tremors were observed at the 24-hour observation period for these animals. Other signs of toxicity in the 0.4 g/kg dose group included piloerection in 1 animal 2 hours and another animal 3 hours after treatment, red extremities in 1 animal 4 and 5 hours after treatment, and perinasal encrustation for 2 animals 3 hours after treatment and for 1 of these animals 4 hours after treatment. Other observations recorded for male animals included salivation for 1 animal in the 0.2 g/kg dose group and

perinasal encrustation for 1 control animal from the 3-hour to the 24-hour observation periods.

A possible decrease in arousal was observed in animals in the 3 highest dose groups at 5 and 24 hours after treatment. All control animals were described as active and alert, while at least 2 of the 4 animals in the 3 highest dose groups were described as inactive and alert.

In the first part of phase II, observations were limited to the females in the highest dose group (0.15 g/kg). Fine tremors were observed in 1 female animal at the 6- and 7-hour observation periods and gait alteration described as "walks on toes" was observed for 1 animal from the 2-hour through the 24-hour observation period. No other observations were recorded for any other female animals in any dose group. In Part 2 of Phase II, tremors were observed for both females at the highest dose (0.2 g/kg) and 1 female each at 0.15 and 0.1 g/kg. These tremors were generally described as fine and were observed for all 3 groups during the 3 and 4-hour observation periods and for the highest dose group during the 2-, 5-, and 6-hour observation periods. One animal in the 0.2 g/kg dose group was hyperactive at the 2-hour through 5-hour observation periods and again at the 24-hour observation period and as having perinasal encrustation at the 4-hour observation.

2. Mortality

In phase I, the males dosed at 1.4 and 2.5 g/kg were found dead within 5.5 hours after dosing. The female dosed at 0.2 g/kg was found dead 4 hours after dosing. No observations, other than tremors, were observed for this animal. The females dosed at 0.4 and 0.8 g/kg were found dead 5 hours after dosing.

No mortality was observed during phase II of the study.

B. BODY WEIGHT

No treatment-related changes in bw were observed in males or females in phase II of the study.

Table 6.7.1/01-2 Summary of observations in phase II part one

Group (g/kg)	Males					Females				
	0.000	0.040	0.100	0.200	0.400	0.000	0.025	0.050	0.100	0.150
1 Hour post-treatment										
Active/alert	4	4	4	3	4	4	4	4	4	4
Inactive/alert	0	0	0	1	0	-	-	-	-	-
Tremors (none)	4	4	4	4	4	4	4	4	4	4
2 Hour post-treatment										
Active/alert	4	4	4	3	4	4	4	3	4	4
Inactive/alert	0	0	0	1	0	0	0	1	0	0
Tremors (none)	4	4	4	4	2	4	4	4	4	4
Tremors (fine)	0	0	0	0	2	-	-	-	-	-
3 Hour post-treatment										
Active/alert	4	2	4	3	4	3	4	3	4	3
Inactive/alert	0	2	0	1	0	1	0	1	0	1
Tremors (none)	4	4	4	4	0	4	4	4	4	4
Tremors (fine)	0	0	0	0	3	-	-	-	-	-
Tremors (coarse)	0	0	0	0	1	-	-	-	-	-
4 Hour post-treatment										
Active/alert	3	3	3	1	4	3	3	2	3	4
Inactive/alert	1	1	1	3	0	1	1	2	1	0
Tremors (none)	4	4	4	4	0	4	4	4	4	4
Tremors (fine)	0	0	0	0	3	-	-	-	-	-
Tremors (coarse)	0	0	0	0	1	-	-	-	-	-
5 Hour post-treatment										
Active/alert	4	3	2	2	2	-	-	-	-	-
Inactive/alert	0	1	2	2	2	-	-	-	-	-
Tremors (none)	4	4	4	4	1	-	-	-	-	-
Tremors (fine)	0	0	0	0	3	-	-	-	-	-
6 Hour post-treatment										
Active/alert	-	-	-	-	-	3	2	2	1	2

Inactive/alert	-	-	-	-	-	1	2	2	3	2
Tremors (none)	-	-	-	-	-	4	4	4	4	3
Tremors (fine)	-	-	-	-	-	0	0	0	0	1
7 Hour post-treatment										
Active/alert	-	-	-	-	-	3	2	2	3	4
Inactive/alert	-	-	-	-	-	1	2	2	1	0
Tremors (none)	-	-	-	-	-	4	4	4	4	3
Tremors (fine)	-	-	-	-	-	0	0	0	0	1
24 Hour post-treatment										
Active/alert	4	4	2	1	2	2	3	2	3	3
Inactive/alert	0	0	2	3	2	2	1	2	1	1
Tremors (none)	4	4	4	4	4	4	4	4	4	4

Table 6.7.1/01-2 Summary of observations in phase II part one

Group (g/kg)	Females					
	0.000	0.025	0.050	0.100	0.150	0.200
1 Hour post-treatment						
Active/alert	2	2	2	2	2	2
Tremors (none)	2	2	2	2	2	2
2 Hour post-treatment						
Active/alert	1	2	2	2	2	1
Inactive/alert	1	0	0	0	0	0
Hyperactive	0	0	0	0	0	1
Tremors (none)	2	2	2	2	2	1
Tremors (fine)	0	0	0	0	0	1
3 Hour post-treatment						
Active/alert	2	2	2	2	2	1
Hyperactive	0	0	0	0	0	1
Tremors (none)	2	2	2	1	1	0
Tremors (fine)	0	0	0	1	1	2
4 Hour post-treatment						
Active/alert	2	1	1	2	2	0
Inactive/alert	0	1	1	0	0	1
Hyperactive	0	0	0	0	0	1
Tremors (none)	2	2	2	1	1	0
Tremors (fine)	0	0	0	1	1	1
Tremors (coarse)	0	0	0	0	0	1
5 Hour post-treatment						
Active/alert	1	1	2	2	0	0
Inactive/alert	1	1	0	0	2	1
Hyperactive	0	0	0	0	0	1
Tremors (none)	2	2	2	2	2	0
Tremors (fine)	0	0	0	0	0	2
6 Hour post-treatment						
Active/alert	2	1	2	2	1	1
Inactive/alert	0	1	0	0	1	1
Tremors (none)	2	2	2	2	2	1
Tremors (fine)	0	0	0	0	0	1
24 Hour post-treatment						
Active/alert	2	1	1	2	2	1
Inactive/alert	0	1	1	0	0	0
Hyperactive	0	0	0	0	0	1
Tremors (none)	2	2	2	2	2	2

Conclusions:

Findings from this study indicate that female rats are more sensitive to the test substance than male rats. Based upon the results of this study, dose levels of 0.04, 0.125, and 0.4g total pyrethrins/kg body weight were selected for males in the definitive acute neurotoxicity study. The doses were to be administered as a solution of pyrethrum extract in corn oil to

contain 10% total pyrethrums. For females, dose levels of 0.02, 0.063, and 0.2 g total pyrethrins/kg bodyweight were selected for use in the definitive acute neurotoxicity study. The doses were to be administered as a solution of pyrethrum extract in corn oil to contain 5% total pyrethrums. Based upon the time course of effects observed in this study, 3 to 5 hours was selected as the time for the first post-exposure evaluation in the definitive acute neurotoxicity study.

ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

This was a GLP compliant study. The study did not follow any OECD guidelines and was used to establish dose levels and select appropriate evaluation intervals for a definitive acute oral neurotoxicity study.

Conclusion:

Appropriate dose levels and evaluations timepoints were selected for the definitive neurotoxicity study based on the findings from this study; i.e. dose levels of 0.04, 0.125, and 0.4 g total pyrethrins/kg body weight for males, and dose levels of 0.02, 0.063, and 0.2 g total pyrethrins/kg body weight for females, and 3 to 5 hours as the time for the first post-exposure evaluation time of 3 to 5 hours.

ASSESSMENT AND CONCLUSION BY RMS

RMS agrees with the assessment and the conclusion of the applicant.

The study did not follow any OECD guidelines but it is considered adequate to establish dose levels and time intervals for evaluation in a definitive acute oral neurotoxicity study.

Based on the results of the study, a dosing solution containing 10% total pyrethrins and dose levels of 0.04, 0.125, and 0.4 g total pyrethrins/kg body weight were selected for males. For females in the definitive acute neurotoxicity study, a dosing solution containing 5% total pyrethrins and dose levels of 0.02, 0.063, and 0.2 g total pyrethrins/kg body weight were selected. In addition, 3 to 5 hours was determined as the time for the first post-exposure evaluation time.

- 9) *Comparative functional observational battery study of twelve commercial pyrethroid insecticides in male rats following acute oral exposure; Weiner M. L., Nemeč M., Sheets L., Sargent D., & Breckenridge C., 2009; Cross reference IIA 5.7.1/03*

Guidelines:

None

GLP:

Yes.

The study is acceptable as supportive information

Executive Summary:

Twelve commercial pyrethroid insecticides (technical-grade active ingredients) were evaluated individually for acute neurobehavioral manifestations of toxicity under conditions suited to assist with determining whether they act by a common mechanism of toxicity. The pyrethroids that were tested reflect a diversity of structures, including six with an α -cyano phenoxybenzyl moiety (bicyfluthrin, l-cyhalothrin, cypermethrin, deltamethrin, esfenvalerate and fenpropathrin) and six without this moiety (bifenthrin, S-bioallethrin, permethrin, pyrethrins, resmethrin and tefluthrin). These chemicals also present a variety of behavioural effects, including ones that are historically classified as causing a T (tremor), CS

(choreoathetosis with salivation) or intermediate syndrome of intoxication, and others that have not previously been classified. Each pyrethroid that was tested consisted of the complement of isomers that occur in commercial products—a key factor for relevance for environmental and human exposure and for comparisons, since the biological activity of the individual isomers can vary tremendously.

Young-adult male Sprague–Dawley rats (10 per dose group) were administered a single dose of pyrethroid by oral gavage, in corn oil, at a volume of 5 ml/kg. Each was tested at a range of two or three dose levels, including a minimally toxic dose, to establish the more sensitive manifestations of toxicity, and a more toxic dose, to establish a more complete spectrum of neurobehavioral manifestations. Animals were evaluated using a functional observational battery (FOB) that was designed to characterize and distinguish effects classically associated with T or CS syndromes of intoxication. The FOB was performed when manifestations of toxicity were most apparent at the time of peak effect (2, 4, or 8 h post-dosing) by observers who were blinded to dose group assignment, thus avoiding possible bias. The results from this study indicate that some pyrethroids clearly exhibit the historic classification symptoms of the T and CS syndromes while others do so less obviously. Use of the statistical technique of Principal Component Analysis (PCA) further helped interpret the study findings.

These results establish manifestations of neurotoxicity in vivo that can be used as weight of evidence to determine whether pyrethroid insecticides act through a common mechanism of toxicity in mammals. Based on a review of the FOB data, analyzed by PCA, and other published data, two common mechanism groups are proposed. Group 1 (T syndrome) would include pyrethrins, bifenthrin, resmethrin, permethrin, S-bioallethrin and tefluthrin. Group 2 (CS syndrome) would include cypermethrin, deltamethrin, esfenvalerate, b-cyfluthrin and l-cyhalothrin. Fenpropathrin exhibited features of both groups.

This summary is focused on the information for pyrethrins only and will refer to the test compounds when necessary.

Material and methods:

A. MATERIALS

1. Test Material: Pyrethrins

Description: Not reported

Lot/Batch: Not reported

Purity: Not reported (Purity and stability verification for the test substances were confirmed by the respective suppliers by Certificates of Analysis or GLP analyses).

2. Vehicle: Mazola corn oil

3. Test Animals

Species: Rat

Strain: Charles River Crl:CD[®](SD)IGS BR

Age: 28-29 days

Sex: Male

Weight at dosing: 219-284 g (males) and 136-182 g (females)

Source: Charles River Laboratories, Inc., Raleigh, NC

Acclimation period: At least 14-days

Diet: Certified Rodent chow[®] #5002 (PMI Nutrition International, Inc.) *ad libitum*

Water: Tap water *ad libitum*

Housing: three/cage by sex for the first three days of the 14-day acclimation period, and housed individually thereafter.

Environmental Conditions: Temperature: 71 ±3°F

Humidity: 50 ±20%

Photoperiod: 12-hour light/dark cycle

B. STUDY DESIGN AND METHODS

1. In-life dates Not reported

2. Animal assignment and treatment

The dose levels and the times-to-peak effect for each test substance were based on data from a rangefinding study at the same laboratory. For pyrethrins, dose levels of 400 and 800 mg/kg body weight were selected and the time to peak effect for clinical signs of toxicity was 4 hours post administration. The rats were allocated to 2 treatment groups and a control group of 10 rats/group, using a computerized randomisation procedure. At the time of group assignment, only rats with bw within ± 20% of the population mean were included. The selected route of administration was oral gavage. The vehicle and test substance formulations were administered orally by gastric intubation via a 16-gauge stainless steel gavage cannula as a single dose. The day of dose administration was termed study day 0 for that animal. Individual dosages were based on the day 0 body weight.

3. Statistics

Each mean was presented with the SD, and the number of animals (N) used to calculate the mean. The numeric data were subjected to statistical analyses by the Dunnett's Test, except for hindlimb resistance and extensor strength in the neuromuscular parameters, which were subjected to the Fisher's Exact Test. In addition, Principal Component Analysis and Factor Analysis were used for a more thorough interpretation of these data (Breckenridge *et al.*, 2009).

C. METHODS

1. Observations

All animals were observed twice daily (morning and afternoon) for mortality and moribundity. Clinical observations were performed daily, except on the day of the Functional Observational Battery.

2. Body weight

Animals were randomized to groups based on body weight 1 week prior to dosing so that body weights were similar on the day of dosing. Individual body weights were recorded at randomization and prior to dose administration on study day 0. Body weights were also recorded during the FOB and prior to terminal euthanasia.

3. Functional Observational Battery (FOB):

The FOB used in this study was based on a standardized procedure developed and used by the laboratory. It is based on procedures in the US EPA OPPTS Health Effects Test Guideline 870.6200 (US EPA, 1996). The FOB was modified to include additional details (e.g., the coarseness of tremor) to distinguish findings particularly associated with pyrethroid intoxication. Tremors were scored on a severity scale of 1–5: 1 indicates no tremors; 2 indicates slight (1.5 mm) tremors; 3 indicates moderately coarse (3 mm) tremors with slight impairment; 4 indicates markedly coarse (4.5 mm) tremors with marked impairment of locomotion and 5 indicates extremely coarse (6 mm) tremors and locomotion impossible. For aerial righting and landing footsplay, the animals landed on a well-cushioned surface and the test was not performed if the animal was judged unable to perform the test. Observations were recorded for all animals at the time of peak effect after test substance administration. Study technicians were given special training to distinguish classical symptoms of pyrethroid intoxication. Testing was performed without the technician's knowledge of dose group assignment and inter-observer reliability was established to verify consistency among the technicians by verification of training in the laboratory with standard pyrethroids.

The FOB was performed in a sound-attenuated room equipped with a white noise generator set to operate at 70 ± 10 dB, while home cage observations were performed in the animal room.

The FOB consisted of six types of observations: home cage, handling, open field, sensory, neuromuscular and physiological observations. Table below summarizes the specific parameters evaluated for each category of the FOB observations.

Type of observation	Parameter	
	Posture	Biting
Home cage	Convulsions/tremors	Palpebral (eyelid) closure
	Faeces consistency	
	Ease of removal from cage	Ease of handling animal in hand
Handling	Lacrimation/chromodacryorrhea	Salivation
	Pilorection	Fur appearance
	Palpebral closure	Respiratory rate/character
	Eye prominence	Mucous membrane/eye/skin colour
	Red/crusty deposits	Muscle tone
Open field	Mobility	Gait
	Rearing	Arousal
	Convulsion/tremors	Urination
	Grooming	Defecation
	Bizarre/stereotypic behaviour	Gait score
	Time to first step (s)	Backing
Sensory	Approach response	Touch response
	Startle response	Tail pinch response
	Pupil response	Eyeblink response
	Forelimb response	Hindlimb response
	Air-righting response	Olfactory response
Neuromuscular	Hind limb extensor strength	Grip strength: hind and forelimb
	Hindlimb foot splay	Rotorod performance
Physiological	Body temperature	Body weight
	Catalepsy	

4. Macroscopic examination (unscheduled deaths)

Animals found dead during the study underwent a gross necropsy examination. This included, but was not limited to, examination of the external surface, all orifices, and the cranial, thoracic, abdominal and pelvic cavities, including viscera.

5. Schedule euthanasia:

Following clinical observations on the day after treatment, surviving animals were euthanized by carbon dioxide inhalation and discarded without necropsy.

Results and discussion:

A. OBSERVATIONS

5. Clinical signs of toxicity

Signs of toxicity in animals dosed pyrethrins had resolved in all surviving animals by the day after treatment.

6. Mortality

All animals dosed pyrethrins survived to scheduled termination.

B. BODY WEIGHT AND BODY WEIGHT GAIN

Body weight for test substance-treated animals was generally similar to the concurrent control group values on the day of dosing and at the scheduled euthanasia.

C. NEUROBEHAVIOURAL EVALUATIONS

3. Functional observation battery (FOB)

FOB findings at the times of peak effect for pyrethrins (Type 1-T syndrome – non- α -cyano substituent in the 3-phenoxybenzyl alcohol moiety of the molecule) are summarized in Table 6.7.1/03-1, respectively. The pyrethrins were considered to be the least potent type I

pyrethroid based on FOB data.

Table 6.7.1/03-1 Summary of FOB findings for Pyrethrins

Observation (Time to peak effect 4h)	Dose level (mg/kg)	
	400	800
Home cage observations		
Sitting, head held low		
Flattened, limbs may be extended		1
Rearing		
Splayed hindlimbs		
Clonic convulsions (repetitive movement of mouth/jaw)		
Clonic convulsions (back twitches)	1	
Clonic convulsions (head/body twitches) myoclonus		1
Clonic convulsions (irregular jerking, limbs)		1
Clonic convulsions (whole body)		1
Slight tremors	1	2
Moderately coarse tremors		
Markedly coarse tremors		
Extremely coarse tremors		
Biting of self		
Handling observations		
Salivation		1 (1)
Ventral staining	1	1
Ventral wetness		
Abdominogenital wetness		
Slightly soiled fur		
Red deposits—nose		2 (1)
Pale mucous membrane		
Pale skin		
Pulsating eyes		
Exophthalmus		
Moderately difficult to remove from cage	1	
High difficulty in handling		
Open field observations		
Ataxia, excessive sway, rocks, lurches		
Slightly impaired mobility		
Moderately impaired mobility		
Walking on tiptoes	1 (1)	3 (1)
Body drags, body sways, abdomen contacts surface		
Hindlimbs splayed or dragging		
Hunched body	1	1
Gait impairment—slight		
Gait impairment—considerable		
Gait impairment—severe, cannot walk without falling		
Clonic convulsions (back twitches)		1
Clonic convulsions (head/body twitches)		2 (1)
Clonic convulsions (irregular jerking, limbs)	2	
Clonic convulsions (whole body)		
Slight tremors	1 (1)	3 (1)
Moderately coarse tremors		
Markedly coarse tremors		1
Extremely coarse tremors		
Low arousal level		
Stereotypic behavior (head flick)		
Sensory observations		
Approach reaction—no response		
Approach reaction—more energetic response (more than slight)		
Touch response—no reaction		
Touch response—more energetic response (more than slight)		

Startle response—more energetic response (more than slight)		
Olfactory orientation—no reaction		
Air-righting reflex—slightly uncoordinated		
Exaggerated hindlimb flexion	2	1
No hindlimb extension		
No forelimb extension		
Neuromuscular observations		
Reduced hindlimb resistance		

Note: Numerals represent the numbers of animals with findings. Occurrences of findings in the control group are indicated in parenthesis next to the group finding. If the finding was not observed in the control, no number has been included.

Conclusion:

Pyrethrins is a Type 1/T syndrome pyrethroid, which is consistent with its structure and other pyrethroids without an α -cyano substituent in the 3-phenoxybenzyl alcohol moiety of the molecule. Although there is no known common toxophore that mediates acute toxicity of pyrethroids, the presence of the α -cyano substituent in the 3-phenoxybenzyl alcohol moiety of the molecule confers greater potency by an estimated order of magnitude in acute lethality studies in rodents. In addition, the manifestations of the particular toxic effect of neurotoxicity can also be related to the presence or absence of the α -cyano substituent, as noted in the early distinctions of the two structural classes of the early pyrethroids (Verschoyle and Aldridge, 1980; Lawrence and Casida, 1982) and by the present study using the FOB.

In the present study the potency of the α -cyano-containing pyrethroids was generally higher than the noncyano pyrethroids. The lowest dose tested ranged from 10 to 65 mg/kg for the α -cyano pyrethroids with l-cyhalothrin (10 mg/kg) > deltamethrin and b-cyfluthrin (12.5 mg/kg) > fenpropathrin and esfenvalerate (15 mg/kg) > cypermethrin (65mg/kg). The potency of the non-cyano pyrethroids was generally lower than the α -cyano pyrethroids based on the lowest dose tested: tefluthrin (10 mg/kg) > bifenthrin (40 mg/kg) > S-bioallethrin (150 mg/kg) > permethrin (200mg/kg) > resmethrin (350 mg/kg) > pyrethrins (400 mg/kg).

ASSESSMENT AND CONCLUSION BY APPLICANT

Assessment:

This was a GLP compliant study. This study was not evaluated for the first EU approval review of pyrethrins (DAR Vol. 3 B6, 2007). The study is not a guideline study, but the functional observations conducted are consistent with those of OECD guideline 424 (1997). The study is conducted at higher dose levels than those tested in Hermansky and Hurley (1993).

The study shows that pyrethrins are of significantly lower neurotoxicity than all other pyrethroid molecules tested. The study is conducted at higher dose levels than those tested in Hermansky and Hurley (1993) and do not therefore have any impact on the dose levels selected for human health risk assessment.

ASSESSMENT AND CONCLUSION BY RMS

RMS agrees with the assessment and the conclusion of the applicant.

According to the study, pyrethrins are considered to be the least potent type I pyrethroid (Type 1-T syndrome – non- α -cyano substituent in the 3-phenoxybenzyl alcohol moiety of the molecule) based on FOB data. Moreover, pyrethrins are of significantly lower neurotoxicity than all other pyrethroid molecules tested.

The study is conducted at higher dose levels than those tested in Hermansky and Hurley (1993) and RMS agrees that does not have any impact on the dose levels selected for human health risk assessment.

DAR - ANNEX B.9 Ecotoxicology

Active substance: Pyrethrins

- 1) *Pyrethrum extract (FEK-99) - acute toxicity to mysid shrimp (Mysidopsis bahia) under flow-through conditions. Machado M.W., 1994e*

Guidelines:

In accordance with U.S.EPA Pesticide Assessment Guidelines, Subdivision E, Section 72-3.

Testing Laboratory and dates:

USA conducted the study during the period December 14, 1993 to December 18, 1993.

GLP:

Yes (self-certified), with the following exceptions:

- Routine water and food contaminant screening analyses for pesticides, PCBs and metals were not collected in accordance with GLP (i.e., no distinct protocol, Study Director, etc.)
- Total organic carbon analyses for filtered seawater were not collected in accordance with GLP
- Documentation of observations made during a single interval of a preliminary exposure was recorded using pencil

These deviations were not considered to have affected the scientific validity of the study or the interpretation of the results.

The study is acceptable

Executive Summary:

In an acute toxicity laboratory study under flow-through conditions, *Mysidopsis bahia* were exposed to total Pyrethrins (57.488% purity) at nominal concentrations of 0, 0.38, 0.75, 1.5, 3.0, 6.0 µg/L over a period of 96 hours. Each concentration was testing using 20 mysid shrimp per treatment level. Dilution water and acetone solvent were also tested as negative and solvent controls, respectively.

At test termination, mean cumulative mortalities of 5%, 10% and 55% were observed among mysids exposed to the 0.34, 0.81, and 1.6 µg total Pyrethrins/L treatment levels. Sub-lethal effects were observed among several of the surviving mysids exposed to 0.81 µg total Pyrethrins/L and among all of the surviving mysid exposed to the 1.6 µg total Pyrethrins/L treatment level.

The 96-hour LC₅₀ (95% confidence interval) was calculated by moving average angle analysis to be 1.4 µg total Pyrethrins/L. The 96-hour NOEC was determined to be 0.34 µg total Pyrethrins/L. Based on these results and on criteria established by Directive 67/548/EEC, pyrethrum extract (FEK-99) would be classified as very toxic to mysid shrimp.

Materials and Methods:

MATERIALS

Test Material:	Pyrethrum extract (FEK-99)
Description:	Brown liquid
Lot/Batch #:	R92-254
Purity:	57.488% total Pyrethrins
CAS #:	8003-34-7
Stability of test compound:	> 5 years at 0°C in the absence of light
Vehicle and/or positive control:	Acetone (solvent control)
	Dilution water (negative control)

Test organisms -

Species: *Mysidopsis bahia*

Source:

Age: ≤24 h

Acclimation conditions:

Acclimation period:	≥ 14 days
Diet:	Mysid cultures were fed live brine shrimp (<i>Adernia salina</i>) nauplii twice daily
Temperature:	24°C to 26°C
Dissolved oxygen:	81% to 87%
Photoperiod:	16 h light, 8 h darkness
5. Exposure conditions:	
Temperature:	25 ± 1°C
Feed:	Mysid cultures were fed live brine shrimp (<i>Adernia salina</i>) nauplii twice daily
Photoperiod:	16 h light, 8 h darkness
Dilution Water:	
Water used in study:	Seawater piped in from the Cape Cod Canal, Bourne, Massachusetts, USA from approximately 4 meters offshore at a depth of 0.5 meters
Salinity:	31 to 32‰
pH:	7.8 to 7.9

Findings:

MORTALITY

Cumulative mortality and sublethal effect data for *Mysidopsis bahia* during 96 hours flow-through exposure to FEK-99 are given in Table B.9.1.1.

Table B.9.1.1: Cumulative mortality and sub-lethal effects for *Mysidopsis bahia* during 96 hours flow-through exposure to FEK-99

Mean measured concentration ^a	Cumulative mortality (%)			
	24 hour (mean)	48 hour (mean)	72 hour (mean)	96 hour (mean)
Control	0	0	0	0
Solvent control	0	0	0	0
0.29	0	0	0	0
0.34	0	5	5	5
0.81	5 ^{bc}	10 ^{bc}	10 ^{bc}	10 ^{fh}
1.6	5 ^d	15 ^{ef}	30 ^{ef}	55 ^{ef}
3.4	20 ^{ef}	95 ^g	100	100

^aConcentrations measured as µg total Pyrethrins/L

^bTwo of the surviving mysids exhibited erratic swimming behavior

^cOne of the surviving mysids exhibited a partial loss of equilibrium.

^dAll of the surviving mysids exhibited a partial loss of equilibrium

^eSeveral of the surviving mysids exhibited a complete loss of equilibrium

^fSeveral of the surviving mysids exhibited a partial loss of equilibrium

^gAll of the surviving mysids exhibited a complete loss of equilibrium

^hSeveral of the surviving mysids exhibited erratic swimming behavior

Conclusions:

The 96-hour LC₅₀ based on mean measured concentrations of FEK-99 with *Mysidopsis bahia* was determined to be 1.4 µg/L total Pyrethrins, and the NOEC was 0.34 µg/L. Utilizing the concentration-effect response observed during this study and criteria established by Directive 67/548/EEC, pyrethrum extract (FEK 99) would be classified as very toxic to *Mysidopsis bahia*. (Machado MW 1994e).

Analytical data on concentrations in the test media

Materials and Methods:

MATERIALS

As per Machado MW 1994e, analysis by GC-ECD.

Findings:

The results of the analysis of the exposure solutions for total Pyrethrins during the in-life portion of the definitive exposure are presented in Table B.9.1.1 in the above section. Mean measured concentrations established for this study defined the exposure levels as 0.29, 0.34, 0.81, 1.6 and 3.4 µg total Pyrethrins/L. The mean measured concentrations averaged 60% of nominal (N=28) with a mean coefficient of variation of 25%. The ratio of the highest measured concentration to the lowest measured concentration at each treatment level was determined and ranged from 1.5 to 2.7.

Analyses of the QC samples resulted in measured concentrations which fell within two standard deviations of the acceptable recovery range, which exceeded the minimum acceptance criteria (i.e., within three standard deviations of the acceptable recovery range). Measured concentrations for the QC samples averaged 114% (N=5) of the nominal fortified levels (0.375 to 6.00 µg total Pyrethrins/L).

Conclusions:

Mean measured concentrations established for this study defined the exposure levels as 0.29, 0.34, 0.81, 1.6 and 3.4 µg total Pyrethrins/L. (Machado MW 1994e)

2) *Pyrethrum extract (FEK-99) - acute toxicity to eastern oyster (Crassostrea virginica) under flow-through conditions. Dionne E., 1994*

Guidelines:

In accordance with U.S.EPA Pesticide Assessment Guidelines, Subdivision E, Section 72-3.

Testing Laboratory and dates:

[REDACTED]

USA conducted the study during the period October 22, 1993 to October 26, 1993.

GLP:

Yes (self-certified), with the following exceptions:

- Routine water and food contaminant screening analyses for pesticides, PCBs and metals were conducted using standard U.S. EPA procedures by Lancaster Laboratories, Lancaster, Pennsylvania, USA. These data were not collected in accordance with GLP procedures (i.e. no distinct protocol, Study Director, etc.).
- Total organic carbon analyses for filtered seawater conducted were not collected in accordance with GLP procedures.

These deviations were not considered to have affected the scientific validity of the study or the interpretation of the results.

The study is acceptable according to US EPA documents. No OECD guidelines were found to be compared to this study

Executive Summary:

In an acute toxicity laboratory study under flow-through conditions, *Crassostrea virginica* were exposed to total Pyrethrins (57.488% purity) at nominal concentrations of 0, 26, 43, 72, 120 and 200 µg/L over a period of 96 hours. The Pacific oyster (*Crassostrea gigas*) and

the Eastern oyster are the preferred species for the shell deposition test as they have demonstrated sensitivity to known toxicants and because a substantial database is available on each species. Eastern oysters were selected for this study based on their availability. Each concentration was tested using 40 oysters per treatment level. Dilution water and acetone solvent were also tested as negative and solvent controls, respectively.

At test termination, growth among dilution water control oysters averaged 2.6 mm. The average shell deposition observed was considered representative for this species and acceptable for establishing the relative toxicity of FEK-99 to Eastern oysters. Reduced feeding and reduced fecal and pseudofecal production were observed among oysters exposed to the 130 µg total Pyrethrins/L concentration. No sublethal effects were observed among oysters exposed to any of the remaining concentrations tested or the controls. Shell growth among oysters exposed to the control and solvent control averaged 2.6 and 2.2 mm, respectively. Shell growth among oysters exposed to 20, 45, 68 and 130 µg total Pyrethrins/L was reduced by 13, 36, 41 and 68%, respectively, and was significantly different when compared to the growth of the pooled control oysters. Shell growth reduction at the 14 µg total Pyrethrins/L treatment level was 12%.

The EC₅₀ (95% confidence intervals) was calculated by linear regression to be 87 µg total Pyrethrins/L. The NOEC was determined to be 14 µg total Pyrethrins/L.

Materials and Methods:

MATERIALS

Test Material:	Pyrethrum extract (FEK-99)
Description:	Brown liquid
Lot/Batch #:	R92-254
Purity:	57.488% total Pyrethrins
CAS #:	8003-34-7
Stability of test compound:	> 5 years at 0°C in the absence of light
Vehicle and/or positive control:	Acetone (solvent control) Dilution water (negative control)
Test organisms -	
Species:	<i>Crassostrea virginica</i> (Eastern oyster)
Source:	P. Cummins Oyster Co., Pasadena, Maryland, USA
Age:	Pre-spawn condition of gonadal development
Mean weight at start of the study:	Not documented
Mean standard length at start of the study:	25 to 50 mm
Acclimation conditions:	
Acclimation period:	23 days
Diet:	Supplementary algal diet of <i>Isochysis galbana</i> , clone T-Iso and <i>Tetraselmis maculate</i> besides natural presence of algae in seawater used in study.
Temperature:	19 to 21°C
Dissolved oxygen:	74 to 99% of saturation
Photoperiod:	16 hours light, 8 hours darkness
Exposure conditions:	
Temperature:	20 to 22°C
Feed:	Supplemental feedings of algae (<i>Isochysis galbana</i>) besides natural presence of algae in seawater used in study.
Photoperiod:	16 hours light, 8 hours darkness
Dilution Water:	
Water used in study:	Natural unfiltered seawater pumped from the Cape Cod Canal, Bourne, Massachusetts, USA from about 4 meters offshore at a depth of approximately 0.5 meters.
Salinity:	32‰
pH:	7.8

Water Hardness: Not documented

Findings:

MEASURED EFFECTS

Effects of FEK-99 exposure to *Crassostrea virginica* are summarised in Table B.9.1.2.

Table B.9.1.2: Effects of FEK-99 on the shell deposition of *Crassostrea virginica* after 96 hours^a

Mean measured concentration ^b	Mean shell deposition ^c (mm)	Mean percentage reduction ^d
Control	2.6(0.9)	NA ^f
Solvent control	2.2(0.8)	NA ^f
Pooled control	2.4(0.9)	NA ^f
14	2.1(0.7)	12
20	2.1(0.9)	13 ^e
45	1.5(0.6)	36 ^e
68	1.4(0.5)	41 ^e
130	0.8(0.4)	68 ^e

^aThe EC₅₀ was calculated to be 87 µg total Pyrethrins/L. The NOEC was estimated to be 14 µg total Pyrethrins/L

^bConcentrations expressed as µg total Pyrethrins/L

^cThe mean shell deposition is presented with the standard deviation in parentheses and represents 40 oysters/treatment

^dThe formula for the calculation of mean percent reduction is presented in the Study Protocol

^eSignificantly different as compared to the performance of the pooled control oysters

^fNA = not applicable

OBSERVATIONS

Observations on sublethal effects were observed at test termination (96-hours), and included reduced feeding and reduced fecal and pseudofecal production among oysters exposed to the 130 µg total Pyrethrins/l concentration. No sublethal effects were observed among oysters exposed to any of the remaining concentrations tested or the controls.

Growth among dilution water control oysters at test termination averaged 2.6 mm. The growth of control organisms during this study exceeded the required minimum and was within the historical range (0.9 - 4.5 mm). Based on these data, the average shell deposition observed during this study is considered representative for this species and acceptable for establishing the relative toxicity of FEK 99 to Eastern oysters. At test termination (96 hours), reduced feeding and reduced fecal and pseudofecal production were observed among oysters exposed to the 130 µg total Pyrethrins/L concentration. No sublethal effects were observed among oysters exposed to any of the remaining concentrations tested or the controls. Shell growth among oysters exposed to the control and solvent control averaged 2.6 and 2.2 mm, respectively. Statistical analysis determined no significant difference between shell deposition in the control and solvent control, therefore, control data were pooled for further analyses. Shell growth among oysters exposed to 20, 45, 68 and 130 µg total Pyrethrins/L was reduced by 13, 36, 41 and 68%, respectively, and was significantly different when compared to the growth of the pooled control oysters. Shell growth reduction at the 14 µg total Pyrethrins/L treatment level was 12%, which was not statistically different than the growth of the pooled control organisms. Effects observed during this test were clearly concentration-dependent.

Conclusions:

Under the conditions of this study, the 96-hour EC₅₀ value for FEK 99 with *Crassostrea virginica* was determined to be 87 µg/L. The NOEC was 14 µg/L total Pyrethrins. (Dionne E 1994)

Analytical data on concentrations in the test media

Materials and Methods:

MATERIALS

As per Dionne E 1994, analysis by GC-ECD.

Findings:

The analyses of the exposure solutions for total Pyrethrins during the in-life portion of the definitive exposure period are presented in Table 9.1.3. Mean measured concentrations established for this study defined the exposure levels as 14, 20, 45, 68 and 130 µg total Pyrethrins/L. The mean measured concentrations averaged 56% of nominal (N = 20) with a mean coefficient of variation of 16%. The ratio of the highest measured concentration to the lowest measured concentration at each treatment level was determined and ranged from 1.3 to 1.5. Analyses of the Quality Control samples resulted in measured concentrations which fell within two standard deviations of the acceptable recovery range, which exceeded the minimum acceptance criteria (i.e., within three standard deviations of the acceptable recovery range). Measured concentrations for the QC samples averaged 88.8% (N = 6) of the nominal fortified levels (25.0 to 200 µg total Pyrethrins/L).

Table B.9.1.3: Mean concentrations of total Pyrethrins measured in exposure solutions during the 96-hour flow-through exposure of Eastern oysters (*Crassostrea virginica*)

Nominal concentration	Measured concentration ^{ab}		% nominal ^d	
	0-hour	96-hour	Mean (CV) ^c	
Control	<3.1	<3.7	NA ^e	NA
Solvent control	<3.1	<3.7	NA	NA
26	15	12	14(14)	52
43	22	19	20(17)	47
72	47	43.5	45(15)	63
120	71.5	65	68(17)	57
200	145	105	130(17)	63
Stock solution ^f	0.42	0.35	0.39	96
(25.0) QC 1 ^g	27.1 (108) ^h	17.5 (70.1)		
(75.0) QC 2	79.0 (105)	55.9 (74.5)		
(200) QC 3	218 (109)	131 (65.5)		

^aConcentrations expressed as µg total Pyrethrins/L

^bMeasured concentrations have been corrected for average QC recovery (i.e., 88.8%)

^cCV = coefficient of variation

^dPercent of nominal was calculated for each treatment level by dividing the mean measured concentration by the nominal concentration and multiplying by 100

^eNA = not applicable

^fConcentrations expressed as mg total Pyrethrins/mL

^gQC = Quality Control sample

^hPercent of nominal for each QC sample is presented in parentheses

Conclusions:

The geometric mean measured concentrations for the study were 14, 20, 45, 68, and 140 µg total Pyrethrins/L. (Dionne E 1994)

B8.9.1 Effects on algal growth and growth rate

- 1) *Refined pyrethrum extract - algal growth inhibition assay. Jenkins C.A., 2003*

Guidelines:

EC Methods for Determination of Ecotoxicity Annex to Directive 92/69/EEC Part C, Method 3 and OECD 201

Testing Laboratory and dates:

conducted the study during the period April 26, 2002 to November 15, 2002.

GLP:

Fully GLP compliant.

The study is acceptable.

Executive Summary:

In an algal growth inhibition assay, *Selenastrum capricornutum* cultures were exposed to water accommodated fractions of refined pyrethrum extract (57.03% total Pyrethrins) dispersed in nutrient media at nominal loading rates of 0, 0.5, 1.1, 2.42, 5.32, 11.7, 25.8 and 56.7 mg/L. The mean measured levels were 0, 0.265, 0.615, 0.889, 1.25, 1.30, 1.95 and 1.78 mg/L.

After 72 hours of exposure, neither the E_{bL50} (median effect loading rate based on area under the growth curve) nor the E_{rL50} (median effect loading rate based on the growth curve) values for *S. capricornutum* could be calculated because <50% inhibition occurred during the definitive test. Similar levels of inhibition were obtained at the three highest nominal loading rates, 11.7 to 56.7 mg/L. The initial measured concentrations of test substance at these concentrations were similar (2.25 to 2.89 mg/L). Consequently, the E_{bL50} and E_{rL50} values were >56.7 mg/L based on the nominal loading rate and >1.95 mg/L expressed in terms of the highest mean measured level. No microscopic abnormalities were noted. The "no observed effect loading rate" (NOELR) for area under the growth curve and for growth rate was 5.32 mg/L (nominal); the NOEC was 1.25 mg/L (measured).

Materials and Methods:

MATERIALS

Test Material:	Refined pyrethrum extract
Description:	Clear amber liquid
Lot/Batch #:	FEK-99
Purity:	57.03% total Pyrethrins
CAS #:	8003-34-7
Stability of test compound:	Stability during the course of this study was demonstrated by analysis.
Vehicle and/or positive control:	Sterile culture medium
Test organisms -	
Species:	<i>Selenastrum capricornutum</i>
Strain:	CCAP 278/4
Source:	Huntingdon Research Centre culture collection of algae and protozoa, Institute of freshwater ecology, Cumbria, UK
Culture medium:	Sterile algal nutrient medium as recommended in OECD Procedure 201 and EC Directive 92/69/EEC Official Journal no L383A, part C3
Pre-culture conditions:	
Light levels:	Not documented
Photoperiod:	Continuous
Temperature:	21 to 25°C (with an occurrence of 18.1°C in the initial 24 h)
Cell density:	0.9 x 10 ⁶ cells/mL, with final aliquot of secondary culture diluted to 1.0 x 10 ⁴ cells/mL before use
Environmental conditions:	test
Light levels:	9800 to 8500 lux
Photoperiod:	Continuous
Temperature:	23 ± 2 °C
Incubation period:	72 hours without renewal

Cell maintenance: Cells were maintained in volumetric flasks. Gaseous exchange and suspension of the cells were ensured by oscillating on an orbital shaker at 150 cycles/min.

Findings:

VALIDITY CRITERIA

After 72 h, the measured levels had decreased, ranging between 2 and 36% of their nominal values and 33 and 58% of their initial values. Failure to achieve the nominal concentrations at the higher levels was attributed to the limit of aqueous solubility of the test substance having been exceeded. After 72 h, a sample without algal cells showed a small decrease in the measured level compared to medium with algal cells. After 72 hours, analysis of a sample of medium containing test substance at 0.5 mg/L, which had been incubated without algal cells, showed a small decrease in the measured level compared to medium that had been incubated with algal cells (22% compared to 63%). A sample taken at 56.7 mg/L, which had been incubated without algal cells indicated that the exposure level was maintained (97%) whereas in the presence of algal cells a loss of 42% was noted. These results indicate that the stability of the test substance was affected by the presence of algal cells.

CELL DENSITY AND GROWTH RATE

Calculated values for the levels of inhibition of growth rate and biomass are given in Table B.9.2.1. Similar levels of inhibition were observed at 11.7 to 56.7 mg/l; this was not unexpected given the similarity of the initial measured levels of the test substance at these concentrations, which approximated the limit of solubility of the test substance under the conditions of the test. The reason for the reduction in cell growth at 1.1 mg/L was unknown but could not be attributed to the presence of test substance because no adverse effects occurred at the next two highest levels.

Although less than 50% inhibition occurred at 56.7 mg/L, further testing at higher concentrations in an attempt to define an EC₅₀ value was not considered necessary because the solubility of the test substance had been exceeded. Mean values of cell densities in algal cultures on each sampling occasion are presented in Table B.9.2.2.

Table B.9.2.1: Inhibition of growth of algae exposed for 72 hours to refined pyrethrum extract

Exposure concentrations (mg refined pyrethrum extract/L)	Mean AUC ^{a,b} 72 h (% inhibition)		Mean growth rate ^c at 0-72 h (% inhibition)
	Nominal ^d	Measured ^e	
Control	ND ^f	3497	7.239
0.5	0.265	3793	7.333
1.1	0.615	3038	6.892
2.42	0.889	3394	7.230
5.32	1.25	3510	7.202
11.7	1.30	3032	6.940
25.8	1.95	2792	6.854
56.7	1.78	2649	6.885

^aAUC = area under the curve

^b × 10⁴

^c × 10⁻²

^dnominal loading rates

^emean measured concentrations

^fND = none detected (<0.01 mg/L)

Table B.9.2.2: Cell density of algae exposed for 72 hours to refined

pyrethrum extract

Exposure concentrations (mg refined pyrethrum extract/L)	Mean cell densities ($\times 10^4$ cells/mL)				
	Nominal ^a	Measured ^b	24 h	48 h	72 h
Control		ND ^c	9.91	46.2	184
0.5		0.265	12.1	50.3	196
1.1		0.615	10.6	47.0	143
2.42		0.889	8.17	44.6	182
5.32		1.25	11.6	47.8	179
11.7		1.30	9.88	45.0	148
25.8		1.95	9.20	40.0	139
56.7		1.78	7.67	34.0	142

^anominal loading rates

^bmean measured concentrations

^cND = none detected (<0.01 mg/L)

Conclusions:

After 72 hours of exposure to Refined Pyrethrum Extract, neither the E_bL₅₀, nor the E_rL₅₀ values for *Selenastrum capricornutum* Strain no (CCAP 278/4) could be calculated because less than 50% inhibition occurred during the definitive test. Similar levels of inhibition were obtained at the three highest nominal loading rates employed in the test (11.7 to 56.7 mg/L); the initial measured concentrations of test substance at these concentrations were similar (2.25 to 2.89 mg/l). Consequently, the E_bL₅₀ and E_rL₅₀ values were >56.7 mg/L based on the nominal loading rate and >1.95 mg/L expressed in terms of the highest mean measured level. The "no observed effect loading rate" (NOELR) for area under the growth curve and for growth rate was 5.32 mg/L (nominal); the NOEC was 1.25 mg/L (measured). (Jenkins CA 2003)

Analytical data on concentrations in the test media

Materials and Methods:

MATERIALS

As per Jenkins CA 2003 analysis by GC-FID.

Findings:

The results of chemical analysis are given in Table B.9.2.3. The overall mean measured levels of refined pyrethrum extract were 0.265, 0.615, 0.889, 1.25, 1.30, 1.95 and 1.78 mg/L (lowest to highest nominal concentrations respectively). Failure to achieve the nominal concentrations at the higher levels was attributed to the limit of aqueous solubility of the test substance having been exceeded.

After 72 hours, analysis of a sample of medium containing refined pyrethrum extract at 0.5 mg/L, which had been incubated without algal cells showed a small decrease in the measured level compared to medium that had been incubated with algal cells (22% compared to 63%). These results indicate that the stability of the test substance was affected by the presence of algal cells.

Table B.9.2.3: Measured concentrations

Nominal concentration refined pyrethrum extract	Measured refined pyrethrum extract concentrations	Overall geometric mean				
		0 h	% N	72 h	% N ^b	% t0 ^c

0	ND ^a	-	ND	-	-	-
0.500	0.432	86	0.162	32	38	0.265
0.500 ^d	-	-	0.339	68	78	-
1.10	0.944	86	0.401	36	42	0.615
2.42	1.48	61	0.534	22	36	0.889
5.32	1.89	35	0.823	15	44	1.25
11.7	2.25	19	0.752	6	33	1.30
25.8	2.89	11	1.31	5	45	1.95
56.7	2.34	4	1.36	2	58	1.78
56.7 ^d	-	-	2.27	4	97	-

^aND = none detected (<0.01 mg/L)

^b% N = measured concentration as a % of nominal concentration

^c% t0 = measured concentration after 72 h as % of starting concentrations

^dculture medium incubated under test condition without algal cells

Conclusions:

At the start of the test, the measured levels of refined pyrethrum extract in samples of the test cultures ranged from 86% of its nominal value at the lowest concentration (0.5 mg/L) to 4% of nominal at the highest concentration (56.7 mg/L). After 72 hours, the measured levels had decreased, ranging between 2 and 36% of their nominal values; between 33 and 58% of their initial values. The overall mean measured levels of refined pyrethrum extract were 0.265, 0.615, 0.889, 1.25, 1.30, 1.95 and 1.78 mg/L. (Jenkins CA 2003)

2) *Natural Pyrethrum: algal inhibition test. Mead C., McKenzie J., 2003*

Guidelines:

EC Methods for Determination of ecotoxicity annex to Directive 92/69/EEC Part C, Method 3 and OECD 201

Testing Laboratory and dates:

conducted the study during the period April 07, 2003 to April 10, 2003.

GLP:

Fully GLP compliant.

The study is acceptable.

Executive Summary:

In an algal growth inhibition test, six replicate flasks with *Scenedesmus subspicatus* cultures were exposed to refined pyrethrum extract (76.54% total Pyrethrins) at a single nominal concentration of 2.32 mg/L for 72 hours under constant illumination and shaking at a temperature of 24 ± 1°C. Samples of the algal populations were removed daily and cell concentrations determined for each control and treatment group.

It could be concluded that EC₅₀ values were greater than 2.32 mg a.s./L, and correspondingly, the No Observed Effect Concentration was 2.32 mg a.s./L if based on nominal concentrations. EC₅₀ values were greater than 1.27 mg/L and correspondingly, the NOEC was 1.27 mg/L if based on mean measured test concentrations.

Materials and Methods:

MATERIALS

Test Material:	Natural pyrethrum
Description:	Slightly cloudy, viscous liquid
Lot/Batch #:	LAB-5
Purity:	76.54% total Pyrethrins
CAS #:	Not documented
Stability of test compound:	Stability during the course of this study was demonstrated by analysis

Vehicle and/or control:	positive	Dimethylformamide (DMF)
Test organisms - Species:		<i>Scenedesmus subspicatus</i>
Strain:		CCAP 276/20
Source:		Culture Collection of Algae and Protozoa (CCAP), Institute of Freshwater Ecology, The Ferry House, Far Sawrey, Ambleside, Cumbria
Culture medium:		Sterile algal nutrient medium as recommended in OECD Procedure 201 and EC Directive 92/69/EEC Official Journal no L383A, part C3
Pre-culture conditions:		
Light levels:		7000 lux (approx.)
Photoperiod:		Continuous
Temperature:		21 ± 1°C
Cell density:		2.11 × 10 ⁶ cells/mL
Environmental conditions:	test	
Light levels:		7000 lux (approx)
Photoperiod:		Continuous
Temperature:		24 ± 1°C
Incubation period:		72 hours
Cell maintenance:		Cells were maintained in volumetric flasks. Gaseous exchange and suspension of the cells were ensured by shaking at approximately 150 rpm

Findings

VALIDITY CRITERIA

Analysis of the test preparations at 0 hours showed the measured concentrations to be 84% and 92% of the nominal value. After 72 hours, there was a marked decline in the measured concentrations to 13% and 29% of the nominal value. Stability analyses conducted indicated that the test material was stable in culture medium for the test period, and hence the decline in measured test concentrations was considered to be due to adsorption to algal cells. Recovery analyses conducted in the presence of algal cells showed that immediate adsorption to algal cells did not occur, however this does not preclude long term adsorption occurring over the test period in the presence of actively growing algal cells. Adsorption was not a factor in the stability analyses conducted, as no algal cells were present.

Given the decline in measured concentrations over the test period, the results of the test were based on the mean measured test concentrations in order to give a "worst case" analysis of the data. The mean measured test concentrations are given in Table B.9.2.4.

Table B.9.2.4: Mean measured test concentrations on replicates 1 to 6

Nominal Concentration (mg/L)	Mean Measured Concentration (mg/L)	Expressed as a Percent of the Nominal Concentration (%)
2.32 (R ₁ -R ₃)	1.12	48
2.32 (R ₄ -R ₆)	1.41	61
R ₁ -R ₆ = Replicates 1 to 6		

The results were based on the mean measured test concentrations of 1.27 mg/L. The use of mean measured test concentrations in the calculation of the results of the test had no significant effect.

CELL DENSITY AND GROWTH RATE

Calculated values for the levels of inhibition of growth rate and biomass are given in Table

B.9.2.5, where it is made clear that no inhibition effect was observed on both parameters. Cell densities in the test are given in Table B.9.2.6 .

The cell concentration of the control cultures increased by a factor of 108 and the cell concentration of the solvent control cultures increased by a factor of 90 during the test in line with the OECD Guideline that states the enhancement must be at least by a factor of 16 after 72 hours.

Table B.9.2.5: Inhibition of growth rate and biomass

Nominal concentration (mg/L)	Area under curve at 72 h	% Inhibition	Growth rate (0 - 72 h)	% Inhibition
Control	2.18×10^7	-	0.065	-
Solvent control	2.04×10^7	-	0.062	-
2.32	2.12×10^7	[4]	0.062	0

[4] increase in growth as compared to the controls

E_bC₅₀ (72 h) >2.32 mg/L

E_rC₅₀ (0 - 72 h) >2.32 mg/L

Table B.9.2.6: Cell densities of algae exposed for 72 hours to the test substance

Nominal Concentration (mg/L)		Cell density ⁽¹⁾ (cells per mL)			
		0 h	24 h	48 h	72 h
Control	R ₁	1.15×10^4	1.26×10^5	2.02×10^5	1.22×10^6
	R ₂	1.01×10^4	1.26×10^5	2.11×10^5	1.21×10^6
	R ₃	1.18×10^4	1.26×10^5	2.11×10^5	1.19×10^6
	Mean	1.11×10^4	1.26×10^5	2.08×10^5	1.21×10^6
Solvent control	R ₁	1.23×10^4	1.24×10^5	2.25×10^5	1.01×10^6
	R ₂	1.19×10^4	1.23×10^5	1.91×10^5	1.30×10^6
	R ₃	1.22×10^4	1.21×10^5	2.31×10^5	9.47×10^6
	Mean	1.21×10^4	1.23×10^5	2.16×10^5	1.09×10^6
2.32	R ₁	1.42×10^4	1.21×10^5	1.86×10^5	1.18×10^6
	R ₂	1.43×10^4	1.22×10^5	2.03×10^5	1.19×10^6
	R ₃	1.33×10^4	1.19×10^5	1.83×10^5	1.18×10^6
	R ₄	1.35×10^4	1.21×10^5	2.08×10^5	1.17×10^6
	R ₅	1.31×10^4	1.25×10^5	2.12×10^5	1.19×10^6
	R ₆	1.32×10^4	1.22×10^5	2.10×10^5	1.21×10^6
	Mean	1.36×10^4	1.22×10^5	2.00×10^5	1.19×10^6

(1) Cell densities represent the mean number of cells per ml calculated from the mean of the cell counts from 3 counts for each of the replicate flasks

R₁ - R₆ = Replicates 1 to 6

Conclusions:

Neither the growth, nor the biomass of *Scenedesmus subspicatus* was affected by the presence of the test material over the 72-hour exposure period. The NOEC was 2.32 mg/L. Based on the mean measured test concentrations of the test media the EC₅₀ values were estimated to be > 1.27 mg/L. (Mead C, McKenzie J 2003)

Analytical data on concentrations in the test media

Materials and Methods:

MATERIALS:

As per Mead C, McKenzie J 2003 analysis by HPLC with detection UV at 230nm.

Findings:

The detection system was found to have acceptable linearity (R² = 0.9999, ranging from 0 to

101 mg/L). The recoveries obtained (mean: 85%) allowed for a consideration of the method as being sufficiently accurate and precise for the purposes of the test. The limit of quantitation was assessed down to 0.020 mg/L and the stability and recovery from test samples prepared as in the trials showed that there was a marked decline in measured test concentrations over the test period (results given in Table B.9.2.7 to Table B.9.2.9), though this decline was considered to be due to adsorption to algal cells.

Table B.9.2.7: Verification of test concentrations

Nominal concentration (mg/L)	Recoveries		
	(mg/L)	(%)	Mean (%)
2.32	2.09	90	90
2.32	2.07	89	
2.32 plus algae	2.07	89	Not applicable

Table B.9.2.8: Stability results from test samples prepared as in the trials after an exposure period of 72 hours in different conditions

Nominal concentration (mg/L)	2.32
Concentration found initially (mg/L)	2.08
Concentration found after storage in light conditions (mg/L)	1.97
Expressed as a percent of the initial concentration	95
Concentration found after storage in dark conditions (mg/L)	2.20
Expressed as a percent of the initial concentration	106
Concentration found after storage in dark conditions (mg/L)-unsonicated sample	2.05
Expressed as a percent of the initial concentration	98

Table B.9.2.9: Recovery results from test samples prepared as in the trials before and after an exposure period of 72 hours

Sample	Nominal concentration (mg/L)	Concentration found (mg/L)	Expressed as a percent of the nominal concentration (%)
0 hours	Solvent control	<LOQ	-
	2.32 R ₁ -R ₃	1.95	84
	2.32 R ₄ -R ₆	2.14	92
72 hours	Solvent control	<LOQ	-
	2.32 R ₁ -R ₃	0.294	13
	2.32 R ₄ -R ₆	0.672	29

Conclusions:

At the start of the test, the measured levels of test item in samples of the test cultures ranged from 84% of its nominal value to 92%. After 72 hours, the measured levels had decreased, ranging between 13 and 29% of the nominal concentration. Although the addition of algae to the medium showed that they did not influence on the immediate recovery of test item, it was considered after a 72-hour exposure period, that there was a process of adsorption of the test item to the algal cells. The EC₅₀ was therefore based on mean measured concentrations. (Mead C, McKenzie J 2003).

Acute fish (DAR renewal)

Data point:	KCA 8.2.1/04
Report author	Teigeler, M.
Report year	2013
Report title	Fish acute toxicity test over 96 h under flow through conditions (OECD TG 203, 1992) Acute toxicity of refined pyrethrum concentrate on the zebrafish (<i>Danio rerio</i>)
Report No	GAB-034/4-32/A
Document No	GAB-034/4-32/A
Guidelines followed in study	OECD Guideline 203 (1992) EEC method C.1 (1992)
Deviations from current test guideline	None
Previous evaluation	No, not previously submitted
GLP/Officially recognised testing facilities	Yes, conducted under GLP/Officially recognised testing facilities
Acceptability/Reliability:	Yes

Executive Summary

The 96-hour acute toxicity of refined pyrethrum concentrate to the zebrafish (*Danio rerio*) was determined under flow-through conditions in a dose response test. The nominal test concentrations were 50.0, 25.0, 12.5, 6.25 and 3.125 µg/L, plus a dilution water control was tested in parallel. Seven fish were tested per test concentration and the control. The mortality and sublethal effects were determined after 3 h, 24 h, 48 h, 72 h and 96 h. The 96-hour LC₅₀ was determined to be 19.8 µg refined pyrethrum concentrate/L. The NOEC based on observations of mortality and clinical signs was determined to be 13.2 µg/L (mean measured concentration).

I. MATERIALS AND METHODS

A. MATERIALS

1. **Test material:** Refined pyrethrum concentrate
Description: Liquid, limpid, pale yellow
Lot/Batch: FEK-99
Content of a.s.: Total Pyrethrins: 57.03% composed as follows: Pyrethrin 1: 37.12%, Pyrethrin 2: 19.91%
Water solubility: emulsifiable

B. STUDY DESIGN AND METHODS

1. **Test animals:** Zebrafish (*Danio rerio*) (Teleostei, Cyprinidae)
Total weight: Mean: 0.071 ± 0.007 g
Total length: Mean: 2.0 + 0.3 cm
Source: Test facility
Acclimation: Minimum of 12 days
Diet: Fed *ad libitum* throughout the holding period with live brine shrimp (*Artemia spp.*) nauplii and ground flake food Tetra Min® (Tetra Werke, Melle, Germany) once daily, except during the test as well as 24 h before test start.
2. **Dilution water:** Purified drinking water
Total hardness: 1.1 mmol/L
Alkalinity: 1.8 mmol/L
pH: 7.75
Conductivity: 261.2 µS/cm
3. **Test vessels:** Glass aquaria with a volume of 25 L
4. **Environmental conditions:**
Temperature: 23.0 ± 2 °C
pH: 7.9 – 8.3
Dissolved oxygen: 77 – 114 % of oxygen saturation
Photoperiod: 12 hours light: 12 hours darkness
5. **Animal assignment and treatment:**

Zebrafish were exposed to nominal concentrations of 50.0, 25.0, 12.5, 6.25 and 3.125 µg/L for a period of 96 hours under flow-through conditions. The test included one control with dilution water only. Seven fish each were used for the test concentrations and for the control. One test vessel per test concentration was installed.

For each replicate vessel, an individual dosage system was used. Dilution water was pumped by a water dosage pump (membrane pump, Prominent, Heidelberg, Germany) into a mixing chamber, placed on a magnetic stirrer. The stock solution was added into the mixing chamber via a stock solution dosage pump (membrane pump with a stainless-steel

head, Prominent, Heidelberg, Germany). The prepared test solution flowed into the test vessels via flexible tubes. The daily water exchange rate was 5 volumes. The dilution water control was served by dilution water only. For every test vessel a water flow rate of 5.21 L/h per vessel was adjusted, resulting in a daily turnover of 5 volumes. The flow-through system was served by test solutions at least 24 h before adding the fish. The test animals were introduced in the test vessels after the concentrations of the test substance were within an acceptable range ($\pm 20\%$ of nominal concentrations). The mean fish weight resulted in a loading of 0.02 g/L test medium.

6. Dose preparation:

For the preparation of the stock solution acetone was used as a solvent. Stock solution was prepared by adding 15.7, 31.3, 62.5, 125 and 250 mg of the test substance into 20 mL of acetone. Pre-warmed brown glass bottles were used. In the pre-warmed bottles 2.2 mL of each stock solution were added, blown with nitrogen and allowed the evaporation of the solvent. The bottles were then filled with 11 L Cu-free water and allowed to stir for 24 hours. This procedure was repeated daily for each concentration.

7. Measurements and observations:

Mortality and abnormal behaviour were recorded after 3 h, 24 h, 48 h, 72 h and 96 h. Dead animals were eliminated from the vessels as soon as they are discovered.

Oxygen concentration, pH and temperature were measured directly before adding the fish and afterwards once per day. Water samples from all test vessels of the treatment levels and the control were taken for analysis at test start, after 48 hours and at test end (after 96 hours). The active substances were analysed using LC-MS. The method validation data are presented in M-CA, Section 4, Point CA 4.1.2/27.

8. Statistics:

The test results were statistically analysed to determine LC₁₀ and LC₅₀ values together with 95 % confidence intervals using Probit analysis assuming log-normal distribution of the values. The computer program ToxRat was used for statistical evaluations.

II. RESULTS AND DISCUSSION

A. VALIDITY CRITERIA

- The mortality in the controls should not exceed 10% at the end of the test (observed: 0%).
- The dissolved oxygen concentration should be at least 60% of the air saturation throughout the test (observed: $\geq 77\%$ of oxygen saturation).
- The concentration of the test substance should be at least 80% of the nominal concentration throughout the test (observed: $\geq 63.9\%$ of nominal concentration). Therefore, the results are based on the mean measured concentrations of the test substance.

All validity criteria for OECD Guideline 203 were met for the control group.

B. MORTALITY AND SUBLETHAL EFFECTS

After three hours exposure with refined pyrethrum concentrate the fish showed already symptoms as inactivity and incoordination at mean measured concentrations of 29.8 and 42.5 $\mu\text{g/L}$. In both concentrations the fish were mainly found to be at the water surface. A mortality of 85.7% at 42.5 $\mu\text{g/L}$ and 14.3 % at 29.8 $\mu\text{g/L}$ occurred after 24 hours. The surviving fish still showed inactivity and incoordination at these concentrations. After 48 hours a mortality of 100 % was found at concentrations of 29.8 and 42.5 $\mu\text{g/L}$.

No clinical signs and no mortality occurred at concentrations of 3.30, 7.09 and 13.2 μg refined pyrethrum concentrate/L over the whole experimental period (Table 9.2.1-10).

Table 9.2.1-10: Effect on mortality and sub-lethal effects of zebrafish (7 fish per concentration) exposed to refined pyrethrum concentrate

Mean measured concentration ($\mu\text{g/L}$)	Cumulative mortality (%)				
	3-hours	24-hours	48-hours	72-hours	96-hours
Control	0	0	0	0	0
3.30	0	0	0	0	0
7.09	0	0	0	0	0
13.2	0	0	0	0	0
29.8	0 ^{abc}	14.3 ^{cd}	100	100	100
42.5	0 ^{ad}	85.7 ^{de}	100	100	100

^a All fish were on water surface

^b Three fish showed incoordination

^c One fish showed inactivity

^d All fish showed incoordination

^e Four surviving fish showed incoordination

^f Six surviving fish were on water surface

^g One surviving fish showed incoordination

^h One surviving fish was on water surface

Based on the results of this study, the 96-hour LC₅₀ was determined to be 19.8 µg refined pyrethrum concentrate/L. The NOEC, based on observation of mortality and clinical signs, was determined to be 13.2 µg/L (mean measured concentrations) (Table 9.2.1-11).

Table 9.2.1-11: LC_x values for zebrafish exposed to refined pyrethrum concentrate after 96 hours

Endpoint	Mean measured concentration (µg refined pyrethrum concentrate/L)
96 hour-LC ₁₀ (99% confidence intervals)	18.2 (n.d.)
96 hour-LC ₂₀ (99% confidence intervals)	18.7 (n.d.)
96 hour-LC ₅₀ (99% confidence intervals)	19.8 (n.d.)
NOEC	13.2

n.d.: not determined due to mathematical reasons or inappropriate data

C. ANALYSIS

The measured concentration of the test substance ranged from 63.9 % to 132.5 % of nominal concentration. Mean measured concentrations were calculated to be 3.30, 7.09, 13.2, 29.8 and 42.5 µg/L. The evaluation of the effects was based on the mean measured concentrations of the test substance (Table 9.2.1-12).

Table 9.2.1-12: Measured concentrations of refined pyrethrum concentrate in the exposure solutions

Nominal concentration (µg test substance/L)	Mean measured concentration (µg test substance/L)	Percent of nominal (%)
Control	n.a.	n.a.
3.125	3.30	105.6
6.25	7.09	113.4
12.5	13.2	105.8
25.0	29.8	119.2
50.0	42.5	85.0

n.a.: Not applicable

D. DEFICIENCIES

None.

III. CONCLUSION

Based on the results of this study, the 96-hour LC₅₀ of refined Pyrethrum concentrate to zebrafish was determined to be 19.8 µg refined Pyrethrum concentrate/L. The NOEC, based on observation of mortality and clinical signs, was determined to be 13.2 µg/L (mean measured concentrations).

Assessment and conclusion by applicant

Assessment:

The deviation of measured test substance deviated by more than 20% from nominal. All test concentrations were analysed at the beginning and at the end of the test and at an additional point of time (48 h). The results are based on mean measured values. Concentrations refer to total pyrethrins, even though referred to as "test item" or "Pyrethrum concentrate". According to guideline OECD 203 (June 2019) fish should be observed twice per day. In the current test they were inspected twice on day 1 and once per day on days 2-4.

Conclusion:

The study complies with the data requirements given in Commission Regulation No 283/2013.

The 96-hour LC₅₀ of refined pyrethrum concentrate to zebrafish was determined to be 19.8 µg refined Pyrethrum concentrate/L. The NOEC, based on observation of mortality and clinical signs, was determined to be 13.2 µg/L (mean measured concentrations).

Assessment and conclusion by RMS

The study has been evaluated by RMS in accordance to OECD guideline 203 (1992). All validity criteria were met despite the following deviation from OECD guideline 203 (2019): a minimum of 2 biological observations on days 2-4 should be inspected twice per day, whilst in this study biological observations were inspected only once per day during days 2-4.

Moreover, OECD guideline 203 (2019) states that « the LC₅₀, the confidence limits (95%) and the slope of the curve should be estimated using appropriate statistical methods », but in this study no confidence limits are reported due to

mathematical reasons or inappropriate data. In addition, it's reported in the KCA 8.2.1/04 (Table 14) that the variance of the slope *b* of probit regression is much more higher than the estimated value of *b*. At this regard, OECD guideline 203 (2019) specifies that « When an experiment results in only one concentration with partial mortality or no concentration with partial mortality, classical maximum likelihood methods cannot be used to estimate the LC₅₀, the slope of the concentration-response curve cannot be estimated, and a confidence interval for the LC₅₀ may not be estimable. In such cases, estimates of the LC₅₀ can be made using various techniques such as the Spearman-Kärber method (Stephan, 1977), the binomial method (USEPA, 2002), the moving average method (ISO, 1996), or as a last resort, the graphical method (USEPA, 2002). These non classical methods can give precise LC₅₀ estimates and are useful to evaluate acute fish studies yielding results that cannot be analysed using classical probit maximum likelihood techniques. » In this study, no concentrations resulted in partial mortality but linear maximum likelihood regression was used to fit the probit model. However, as response to the issue raised by zRMS, the applicant provided a statistical re-analysis of LC₅₀ value based on binomial method (Report N. 1504359.UK0 - 6297) that confirmed the previously derived endpoint.

The recalculation of endpoints using the binomial method according to OECD 203 (2019) resulted in LC₅₀ value of 19.83 µg/L (95% CI: 13.2 – 29.8 µg/L) for *Danio rerio* (mean measured concentrations). The NOEC, based on observation of mortality and clinical signs, was determined to be 13.2 µg/L (mean measured concentrations).

Data point:	KCA 8.2.1/05
Report author	Teigeler, M.
Report year	2013
Report title	Fish acute toxicity test over 96 h under flow through conditions (OECD TG 203, 1992) Acute toxicity of refined pyrethrum concentrate on the three-spined stickleback (<i>Gasterosteus aculeatus</i>)
Report No	GAB-034/4-32/G
Document No	GAB-034/4-32/G
Guidelines followed in study	OECD Guideline 203 (1992) EEC method C.1 (1992)
Deviations from current test guideline	None
Previous evaluation	No, not previously submitted
GLP/Officially recognised testing facilities	Yes, conducted under GLP/Officially recognised testing facilities
Acceptability/Reliability:	Yes

Executive Summary

The 96-hour acute toxicity of refined pyrethrum concentrate to the three-spined stickleback (*Gasterosteus aculeatus*) was determined under flow-through conditions in a dose response test. The nominal test concentrations were 50.0, 25.0, 12.5, 6.25 and 3.125 µg/L, plus a dilution water control was tested in parallel. Seven fish were tested per test concentration and the control. The mortality and sublethal effects were determined after 3 h, 24 h, 48 h, 72 h and 96 h. The 96-hour LC₅₀ was determined to be 10.9 µg refined pyrethrum concentrate/L. The NOEC based on observations of mortality and clinical signs was determined to be 6.57 µg/L (mean measured).

I. MATERIALS AND METHODS

A. MATERIALS

1. Test material: Refined pyrethrum concentrate
- Description: Liquid, limpid, pale yellow
- Lot/Batch: FEK-99
- Content of a.s.: Total Pyrethrins: 57.03% composed as follows: Pyrethrin 1: 37.12%, Pyrethrin 2: 19.91%

B. STUDY DESIGN AND METHODS

1. Test animals: Three-spined stickleback (*Gasterosteus aculeatus*) (Teleostei, Gasterosteidae)
- Total weight: Mean: 0.66 ± 0.21 g
- Total length: Mean: 4.1 ± 0.2 cm
- Source: Collected from a pond plant at Kirchhundem-Albaum. The plant was fed with a steady stream of freshwater from a near runnel.
- Acclimation: Minimum of 12 days

Diet:	Fed <i>ad libitum</i> throughout the holding period with live brine shrimp (<i>Artemia spp.</i>) nauplii and ground flake food Tetra Min® (Tetra Werke, Melle, Germany) once daily, except during the test as well as 24 h before test start.
2. Dilution water:	Purified drinking water
Total hardness:	1.1 mmol/L
Alkalinity:	1.8 mmol/L
pH:	7.75
Conductivity:	261.2 µS/cm
3. Test vessels:	Glass aquaria with a volume of 25 L
4. Environmental conditions:	
Temperature:	14.9 – 15.2 °C
pH:	7.8 – 8.4
Dissolved oxygen:	88 – 98 % of oxygen saturation
Photoperiod:	12 hours light: 12 hours darkness

5. Animal assignment and treatment:

The test fish were exposed to nominal concentrations of 50.0, 25.0, 12.5, 6.25 and 3.125 µg pyrethrins/L for a period of 96 hours under flow-through conditions. The test included one control with dilution water only. Seven fish each were used for the test concentration and for the control. One test vessel per test concentration was installed.

For each replicate vessel, an individual dosage system was used. Dilution water was pumped by a water dosage pump (membrane pump, Prominent, Heidelberg, Germany) into a mixing chamber, placed on a magnetic stirrer. The stock solution was added into the mixing chamber via a stock solution dosage pump (membrane pump with a stainless steel head, Prominent, Heidelberg, Germany). The prepared test solution flowed into the test vessels via flexible tubes. The daily water exchange rate was 5 volumes. The dilution water control was served by dilution water only. For every test vessel a water flow rate of 5.21 L/h per vessel was adjusted, resulting in a daily turnover of 5 volumes. The flow-through system was served by test solutions at least 24 h before adding the fish. The test animals were introduced in the test vessels after the concentrations of the test substance were within an acceptable range (± 20 % of nominal concentrations). The mean fish weight resulted in a loading of 0.18 g/L test medium.

6. Dose preparation:

For the preparation of the stock solution acetone was used as a solvent. Stock solution was prepared by adding 15.7, 31.3, 62.5, 125 and 250 mg of the test substance into 20 mL of acetone. Pre-warmed brown glass bottles were used. In the pre-warmed bottles 2.2 mL of each stock solution were added, blown with nitrogen and allowed the evaporation of the solvent. The bottles were then filled with 11 L Cu-free water and allowed to stir for 24 hours. This procedure was repeated daily for each concentration.

7. Measurements and observations:

Mortality and abnormal behaviour were recorded after 3 h, 24 h, 48 h, 72 h and 96 h. Dead animals were eliminated from the vessels as soon as they are discovered.

Oxygen concentration, pH and temperature were measured directly before adding the fish and afterwards once per day. Water samples from all test vessels of the treatment levels and the control were taken for analysis at test start, after 48 hours and at test end (after 96 hours). The active substances were analysed using LC-MS. The method validation data are presented in M-CA, Section 4, Point CA 4.1.2/27.

8. Statistics:

The test results were statistically analysed to determine LC₁₀ and LC₅₀ values together with 95 % confidence intervals using Probit analysis assuming log-normal distribution of the values. The computer program ToxRat was used for statistical evaluations.

II. RESULTS AND DISCUSSION

A. VALIDITY CRITERIA

- The mortality in the controls should not exceed 10% at the end of the test (observed: 0%).
- The dissolved oxygen concentration should be at least 60% of the air saturation throughout the test (observed: ≥ 88 % of oxygen saturation).
- The concentration of the test substance should be at least 80% of the nominal concentration throughout the test (observed: ≥ 51.8 % of nominal concentration). Therefore, the results are based on the mean measured concentrations of the test substance.

All validity criteria for OECD Guideline 203 were met for the control group.

B. MORTALITY AND SUBLETHAL EFFECTS

After three hours exposure with refined pyrethrum concentrate the fish showed already symptoms as incoordination and a lateral body position at mean measured concentrations of 30.3 and 45.0 µg/L. In both concentrations the fish position was mainly on the bottom of the aquarium. A mortality of 71.4% at 45.0 µg/L, 28.6 % at 30.3 µg/L and 14.3% at 18.0 µg/L occurred after 24 hours. The surviving fish still showed incoordination and a lateral body position at these concentrations. After 48 hours a mortality of 100 % was found at a concentration of 45.0 µg/L. At concentrations of 18.0 and 30.3 µg/L the mortality was determined to be 42.9 % and 71.4 %. A 100 % mortality was found at 30.3 µg/L after 72

hours, whereas at 18.0 µg/L the mortality was 85.7 %. At the end of the test (96 hours) also the last fish in the treatment of 18.0 µg/L died and a mortality of 100 % was found.

No clinical signs and no mortality occurred at concentrations of 2.10 and 6.57 µg refined pyrethrum concentrate/L over the whole experimental period (Table 9.2.1-13).

Table 9.2.1-13: Effect on mortality and sub-lethal effects of three-spined stickleback (7 fish per test concentration) exposed to refined pyrethrum concentrate

Mean measured concentration (µg/L)	Cumulative mortality (%)				
	3-hours	24-hours	48-hours	72-hours	96-hours
Control	0	0	0	0	0
2.10	0	0	0	0	0
6.57	0	0	0	0	0
18.0	0	14.3 ^c	42.9 ^d	85.7 ^f	100
30.3	0 ^a	28.6 ^b	71.4 ^e	100	100
45.0	0 ^b	71.4 ^b	100	100	100

^a Two surviving fish showed incoordination, a lateral body position and were mainly on the bottom of the aquarium

^b All surviving fish showed incoordination, a lateral body position and were mainly on the bottom of the aquarium

^c Two surviving fish showed incoordination

^d One surviving fish showed incoordination and two surviving fish showed a lateral body position and were mainly on the bottom of the aquarium

^e All surviving fish showed a gasping respiration, a lateral body position and were mainly on the bottom of the aquarium

^f The surviving fish showed a lateral body position and was mainly on the bottom of the aquarium

Based on the results of this study, the 96-hour LC₅₀ was determined to be 10.9 µg refined pyrethrum concentrate/L. The NOEC, based on observation of mortality and clinical signs, was determined to be 6.57 µg/L (mean measured concentrations) (Table 9.2.1-14).

Table 9.2.1-14: LC_x values for three-spined stickleback exposed to refined pyrethrum concentrate after 96 hours

Endpoint	Mean measured concentration (µg refined pyrethrum concentrate/L)
96 hour-LC ₁₀ (95% confidence intervals)	9.75 (n.d.)
96 hour-LC ₂₀ (95% confidence intervals)	10.1 (n.d.)
96 hour-LC ₅₀ (95% confidence intervals)	10.9 (n.d.)
NOEC	6.57

n.d.: not determined due to mathematical reasons or inappropriate data

C. ANALYSIS

The measured concentration of the test substance ranged from 51.8 % to 159.9 % of nominal concentration. Mean measured concentrations were calculated to be 2.10, 6.57, 18.0, 30.3 and 45.0 µg/L. The evaluation of the effects was based on the mean measured concentrations of the test substance (Table 9.2.1-15).

Table 9.2.1-15: Measured concentrations of refined pyrethrum concentrate in the exposure solutions

Nominal concentration (µg test substance/L)	Mean measured concentration (µg test substance/L)	Percent of nominal (%)
Control	n.a.	n.a.
3.125	2.10	67.2
6.25	6.57	105.1
12.5	18.0	144.4
25.0	30.3	121.1
50.0	45.0	90.1

n.a.: Not applicable

D. DEFICIENCIES

None.

III. CONCLUSION

Based on the results of this study, the 96-hour LC₅₀ of refined pyrethrum concentrate to three-spined stickleback was determined to be 10.9 µg refined pyrethrum concentrate/L and the NOEC, based on observation of mortality and clinical signs, was determined to be 6.57 µg/L (mean measured concentrations).

Assessment and conclusion by applicant

Assessment:

The deviation of measured test substance deviated by more than 20% from nominal. All test concentrations were analysed at the beginning and at the end of the test and at an additional point of time (48 h). The results are based on mean measured values of total pyrethrins. According to guideline OECD 203 (June 2019) fish should be observed twice per day. In the current test they were inspected twice on day 1 and once per day on days 2-4.

Conclusion:

The study complies with the data requirements given in Commission Regulation No 283/2013.

The 96-hour LC₅₀ of refined pyrethrum concentrate to three-spined stickleback was determined to be 10.9 µg pyrethrins/L. The NOEC, based on observation of mortality and clinical signs, was determined to be 6.57 µg/L (mean measured concentrations).

Assessment and conclusion by RMS

The study has been evaluated by RMS in accordance to OECD guideline 203 (1992). All validity criteria were met despite the following deviations from OECD guideline 203 (2019) : (I) The mean total length of tested animals in the current study is 4.1 mm, in contrast with the recommended length range of 1-2 cm, (II) a minimum of 2 biological observations should be conducted on days 2-4, whilst in this study biological observations were inspected only once per day on days 2-4.

Moreover, OECD guideline 203 (2019) states that « the LC₅₀, the confidence limits (95%) and the slope of the curve should be estimated using appropriate statistical methods », but in this study no confidence limits are reported due to mathematical reasons or inappropriate data. In addition, it's reported in the KCA 8.2.1/04 (Table 14) that the variance of the slope b of probit regression is much more higher than the estimated value of b. At this regard, OECD guideline 203 (2019) specifies that « When an experiment results in only one concentration with partial mortality or no concentration with partial mortality, classical maximum likelihood methods cannot be used to estimate the LC₅₀, the slope of the concentration-response curve cannot be estimated, and a confidence interval for the LC₅₀ may not be estimable. In such cases, estimates of the LC₅₀ can be made using various techniques such as the Spearman-Kärber method (Stephan, 1977), the binomial method (USEPA, 2002), the moving average method (ISO, 1996), or as a last resort, the graphical method (USEPA, 2002). These non classical methods can give precise LC₅₀ estimates and are useful to evaluate acute fish studies yielding results that cannot be analysed using classical probit maximum likelihood techniques. » In this study, no concentrations resulted in partial mortality but linear maximum likelihood regression was used to fit the probit model. However, as response to the issue raised by zRMS, the applicant provided a statistical re-analysis of LC₅₀ value based on binomial method (Report N. 1504359.UK0 - 6297) that confirmed the previously derived endpoint.

The recalculation of endpoints using the binomial method according to OECD 203 (2019) resulted in LC₅₀ value of 10.88 µg/L (95% CI: 6.57 – 18.0 µg/L) for *Gasterosteus aculeatus* (mean measured concentrations).

The NOEC is 6.57 µg/L (mean measured concentrations).

Chronic fish (DAR renewal)

Data point:	KCA 8.2.2.1/02
Report author	Lee, M. R.
Report year	2012
Report title	Pyrethrins TGAI - Early Life-Stage Toxicity Test with Sheepshead Minnow (<i>Cyprinodon variegatus</i>) Following OPPTS Draft guideline 850.1400
Report No	13513.6106
Document No	N/A
Guidelines followed in study	OPPTS Draft guideline 850.1400
Deviations from current test guideline	Due to an error, one replicate in one concentration contained 25 instead of 30 organisms. In the same replicate 18 instead of 15 organisms were exposed after hatch until day 5. Days to hatch not reported.
Previous evaluation	No, not previously submitted
GLP/Officially recognised testing facilities	Yes, conducted under GLP/Officially recognised testing facilities
Acceptability/Reliability:	Yes

Executive Summary

The effects of Pyrethrins TGAI to embryos and larvae of the sheepshead minnow (*Cyprinodon variegatus*) were determined under flow-through conditions during 33 days. The exposure period included a 5-day incubation period and a 28-day post hatch exposure period. The nominal test concentrations were 1.3, 2.5, 5.0, 10 and 20 µg a.s./L, plus a dilution water and a solvent (DMF) control were tested in parallel. 120 organisms in four replicates were tested per test concentration and controls. The endpoints evaluated were embryo hatching success, percentage of embryos that produce live, normal larvae at hatch, larval survival and larval growth (total leech length and dry weight). The LOEC was determined to be 7.0 µg a.s./L and the NOEC was determined to be 3.5 µg a.s./L (mean measured).

I. MATERIALS AND METHODS

A. MATERIALS

1. **Test material:** Pyrethrins TGAI
Description: Brown liquid
Lot/Batch: 230-089
Content of a.s.: 53.48% total Pyrethrins

B. STUDY DESIGN AND METHODS

1. **Test animals:** Sheepshead minnow (*Cyprinodon variegatus*)
Age: Embryos of approx. 36 hours old
Source: Aquatic Biosystems, Inc., Fort Collins, Colorado, USA
Acclimation: The embryos were allowed to acclimate to test temperature over one hour.
Diet: On day 5, larvae were fed once with live brine shrimp nauplii (*Artemia salina*) three times daily except during the 24 hours prior to testing.
2. **Dilution water:** Dilute, filtered natural seawater from the Cape Cod Canal, Bourne, Massachusetts.
Salinity: 19 – 20‰
Total organic carbon: 1.1 – 1.3 mg/L
pH: 7.8 – 8.1
3. **Test vessels:** Glass aquaria (39 x 10 x 20 cm) with a 14.5 cm high side drain which maintained a constant volume of 6.5 L.

Embryo incubation cups were glass jars (5 cm diameter, 8 cm high) with 475- μ m nylon screen bottoms.

4. **Environmental conditions:**

Temperature:	24 – 27°C
pH:	7.7 – 8.0
Dissolved oxygen:	49 - 99% of oxygen saturation
Salinity:	19 – 22‰
Photoperiod:	16 hours light: 8 hours darkness (830 – 1000 lux)

5. **Animal assignment and treatment:**

Following the acclimation period, the embryos were impartially distributed to the embryo incubation cups five a time until all cups contained 30 embryos, and they were microscopically examined for viability. The incubation cups were then suspended in the respective exposure aquaria (one cup per replicate vessel) and exposed to nominal concentrations of 1.3, 2.5, 5.0, 10 and 20 μ g a.s./L during 33 days under flow-through conditions. The test included one control with dilution water only and one solvent control with DMF. There were 60 organisms per treatment level in four replicate exposure aquaria at the beginning of the test. On day 5 (completion of hatch), the surviving larvae were thinned to 515 organisms per replicate, resulting in 60 organisms per concentration. Test aquaria were impartially positioned in a water bath containing circulating water.

Prior to exposure initiation, a Harvard Apparatus syringe pump in conjunction with a 10-mL Spectrum Chromatography gas-tight syringe was calibrated to deliver 0.0097 mL/cycle of the 4.0 mg a.s./mL stock solution into the diluter system's chemical mixing chamber which also received 1.94 L/cycle of dilution water.

The exposure system consisted of an intermittent flow proportional diluter, a temperature-controlled water bath and a set of 28 exposure aquaria. Flow-splitting cells were employed to equally distribute the solutions to the replicate vessels at a rate of 250 mL of control and test solution vessel per cycle.

From exposure initiation to day 26, the diluter system was calibrated to deliver the control and test solutions to the exposure aquaria (49.5 L/aquarium/day) at a rate sufficient to provide approx. 7.6 aquarium volumes per 24-hour period, with a 90% replacement time of approx. 7 hours. From day to test termination (day 33), the diluter system was calibrated to deliver control and test solutions to the exposure aquaria (64.5 L/aquarium/day) at a rate sufficient to provide approx. 9.9 aquarium volumes per 24-hour period, with a 90% replacement time of approx. 5 hours.

During the 28-day post-hatch exposure period, biomass loading did not exceed 0.059 g/L flowing solution per day or 0.45 g/L of solution at any time, in any exposure aquarium.

6. **Dose preparation:**

Prior to exposure initiation and every three to four days thereafter throughout the exposure, a 4.0 mg a.s./mL stock solution was prepared by placing approx. 0.0748 g of Pyrethrins TGAI (0.0400 g as active ingredient) in a 10 mL volumetric flask and bringing it to volume with dimethylformamide (DMF). The concentration of Pyrethrins TGAI in the solution contained within the mixing chamber was equivalent to that of the highest nominal test concentration (20 μ g a.s./L) and was proportionally diluted (50%) to produce the remaining nominal test concentrations (10, 5.0, 2.5 and 1.3 μ g a.s./L).

Prior to exposure initiation and for the first 25 days of exposure, a 36 μ L/mL solvent stock solution was prepared by diluting 36 mL of DMF to 1000 mL with purified reagent water. At exposure day 26 and for the remainder of the exposure a 120 μ L/mL solvent stock solution was prepared by diluting 120 mL of DMF to 1000 mL with purified reagent water.

The concentration of DMF in the solution in the mixing chamber, as well as the high test concentration, constituted the highest DMF concentration (5.0 μ g/L). A Fluid Metering, Inc. (FMI) pump was calibrated to deliver the requisite amount of the DMF stock solution per cycle to the requisite amount of dilution water per cycle which was subsequently delivered to the solvent control vessels and the treatment level solutions. For the first 25 days of exposure, an FMI pump was calibrated to deliver 0.704 mL per cycle of a 36 μ L/mL DMF stock solution to 5.03 L of dilution water per cycle. From day 26 until exposure termination, an FMI pump was calibrated to deliver 0.224 mL per cycle of a 120 μ L/mL DMF stock solution to 5.45 L of dilution water per cycle. Throughout the entire exposure, the DMF concentration in the solvent control and the treatment levels was 5.0 μ L/mL, which was equal to the DMF concentration in the high pyrethrins test concentration

7. **Measurements and observations:**

Embryo mortalities were counted daily until hatching was complete (exposure day 5). The 28-day post-hatch larval exposure was initiated following completion of hatch on study day 5 and placed into their respective exposure aquaria. During the post-hatch exposure period, observations of larval survival, behaviour and appearance were made and recorded daily. Larval survival was estimated daily during the post-hatch period. At 28 days post-hatch exposure (study termination) the percent larval survival was determined. The surviving larvae were anaesthetised, measured to determine the total length, dried in an oven and weighed individually to determine the dry weight.

Dissolved oxygen concentration, pH, salinity and temperature were measured daily in each aquarium at exposure initiation and in alternating replicates daily thereafter until test termination.

Water samples were removed from alternating replicate solutions of each treatment level and the controls on test days 0, 5, 11, 14, 18, 21, 25 and 33 and analysed for the concentration of pyrethrins TGAI. All exposure solution samples were analysed using gas chromatography with micron electron capture detection (GC- μ ECD). The method validation data are presented in M-CA, Section 4, Point CA 4.1.2/28.

8. Statistics:

Data obtained on percent embryo hatch and larval survival and growth at study termination were analysed to identify significant differences between the treatment and control organisms. Analyses were performed using the mean organism response in each aquarium group rather than individual response values. All statistical analyses were conducted at the 95% level of certainty except in the case of Shapiro-Wilks', modified Levenes' Equality of Variance and Bartlett's Tests in which the 99% level of certainty was applied. The following procedures were used:

- An equal variance t-Test was used to evaluate the endpoints and to compare the performance the dilution water control organisms with that of the solvent control organisms. For embryo hatching success and percent normal larvae at hatch, no significant difference was determined between negative control and solvent control data. For percent survival, length and dry weight at test termination, a significant difference was determined between negative control and solvent control data.
- Statistical analysis of percentage hatching success, percent normal larvae at hatch and percentage larval survival was performed following arc-sin square-root percentage transformation of data.
- The Shapiro-Wilks' Test for normality was used to compare the observed sample distribution with a normal distribution for all endpoints. All data with the exception of percent normal larvae at hatch were normally distributed when treatment data were evaluated.
- As a check on the assumption of homogeneity of variance, data for each endpoint were analysed using Bartlett's Test or modified Levenes' Equality of Variance Test. All data with the exception of percent normal larvae at hatch met the assumption of homogeneity of variance when treatment data were evaluated.
- Percent normal larvae at hatch were evaluated using Steel's Many-One-Rank Test, a non-parametric procedure to evaluate treatment effects. Embryo hatching success and larval survival at test termination data were evaluated using Dunnett's Multiple comparison Test, a parametric procedure to evaluate treatment effects.
- Length and dry weight data were evaluated using Bonferroni's Adjusted t-Test, a parametric procedure to evaluate treatment effects for these endpoints.

A computer program CETIS Version 1.8.1.1 (Ives, 2009) was used to perform the statistical computations.

II. RESULTS AND DISCUSSION

A. VALIDITY CRITERIA

- The overall survival of fertilised eggs and post-hatch success in the control and in the solvent control should be ≥ 70 and 75%, respectively (observed: ≥ 95 and 82%, respectively).
- The dissolved oxygen concentration should be $> 60\%$ of the air saturation throughout the test (observed: $\geq 49\% - 99\%$ of oxygen saturation).
- The water temperature should be within $25 \pm 1.5^\circ\text{C}$ between test chambers or between successive days at any time during the test (observed: $24 - 27^\circ\text{C}$).
- The concentrations of the test substance were measured analytically.

The dissolved oxygen concentration was below 60% of the air saturation but it was not allowed to remain below 60% for more than 8 hours. All other validity criteria for OECD Guideline 210 were met for the control groups.

B. HATCHABILITY AND SURVIVAL

The negative control and solvent control were significantly different for larval survival, total body length and dry body weight. Therefore separate analyses were performed to assess treatment effects, i.e. first treatment groups were compared to the solvent control and then to the negative control. When comparing the treatment groups to the solvent control, the most sensitive endpoint was larval survival (LOEC = $15 \mu\text{g a.s./L}$). When comparing the treatment groups to the dilution water control, the most sensitive endpoint was larval growth (LOEC = $7.0 \mu\text{g a.s./L}$). Therefore treatment groups were compared against the dilution water control; data in order to determine the most conservative estimate of toxicity.

At the completion of the hatching period (day 5), embryo hatching success in the dilution water control and the solvent control averaged 95 and 98%, respectively. Embryo hatching success in in the 0.70, 1.7, 3.5, 7.0 and $15 \mu\text{g a.s./L}$ treatment levels averaged 93, 100, 96, 97 and 98% respectively, and was not statistically reduced from the dilution water control.

At the completion of the hatching period (day 5), the percent of live normal larvae in both controls averaged 100%. Percent of live normal larvae in the 0.70, 1.7, 3.5, 7.0 and $15 \mu\text{g a.s./L}$ treatment level averaged 100, 99, 100, 100 and 99% respectively, and was not statistically reduced from the dilution water control.

Following 28-days post-hatch exposure (study day 33), larval survival in the dilution water control and solvent control averaged 82 and 97%, respectively. Larval survival in the 0.70, 1.7, 3.5, 7.0 and $15 \mu\text{g a.s./L}$ treatment levels averaged 88, 93, 93, 85 and 12%, respectively. There was a significant reduction ($p < 0.05$) in larval survival among fish exposed to the $15 \mu\text{g a.s./L}$ treatment level compared to the negative control (82%). Hatchability and survival data are presented in Table 9.2.2.1-4.

Table 9.2.2.1-4: Percent embryo hatching success and normal larvae at completion of hatch (day 5) and survival of sheepshead minnow larvae following 33-day exposure (28 days post-hatch) to pyrethrins TGAI

Mean measured concentration (µg a.s./L)	Mean embryo hatching success (%)	Mean normal larvae at hatch (%)	Mean larval survival at Day 28 (%)
Control	95	100	82
Solvent control	98	100	97 ^a
0.70	93	100	88
1.7	100	99	93
3.5	96	100	93
7.0	97	100	85
15	98	99	12 ^b

^a Significantly difference between the dilution water control and solvent control ($p < 0.05$)

^b Significantly reduced compared to the dilution water control ($p < 0.05$)

C. GROWTH

At test termination, total length of larvae averaged 23.8 and 22.9 mm in the dilution water control and solvent control, respectively. Total length of larvae exposed to the 0.70, 1.7, 3.5, 7.0 and 15 µg a.s./L treatment levels averaged 23.4, 23.0, 22.4 and 22.3 mm, respectively. Statistical analysis determined a significant reduction ($p < 0.05$) in total length among fish exposed to the 7.0 and 15 µg a.s./L treatment levels tested compared to the dilution water control (23.8 mm). The dry weight of larvae in the dilution water control and solvent control averaged 0.0536 and 0.0451 g, respectively. Dry weight of larvae exposed to the 0.70, 1.7, 3.5, 7.0 and 15 µg a.s./L treatment levels averaged 0.0501, 0.0469, 0.0467, 0.0443 and 0.0457 g, respectively. Statistical analysis determined a significant reduction ($p < 0.05$) in dry weight among fish exposed to the 7.0 treatment level tested compared to the dilution water control (0.0536 g). Growth data are presented in Table 9.2.2.1-5.

Table 9.2.2.1-5: Total length and dry weight of sheepshead minnow larvae following 33-day exposure (28 days post-hatch) to pyrethrins TGAI

Mean measured concentration (µg a.s./L)	Mean total length (mm) (SD)	Mean dry weight (g) (SD)
Control	23.8 (0.64)	0.0536 (0.0048)
Solvent control	22.9 (0.32) ^a	0.0451 (0.0010) ^a
0.70	23.4 (0.48)	0.0501 (0.0039)
1.7	23.0 (0.50)	0.0469 (0.0028)
3.5	23.0 (0.24)	0.0467 (0.0012)
7.0	22.4 (0.58) ^b	0.0443 (0.0039) ^b
15	22.3 (1.55) ^b	0.0457 (0.0105)

SD: standard deviation

^a Significantly difference between the dilution water control and solvent control ($p < 0.05$)

^b Significantly reduced compared to the dilution water control ($p < 0.05$)

Based on the effects observed during this study, larval growth was the most sensitive indicator of toxicity, the No-Observed-Effect Concentration (NOEC) was determined to be 3.5 µg a.s./L and the Lowest-Observable-Effect Concentration (LOEC) was determined to be 7.0 µg a.s./L (mean measured).

D. ANALYSIS

The mean measured concentrations ranged from 53 to 74% of nominal with a coefficient of variation $\leq 10\%$ and defined the exposure levels as 0.70, 1.7, 3.5, 7.0 and 15 µg a.s./L (Table 9.2.2.1-6).

Table 9.2.2.1-6: Measured concentrations of pyrethrins TGAI in the exposure solutions

Nominal concentration (µg a.s./L)	Mean measured concentration (µg a.s./L) (%CV)	Percent of nominal (%)
Control	n.a.	n.a.
Solvent control	n.a.	n.a.
1.3	0.70 (6.4)	53
2.5	1.7 (10)	69
5.0	3.5 (7.8)	70
10	7.0 (4.4)	70
20	15 (8.1)	74

CV: coefficient of variation

n.a.: Not applicable

D. DEFICIENCIES

Organism assignment: In the concentration of 5 µg a.s./L replicate B contained 25 eggs instead of 25. Thus 115 eggs instead of 120 eggs were distributed to the four replicates of this treatment level. Calculations of hatching success were adjusted to account for this loading error. Further 18 instead of 15 larvae were transferred to this replicate at thinning. The three additional larvae were removed on day 5.

Time to hatch was not recorded.

Food: live brine shrimp fed instead of flake food. The different food is not considered to be a deficiency.

III. CONCLUSION

Based on the results of this study, larval growth was the most sensitive indicator of the toxicity of pyrethrins TGAI to sheepshead minnow. The LOEC was determined to be 7.0 µg a.s./L and the NOEC was determined to be 3.5 µg a.s./L (mean measured).

Assessment and conclusion by applicant

Assessment:

The test was performed according to OPPTS Draft guideline 850.1400. Even though in one replicate of the 5 µg a.s./L group loading errors occurred, the study was over-all well performed and is compliant with the OECD guideline 210. It is noted that the time to hatch was not recorded. The study is suited to determine lethal concentrations and the No-Observed-Effect Concentration based on sub-lethal parameters. An EC₁₀ was not calculated, as no clear dose-response was shown in the investigated parameters.

Conclusion:

Based on the results of this study, larval growth was the most sensitive indicator of the toxicity of pyrethrins TGAI to sheepshead minnow. The LOEC was determined to be 7.0 µg a.s./L and the NOEC was determined to be 3.5 µg a.s./L (mean measured).

Assessment and conclusion by RMS

The study has been evaluated by RMS in accordance to OPPTS Draft guideline 850.1400. A minor deviation from this study to the current OECD guideline regards the food for tested organisms : in this study live brine shrimp was used in contrast to the recommended frozen brine shrimp.

As reported in OPPTS Draft guideline 850.1400 and in OECD guideline 210, for a test to be valid, the dissolved oxygen concentration must be between 60 and 100 percent of the air saturation value throughout the test, but in this study the dissolved oxygen varies from 49% - 99% of oxygen saturation. The applicant reports that the saturation was not allowed to remain below 60% for more than 8 hours. During the commenting period, the coRMS commented on this issue indicating that the drop of O₂ levels could compromise test results. In order to statistically control for that, the most conservative assumption would be it negatively affected only controls or low concentrations and not higher concentrations. Thus, control values would be underestimated. To give a better estimate, the highest value of all tests should be used as control. E.g. control values would be for mean embryo hatching success 100 %, mean normal larvae at hatch 100 %, mean larval survival at day 28 97 %, mean total length (mm) (SD) 23.8 (0.64), mean dry weight (g) (SD) 0.0536 (0.0048). As for the endpoints larval growth (weight and length), the NOEC would be the same. But for the other endpoints, a change could occur. Re-calculations could only be done with the original data.

The LOEC is 7.0 µg a.s./L and the NOEC is 3.5 µg a.s./L (mean measured).

Chronic invertebrates (DAR renewal)

Data point:	KCA 8.2.5.2/01
Report author	Lee, M.R.
Report year	2013
Report title	Pyrethrum Stewardship Blend – Life Cycle Toxicity Test with Mysids (<i>Americamysis bahia</i>) Following Draft OPPTS Guideline 850.1350
Report No	13513.6105
Document No	N/A
Guidelines followed in study	Draft OPPTS Guideline 850.1350 (1996)
Deviations from current test guideline	None
Previous evaluation	No, not previously submitted
GLP/Officially recognised testing facilities	Yes, conducted under GLP/Officially recognised testing facilities
Acceptability/Reliability:	Yes

Executive Summary

The chronic toxicity (full life-cycle) of Pyrethrum Stewardship Blend to the mysid, *Americamysis bahia* was determined under flow-through conditions, over 28 days. The nominal test concentrations of 0.031, 0.063, 0.13, 0.25 and 0.50 µg/L (mean measured concentrations: 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L) plus a dilution water and a solvent (acetone) control were tested in parallel. Four replicates, each containing 10 Mysids, were tested per test concentration and the controls. Based on mean measured concentrations and male and female growth (total body length and dry weight, the most sensitive indicators of toxicity), the No-Observed-Effect Concentration (NOEC) was determined to be 0.25 µg/L. The Lowest-Observed-Effect Concentration (LOEC) for mysids was determined to be 0.63 µg/L. Therefore, the Maximum-

Acceptable-Toxicant Concentration (MATC) was calculated to be 0.40 µg/L. Based on linear interpolation, the 7, 14, 21 and 28-day LC₅₀ values were estimated to be > 1.1 µg/L, the highest mean measured concentration tested.

I. MATERIALS AND METHODS

A. MATERIALS

- | | | |
|----|-------------------------|---|
| 1. | Test material: | Pyrethrum Stewardship Blend |
| | Description: | Not specified |
| | Lot/Batch: | 230-089 |
| | Content of a.s.: | 53.48% as total pyrethrins (29.46% as pyrethrins I and 24.02% as pyrethrins II) |

B. STUDY DESIGN AND METHODS

- | | | |
|----|----------------------------------|---|
| 1. | Test animals: | Mysid (<i>Americamysis bahia</i>) |
| | Age: | ≤ 20 hours old at test initiation |
| | Source: | Smithers Viscient culture facility |
| | Acclimation: | not specified, in-house culture |
| | Diet: | Live brine shrimp (<i>Artemia salina</i>) twice daily. |
| 2. | Dilution water: | Filtered natural seawater from the Cape Cod Canal, Bourne, Massachusetts |
| | Salinity: | 20 - 21‰ |
| | pH: | 7.6 - 8.0 |
| 3. | Test vessels: | Glass aquarium (30 x 15 x 20 cm) with a 10-cm high side drain that maintained a constant exposure solution volume of approximately 4.5 L. |
| 4. | Environmental conditions: | |
| | Temperature: | 24 – 27 °C |
| | pH: | 7.6 – 8.0 |
| | Dissolved oxygen: | 5.02 – 7.80 mg/L (70.3 – 105% saturation) |
| | Salinity: | 20 - 21‰ |
| | Photoperiod: | 16 hours light: 8 hours darkness (230 - 310 lux) |

5. Animal assignment and treatment:

A total of 20 mysids (5 mysids per replicate, 4 replicates per concentration and controls) were exposed to nominal test concentrations of 0.031, 0.063, 0.13, 0.25 and 0.50 µg/L, a dilution water control and a solvent (acetone) control for 28 days under flow-through conditions.

The exposure system consisted of a calibrated intermittent-flow proportional diluter (Mount and Brungs, 1967) a temperature-controlled water bath, and a set of 28 exposure aquaria. During each cycle of the diluter system, approximately 500 mL of exposure solution was delivered to each replicate test vessel with a flow-splitting accuracy of 5%. During the study, the diluter provided the exposure solutions to each test vessel at a rate of approximately 7.7 aquarium volume additions per day to provide a 90% test solution replacement rate of approximately 7 hours (Sprague, 1969).

For the first 11 days of exposure, each exposure aquarium contained two retention chambers, used to retain sexually immature mysids, constructed of glass petri dishes, 10 cm in diameter, 2 cm deep, to which a 14 cm high Nitex® screen collar (350-µm mesh size opening) was attached with silicone sealant. The solution volume within the retention chambers was approximately 785 mL. Once all mysids appeared to be sexually mature (test day 12), male and female pairs were transferred to separate pairing chambers. Following this distribution, each exposure aquarium contained one retention chamber and five pairing chambers, used to retain sexually mature male and female organisms, constructed of 6-cm diameter, 1.5 cm deep petri dishes, to which a 14 cm high Nitex® screen collar (350-µm mesh size opening) was attached with silicone sealant. Solution volume within the pairing chambers was approximately 250 mL. The study was conducted in a water bath designed to maintain the test solution temperatures at 25 ± 2 °C.

During the 28-day exposure period, biomass loading did not exceed 0.0026 g/L flowing solution per day or 0.020 g/L of solution at any time, in any exposure aquarium.

6. Dose preparation:

A primary stock solution at approximately 4.0 mg/mL was prepared by adding approximately 0.07615 g of Pyrethrum Concentrate (Stewardship Blend, 0.04073 g as active ingredient) to a 10-mL volumetric flask and bringing it to volume with acetone. Each stock solution was manually mixed by inverting the flask until the solution appeared to be homogenous. A 5.8 µL/mL solvent stock solution was prepared by diluting 5.8 mL of acetone to a final volume of 1000 mL with reagent grade water in a graduated cylinder.

Prior to exposure initiation, a Harvard Apparatus syringe pump in conjunction with a 2.5-mL Hamilton gas-tight syringe was calibrated to deliver 0.00144 mL/cycle of the 4.0 mg/mL diluter stock solution into the diluter's chemical mixing chamber which also received 1.94 L of dilution water per cycle. The mixing chamber was positioned over a magnetic stir plate and was partially submerged within an ultrasonic water bath, which aided in the solubilisation of the test substance into the dilution water. The solution contained in the mixing chamber constituted the highest target test concentration (0.50 µg/L) and was subsequently diluted (50%) to provide the remaining nominal exposure concentrations (0.25, 0.13, 0.063 and 0.031 µg/L).

The concentration of acetone in the solution in the mixing chamber and the high test concentration constituted the highest acetone concentration (0.74 µL/L). An FMI pump, in conjunction with a 500-mL graduated stock bottle, was calibrated to deliver 0.64 mL/cycle of the 5.8 µL/mL solvent stock solution to 5.0 L of dilution water per cycle which was subsequently delivered to the solvent control and treatment vessels. The acetone concentration in the solvent control and the treatment levels was 0.74 µL/L, which was equal to that of the high test concentration.

7. Measurements and observations:

Observations of stress, abnormal behaviour (including discoloration, immobilization and inability to maintain position in the water column), and survival were made at the time an F1 generation pairing chamber was established and daily thereafter for 96 hours. Dead mysids were recorded and removed from each replicate test vessel daily. Missing mysids were considered dead.

Reproductive success was calculated for each replicate aquarium (treatments and the controls) as the total number of offspring produced per female. In addition, the percentage of actively reproducing females in each replicate of each treatment and the controls was determined.

At test termination (day 28), individual lengths and weights of all surviving males and females were recorded separately for each replicate of each concentration and the controls.

Temperature, dissolved oxygen concentration, pH and salinity were measured in each replicate on day 0 and alternated between replicates daily thereafter throughout the exposure period, for each treatment level and the controls.

Water samples were removed from alternating replicate solutions of each treatment level and the control on days 0, 7, 14, 21 and 28 and analysed for pyrethrins concentration (as pyrethrins I) using liquid chromatography with mass spectrometry (LC/MS/MS). The method validation data are presented in M-CA, Section 4, Point CA 4.1.2/31.

8. Statistics:

Data obtained from the test organisms were statistically analysed to establish treatment level effects. The endpoints used for determination of significant adverse effect on F0 organisms included 28-day survival, male and female survival, growth (average dry body weight and average total body length) of both male and female mysids and reproduction (number of young released per female). All statistical conclusions were made at the 95% level of certainty except in the case of the basic assumption tests, e.g., Shapiro-Wilks' Test and Bartlett's Test, in which the 99% level of certainty was applied. The following procedures were used:

- An Equal Variance Two-Sample Test was conducted to statistically compare control to the solvent control data. For this study, no significant differences were determined between the dilution water control and solvent control, therefore the dilution water control and the solvent control were pooled to establish treatment effects for all endpoints.
- Binominal endpoints (e.g., 28-day survival, male and female survival and F1 survival) were analysed using Fisher's Exact Test with Bonferroni-Holm's Adjustment.
- The Shapiro-Wilks Test for normality was conducted and compared the observed sample distribution with a normal distribution. For this study, all continuous data met this assumption.
- As a check on the assumption of homogeneity of variance, data for each endpoint were analysed using Bartlett's Test. For this study, all continuous data met this assumption.
- All continuous endpoints met the assumptions of normal distribution and homogeneity; therefore, Dunnett's Multiple Comparison Test, a parametric procedure, was used to evaluate the data.

CETIS™ was used to perform the statistical computations.

II. RESULTS AND DISCUSSION

A. VALIDITY CRITERIA

- The F0 post-pairing survival in the negative control should be > 70% at the end of the test (observed: 79%).
- The reproductively active females in the negative control should be ≥ 75% at the end of the test (observed: 100%).
- The 10th and 90th percentile of the development stage distribution in the negative control should not differ by more than 4 stages (observed: approximately 2 stages).
- The mean number of offspring produced per female surviving in the negative control should be ≥ 3 (observed: ≥ 16.5 offspring per female)

All validity criteria of the OPPTS 850.1350 Guideline were met for the control group.

B. BIOLOGICAL RESULTS

Survival and reproductive success

The biological results have been reported based on mean measured concentrations. No behavioural abnormalities were observed during the exposure period. At test termination, mean survival of 66 and 80% was observed among male mysids in the control and solvent control, respectively (pooled control = 73%). Mean survival of 83, 88, 72, 90 and 73% was observed among male mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels, respectively. Fisher's Exact Test with Bonferroni-Holm's Adjustment determined no significant difference in male survival among organisms exposed to any of the treatment levels tested compared to the pooled control data. At test termination, mean survival of 85 and 91% was observed among female mysids in the control and solvent control, respectively (pooled control = 88%). Mean survival of 90, 89, 85, 74 and 89% was observed among female mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels, respectively. Fisher's Exact Test with Bonferroni-Holm's Adjustment determined no significant

difference in female survival among organisms exposed to any of the treatment levels tested compared to the pooled control data.

Following 28 days of exposure, mean survival of 60 and 69% was observed among organisms in the control and solvent control, respectively (pooled control = 65%). Mean survival of 80, 67, 68, 69 and 41% was observed among mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels, respectively. Fisher's Exact Test with Bonferroni-Holm's Adjustment determined a significant difference in survival among organisms exposed to the 1.1 µg/L treatment level compared to the pooled control data.

At test termination, the mean number of offspring per female for organisms in the control and solvent control was 16.5 and 17.8, respectively (pooled control = 17.2 number offspring per female). The mean number of offspring per female was 16.3, 17.0, 13.3 and 3.1 among mysids exposed to the 0.044, 0.12, 0.25 and 0.63 µg/L treatment levels, respectively. Dunnett's Multiple Comparison Test determined a significant difference in the mean number of offspring per female among organisms exposed to 0.63 µg/L treatment level tested compared to the pooled control data. Since females exposed to the 1.1 µg/L treatment level did not produce any young, the 1.1 µg/L treatment level was excluded from statistical analysis.

Significant adult mortalities were observed at these doses; therefore young were not collected for the 1.1 µg/L treatment level or for two replicates of the 0.63 µg/L treatment level. Following the 96-hour observation period, mean percent survival of 93 and 98% was observed among F1 mysids in the control and solvent control, respectively (pooled control = 96%). Mean percent survival of 100, 98, 98 and 95% was observed among F1 mysids exposed to the 0.044, 0.12, 0.25 and 0.63 µg/L treatment levels, respectively. Fisher's Exact Test with Bonferroni-Holm's Adjustment determined no significant difference in F1 mysid survival among organisms exposed to any of the treatment levels statistically analysed compared to the pooled control data.

A summary of the first generation (F₀) survival, reproductive success data and the F1 generation 96-hour post-release survival is presented in Table 9.2.5.2-1.

Table 9.2.5.2-1: Mean percent first generation (F₀) survival, mean number of offspring produced per female mysids exposed to Pyrethrum Stewardship Blend during 28 days and F₁ generation 96-hour post-release survival

Mean measured concentration (µg/L)	Mean % 28-day survival (SD)	Mean no. of offspring/female (SD)	Mean % 96-hour post-release survival (SD)
Control	6 (8.9)	16.5 (5.3)	93 (9.6)
Solvent control	69 (17.0)	17.8 (4.0)	98 (5.0)
Pooled control	65	17.2 (1.6)	96
0.044	80 (7.7)	16.3 (2.4)	100 (0)
0.12	67 (19)	17.0 (1.0)	98 (5.0)
0.25	68 (7.1)	13.3 (2.9)	98 (5.0)
0.63	69 (11)	3.1 (2.0) ^b	95 (7.1)
1.1	41 (19) ^a	0 (0) ^c	n.a. ^d

SD: Standard deviation

n.a.: Not applicable

^a Significantly reduced compared to the pooled control based on Fisher's Exact Test with a Bonferroni-Holm Adjustment.

^b Significantly reduced compared to the pooled control based on Dunnett's Multiple Comparison Test.

^c Due to the survival effect observed, statistical analysis was not conducted.

^d Data excluded from statistical analysis due to significant adult mortalities.

Growth

The average total body length of male mysids in the control and solvent control was 7.09 and 7.13 mm, respectively (pooled control = 7.11 mm). The average total body length of male mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels was 7.04, 7.05, 7.06, 6.28 and 5.85 mm, respectively. Dunnett's Multiple Comparison Test determined a significant difference in the total body length of male mysids exposed to the 0.63 and 1.1 µg/L treatment levels compared to the pooled control data.

The average total body length of female mysids in the control and solvent control was 7.29 and 7.42 mm, respectively (pooled control = 7.36 mm). The average total body length of female mysids exposed to 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels was 7.46, 7.30, 7.41, 6.77 and 6.31 mm, respectively. Dunnett's Multiple Comparison Test determined a significant difference in the total body length of female mysids exposed to the 0.63 and 1.1 µg/L treatment levels compared to the pooled control data.

The average dry body weight of male mysids in the control and solvent control was 0.81 and 0.83 mg, respectively (pooled control = 0.82 mg). The average dry body weight of male mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels was 0.82, 0.85, 0.86, 0.66 and 0.62 mg, respectively. Dunnett's Multiple Comparison Test determined a significant difference in the dry body weight of male mysids exposed to the 0.63 and 1.1 µg/L treatment levels compared to the pooled control data.

The average dry body weight of male mysids in the control and solvent control was 0.81 and 0.83 mg, respectively (pooled control = 0.82 mg). The average dry body weight of male mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels was 0.82, 0.85, 0.86, 0.66 and 0.62 mg, respectively. Dunnett's Multiple Comparison Test determined a significant difference in the dry body weight of male mysids exposed to the 0.63 and 1.1 µg/L treatment levels compared to the pooled control data. The average dry body weight of female mysids in the control and solvent control was 1.15 and 1.27 mg, respectively (pooled control = 1.21 mg). The average dry body weight of female mysids exposed to the 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L treatment levels was 1.19, 1.17, 1.20, 0.87 and 0.67 mg, respectively. Dunnett's Multiple Comparison Test determined a significant difference in the dry body weight of female mysids exposed to the 0.63 and 1.1 µg/L treatment levels compared to the pooled control data.

Measurements of growth, as average total body length and average dry body weight, for all surviving adult mysids at test termination are presented in Table 9.2.5.2-2.

Table 9.2.5.2-2: Mean total body length and dry body weight of first generation (F₀) male and female mysids exposed to Pyrethrum Stewardship Blend during 28 days

Mean measured concentration (µg/L)	Mean total body length in mm (SD)		Mean dry weight in mg (SD)	
	Males	Females	Males	Females
Control	7.09 (0.28)	7.29 (0.37)	0.81 (0.11)	1.15 (0.18)
Solvent control	7.13 (0.14)	7.42 (0.10)	0.83 (0.024)	1.27 (0.072)
Pooled control	7.11 (0.07)	7.36 (0.09)	0.82 (0.03)	1.21 (0.05)
0.044	7.04 (0.13)	7.46 (0.12)	0.82 (0.046)	1.19 (0.14)
0.12	7.05 (0.049)	7.30 (0.17)	0.85 (0.030)	1.17 (0.051)
0.25	7.06 (0.21)	7.41 (0.20)	0.86 (0.063)	1.20 (0.089)
0.63	6.28 (0.15) ^a	6.77 (0.23) ^a	0.66 (0.017) ^a	0.87 (0.066) ^a
1.1	5.85 (0.30) ^a	5.31 (0.06) ^a	0.62 (0.035) ^a	0.67 (0.081) ^a

SD: Standard deviation

^a Significantly reduced compared to the pooled control, based on Dunnett's Multiple Comparison Test.

Based on linear interpolation the 7, 14, 21 and 28-day LC₅₀ values were estimated to be > 1.1 µg/L, the highest mean measured concentration tested. Based on mean measured concentrations and male and female growth (total body length and dry weight), the NOEC was determined to be 0.25 µg/L and the LOEC was determined to be 0.63 µg/L. Therefore, the MATC was calculated to be 0.40 µg/L.

Based on linear interpolation, the 7, 14, 21 and 28-day LC₅₀ values were estimated to be > 1.1 µg/L, the highest mean measured concentration tested.

D. ANALYSIS

Measured concentrations of Pyrethrum Stewardship Blend were slightly variable, but maintained the expected concentration gradient (50% dilution series) throughout the exposure. The coefficient of variation for all measured concentrations ranged from 13 to 31% and defined the exposure levels as 0.044, 0.12, 0.25, 0.63 and 1.1 µg/L (Table 9.2.5.2-3).

Table 9.2.5.2-3: Measured concentrations of Pyrethrum Stewardship Blend in the exposure solutions during the 28-day life-cycle exposure of mysids

Nominal concentration (µg/L)	Mean measured concentration (µg/L) (SD)	% CV	Percent of nominal (%)
Control	n.a.	n.a.	n.a.
Solvent control	n.a.	n.a.	n.a.
0.031	0.044 (0.0096)	22	140
0.063	0.12 (0.021)	17	190
0.13	0.25 (0.033)	13	190
0.25	0.63 (0.20)	31	250
0.50	1.1 (0.25)	23	220

SD: standard deviations

CV: coefficient of variation

n.a.: Not applicable

E. DEFICIENCIES

The protocol states that during the test the diluter will be visually inspected at least twice daily and complete check of diluter functioning will be made once daily. For the entire duration of the exposure, the diluter function was completely checked twice daily. The diluter system was visually inspected and a complete check of diluter function was made twice daily to more effectively prevent malfunctions. This deviation does not have a negative impact on the results or interpretation of the study.

III. CONCLUSION

Based on mean measured concentrations and male and female growth (total body length and dry weight, the most sensitive indicators of toxicity), the NOEC of Pyrethrum Stewardship Blend to *Americamysis bahia* was determined to be 0.25 µg/L. The LOEC was determined to be 0.63 µg/L. Therefore, the MATC was calculated to be 0.40 µg/L. Based on linear interpolation, the 7, 14, 21 and 28-day LC₅₀ values were estimated to be > 1.1 µg/L, the highest mean measured Pyrethrum Stewardship Blend concentration tested.

Assessment and conclusion by applicant

Assessment:

This study was performed in 2013 according to US-requirements. The study is considered to be acceptable.

Conclusion:

The No-Observed-effect concentration (NOEC) was determined to be 0.25 µg pyrethrins/L.

Assessment and conclusion by RMS

The study has been evaluated by RMS in accordance with OPPTS Guideline 850.1350 (1996).

The following minor deviations are reported : the photoperiod of this study (16 hours light: 8 hours darkness) is not the recommended photoperiod in the OPPTS guidance (14 hours light: 10 hours darkness) ; the body length of F0 mysids should be recorded at the first observation day (depending on time of sexual maturation) and on day 28 but in this study body length was recorded only at test termination.

All validity criteria were met and the study can be considered acceptable for riskassessment.

The NOEC is 0.25 µg pyrethrins/L (mean measured concentrations).

As regards analytical methods, the expert concluded that the method of this study is not acceptable (method is not fully validated - results are not in accordance with RD – two different reference materials were used - materials are not compliant with RD). Moreover, the composition of batch remained uncharacterized. Please refer to Volume 4 comments.

Water/sediment organisms (DAR renewal)

B.9.2.8.1. Freshwater Amphipods

Data point:	KCA 8.2.8/01
Report author	Bradley, M.W.
Report year	2013
Report title	Pyrethrum Stewardship Blend - Acute Toxicity to Freshwater Amphipods (<i>Hyalella azteca</i>) Under Flow-Through Conditions
Report No	13513.6133
Document No	N/A
Guidelines followed in study	US EPA Draft Guideline 850.1020 (1996)
Deviations from current test guideline	None. Study not an EU data requirement.
Previous evaluation	No, not previously submitted
GLP/Officially recognised testing facilities	Yes, conducted under GLP/Officially recognised testing facilities
Acceptability/Reliability:	Yes

Executive Summary

The acute toxicity of Pyrethrum Stewardship Blend to the freshwater amphipod *Hyalella azteca* was determined under flow-through conditions for 96 hours. The nominal test concentrations were 0.25, 0.50, 1.0, 2.0 and 4.0 µg/L (mean measured concentrations: 0.10, 0.24, 0.54, 0.88 and 2.2 µg/L) plus a dilution water and a solvent (acetone) control were tested in parallel. One replicate aquarium containing three retention chambers, each with ten amphipods, was included for each test concentration and the controls.

Based on nominal concentrations, the 96-hour LC₅₀ value was determined by the Trimmed Spearman-Kärber Method to be 1.5 µg/L, with 95% confidence intervals of 1.3 to 1.8 µg/L. The No-Observed-Effect Concentration (NOEC) and Lowest-Observed-Effect Concentration (LOEC) were determined to be 0.50 and 1.0 µg/L, respectively. Based on mean measured concentrations, the 96-hour LC₅₀ value was by the Trimmed Spearman-Kärber Method to be 0.76 µg/L, with 95% confidence intervals of 0.64 to 0.92 µg/L. The NOEC and LOEC were determined to be 0.24 and 0.54 µg/L, respectively.

I. MATERIALS AND METHODS

A. MATERIALS

1. **Test material:** Pyrethrum Stewardship Blend
Description: Not specified
Lot/Batch: 230-089
Content of a.s.: 53.40% as total pyrethrins (29.88% as pyrethrins I and 23.52% as pyrethrins II)

B. STUDY DESIGN AND METHODS

1. **Test animals:** Freshwater amphipod (*Hyalella azteca*)
Age: 9 days old at test initiation
Source: Smithers Viscient cultures
- Acclimation:** 48 hours
Diet: Flaked fish food suspension (YCT) three times daily
2. **Dilution water:** Laboratory well water
Total hardness: 74 – 78 mg/L as CaCO₃
Total alkalinity: 28 mg/L as CaCO₃
pH: 6.9
Conductivity: 320 – 400 µmhos/cm
3. **Test vessels:** Glass aquaria (30 x 15 x 20 cm) with a 15-cm high side drain that maintained a constant exposure solution volume of 6.8 L.
4. **Environmental conditions:**
Temperature: 22 – 25°C
pH: 7.0 – 7.7
Dissolved oxygen: 6.4 – 8.9 mg/L
Photoperiod: 16 hours light: 8 hours darkness (220 - 490 lux)
5. **Animal assignment and treatment:**

Thirty amphipods (three chambers per treatment aquarium, each containing ten amphipods) were exposed to nominal concentrations of 0.50, 1.0, 2.0, 4.0 and 8.0 µg/L, a dilution water control and a solvent (acetone) control, in a flow-through study for a duration of 96 hours. Each exposure aquarium contained three retention chambers consisting of 250-µm mesh Nitex® screen (14 cm long) attached to a 6-cm diameter petri dish (1.5 cm high) using silicone, which remained partially submerged throughout the exposure. To initiate the study, ten amphipods were added impartially to retention chambers.

The flow-through test was conducted using an exposure system consisting of an intermittent-flow proportional diluter (Mount and Brungs, 1967), a temperature-controlled water bath and a set of seven exposure aquaria. The test system was designed to provide five concentrations of the test substance, a dilution water control and a solvent control. One replicate aquarium was included for each test concentration and the controls. The diluter delivered the control and test solutions to the exposure aquaria (68 L/aquarium/day) at a rate sufficient to provide approximately 10 aquarium volumes per 24-hour period, with a 90% replacement time of approximately 6 hours.

Test aquaria were positioned in a water bath designed to maintain the test solution's temperature 23 ± 1 °C.

6. Dose preparation:

A 40 µg/mL diluter stock solution was prepared prior to exposure initiation by bringing 0.01503 g of Pyrethrum Stewardship Blend (0.00803 g as active ingredient) to a volume of 200 mL with acetone. In addition, a 510 µL/mL solvent stock solution was prepared by bringing 102 mL of acetone to a volume of 200 mL with deionized water.

Prior to exposure initiation, a Harvard Apparatus syringe pump in conjunction with a 50-mL Glenco gas-tight syringe was calibrated to deliver 0.0785 mL/cycle of the 40 µg/mL diluter stock solution to the diluter's mixing chamber which also received 0.785 L of dilution water per cycle. The mixing chamber was positioned over a magnetic stir plate and was partially submerged within an ultrasonic water bath which aided in the solubilisation of the test substance into the dilution water. The solution contained in the mixing chamber constituted the highest nominal test concentration (4.0 µg/L) and was subsequently diluted (50%) to provide the remaining nominal exposure concentrations (2.0, 1.0, 0.50 and 0.25 µg/L). The concentration of acetone in the solution in the mixing chamber and the high test concentration constituted the highest acetone concentration (0.10 mL/L). A similar system was calibrated to deliver 0.0785 mL/cycle of the 510 µL/mL solvent stock solution to 0.40 L of dilution water per cycle which was subsequently delivered to the solvent control chambers. This delivery method ensured the acetone concentration in the solvent control and the treatment levels was 0.10 mL/L, which was equal to that of the high test concentration.

7. Measurements and observations:

The number of dead amphipods, biological observations and observations of the physical characteristics in each retention chamber was recorded at test initiation and after 24, 48, 72 and 96 hours of exposure.

The pH, dissolved oxygen concentration and temperature were measured in retention chamber A of each treatment level and the control at test initiation and in alternating retention chamber replicates daily thereafter. Continuous temperature monitoring was performed in retention chamber B of the 4.0 µg/L nominal treatment level throughout the exposure period.

One water sample was removed from each test solution and the controls for analysis of pyrethrins concentration at 0 hour (test initiation), 48 hours and 96 hours (test termination). Samples were collected from the approximate midpoint of the test vessel by pipet. All exposure solution samples were analysed for pyrethrins concentration (as pyrethrins I) using liquid chromatography with mass spectrometry (LC/MS/MS). The method validation data are presented in M-CA, Section 4, Point CA 4.1.2/32.

8. Statistics:

If at least one test concentration caused mortality of $\geq 50\%$ of the test population, then CETIS® Version 1.8.4.20 (Ives, 2011) was used to calculate the LC₅₀ values and 95% confidence intervals.

II. RESULTS AND DISCUSSION

A. VALIDITY CRITERIA

- The mortality in the controls should not exceed 10% at the end of the test (observed: 0%).
- The dissolved oxygen concentration should be between 60 and 105% of the air saturation throughout the test (observed: $\geq 75\%$ of oxygen saturation).
- The concentration of solvent should not exceed 0.1 mL/L (observed: 0.10 mL acetone/L).
- Dissolved oxygen concentration, pH, temperature, and the concentration of test substance in test chambers should be measured at specified intervals (observed: Dissolved oxygen concentration, pH, temperature were measured daily and pyrethrins concentration at 0 hour, 48 hours and 96 hours).

All validity criteria for US EPA Draft Guideline OPPTS 850.1020 (1996) were met.

B. BIOLOGICAL EFFECTS

Following 96 hours of exposure, 30, 60 and 97% mortality was observed among amphipods exposed to mean measured concentrations of 0.54, 0.88 and 2.2 $\mu\text{g/L}$ (1.0, 2.0 and 4.0 $\mu\text{g/L}$ nominal), respectively. All surviving amphipods exposed to the 0.88 and 2.2 $\mu\text{g/L}$ (2.0 and 4.0 $\mu\text{g/L}$ nominal, respectively) treatment levels were observed to be lethargic. No mortality or adverse effects were observed for amphipods exposed to the remaining treatment levels tested (0.10 and 0.24 $\mu\text{g/L}$ mean measured; 0.25 and 0.50 $\mu\text{g/L}$ nominal) or the controls (Table 9.2.8.1-1).

Table 9.2.8.1-1: Mean cumulative mortality and biological observations of *Hyalella azteca* exposed to Pyrethrum Stewardship Blend

Mean measured concentration (μg total pyrethrins/L)	Mean cumulative mortality of organisms (%)			
	24 hours	48 hours	72 hours	96 hours
Control	0	0	0	0
Solvent control	0	0	0	0
0.10	0	0	0	0
0.24	0	0	0	0
0.54	0	3.3	20	30
0.88	6.7 ^a	30 ^a	47 ^a	60 ^a
2.2	23 ^a	47 ^a	97 ^a	97 ^a

^a All surviving amphipods were observed to be lethargic.

Based on nominal concentrations, the 96-hour LC₅₀ value was determined by the Trimmed Spearman-Kärber Method to be 1.5 $\mu\text{g/L}$, with 95% confidence intervals of 1.3 to 1.8 $\mu\text{g/L}$. The NOEC and LOEC were determined to be 0.50 and 1.0 $\mu\text{g/L}$, respectively. Based on mean measured concentrations, the 96-hour LC₅₀ value was determined by the Trimmed Spearman-Kärber Method to be 0.76 $\mu\text{g/L}$, with 95% confidence intervals of 0.64 to 0.92 $\mu\text{g/L}$. The NOEC and LOEC were determined to be 0.24 and 0.54 $\mu\text{g/L}$, respectively (Table 9.2.8.1-2).

Table 9.2.8.1-2: LC₅₀, NOEC and LOEC values for *Hyalella Azteca* exposed to Pyrethrum Stewardship Blend

Endpoint	Nominal concentration (μg total pyrethrins/L)	Mean measured concentration (μg total pyrethrins/L)
96-hour NOEC	0.50	0.24
96-hour LOEC	1.0	0.54
96-hour LC ₅₀ ^a (95% confidence intervals)	1.5 (1.3 – 1.8)	0.76 (0.64 – 0.92)

^a LC₅₀ values and corresponding 95% confidence intervals were determined using the Trimmed Spearman-Kärber Method

C. ANALYSIS

The mean measured concentrations ranged from 42 to 54% of nominal concentrations and defined the treatment levels tested as 0.10, 0.24, 0.54, 0.88 and 2.2 $\mu\text{g/L}$ (Table 9.2.8.1-3).

Table 9.2.8.1-3: Measured concentrations of total pyrethrins in the exposure solutions

Nominal concentration (µg total pyrethrins/L)	Mean measured concentration (µg total pyrethrins/L)	Percent of nominal (%)
Control	n.a.	n.a.
Solvent control	n.a.	n.a.
0.25	0.10	42
0.50	0.24	48
1.0	0.54	54
2.0	0.88	44
4.0	2.2	54

n.a.: not applicable

D. DEFICIENCIES

During this exposure, the minimum/maximum thermometer recorded a maximum temperature of 25 °C on test day 3 whereas the protocol stated that the test temperature would be maintained at 23 ± 1 °C. Since this temperature is within the tolerance of the test species, this deviation did not have a negative impact on the results or interpretation of the study.

III. CONCLUSION

Based on nominal concentrations, the 96-hour LC₅₀ value for Pyrethrum Stewardship Blend and *Hyalella azteca* was determined by the Trimmed Spearman-Kärber Method to be 1.5 µg/L, with 95% confidence intervals of 1.3 to 1.8 µg/L. The No-Observed-Effect Concentration (NOEC) and Lowest-Observed-Effect Concentration (LOEC) were determined to be 0.50 and 1.0 µg/L, respectively. Based on mean measured concentrations, the 96-hour LC₅₀ value was determined by the Trimmed Spearman-Kärber Method to be 0.76 µg/L, with 95% confidence intervals of 0.64 to 0.92 µg/L. The NOEC and LOEC were determined to be 0.24 and 0.54 µg/L, respectively.

Assessment and conclusion by applicant

Assessment:

The study was performed according to US EPA Draft Guideline 850.1020 (1996) and is not a data requirement for EU registration

Conclusion:

The 96-hour LC₅₀ value was determined to be 0.76 µg/L. The NOEC and LOEC were determined to be 0.24 and 0.54 µg/L, respectively.

Assessment and conclusion by RMS

The study was performed in accordance with US EPA Draft Guideline 850.1020 (1996) and respected the validity criteria.

The following minor deviation is highlighted : in this study the temperature was maintained between 22 – 25°C and the EPA guideline recommends to maintain a temperature of 18 ± 1 °C. However, many other studies show that *Hyalella azteca* can be successfully reared at higher temperature and then this deviation does not invalidate the study.

The 96-hour LC₅₀ is 0.76 (95% confidence limits : 0.64 to 0.92 µg/L) based on mean measured concentration. The NOEC and LOEC is 0.24 and 0.54 µg/L, respectively, based on mean measured concentration.

As regards analytical methods, the expert concluded that the method of this study is not acceptable (method is not fully validated - results are not in accordance with RD – two different reference materials were used - materials are not compliant with RD). Moreover, the composition of batch remained uncharacterized and the study is not accepted by method experts. Please refer to Volume 4 comments.