

Annex XV report

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57

Substance Name: Perfluorobutane sulfonic acid (PFBS) and its salts

EC Number: -

CAS Number: -

Submitted by: Norway

Date: 05.08.2019

This document has been prepared according to template: TEM-0049.03

CONTENTS

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57	11
PART I	14
JUSTIFICATION	14
1. IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES	14
1.1. Name and other identifiers of the substance.....	15
1.2. Composition of the substances	17
1.3. Identity and composition of structurally related substances.....	17
1.3.1. Salt forms - grouping.....	17
1.3.2. Perfluoroalkane sulfonic acids used in read-across	20
1.3.3. PFBS related substances.....	22
1.4. Other PFASs.....	23
1.5. Physicochemical properties.....	24
2. HARMONISED CLASSIFICATION AND LABELLING	28
3. ENVIRONMENTAL FATE PROPERTIES	28
3.1. Degradation	28
3.1.1. Abiotic degradation	29
3.1.2. Biodegradation in water.....	31
3.1.3. Biodegradation in soil.....	33
3.1.4. Read-across to similar PFASs.....	33
3.1.5. Summary and discussion of degradation	35
3.2. Environmental distribution.....	36
3.2.1. Sorption to the organic fraction of soil.....	36
3.2.2. Sorption to the mineral fraction of soil	38
3.2.3. Volatilisation	39
3.2.4. Distribution modelling.....	40
3.2.5. Field data	42

3.2.6. <i>Summary and discussion of environmental distribution</i>	59
3.3. Removal from the environment, decontamination and purification.....	60
3.4. Data indicating potential for long-range transport.....	61
3.4.1. <i>Mobility and the potential for long-range transport</i>	61
3.4.2. <i>Estimation of long-range transport potential</i>	63
3.4.3. <i>Occurrence of PFBS in remote areas</i>	64
3.4.4. <i>Long-range transport of precursors and degradation to PFBS</i>	70
3.4.5. <i>Summary of data indicating potential for long-range transport</i>	71
3.5. Bioaccumulation.....	71
3.5.1. <i>Laboratory studies on bioaccumulation</i>	72
3.5.2. <i>Persistence and mobility in relation to bioaccumulation</i>	76
3.5.3. <i>Field studies</i>	76
3.5.4. <i>Summary of bioaccumulation</i>	83
3.6. Protein binding.....	84
3.7. Enrichment in plants.....	86
3.7.1. <i>Measured levels of PFBS in plants</i>	92
3.8. Temporal trends and comparison with other PFASs.....	94
3.9. Case study: Environmental distribution of PFBS – Fuxin, China.....	95
3.10. Summary and discussion of environmental fate properties.....	97
4. HUMAN HEALTH HAZARD ASSESSMENT	99
4.1. Toxicokinetics (absorption, metabolism, distribution and elimination).....	99
4.1.1. <i>Non-human information</i>	99
4.1.2. <i>Human information</i>	101
4.1.3. <i>Summary</i>	101
4.2. Human exposure routes.....	102
4.3. Levels of PFBS in humans.....	103
4.3.1. <i>Summary</i>	108
4.4. Repeated dose toxicity.....	109

4.4.1. Non-human information	109
4.4.2. Human information	110
4.4.3. Summary and discussion of repeated dose toxicity	111
4.5. Mutagenicity	111
4.5.1. Non-human information	111
4.6. Carcinogenicity	112
4.6.1. Non-human information	112
4.7. Toxicity for reproduction.....	112
4.7.1. Effects on fertility	112
4.8. Developmental toxicity.....	113
4.8.1. Non-human information	113
4.8.2. Human information	114
4.8.3. Summary and discussion of reproductive toxicity	114
4.9. Other effects: Neurotoxicity	115
4.9.1. Non-human information	115
4.9.2. Summary and discussion of neurotoxicity	116
4.10. Other effects: Immunotoxicity.....	116
4.10.1. Non-human information	116
4.10.2. Human information.....	117
4.10.3. Summary and discussion of immunotoxicity.....	117
4.11. Other effects: Endocrine disruption	117
4.11.1. In vitro.....	117
4.11.2. In vivo	118
4.11.3. Summary.....	119
4.12. Summary and discussion of human exposure and health hazard assessment....	120
5. ENVIRONMENTAL HAZARD ASSESSMENT	123
5.1. Aquatic compartment (including sediment)	123
5.1.1. Fish.....	123

5.1.2. Aquatic invertebrates.....	129
5.1.3. Algae and aquatic plants.....	130
5.1.4. Sediment organisms.....	131
5.1.5. Other aquatic organisms.....	131
5.2. Terrestrial compartment.....	132
5.2.1. Toxicity to soil micro-organisms.....	132
5.2.2. Toxicity to terrestrial plants.....	133
5.3. Microbiological activity in sewage treatment systems	133
5.4. Toxicity to birds.....	134
5.5. Mammalian wildlife.....	135
5.6. Other effects: Endocrine disruption.....	135
5.7. Summary and discussion of the environmental hazard assessment.....	139
6. CONCLUSIONS ON THE SVHC PROPERTIES.....	141
6.1. CMR assessment.....	141
6.2. PBT and vPvB assessment	141
6.3. Assessment under Article 57 (f)	141
6.3.1. Summary of the data on the hazardous properties	141
6.3.2. Concerns arising from the substance properties.....	146
6.3.3. Overall concern and assessment of the level of concern	155
6.3.4. Conclusion on the hazard properties and equivalent level of concern assessment	160
REFERENCES FOR PART I.....	163
PART II.....	188
7. REGISTRATION AND C&L NOTIFICATION STATUS	188
7.1. Registration status	188
7.2. CLP notification status	190
8. INFORMATION ON USES OF THE SUBSTANCES	195
9. ADDITIONAL INFORMATION	197
9.1. Substances with similar hazard and use profiles on the Candidate List.....	197
9.2. Existing EU legislation.....	200

9.3. Previous assessments by other authorities	200
REFERENCES FOR PART II	202
ANNEX I: PHYSICAL-CHEMICAL PROPERTIES OF SUBSTANCES USED IN READ-ACROSS	203
ANNEX II: MODELLING NOMINAL BIOTA CONCENTRATION DEVELOPMENT OF PFBS ACCORDING TO THE CROOKES AND FISK MODEL	204

TABLES

Table 1: Substance identity, PFBS.....	15
Table 2: Substance identity, KPFBS – the most widely used and studied salt form of PFBS.....	16
Table 3: Salt forms of PFBS in ECHA's database.....	18
Table 4: Identity and composition of substances used in read-across...	21
Table 5: Physicochemical data for PFBS, as well as its potassium salt KPFBS.....	25
Table 6: Results from BIOWIN estimations of PFBS biodegradation.	31
Table 7: Percent biodegradation of KPFBS.....	32
Table 8: Stability data for C1, C4 and C8 perfluoroalkane sulfonic acids reported at the ECHA dissemination page.....	34
Table 9: Physiochemical properties needed to estimate environmental distribution for PFBS and homologues.....	41
Table 10: Environmental concentrations of PFBS in surface fresh water samples (ng/L).....	43
Table 11: Concentrations of PFBS in surface water near PFAS production facilities or known hotspots (ng/L).....	47
Table 12: Environmental concentrations of PFBS in groundwater (ng/L).	49
Table 13: Concentrations of PFBS in drinking water (ng/L).....	51
Table 14: Concentrations of PFBS in WWTP effluent and in landfill leachate (ng/L).	55
Table 15: Environmental concentrations of PFBS in marine water samples (ng/L).....	56
Table 16: Environmental concentrations of PFBS in soil and sediment samples (ng/g dw).....	58
Table 17: Environmental concentrations of PFBS in remote areas; fresh water, air, snow and ice samples (ng/L).	65
Table 18: Environmental concentrations of PFBS in marine water samples from remote regions (ng/L).....	67
Table 19: Environmental concentrations of PFBS in Arctic biota (ng/g ww).	68

Table 20: Environmental concentrations of PFBS in marine biota from other regions than remote arctic areas (ng/g ww).....	79
Table 21: Environmental concentrations of PFBS in freshwater biota (ng/g ww).....	81
Table 22: Environmental concentrations in other biota samples (ng/g ww)	82
Table 23: Uptake and concentration of PFBS in plants	88
Table 24: Uptake factors and concentrations of PFBS in terrestrial plants near hot spots.....	91
Table 25: PFBS concentrations in environmental and dietary samples..	96
Table 26: Estimated daily intakes of PFBS (ng/kg bw/day).....	96
Table 27: Serum t1/2 of PFBS in different species.	101
Table 28: Concentrations of PFBS in human tissues (ng/g ww) and blood (ng/mL)	106
Table 29: Overview of PFBS-induced health effects	121
Table 30: Overview of short-term effects on fish	124
Table 31: Summary of long-term effects on fish	127
Table 32: Summary of short-term effects on aquatic invertebrates	129
Table 33: Summary of long-term effects in aquatic invertebrates.....	130
Table 34: Summary of effects on algae	130
Table 35: Summary of effects on sediment organisms	131
Table 36: Summary of effects on marine microorganisms.....	131
Table 37: Summary of effects on birds.....	134
Table 38: Overview of transcriptional effect in F1 and F2 offspring (larvae). Modified from Table S4 in the supplementary material. Significant upregulation is shown by upwards arrow and significant downregulation is shown by downwards arrow. A dash indicates non-significant results.	137
Table 39: Summary of endocrine disruptive effects	138
Table 40: Overview of the qualitative components relevant for assessing the level of concern	156
Table 41: Registration status for PFBS and some salt forms	188

Table 42: Notified self-classifications for perfluorobutane sulfonic acid, PFBS, according to CLP criteria	190
Table 43: Notified self-classifications for potassium perfluorobutane sulfonate, KPFBS, according to CLP criteria	191
Table 44: Notified self-classifications and harmonised classifications of additional salts of PFBS that are registered under REACH	192
Table 45: Notified self-classifications according to CLP criteria of salts of PFBS that are not registered under REACH	194
Table 46: PFASs subjected to regulatory activities under REACH	198
Table 47: Physical-chemical properties of substances used in read-across (from OECD QSAR toolbox v4.2)	203

Abbreviations related to per- and polyfluoroalkyl substances (PFASs)

KPFBS	Potassium perfluorobutane sulfonate
PBSF	Perfluorobutane sulfonyl fluoride
PFAAs	Perfluoroalkyl acids (includes carboxylic and sulfonic acids)
PFAS	Per- and polyfluoroalkyl substance
PFBS	Perfluorobutane sulfonic acid
PFCAs	Perfluoroalkyl carboxylic acid
PFDA	Perfluorodecanoic acid
PFDoDA	Perfluorododecanoic acid
PFHxA	Perfluorohexanoic acid
PFHxS	Perfluorohexane sulfonic acid
PFNA	Perfluorononanoic acid
PFOA	Perfluorooctanoic acid
PFOS	Perfluorooctane sulfonic acid
PFSAs	Perfluoroalkane sulfonic acids
PFTeDA	Perfluorotetradecanoic acid
PFTrDA	Perfluorotridecanoic acid
PFUnDA	Perfluoroundecanoic acid

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57

Substance name: Perfluorobutane sulfonic acid (PFBS) and its salts

EC Number: -

CAS Number: -

- It is proposed to identify the substances as substances of equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of Regulation (EC) No 1907/2006 (REACH) according to Article 57(f) of REACH Regulation.

Summary of how the substance meets the criteria set out in Article 57 of the REACH Regulation

Perfluorobutane sulfonic acid (PFBS) and its salts are proposed to be identified as substances of very high concern in accordance with Article 57(f) of Regulation (EC) 1907/2006 (REACH) as there is scientific evidence of probable serious effects to the environment and human health which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of REACH.

PFBS-salts will be fully indistinguishable from PFBS in the environment as the salts will exist in their dissociated anionic sulfonate form, just like PFBS itself, and they will all be a part of the same equilibrium. Hence, all conclusions on different end-points apply to any and all salt forms, as well as PFBS itself.

Several concerns are caused by the intrinsic properties of PFBS. Elements of concern are triggered by individual intrinsic properties or by different combinations of the properties.

PFBS is persistent in the environment. No abiotic or biotic degradation of PFBS at relevant environmental conditions has been shown. This is supported by read-across to perfluoroalkane sulfonic acids with both shorter and longer chain lengths, which also have a very high environmental stability. PFBS shows a preference for distribution to the aqueous phase due to its high solubility in water (fully miscible with water at 20 °C) and low sorption potential ($\log K_{oc}$ 1.2 to 2.7) and is considered very mobile in the environment.

Overall, PFBS has a very high potential to cause widespread effects in wildlife and in humans exposed via environment, due to its persistence, mobility, potential for long-range transport (CTD 17616 km, $P_{OV} = 220$ days), observed adverse effects that may be relevant for human health and the environment, and exposure via drinking water and food. It is considered that the T criteria for ecotoxicity in Annex XIII of REACH are fulfilled for PFBS based on its documented effects on marine medaka that include significant reduction in body length and weight (NOEC 1.0 µg/L), reduced egg production (NOEC 1.0 µg/L), delayed hatching (LOEC 1.0 µg/L), a skewed sex ratio toward male dominance (NOEC 2.9 µg/L). Disturbances of the visual system (NOEC 1.0 µg/L) of marine medaka represent a supporting argument in the ecotoxicity assessment.

It is particularly concerning that PFBS prefers and persists in the aqueous environment, while severe effects of PFBS have been demonstrated for fish, which belong to this particular environmental compartment.

The very high persistence, together with low adsorption potential and high mobility, imply a high potential for increasing environmental concentrations and irreversible exposures of wildlife and of humans via the environment. Hence, PFBS is bioavailable via the aqueous environment. Furthermore, the low adsorption potential and high water solubility leads to difficulties and high costs in removing PFBS using end-of-pipe treatment. Together, these elements of concern lead to a high potential for irreversible effects once effect levels have been reached.

The high global transport potential is demonstrated by detection of PFBS in samples of surface water, snow, ice, air and marine water from remote areas such as the Arctic and the Antarctic. PFBS has also been found in biota like dolphins and whales, as well as green turtles and polar bears, which are both threatened species. This shows that PFBS is bioavailable, and that exposure may occur throughout the food chain and via drinking water and that this is already taking place worldwide.

In pigs BMF values > 1 have been reported (BMF 1.2, whole pig), indicating that PFBS might be bioaccumulative in pigs. A serum elimination half-life of around one month (up to 46 days) has been measured in humans, which is considerably longer than the half-lives measured for rodents (less than 1 day).

The continuous and increasing exposure in human populations cannot be avoided if releases are not minimised. Similarly, wildlife populations cannot be protected from the total quantity of the substance released.

The level of concern is considered very high due to the combination of:

- high potential for irreversible exposure due to very high persistence and mobility and, in the case of human exposure via the environment, difficulties and costs to decontaminate drinking water,
- high potential for increasing contamination and increasing, bioavailable exposures as the intrinsic properties cause a difficulty to remove the substance after release,
- high potential for wide geographic scale contamination,
- ecotoxic effects fulfilling the T criteria in Annex XIII of REACH based on several different effects on marine medaka, including reduced egg production, delayed hatching of larvae, significant reduction in body length and weight, and a skewed sex ratio toward male dominance, supported by disturbances in the visual system,
- thyroid hormonal disturbances observed in rodents and fish and effects on expression of sex hormone receptor activity in tadpoles,
- effects on thyroid hormones together with deficits in perinatal growth, pubertal onset and reproductive organ development in mice,
- observed effects in experimental toxicity studies are of such nature that in combination with the above aspects, they lead to a high potential for serious effects on humans and the environment on a global scale,
- potential for causing serious effects that are not observed in standard tests or effects caused by life-long exposure or inter-generational effects following transfer from mother to offspring in several species,
- high societal concern.

PFBS has been detected in humans worldwide and in different species of wild life, including in endangered species and in remote areas. The substance has been found to transfer from mother to offspring in humans, whales and in birds and may disturb development at sensitive life stages and in vulnerable populations. It may be difficult in practice to manage exposures due to the high mobility of PFBS and the fact that exposures may take place at a different location than where releases occurred and at a different moment in time. The high persistence and high mobility of PFBS together lead to a concern for co-exposure with other contaminants with similar effects on human health and the environment. Co-exposure may last for a very long time, because natural degradation processes for these substances are slow or negligible. This is brought into the weight-of-evidence as supportive information.

Limitations of the available remediation techniques raise a concern that the removal of PFBS from drinking water may only be possible with high societal costs. Remediation of environmental pollution may even be practically impossible due to the high mobility of the substance. Furthermore, PFBS will quickly diffuse from contaminated sites.

Taken together in a weight-of-evidence consideration, the above arguments demonstrate that there is scientific evidence of probable serious adverse effects of PFBS and its salts to the human health and the environment, which gives rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of REACH Regulation.

Registration dossiers submitted for the substance? Yes

PART I

Justification

1. Identity of the substance and physical and chemical properties

PFBS belongs to the group of per- and polyfluoroalkyl substances (PFAS). It consists of a C4 perfluoroalkyl chain attached to a sulfonic acid group. PFBS belongs to the short-chain PFASs. PFBS is similar to the more well-known perfluorooctane sulfonic acid (PFOS), but PFOS carries a C8 perfluoroalkyl chain. In its pure form, PFBS is a corrosive liquid that causes severe skin burns and eye damage.

As a sulfonic acid, PFBS is a strong acid that readily forms sulfonate salts with bases, e.g. sodium, potassium and ammonium salts. In aqueous solution the salt forms will exist in equilibrium with PFBS itself. In the environment PFBS will quickly be deprotonated by available bases, and it will be present dissolved on its anionic form. Salts of PFBS are also dissolved in the water phase and the PFBS-part will be present in its anionic sulfonate ion form. It is usually not differentiated between PFBS and its anionic form or salt forms in analyses and quantifications of PFBS. In the literature, the concentrations reported in environmental and human monitoring studies will always be a sum of the acid PFBS, its anionic sulfonate form and its salt forms, which are all part of the same equilibrium.

For clarity, it is usually referred to PFBS in the discussions and conclusions on health and environmental effects in this document. However, based on the reasoning above, the conclusions are considered equally valid for any PFBS salt as well.

As the scope of the present SVHC dossier we have selected PFBS and any/all of its salts. These entities are indistinguishable in the environment and will all contribute to the PFBS levels.

It should be noted that the potassium salt, KPFBS, is the salt form most widely manufactured, used and studied among the PFBS salts. In many experimental studies the test material has been KPFBS, which is available and convenient to employ in studies. In the physicochemical properties section, Section 1.5, we have included the properties of KPFBS in addition to PFBS itself, in order to illustrate the differences between the free acid and a salt form.

1.1. Name and other identifiers of the substance

Data on the substance identity of PFBS and its potassium salt KPFBS are presented in Table 1 and Table 2, respectively.

Table 1: Substance identity, PFBS.


EC number:	206-793-1
EC name:	1,1,2,2,3,3,4,4,4-Nonafluorobutane-1-sulfonic acid
CAS number (in the EC inventory):	375-73-5
CAS name:	--
IUPAC name:	Nonafluorobutane-1-sulfonic acid
Index number in Annex VI of the CLP Regulation:	
Molecular formula:	C ₄ HF ₉ O ₃ S
Molecular weight:	300.10 g/mol
Synonyms:	Perfluorobutane sulfonic acid, PFBS
Chemical structure:	
SMILES notation:	OS(=O)(=O)C(F)(F)C(F)(F)C(F)(F)C(F)(F)F

Table 2: Substance identity, KPFBS – the most widely used and studied salt form of PFBS.

EC number:	249-616-3
EC name:	Potassium 1,1,2,2,3,3,4,4,4-nonafluorobutane-1-sulfonate
CAS number (in the EC inventory):	29420-49-3
CAS name:	--
IUPAC name:	Potassium nonafluorobutane-1-sulfonate
Index number in Annex VI of the CLP	
Regulation:	
Molecular formula:	C ₄ F ₉ O ₃ SK
Molecular weight:	338.19 g/mol
Synonyms:	Potassium perfluorobutane sulfonate, Perfluorobutane sulfonic acid potassium salt, KPFBS, K-PFBS, PFBSK+,
Chemical structure:	
SMILES notation:	[K+].[O-]S(=O)(=O)C(F)(F)C(F)(F)C(F)(F)C(F)(F)F

1.2. Composition of the substances

This dossier applies to PFBS and any/all of its salts. The salt forms included in ECHA's database, either registered or notified, are listed in Table 3, Section 1.3. However, the dossier is not limited to the currently registered and notified salt forms. It applies to any salt form of PFBS, either currently existing or developed in the future.

Name: Perfluorobutane sulfonic acid, PFBS

Substance type: mono-constituent

1.3. Identity and composition of structurally related substances

1.3.1. Salt forms - grouping

Salt forms of strong acids, such as the sulfonic acid PFBS, are closely related to its parent compound, in this case the free acid PFBS. The acid, the corresponding deprotonated, anionic sulfonate ion and the salt forms are all part of the same acid/base equilibrium. All salt forms of PFBS released, contribute to the overall levels of PFBS in the environment, and therefore to the overall risk for health effects, when exposure is via the environment.

In the present dossier PFBS and its salt forms are grouped together in a common proposal for listing on the Candidate List. The salt forms of PFBS included in ECHA's database, either registered or notified, are listed in Table 3. These PFBS salts are members of the group of substances addressed in this dossier, but any other salt form of PFBS is included as well. According to the above argument, any salt form of PFBS will contribute to PFBS-concentrations through the acid-base equilibrium.

Table 3: Salt forms of PFBS in ECHA's database.

Perfluorobutane sulfonic acid (PFBS)				
CAS No	375-73-5		EC No	206-793-1
Molecular formula	C ₄ HF ₉ O ₃ S		Molecular weight (g/mol)	300.10
Potassium perfluorobutane sulfonate (KPFBS)				
CAS No	29420-49-3		EC No	249-616-3
Molecular formula	C ₄ F ₉ O ₃ SK		Molecular weight (g/mol)	338.19
Tetraethylammonium perfluorobutane sulfonate				
CAS No	25628-08-4		EC No	NA
Molecular formula	C ₁₂ H ₂₀ F ₉ NO ₃ S		Molecular weight (g/mol)	429.34
Tetrabutylphosphonium perfluorobutane sulfonate				
CAS No	220689-12-3		EC No	444-440-5
Molecular formula	C ₂₀ H ₃₆ F ₉ O ₃ PS		Molecular weight (g/mol)	558.52
Triphenylsulfanium perfluorobutane sulfonate				
CAS No	144317-44-2		EC No	478-340-8
Molecular formula	C ₂₂ H ₁₅ F ₉ O ₃ S ₂		Molecular weight (g/mol)	562.47
Dimethyl(phenyl)sulfanium perfluorobutane sulfonate				
CAS No	220133-51-7		EC No	452-310-4
Molecular formula	C ₁₂ H ₁₁ F ₉ O ₃ S ₂		Molecular weight (g/mol)	438.33
Bis(4-t-butylphenyl) idonium perfluorobutane sulfonate				

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

CAS No	NA		EC No	432-660-4
Molecular formula	C ₂₄ H ₂₆ F ₉ IO ₃ S		Molecular weight (g/mol)	692.42
Ammonium perfluorobutane sulfonate				
CAS No	68259-10-9		EC No	269-513-7
Molecular formula	C ₄ H ₄ F ₉ NO ₃ S		Molecular weight (g/mol)	317.13
Lithium perfluorobutane sulfonate				
CAS No	131651-65-5		EC No	NA
Molecular formula	C ₄ F ₉ LiO ₃ S		Molecular weight (g/mol)	306.03
Magnesium perfluorobutane sulfonate				
CAS No	507453-86-3		EC No	NA
Molecular formula	C ₈ F ₁₈ MgO ₆ S ₂		Molecular weight (g/mol)	622.49
Morpholinium perfluorobutane sulfonate				
CAS No	503155-89-3		EC No	NA
Molecular formula	C ₈ H ₁₀ F ₉ NO ₄ S		Molecular weight (g/mol)	387.22
1-(4-Butoxy-1-naphthalenyl)tetrahydrothiophenium 1,1,2,2,3,3,4,4,4-nonafluoro-1-butanesulfonate				
CAS No	NA		EC No	468-770-4
Molecular formula	C ₂₂ H ₂₃ F ₉ O ₄ S ₂		Molecular weight (g/mol)	586.53




In the environment all the salts will exist in their dissociated anionic sulfonate form, just like PFBS itself, and they will all be indistinguishable. Hence, all conclusions on different end-points apply to any and all salt forms, as well as PFBS itself.

1.3.2. Perfluoroalkane sulfonic acids used in read-across

In the assessment of the persistence of PFBS read-across to structurally related perfluoroalkane sulfonic acids (PFSA) have been applied. PFBS, a C4 PFSA, has been compared to one shorter (C1) and one longer (C8) homologue PFSA, trifluoromethane sulfonic acid and perfluorooctane sulfonic acid (PFOS), respectively. In ECHA's database trifluoromethane sulfonic acid, the potassium salt of PFBS, and the tetraethylammonium salt of PFOS are registered, and information on their degradation is found on the ECHA dissemination website. These data have been compared in a read-across approach in order to investigate the influence of perfluoroalkyl chain length on persistence of PFSA.

In Table 4 information on the identity and composition of these substances are presented. The read-across assessment is found in Section 3.1.4.

Table 4: Identity and composition of substances used in read-across

Chain length	C1	C4		C8	
Molecular formula	CF ₃ SO ₃ H	C ₄ F ₉ SO ₃ H	C ₄ F ₉ SO ₃ K	C ₈ F ₁₇ SO ₃ H	C ₈ F ₁₇ SO ₃ N(C ₂ H ₅) ₄
Name	Trifluoromethane sulfonic acid	Perfluorobutane sulfonic acid	Potassium perfluorobutane sulfonate	Perfluorooctane sulfonic acid	Tetraethylammonium perfluorooctane sulfonate
Acronym	Triflic acid	PFBS	KPFBS	PFOS	PFOS-NEt ₃
CAS No	1493-13-6	375-73-5	29420-49-3	1763-23-1	56773-42-3
EC No	216-087-5	206-793-1	249-616-3	217-179-8	260-375-3
Molecular weight (g/mol)	150.08	300.10	338.19	500.13	629.37
Structure		 <p style="text-align: center;">PFBS</p>		 <p style="text-align: center;">PFOS</p>	

1.3.3. PFBS related substances

PFBS is the final degradation product of a range of PFBS-precursors, or PFBS-related substances, that may degrade to PFBS during use, in the waste stage, or in the environment. Hence, contributions to the environmental concentrations of PFBS will come from the production and use of PFBS itself, as well as production and use of PFBS-related substances for various applications. Our concern for PFBS relates to the overall concentrations of PFBS in the environment, and therefore to the combined use and emissions of PFBS, its salts and PFBS-related substances. The manufacture and use of PFBS is limited compared to the related substances.

A comprehensive evaluation of the degradation of PFBS-related substances to PFBS was included in a literature study performed at the University of Oslo (Nielsen, 2017) and elaborated upon in the RMO analysis for PFBS, its salts and related substances (Norwegian Environment Agency, 2018). The degradation study identified several classes of substances that should be considered as precursors to PFBS. Furthermore, COWI A/S performed an investigation, commissioned by the Norwegian Environment Agency in 2017, of sources to PFBS in the environment (Lassen et al., 2017). Different applications of PFBS and related substances are discussed in this study.

Perfluorobutane sulfonyl fluoride (PBSF) is a key intermediate in the production of PFBS-related substances. Most PFBS-related substances may be prepared from PBSF. PBSF is a reactive substance used in manufacture. When PBSF is treated with alcohols or amines in a production process, sulfonic esters and sulfonamides are formed, respectively. PBSF reacts with water in a hydrolytic process with the formation of PFBS. The relationship between PBSF, PFBS-related substances and PFBS is illustrated in Figure 1.




Figure 1: Relationship between PBSF, PFBS-related substances and PFBS

The PFBS-related substances identified in the study by Nielsen, 2017, include PFBS salts, sulfonic acid halides, sulfonic alkyl/olefinic/aryl esters, sulfonamides, sulfones and side-chain fluorinated polymers containing the PFBS moiety. Perfluorobutane sulfinic acid also represents a precursor to PFBS through oxidation to the required sulfonic acid group (Nielsen, 2017). The chemical structures of PFBS-related substances are shown in Figure 2.



R = any chemical group, in particular R = -OH, -F, -Cl, -OR', -NR'R'' where R' and R'' represent any chemical group.

Related substances include e.g. sulfonic halides, sulfonic acid salts, sulfonic esters, sulfonamides, sulfones and sulfinic acid.

Figure 2: Chemical structure of PFBS-related substances

During manufacture, PBSF may be converted into various PFBS-related substances. Different uses of PFBS-related substances are briefly described in Chapter 8 in Part II of this dossier.

In summary, there is a concern related to concentrations of PFBS in the environment. Contributions to the overall amount of PFBS in the environment come from production, use and emissions of PFBS and its salts, as well as from degradation of PFBS-related substances during use, in the waste stage, or in the environment. However, the concern is first and foremost associated with the ultimate degradation product PFBS. On this basis the scope of the present SVHC-dossier has been selected to be PFBS and its salts.

1.4. Other PFASs

Kotthoff and Bücking (2018) have looked at the trends in PFAS chemistry over the last years. They found that the portfolio of detected anthropogenic PFAS in products encountered in daily life has expanded. Yet no clear picture of the full range of individual substances that comprise PFASs is available and this challenges authorities who struggle to cope with uncertainties in managing risk of harm posed by PFASs. This is said to be a result of an incomplete understanding of the range of compounds that are used in different products. Yeung and Mabury (2016) found that after the year 2000, the fraction of unidentified organofluorine in plasma samples is increasing. This suggests that humans are being exposed to an increasing number of new and unidentified fluorinated products.

The number of PFASs on the world market was estimated at more than 3000 in a survey by Swedish authorities in 2015 (Swedish Chemicals Agency, 2015). The OECD/UNEP Global PFC Group (2018) performed a new count in 2018 and identified 4730 different PFAS-related CAS-numbers.

Several PFASs have been identified as substances of very high concern (SVHC), including perfluorohexane sulfonic acid (PFHxS), and the C8 to C14 perfluorinated carboxylic acids: perfluorooctanoic acid (PFOA), perfluorononanoic acid (PFNA), perfluorodecanoic acid (PFDA), perfluoroundecanoic acid (PFUnDA), perfluorododecanoic acid (PFDoDA), perfluorotridecanoic acid (PFTrDA) and perfluorotetradecanoic acid (PFTeDA). In addition, PFOS is identified as a Persistent Organic Pollutant under the Stockholm Convention (UNEP, 2006).

1.5. Physicochemical properties

PFBS is a strong acid, and it is predominately present as a dissociated anion in the environment (Arp and Slinde, 2018). The substance has excellent surfactant properties, having a hydrophilic anionic sulfonate head group, and a hydrophobic perfluoroalkyl end group.

As is always the case for acids/salts, the pure phase properties, like physical state, melting point, boiling point, density, vapour pressure and water solubility, are intrinsically dependant on whether they are determined for the neutral species or the salt. All properties involving water, such as water solubility, K_{ow} and K_{aw} are dependent on the pH-value. PFBS is an acid, and in water it dissociates from the neutral form $C_4F_9SO_3H$ to the anionic $C_4F_9SO_3^-$ form. Over ambient pH ranges, there is a convention to use the symbol "D" instead of "K" for acids and bases for partitioning processes to indicate the pH dependence of the property. The pH dependence is governed by the acid-dissociation constant, pK_a . For monoprotic acids like PFBS, the relationship between K_{ow} and the pH-dependant D_{ow} follows from Equation 1:

Equation 1
$$D_{ow} = K_{ow} / (1 + 10^{(pH - pK_a)})$$

PFBS is a perfluorinated surfactant with a hydrophobic perfluoroalkyl group and a hydrophilic anionic sulfonate group. Perfluorinated surfactants are much more surface active than hydrocarbon surfactants. The substitution of fluorine atoms for hydrogen atoms decreases their surface activity for aqueous solutions, which promotes micellization at lower concentrations and lowers the surface tension relative to that of other hydrocarbon analogues (Kissa, 2001; Moody and Field, 2000). However, it has been found that among perfluorinated surfactants, the surface film of PFBS has a lower stability due to the greater water solubility (Campbell et al., 2009). The aqueous surface tension as a function of the PFBS concentration was measured and compared with PFBA, PFHxA and PFHxS and is illustrated in the article. Furthermore, it was found that PFBS, PFHxS and PFOS become insoluble prior to reaching their critical micelle concentration, Campbell et al. (2009); Solubility of KPFBS: 52.6 g/L at 22.5-24 °C. Hence, PFBS is not expected to form micelles in water.

Physicochemical data for PFBS as compiled by Arp and Slinde (2018) are shown in Table 5. For information, the table also includes data for the potassium salt, KPFBS, as the most widely used and studied PFBS salt.

Table 5: Physicochemical data for PFBS, as well as its potassium salt KPFBS

Property	Salt/Neutral Form	Description of key information	Reference/source of information
Physical state at 20°C and 101.3 kPa	KPFBS	White powder	Experimental, ECHA dissemination website
	PFBS	Liquid	Elvers, 2011
Melting/freezing point	KPFBS	> 280 °C	Experimental, ECHA dissemination website
	PFBS	-21 °C	Experimental, ECHA dissemination website
Boiling point	KPFBS	Decomposes before boiling	Experimental, ECHA dissemination website
	PFBS	198 °C	Experimental, ECHA dissemination website; Grayson, 1978
Vapour pressure	KPFBS	< 1.22E-05 Pa at 20°C ± 1°C	Experimental, ECHA dissemination website
	PFBS	7 Pa at 20°C	Experimental, ECHA dissemination website
		2.8 Pa	Wang et al., 2011; Estimated using COSMOtherm
Density	KPFBS	2.248 g/cm ³ at 20 °C	Experimental, ECHA dissemination website
	PFBS	1.824 g/cm ³ at 20°C	Experimental, ECHA dissemination website

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Property	Salt/Neutral Form	Description of key information	Reference/source of information
Water solubility	KPFBS	52.6 g/L at 22.5-24 °C	ECHA dissemination website;
	PFBS	Fully miscible at 20°C	Experimental, ECHA dissemination website; This conclusion is also expected over the entire ambient pH range based on an estimated water solubility of the neutral species (30 g/L) and the estimated pKa of -3.94
Partitioning coefficient soil/water (log K_{oc})	KPFBS	< 2	ECHA dissemination website, following OECD Guideline 106
	PFBS	1.2 1.79 2.2 2.7 1.2	Pereira et al., 2018; Guelfo and Higgins, 2013; Kwadijk et al., 2010; Vierke et al., 2014; Milinovic et al., 2015;
Dissociation constant pK_a	PFBS	-3.94	Wang et al., 2011; Estimated using COSMOtherm

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Property	Salt/Neutral Form	Description of key information	Reference/source of information
Partition coefficient n-octanol/water (log Kow / pH dependent Dow value)	KPFBS	-1.8 at 23 °C	Experimental, ECHA dissemination website
	PFBS	3.9 (neutral form) -4.0 to 0.0 (pH 4) -7.0 to -3.0 (pH 7) -8.0 to -4.0 (pH 8)	Wang et al., 2011; Estimated using COSMOtherm One experimental value is -0.34 in a 0.01 mol/L concentration (pH 1.7) at 23°C, from Experimental, ECHA dissemination website
Partition coefficient air/water (log Kaw/ pH dependant Daw value)	PFBS	-2.59 (pH << 0)	Wang et al., 2011; Estimated for neutral (nonionised species) using COSMOtherm
		-2.6 (neutral form) -10.5 to -6.5 (pH 4) -13.5 to -9.5 (pH 7) -14.5 to -10.5 (pH 8)	Estimated using log Kaw of -2.6 for the neutral species and pKa -3.94 and 0.14, respectively, and the equation $Daw = (1/(1+10^{(pH - pKa)}))Kaw$ (for monoprotic acids)
Partitioning coefficient n-octanol/air (log Koa)	PFBS	6.49	Wang et al., 2011; Estimated using COSMOtherm

2. Harmonised classification and labelling

There is no harmonised classification available for PFBS or any of its salts. However, there are four different self-classifications for PFBS and six for KPFBS. These are presented in Section 7.2, together with CLP data for the other registered and notified PFBS salts.

The free acid and the salt forms may have quite different physicochemical properties when they are still in their respective forms. However, in the environment or when mixed in a chemical system (e.g. water with buffer salts), the substances will be indistinguishable.

3. Environmental fate properties

The Norwegian Geotechnical Institute (NGI) carried out a study of PFBS in the environment with emphasis on monitoring data and the physical-chemical properties of the substance (Arp and Slinde, 2018). The study was funded by the Norwegian Environment Agency and is an important part of the basis for the evaluation of the environmental fate of PFBS.

The studies referred to in this chapter are according to standard tests, OECD test guidelines or GLP compliant if stated so. Otherwise they are studies published in scientific peer-reviewed journals or public reports.

3.1. Degradation

The perfluorinated substances are among the most stable organic compounds. According to Kissa (2001), this is due to the high electronegativity and low polarisability of fluorine, which results in a high bond energy of the C-F bond. It is not expected that the length of the perfluoroalkyl chain has a major impact on the inherent stability of PFASs. Hence, it is unlikely that a C4 perfluorobutyl compound should be considerably less stable as compared to e.g. the corresponding C8 perfluorooctyl compound (PFBS vs. PFOS in this case).

The C-F bond is among the strongest covalent bonds known, and it is resistant to acids, bases, oxidation and reduction, even at high temperatures. The strength of the C-F bond increases with increasing fluorine substitution at the carbon atom. Perfluoroalkane sulfonic acids (PFASs) are remarkably stable, with an outstanding thermal and chemical stability. Anhydrous PFASs are stable at 400 °C in the absence of air, but they may form hydrogen fluoride at this temperature when moisture is present. The sulfur atoms in PFASs are at their maximum oxidation state, and cannot be oxidised further (Arp and Slinde, 2018).

The Global PFC Group refers to PFASs as highly persistent in the environment, while their potential precursors are transformed into PFASs abiotically or biotically (OECD/UNEP Global PFC Group, 2013). Due to the high resistance to heat and chemical agents, the perfluoroalkyl substances have been frequently used in products with high versatility, strength, resilience and durability. However, the high persistence allows for a wide distribution in the environment, and many PFASs have been detected globally in the environment.

Brendel et al. (2018) recently summarised the current knowledge on the environmental stability of short chain PFASs and concluded that perfluoroalkyl acids (including PFBS) are extremely persistent, and that they do not undergo abiotic or biotic degradation at all under environmental conditions. It was further pointed out that the extreme stability in the

environment is regarded as an incalculable hazard itself, as short-chain perfluoroalkyl acids will stay in the environment for decades to centuries.

Ateia et al. (2019) point out that the short-chain PFASs are equally persistent as their long-chain counterparts, and that the high solubility of short-chain PFAS in water, low/moderate sorption to soils and sediments and resistance to biological and chemical degradation has resulted in their widespread presence in various aquatic environments.

In a review paper Cousins et al. (2016) look at the precautionary principle and chemicals management in relation to perfluoroalkyl acid contamination of groundwater. It is pointed out that all PFASs entering groundwater, irrespective of their perfluoroalkyl chain length and bioaccumulation potential, will result in poorly reversible exposures and risks, as well as further clean-up costs for society. In order to protect groundwater resources for future generations, the authors call for a precautionary approach and prevent the use and release of highly persistent and mobile chemicals such as PFASs. In a different study it is proposed that high persistence alone should be established as a sufficient basis for regulation of a chemical (Cousins et al., 2019). The authors argue that if a chemical is highly persistent, its continuous release will lead to continuously increasing contamination irrespective of the chemical's other physical and chemical properties. The increasing concentrations will result in increasing probabilities for known and unknown effects. Once adverse effects are identified, it will take decades, centuries or even longer to reverse contamination and therefore effects. PFASs are one of three classes of highly persistent compounds addressed in the study.

Parsons et al. (2008) reviewed the biodegradation of perfluorinated compounds. The authors pointed out that the most theoretically plausible degradation pathway for PFASs, such as PFBS, is through reductive defluorination, which could occur under anaerobic conditions. The same authors reported for PFOS that no biodegradation was observed under aerobic conditions, while there were some observations of degradation of PFOS under anaerobic conditions though no metabolites were measured in these studies. In principle, it cannot be ruled out that some degradation of PFBS under anaerobic conditions can occur (e.g. in hypoxic groundwater, marine water or sediments), or will occur in the future if bacteria adapt to utilise the energy present in the PFAS substrates. Indications for such bacterial behaviour were found when a PFOA-degrading bacterial strain was isolated from soil near a PFAS production plant (Yi et al., 2016). The PFOA-degradation has been demonstrated at lab conditions with a low degradation efficiency only. The findings are promising with regards to a future bioremediation technique for PFOA-polluted areas. However, the rates of these processes under environmental conditions are unknown, and potentially very slow or negligible, and have yet to be observed in the environment (Arp and Slinde, 2018).

3.1.1. Abiotic degradation

3.1.1.1. Hydrolysis

A study of KPFBS hydrolysis as a function of pH is reported on ECHA's dissemination website for KPFBS (according to EU method C.7, Degradation: Abiotic Degradation: Hydrolysis as a Function of pH). The substance was studied in pH 4.0, 7.0, and 9.0 buffer systems and at 50 °C. The study is GLP compliant with a Klimisch reliability 2. No changes in the concentration of the substrate were observed after 5 days at any of the different pH values. Since less than 10% hydrolysis occurred, the test substance is considered hydrolytically stable with a hydrolytic half-life, $t_{1/2}$ at 25 °C of more than 1 year.

Comparison of the hydrolytic stability of the C1, C4 (PFBS) and the C8 PFASs based on experimental data in a read-across approach is presented in Section 3.1.4.

PFOS has been concluded to be hydrolytically stable with a hydrolytic half-life of more than 41 years (UNEP, 2006). PFHxS is also considered to be hydrolytically stable under environmental conditions based on read-across to PFOS and PFOA (ECHA, 2017).

3.1.1.2. Phototransformation in air

The QSAR model AOPWIN v1.92 of the EPISuite tool was used to assess degradation in the atmosphere. The model estimates the rate constant for the atmospheric, gas-phase reaction between hydroxyl radicals and organic chemicals. The rate constants estimated by the program are then used to calculate atmospheric half-lives for organic compounds. The model is preset with the following: 1.5×10^6 molecules (radicals)/cm³ per 12 hours of daylight, which was also used for this assessment. The predicted degradation rate constant for PFBS was 0.1400×10^{-12} cm³/(molecule*sec), which is equal to an atmospheric half-life of 76.4 days. It should be emphasised that perfluoroalkyl substances are not fully covered by the EPISuite models, and the estimated overall OH rate constant and half-life are based only on "reaction with N, S and OH-".

Thus, the results should be interpreted with caution. However, the data can be used in a weight of evidence approach indicating that the degradation half-life of PFBS in the atmosphere is above the threshold of two days, which proves that the substance has a potential for long-range transport.

Photolysis of perfluoroalkane sulfonates in the atmosphere is generally considered negligible under environmental conditions (Vecitis et al., 2009). An outlying study is one by Taniyasu et al. (2013) which reported observing photolysis in air at high altitudes (>2500 m) of PFOS, which reacted via dealkylation to form PFBS. It was concluded that PFBS was more recalcitrant and stable. It should be noted that this photodegradation of PFOS has been disputed by Wang et al. (2015a), who argue that PFOS is too stable to undergo atmospheric photolysis. Nevertheless, PFBS, as such, is considered to remain stable to photolysis under environmental conditions, and this has been supported by all studies thus far.

PFOS has an expected half-life in the atmosphere greater than two days and fulfils the POP-criteria (UNEP, 2006). For PFHxS read across to PFOS and PFOA has been used, concluding on an atmospheric half-life which also exceeds two days (ECHA, 2017).

3.1.1.3. Phototransformation in water

A study on the photolysis of PFOS in water has been conducted, where no evidence of direct or indirect photolysis was observed under any of the conditions tested. The indirect photolytic half-life of PFOS at 25°C was calculated to be more than 3.7 years (UNEP, 2006).

Similar data for PFBS have not been found, but based on the structural similarities, PFBS would be expected to have a similar behaviour as PFOS.

3.1.1.4. Oxidation

PFBS was included in a study of the degradation of new substitutes for perfluorinated surfactants (Quinete et al., 2010). The study tested for the abiotic degradation of PFBS

using advanced oxidation processes through ultraviolet radiation, hydrogen peroxide, or both. No significant oxidative degradation of PFBS was observed in the tests.

In summary, a hydrolytic half-life for PFBS in water of more than 1 year has been determined experimentally. An atmospheric half-life of 76.4 days has been predicted, which by far exceeds the criterion in the Stockholm Convention of > 2 days. Although the value was not from a model validated for such chemistry, it is indicative of a potential for long-range transport. The available information demonstrates that PFBS is stable towards hydrolysis, oxidation and photodegradation and very persistent under environmentally relevant conditions.

3.1.2. Biodegradation in water

3.1.2.1. Estimated data

Biodegradation of PFBS was modelled using the EpiSuite BIOWIN v4.10 model in EpiSuite (US EPA, 2000-2017). BIOWIN estimates the probability of rapid aerobic and anaerobic biodegradation of an organic compound in the presence of mixed populations of microorganisms and contains seven separate models. The screening criteria for persistence in the environment are BIOWIN2 <0.5 or BIOWIN6 <0.5 and BIOWIN3 <2.25. For substances where BIOWIN3 indicates a value between 2.25 and 2.75 more degradation relevant information is generally required. The outcome of the estimation is provided in Table 6.

Table 6: Results from BIOWIN estimations of PFBS biodegradation.

Model	Result	Limit	Conclusion
BIOWIN 1, linear probability model	-0.3592		Does not biodegrade fast
BIOWIN 2, non-linear probability model	0.0052	<0.5	Does not biodegrade fast
BIOWIN 3, expert survey ultimate biodegradation model	1.5793	<2.25	Recalcitrant
BIOWIN 4, expert survey primary biodegradation model	2.8572		Weeks-months
BIOWIN 5, MITI linear model	0.1786	≥0.5	Does not biodegrade fast
BIOWIN 6, MITI non-linear model	0.0000	<0.5	Does not biodegrade fast
BIOWIN 7, anaerobic biodegradation model	-0.1666		Does not biodegrade fast
Ready biodegradability prediction			NO

The QSAR estimates for PFBS are all below the screening criteria for being rated as "readily biodegradable" and indicate that PFBS is potentially persistent or very persistent according to ECHA Guidance on PBT/vPvB assessment (ECHA, 2017a). PFBS and PFASs are not always represented well in QSAR estimations, and the results should be interpreted with caution. For example, the main structural difference in the perfluoroalkane sulfonic acid series is the number of "carbons with 4 single bonds and no hydrogens", and in BIOWIN 1-4 these contribute to a more stable structure, while in BIOWIN 5-6 they provide less stability. Hence, comparison with shorter or longer homologues in the perfluoroalkane sulfonic acid group, will give different results for BIOWIN 1-4 and 5-6. However, although the small differences in the estimates among closely related PFAS substances are uncertain, all models clearly suggest that PFBS does not biodegrade fast. This is taken as a strong indication that PFBS will not fulfil the screening criteria for being rated as "readily biodegradable" and PFBS can be regarded as potentially persistent or very persistent according to ECHA Guidance on PBT/vPvB assessment (ECHA, 2017a). In any case, BIOWIN predictions are used only as support for other evidence for persistence of PFBS.

3.1.2.2. Screening studies in aquatic compartment

A screening test on ready biodegradability of KPFBS is provided as key study on ECHAs dissemination webpage. The test was performed according to the test guideline OECD 301E (Modified OECD Screening Test), was GLP compliant, and assessed by registrants with Klimisch reliability 2. The purity of the test substance was 98.7 % and a test concentration of 20 mg/L dissolved organic carbon (DOC) was incubated with secondary effluent of a sewage treatment plant (STP) receiving predominantly domestic sewage. The test was carried out at 22 ± 2 °C, in darkness and under aerobic conditions for 28 days.

Based on DOC removal 14% biodegradation after 14 and 28 days were observed which declined to 7 % on day 21, and this progress is rather unexpected, see Table 7. There was an overall decrease in DOC in the test bottles of 2.5 to 3 mg/L from day 0 to day 28. The blank control had first a decrease in DOC from 3 mg/L to 1 mg/L from day 0 to day 14, but then an increase to 3 mg/L on day 28, which is not normal. Since biodegradation of the test substance is calculated as the decrease in DOC in the test bottles, corrected for the DOC in the blank control, the unusual values from the blank control can explain what initially looks like biodegradation of KPFBS, which justifies for a Klimisch reliability 3. In conclusion, there is no reliable evidence for biodegradation of KPFBS in this study.

Table 7: Percent biodegradation of KPFBS

	Replicate	Day 0	Day 7	Day 14	Day 21	Day 27	Day 28
Blank values reported							
mg/L DOC removal		3	2	1	2	2	3
KPFBS % biodegradation	A		-3 %	0 %	5 %	10 %	15 %
	B		-8 %	-29 %	8 %	8 %	13 %
	Average		-5 %	-14 %	7 %	9 %	14 %

Another experimental study is available on ECHAs dissemination webpage for KPFBS where Quinete et al. (2010) tested KPFBS for biodegradation in conventional screening tests. Biodegradation of <1% after 40 days based on oxygen consumption was observed in a manometric respirometry test (OECD 301 F; OxiTop) using domestic non-adapted sludge of a sewage treatment plant (STP). According to the registrants, the study was deemed as not reliable due to significant methodological deficiencies. Furthermore, a closed-bottle test (OECD 301 D) using inoculum from the Rhine River demonstrated < 3% biodegradation (based on oxygen consumption) of PFBS within 28 days. This study was deemed as reliable with restrictions by registrants since detailed documentation was missing. In buffered media, biodegradation testing on the free acid is equivalent to testing on the salt. Therefore, results for PFBS are directly applicable to KPFBS. In a standardised biological fixed-bed bioreactor test using Rhine River inoculum, no biodegradation of PFBS was observed. The authors concluded that KPFBS is not readily biodegradable.

Comparison of ready biodegradability of the C1, C4 (PFBS) and the C8 PFSA based on experimental data in a read-across approach is presented in Section 3.1.4. No information is available for the short-chain PFASs perfluorobutanoic acid (PFBA) and perfluoropentanoic acid (PFPeA) since they are not registered.

The C8 homologue PFOS has been established as not readily biodegradable (UNEP, 2006), and a read-across to PFOS has also been used to conclude on no "ready biodegradability" for PFHxS (ECHA, 2017).

3.1.2.3. Simulation tests (water and sediments)

No simulation tests have been reported for PFBS.

3.1.3. Biodegradation in soil

There are no experimental soil degradation tests available for PFBS or KPFBS. The C8 homologue PFOS has been found to be persistent in soil cultures (UNEP, 2006) and read-across to PFOS has been used to conclude that also PFHxS is very persistent in soil (ECHA, 2017).

3.1.4. Read-across to similar PFSA

In general, the read-across approach can be applied in the assessment of a property when the substances compared have physicochemical and/or toxicological and/or ecotoxicological properties that are likely to be similar or follow a regular pattern as a result of structural similarity. According to ECHA's practical guide 6 "How to report read-across and categories" similarities may be due to a common functional group, common precursor or breakdown products, constant pattern in changing potency or common constituents or chemical class.

The group of perfluoroalkane sulfonic acids (PFSA) consists of closely related substances with very similar structures and properties. The PFSA include a perfluoroalkyl group of varying carbon chain length, attached to a sulfonic acid group. Hence, the different substances in the group differ only in the number of CF₂-units in the perfluoroalkyl chain, whereas the other structural features are the same. The perfluoroalkyl chain is persistent due to the high stability of the C-F-bond. The sulfonic acid group is at its highest oxidation state and cannot be oxidised further in the environment. It is not expected that the sulfonic

acid group will make the attached perfluoroalkyl chain more susceptible to chemical transformations. If the sulfonic acid group was to have an influence on the perfluoroalkyl chain, it would be on the neighbouring CF₂-group, and this is common for all the PFSA. Hence, the persistence of the members in this class of substances is highly comparable, and an evaluation of the persistence of PFBS in comparison with trifluoromethane sulfonic acid and PFOS in a read-across approach is justified.

In general, when comparing experimental and estimated data, it can be assumed that with increasing chain length, water solubility decreases and the sorption potential increases (ECHA, 2015). Consequently, a short-chain PFSA with a higher solubility in water could potentially be more readily available for aqueous organisms and biodegradation. However, no biodegradation has been observed for trifluoromethane sulfonic acid in screening tests, see Table 8. The persistence of PFSA similar to PFBS has been confirmed. PFOS was in 2006 identified as a Persistent Organic Pollutant and listed in the Stockholm convention (UNEP, 2006). PFOS is considered extremely stable in the environment and does not hydrolyse, photolyse or biodegrade under any environmental conditions tested (OECD, 2002). PFHxS was established as P and vP by the Member State Committee in 2017 when PFHxS was identified as a substance of very high concern based on read-across to PFOS, (ECHA, 2017).

At the ECHA dissemination page there is data for the hydrolytic stability and biodegradability of the C1, C4 and C8 perfluoroalkane sulfonic acids from the registration dossiers for the following substances: trifluoromethane sulfonic acid (C1), PFBS potassium salt (C4) and PFOS triethylammonium salt (C8). The three substances belong to the same PFAS subclass and are different only in the length of the perfluoroalkyl chain. There is no difference in testing the persistence of the free acid or a salt form, as the salts will be in equilibrium with the corresponding acid in the aqueous phase.

The identity and composition of the substances used in the read-across assessment is found in Section 1.3.2, while physical-chemical data for the substances are presented in Annex I. A comparison of the stability data found on the ECHA dissemination webpage for the three substances is shown in Table 8.

Error! Reference source not found.

Table 8: Stability data for C1, C4 and C8 perfluoroalkane sulfonic acids reported at the ECHA dissemination page.

	C1: CF₃SO₃H	C4: KPFBs	C8: PFOS NEt₄-salt
Hydrolytic stability	OECD Guideline 111, Hydrolysis as a Function of pH (pH 4.0, 7.0 and 9.0): Hydrolytically stable at all pH values with estimated half-life greater than 1 year at 25 °C	KPFBs: Hydrolysis as a Function of pH (4.0, 7.0 and 9.0), EU Method C.7 (EG-guideline 92/69): Less than 10% hydrolysis observed. Hydrolysis half life considered to be > 1 year under environmental conditions (25°C).	Study not conducted as the substance shows no chemical moieties that may be subject to hydrolytical reactions.
Biodegradability	OECD Guideline 301 D, Ready Biodegradability: Closed Bottle Test: There was 0% degradation of test item after 28 days.	OECD Guideline 301 E, modified OECD screening test for ready biodegradability: 14% biodegradation observed. Substance concluded not	Ready biodegradation according to OECD Guideline 301E: Biodegradation after 27 days is given as 0%.

	C1: CF₃SO₃H	C4: KPFBS	C8: PFOS NEt₄-salt
	Substance not readily biodegradable.	readily biodegradable. OECD Guideline 301 F, ready biodegradability according to Manometric Respirometry Test: Less than 1% degradation, KPFBS not readily biodegradable. OECD Guideline 301 D, ready biodegradability, Closed Bottle Test: Less than 3% degradation observed. PFBS is not readily biodegradable.	

With regards to the hydrolytic stability, PFBS and the C1 compound were found to be stable at all tested pH values with an estimated half-life greater than 1 year at 25 °C, while PFOS was considered to have no chemical moieties that may be subject to hydrolytical reactions. All three compounds were concluded to be not readily biodegradable in the tests performed. For PFBS some degradation (14%) was reported in one of the tests, but as described in Section 3.1.2.2 this result is considered unreliable and should not affect the conclusion. As the other biodegradability tests for PFBS, as well as the tests for the shorter (C1) and longer (C8) PFASs demonstrate no biodegradation, the conclusion that PFBS is not readily biodegradable is considered reliable.

In summary, based on experimental results from screening tests on KPFBS, which are supported by BIOWIN predictions, and read-across between the C1, C4 (PFBS) and the C8 PFASs in tests of hydrolytic stability and ready biodegradability, it can be concluded that PFBS is not readily biodegradable and is meeting the screening criteria for "potentially P/vP" according to ECHA Guidance on PBT/vPvB assessment (ECHA, 2017a).

3.1.5. Summary and discussion of degradation

The perfluorinated substances are among the most stable organic compounds due to the high bond energy of the C-F bond, which makes the substances resistant to acids, bases, oxidation and reduction, even at high temperatures. The sulfonic acid group is at its highest oxidation state and cannot be oxidised further under environmentally relevant conditions. The high stability of PFASs, combined with their mobility, allows for a wide distribution in the environment, and many PFASs have been detected globally in the environment.

PFBS is stable towards hydrolysis, oxidation and photodegradation in the atmosphere, and no abiotic degradation has been reported at environmental conditions. A study on hydrolysis of KPFBS in water demonstrated a hydrolytic half-life of more than 1 year. Reductive defluorination has been suggested as a plausible degradation pathway for PFASs, but evidence that PFBS undergoes such processes in the environment has not been found.

Based on experimental results from screening tests on KPFBS, read-across to similar measurements for the corresponding C1 and C8 sulfonic acids, as well as supportive evidence from BIOWIN predictions, it can be concluded that PFBS is not "readily biodegradable" but meets the screening criteria for "potentially persistent/very persistent" according to ECHA Guidance on PBT/vPvB assessment (ECHA, 2017a).

No abiotic or biotic degradation of PFBS at relevant environmental conditions has been shown. PFBS has been demonstrated to have a half-life in water of more than 1 year.

3.2. Environmental distribution

3.2.1. Sorption to the organic fraction of soil

Information from the ECHA dissemination website for KPFBS refers to a study of adsorption - desorption using a Batch Equilibrium Method (OECD Guideline 106). The study was performed on the free acid form of PFBS in the pH-range 5.96 to 7.77, as specified by test guideline. Adsorption and desorption testing on the acid is equivalent to testing on the potassium salt. The measured adsorption coefficient was 0 for the tested soils and sediment, and 0.5 for the tested sludge. The low sorption of PFBS to all five matrices tested (three soils, sediment, and sludge) placed PFBS in the very high mobility class ($K_{oc} = 0 - 50$).

When conceptualising sorption/desorption to soil for neutral organic molecules, one has to consider mainly van der Waal type interactions, i.e. non-ionic sorption interactions (Arp and Slinde, 2018). For organic ions, like PFBS, this is more complex, and one has to additionally consider ionic interactions between the substance and soil. Soils exhibit wide variation in their anion exchange capacity (AEC), and hence their ability to retain negatively charged molecules like PFBS.

The complexity of the PFAS chemistry is high, and it is difficult to predict the sorption of the substances from a single sorbent bulk property, such as for example organic carbon (OC) content (Pereira et al., 2018). How various sorbent-specific properties, such as pH and surface-bound cations, interact to determine the binding of PFASs to soils is still not fully understood. At environmental pH, natural organic matter (NOM) carries a negative net charge due to the presence of dissociated carboxylic and phenolic acid groups. The sorption of anionic, organic contaminants is promoted by a decrease in pH and by an increase in the concentration of cations.

Pereira et al. (2018) investigated the effect of solution pH and concentrations of Al^{3+} , Ca^{2+} and Na^{+} on the sorption of PFASs in soils. They found that perfluoroalkane sulfonic acids sorbed more strongly than perfluoroalkyl carboxylic acids. The PFAS sorption was further found to increase with increasing perfluorocarbon chain length with 0.60 log K_{oc} units per CF_2 moiety for C_3 - C_{10} PFCAs and 0.83 log K_{oc} units per CF_2 moiety for C_4 , C_6 , and C_8 PFSAs. Short-chained PFASs, including PFBS, were weakly sorbed (less than 10% on average), while long-chained PFASs sorbed strongly (on average, 99-100%). Log K_{oc} for PFBS was determined to be in the interval -0.7 to 2.2 at different pH-values and cation concentrations, with an average log K_{oc} at 1.2.

Though adsorption/desorption to soils is commonly normalised to the organic carbon fraction, i.e. K_{oc} value, a common discussion point regarding the adsorption/desorption of ions is the role of clays and minerals (Droge and Goss, 2013). Clays and minerals can have widely differing available surface areas for sorption and AEC values. It is therefore challenging to include a generic parameter to account for clay sorption.

Guelfo and Higgins (2013) studied the subsurface transport potential of perfluoroalkyl acids in batch sorption experiments with various soils and in the presence of co-contaminants relevant to aqueous film-forming foam (AFFF)-impacted sites. The results indicated that PFAA groundwater transport will depend on the solid phase characteristics at the site as well as PFAA concentration and chain length. Detailed site-specific information will likely be needed to accurately predict PFAA transport at AFFF-impacted areas. A log K_{OC} value for PFBS at 1.79 ± 0.10 was reported, together with a range of log K_d values for different soil types, -0.55 to 0.21 (K_d = concentration-specific solid-water distribution coefficient, L/Kg).

Kwadijk et al. (2010) examined the sorption of PFAS contaminated areas in the Netherlands and derived a log K_{OC} of 2.2 for PFBS, and a log K_d at 1.42 ± 0.50 . Vierke et al. (2014) investigated the transport of perfluoroalkyl acids in a water-saturated sediment column under near-natural conditions. The authors concluded that short-chain PFCAs and PFSAAs, like PFBS, may be problematic if contaminated surface waters are to be used for drinking water production via riverbank filtration as the short-chain substances will pass through the sediments. A log K_{OC} of 2.7 for PFBS was reported.

In a study of the sorption behaviour of three perfluoroalkyl substances (PFBS, PFOS and PFOA) in six different soils with contrasting characteristics, it was found that PFBS and PFOA showed a considerably lower sorption than PFOS in soils (Milinovic et al., 2015). This was attributed to the varying hydrophobicity of the substances, which is derived from the length of their perfluorinated carbon chain and the functional hydrophilic group, i.e. sulfonic vs. carboxylic acid. Furthermore, the sorption of PFOS was found to be highly irreversible, while PFOA and PFBS showed much higher desorption yields. For PFBS a log K_{OC} of 1.2 was determined.

The Reach Guidance, Chapter R.7b: Endpoint Specific Guidance defines a log K_{OC} of 3.0 as threshold when a substance can be considered to have a high potential for adsorption. At lower log K_{OC} , the adsorption is low, while the preference for the water phase is correspondingly higher. Hence, with log K_{OC} in the range 1.2 to 2.7, PFBS will be distributed relatively easily within and between environmental compartments and should be considered as mobile in the environment.

Arp and Slinde (2018) considered a log K_{OC} of 2.2 (equal to a K_{OC} of 158 L/Kg) and pointed out that this indicates that PFBS is quite mobile in the aquatic environment. The definition of K_{OC} is given in Equation 2:

$$\text{Equation 2} \quad K_{OC} = C_{soil} / (C_{water} * f_{OC})$$

where C_{soil} (ng/Kg) is the soil concentration, C_{water} (ng/L) is the water concentration and f_{OC} the fraction of organic carbon. At 2% f_{OC} one can calculate as presented in Equation 3:

$$\text{Equation 3} \quad C_{soil} \text{ (ng/kg)} = K_{OC} * C_{water} * 0.02 = 3.16 * C_{water} \text{ (ng/L)}$$

Hence, the concentration of PFBS in soil or sediment per kg is expected to be roughly three times the concentration in water per L. When water flows through soil, PFBS can be more readily transported compared to substances with a higher log K_{OC} value.

In summary, the high complexity of the PFAS chemistry makes it difficult to predict the sorption of the substances from a single sorbent bulk property. Properties such as pH, identity of the soil/sediment and surface-bound cations interact to determine the binding of PFASs. The sulfonic acids tend to sorb more strongly than carboxylic acids, and the PFAS sorption tends to increase with increasing perfluorocarbon chain length. However, PFBS has been found to be a weakly sorbing substance with a high mobility in the environment. PFBS has a preference for distribution to the aqueous phase and is relatively

readily transported when water flows through soil. Log K_{oc} for PFBS has been reported in the range 1.2 to 2.7.

3.2.2. Sorption to the mineral fraction of soil

In addition to the highly complex PFAS chemistry that makes it difficult to predict their sorption, the highly complex nature of the soil complicates matters further. Soils consist of organic matter, minerals and pore spaces filled with air and water (Brady and Weil, 2010). Sand, silt and clay all provide minerals and surface area for the sorption of PFAS. Sand, silt and clay differ in their particle size, and smaller clay particles have colloidal properties carrying positive and/or negative charges. Both silicates, iron and aluminium oxides drive the soils chemical and physical activity (Brady and Weil, 2010).

Equation 4 describes the sorption (represented by K_D) of ionic species to minerals (e.g. Milinovic et al., 2015):

$$\text{Equation 4} \quad K_D = f_{oc}K_{oc} + f_{\text{minerals/clays}} K_{\text{minerals/clays}}$$

Where f_{oc} is the fraction of organic carbon in the soil, $f_{\text{minerals/clays}}$ is the fraction of minerals and clays in the soil, and the corresponding K values represent partitioning coefficients for the sorption of ionic species to both the organic carbon (K_{oc}) and the minerals and clays ($K_{\text{minerals/clays}}$).

A more specific model for ion exchange sites on minerals and clays has been developed for cations (Droge and Goss, 2013a), and this may be extended to anions, Equation 5.

$$\text{Equation 5} \quad K_D = f_{oc}K_{oc} + f_{IEC} K_{IEC-clays}$$

Where f_{IEC} is the fraction of ion exchange capacity sites of a soil per volume, and $K_{IEC-clays}$ the compound specific sorption to the ion exchange sites.

An important consideration for sorption of ions like PFBS, is that the soil solution chemistry will also play a role. This is especially true if ion exchange mechanisms dominate sorption, as the pH and salt type and concentration can affect the availability of ion exchange sites on the soil or mineral surfaces (Higgins and Luthy, 2006), as well as potentially influencing the ionic state of PFBS via protonation. Both Equation 4 and Equation 5 indicate that sorption to a soil is the sum of sorption to the organic carbon and to mineral component. In cases where sorption to minerals is very strong, or at very low organic carbon content (for example freshwater and marine sediments), electrostatic interactions with mineral phases increases and may become the dominate sorption phase (Barzen-Hanson et al., 2017).

Despite the organic matter comprising between 2 and 5 % of the soils volume and the minerals making up between 45 and 58 % (Brady and Weil, 2010; ECHA Guidance Chapter R.16 on Environmental Exposure Assessment), previous studies investigating the sorption of PFAS to soil have often concluded that sorption is most strongly correlated with the organic matter fraction. For example, Qian et al. (2017) concluded that soil organic matter enhanced the sorption of PFOS to the greatest extent in soils with varying clay, and therefore oxide and silicate contents (soil organic matter contents of 4.91-11.83 g/kg). In another study investigating the sorption of PFOS to soil, three Ferrsols, two Cambosols and one Isohumosol were selected (Wei et al., 2017). These soils varied in terms of pH, organic carbon content, Fe_2O_3 content, Al_2O_3 content, soil texture, specific surface area, average pore diameter and total pore volume. The results from a linear regression analysis revealed a positive correlation of PFOS sorption and Al_2O_3 , Fe_2O_3 and organic carbon contents (most significant relationship being with Al_2O_3 content). Li et al. (2018) conducted

a meta-analysis in order to investigate whether sorption to soils and sediments could be explained by the physicochemical properties of the soils or sediments. These authors concluded that a combination of organic carbon, pH and clay content had a significant role on sorption. A previous study investigating the sorption of PFBS to the aluminium oxide mineral boehmite (AlOOH) and humic acid found that sorption to humic acid and even humic acid coated boehmite was much stronger than to boehmite. This study nicely shows that even very low organic matter contents can dominate PFBS sorption (Wang et al., 2015b).

Milinovic et al. (2015) characterised sorption of PFAS to soils with a broad range of organic carbon (dissolved organic carbon from 180 to 7250 mg/kg) contents. The authors concluded that the soil mineral fraction had a negligible role in the sorption of PFBS to the soils tested, and that the organic matter dominated. In direct contrast to this, Li et al. (2019) reported that the sorption of PFBS to six soils with distinctly different physicochemical properties was not driven by the total organic carbon content of the soil which was between 2580 and 32800 mg/kg, but by the content of proteins, anion exchange capacity and the content of iron oxides (Fe₂O₃ content 1.51 – 9.21 mg/g). In a study specifically focusing on fluorotelomer sulfonates, correlations of sorption with pH, CEC, AEC and organic carbon content were investigated (Barzen-Hanson et al., 2017). These authors reported a contribution to sorption from the mineral phase for the soils with the lowest organic carbon content (percent organic carbon 0.098-7.7).

A simplistic way to describe the sorption of PFBS to minerals is to consider sorption to the silicates which are generally negatively charged and sorption to the iron and aluminium oxides which can be positively charged depending on the soil water chemistry. Silicates repel organic anions like PFBS, while positively charged metal oxides could potentially sorb PFBS. Sorption to metal oxides occurs via ion exchange interactions (as described above), with a contribution from hydrogen bonding ligand exchange and surface complexation (Li et al., 2019a). This has been demonstrated for PFOS which sorbed to alumina but not to silica (Hellsing et al., 2016). Further, the sorption behaviour and mechanisms of PFAA to the clays: montmorillonite, kaolinite and hematite were investigated by Zhao et al. (2014). The authors reported sorption to all three minerals, with the sorption strength order being hematite (an iron oxide) > kaolinite (an alumino-silicate) > montmorillonite (a phyllosilicate). Adsorption to alumina (Al₂O₃) has also been demonstrated for PFOS and PFOA with the authors of the study postulating that the protonated alumina surface rendered it accessible for compounds with negative charges to attach themselves as a result of electrostatic interactions. Sorption was sensitive to changes in pH and ionic strength, particularly the presence of Ca²⁺ ions (Wang and Shih, 2011). PFOS and PFOA have also been reported to sorb to organo-montmorillonites (Na₃(Al, Mg)₂Si₄O₁₀(OH)₂.4H₂O) (Zhou et al., 2010), goethite (Lath et al., 2018) and boehmite (AlOOH) (Wang et al., 2012).

In summary, the influence of minerals and soil chemistry on the sorption of PFBS is complex, and modelling of the processes and mechanisms involved is difficult. High organic carbon contents in soils most often dominate sorption, but even trace amounts of organic carbon can have the same effect. The contribution of minerals to sorption occurs in addition to that of organic carbon, and this only becomes substantial when mineral oxides dominate the clay fraction and the appropriate solution chemistry is present. In the complete absence of organic carbon in soils or sediments, sorption to minerals would be a function of the complex solution chemistry.

3.2.3. Volatilisation

PFBS is a strong acid which will be fully deprotonated at environmentally relevant conditions (Arp and Slinde, 2018). The anion in solution, as well as dissolved PFBS salts

do not volatilise. However, the non-deprotonated, neutral PFBS does volatilise to ca. 2.8-7 Pa. This implies that traces of PFBS fumes would be present in a room containing neutral PFBS and poor ventilation.

The pH dependant air-water partition coefficient, $\log D_{aw}$, was estimated at -6.5 to -14.5 (see

Table 9). This indicates that volatilisation from water is negligible. At pH 8, essentially all PFBS is ionic, and the substance would remain exclusively in water.

Because of negligible volatilisation of PFBS, its presence in the air would be primarily due to direct emissions of the substance or its salts into the air. Another possibility is through emissions of contaminated particles or water droplets (e.g. contaminated marine aerosols). When in the atmosphere, PFBS is expected to readily partition with surface water and water droplets (rain, cloud droplets, fog droplets, etc.) based on the very low $\log D_{aw}$ value. Therefore, it will undergo efficient removal from the atmosphere via wet-deposition.

The Henry's Law constant describes the tendency of a substance to volatilise from water. It can be calculated by Equation 6 (ECHA Guidance Chapter R.16 on Environmental Exposure Assessment):

$$\text{Equation 6} \quad \text{HENRY} = \text{VP} * \text{MOLW} / \text{SOL}$$

Where HENRY is the substance specific Henry's Law constant, VP is its vapour pressure, MOLW is molecular weight and SOL is water solubility.

Using the physical chemical properties reported in Section 1.5 (measured VP = 7 Pa; MOLW 300.1 g/mol; measured SOL = 52.6 g/L as reported on the ECHA dissemination page) the Henry's Law constant of PFBS is 0.040 Pa * m³/mol.

The calculated value for the Henry's Law constant for PFBS, 0.040 Pa * m³/mol, is far below the limit value for volatile chemicals, >250 Pa * m³/mol. This shows that PFBS has a low tendency to volatilise from water and that the aqueous environmental compartment is preferred for PFBS, compared to the atmosphere.

3.2.4. Distribution modelling

Distribution modelling for PFBS in comparison with other perfluoroalkane sulfonic acids (PFHxS, PFOS and PFDS) was performed by the Norwegian Geotechnical Institute (Arp and Slinde, 2018).

Distribution modelling of anionic substances like PFBS and other PFAS is not straightforward. Most distribution models were developed for neutral substances, wherein distribution modelling is generally described using three main partitioning coefficients: K_{ow} , K_{oa} and K_{aw} . However, octanol as a surrogate phase for soil/sediment/aerosol organic matter, or biological membranes, is inappropriate for anionic substances like PFBS, because octanol cannot make ionic interactions. One approach that has been used to solve this for PFOS is to apply the experimental K_{oc} value in distribution modelling, rather than the K_{ow} (Armitage et al., 2009), as the experimental K_{oc} inherently includes some ionic interactions to a naturally occurring phase. In this case, basic distribution modelling can be done using the pH-dependant K_{aw} , K_{oc} and the organic carbon-air partitioning coefficient $K_{oc, a}$. For the selected perfluoroalkane sulfonates, values for these terms are presented in

Table 9.

Table 9: Physiochemical properties needed to estimate environmental distribution for PFBS and homologues.

Parameter	PFBS	PFHxS	PFOS	PFDS	References
log K_{aw} (neutral)	-2.6	-2.4	-1.7	-1.2	Wang et al., 2011
log D_{aw} (pH 8)	-14.5	-13.8	-13.1	-12.0	Derived from Wang et al., 2011
					PFBS: Kwadijk et al., 2010
log K_{oc}	2.2	2.6	3.0	3.4	PFOS: Zareitalablad et al., 2013 PFHxS and PFDS extrapolated (read-across)
log $K_{oc, a}$	4.8	5.0	4.7	4.6	Calculated from the thermodynamic triangle: $\log K_{oc, a} = \log K_{oc} - \log K_{aw}$

The log K_{oc} values spread from 2.2 (PFBS) to 3.4 (PFDS), which would imply PFDS, PFOS and PFHxS sorb stronger than PFBS by factors of 16, 6 and 2.5 respectively.

The Globo-POP model (Wania, 2003) has been established to model the global distribution of neutral substances. In Figure 3 below, output from the Globo-POP model is used to illustrate the partitioning properties of substances that, globally, are predominantly in the air, water or soil phases, based on the following assumptions: the substance is non-degradable (which is appropriate for PFBS), 10 years of continuous emissions has occurred, emissions occur equally to soil, water, air (1/3rd each), and the global distribution of emissions is zonally distributed similarly to the human population. In this figure, lines are overlaid that represented the physical chemical properties of PFBS, PFHxS, PFOS and PFDS (Table 9), though using the pH-dependant K_{aw} , K_{oc} and the organic carbon-air partitioning coefficient $K_{oc, a}$ which is more appropriate for negatively charged substances.

As is indicated by Figure 3, PFBS is tightly clustered along with PFHxS, PFOS and PFDS, which indicates that they are all primarily in the water phase, while PFBS shows the highest preference for water.




Figure 3: Global POP chemical space distribution for non-degradable substances being emitted into air-water-soil at equal levels for the 10 years, with a similar zonal distribution the human population. The location of PFBS, PFHxS, PFOS and PFDS is overlaid, using $\log K_{OC}$ for $\log K_{OW}$, $\log K_{OC, a}$ instead of $\log K_{Oa}$ and considering the pH dependence of $\log K_{aw}$. Note that at ambient pH, the $\log K_{aw}$ is even lower than presented in this chart, and the substances would be strongly within the sea and freshwater media area of the chart. Other substances presented in this figure for comparison are CBzs (chlorobenzenes), PCBs (polychlorinated biphenyls) and PCDDs and PCDFs (dioxins). This Figure is modified and reprinted with permission from Wania and the American Chemical Society.

3.2.5. Field data

There are numerous studies reporting PFBS data in the environment compiled from peer-reviewed scientific research papers, thesis and reports. It has to be noticed that monitoring data encompass uncertainties and variability, but the reliability of the published data is considered to be of sufficient quality. Arp and Slinde (2018) gathered a non-exhaustive list of monitoring data for PFBS in surface water, ground water, drinking water, wastewater and leachate from landfill.

3.2.5.1. PFBS in surface fresh water

PFBS has been found in surface fresh waters throughout Europe, Asia and North America. In Table 10 PFBS concentrations from surface water not connected to known point sources have been compiled.

Table 10: Environmental concentrations of PFBS in surface fresh water samples (ng/L).

Location	Date	Mean +/- SD	Min. – Max. (Median)	n (det. Freq), LOD	Reference
Baltic sea	2013-2014	0.098		4/4 (100 %) LOD = 0.04	Gebbink et al., 2016a
Elbe River	2014	2.3	0.80 - 3.60 (2.40)	(100 %) LOD = 0.13	Heydebreck et al., 2015
France	2009	<1	<1 - 3 (<1)	(1 %) LOD = 4	Boiteux et al., 2012
France	2012		<0.02 - 29.00	(65 %) LOD = 0.02	Munoz et al., 2015
Germany Bad Godesberg	2008-2009	<0.01	<0.01	0 % LOD = 0.01	Wilhelm et al., 2010
Germany Bimmen-Lobith	2008-2009		0.00 - 0.08 (0.02)	(89 %) LOD = 0.01	Wilhelm et al., 2010
Germany Düsseldorf-Flehe	2008-2009		0.00 - 0.10 (0.03)	(91 %) LOD = 0.01	Wilhelm et al., 2010
Germany Möhne river	2008-2009		0.017 - 0.032 (0.023)	35 (100 %) LOD = 0.01	Wilhelm et al., 2010
Germany WkSt Rhein-Nord Kleve	2008-2009		0.00 - 0.06 (0.03)	(92 %) LOD = 0.01	Wilhelm et al., 2010
Germany WkSt Süd/Bad Honnef	2008-2009	<0.01	<0.01 - 0.03 (<0.01)	(7 %) LOD = 0.01	Wilhelm et al., 2010
Germany, Moehne river	2006	319	<LOD - 1,450 (83)	12 LOD = 2	Skutlarek et al., 2006
Germany, Rhine river	2006	14.3	<LOD - 46.00 (12.50)	38 LOD = 2	Skutlarek et al., 2006
Germany, Rhine watershed	2008		<0.08 - 181.00	LOD = Not reported	Möller et al., 2010

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min. – Max. (Median)	n (det. Freq), LOD	Reference
Germany, Rhine River	2013	15.6	1.40 - 40.00 (15.00)	(100 %) LOD = Not reported	Heydebreck et al., 2015
Germany, Rhine River	2016	21.9	0.46 - 146.00 (20.1)	20 (100 %) LOD = 0.03	Pan et al., 2018
Germany, Ruhr river	2006	16.7	<LOD - 71.00 (13.00)	27 LOD = 2	Skutlarek et al., 2006
Germany, River Elbe	2007	6.3	3.40 - 17.70 (5.40)	LOD = 0.50 (LOQ= 1.67)	Ahrens et al., 2009a
Italy (river basin survey)	2010-2013	8.90 ± 8.90	<LOD - 31.40 (5.60)	104 (74 %) LOD = 1	Valsecchi et al., 2015
North Sea, Baltic Sea and Norwegian Sea	2009		6.51	LOD = 0.05	Ahrens et al., 2010
Norway, lake	2004	nd		5 LOD = 0.03	Kallenborn et al., 2004
Norway, lake	2004	42.8	5.64 - 112.00	6 LOD = 0.03	Kallenborn et al., 2004
Norway, River Alna River Drammens-elva River Glomma	2016		0.05- 0.74 0.05-0.1 0.2-0.5	LOD = Not reported	Skarbøvik et al., 2017
Nordic countries, rivers and lakes	2017		<0.02-1.62	(36 %) LOD = Not reported	Kärroman et al., 2019

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min. – Max. (Median)	n (det. Freq), LOD	Reference
Spain	2009	1.22 ± 1.55	n.d. - 5.50	12 (17 %) LOD = 0.02-1.5	Pico et al., 2012
Spain	2009	1.33 ± 3.01	<0.10 - 10.10 (0.20)	30 (57 %) LOD = Not reported	Domingo et al., 2012
SW along River Elbe	2007	2.2	0.00 - 3.40 (2.30)	30 (100 %) LOD = 0.5	Ahrens et al., 2009b
Sweden, Mälaren Lake	2016	1.43	0.75 - 1.92	10 (100 %) LOD = 0.05	Pan et al., 2018
Sweden, river	2015	0.24 ± 0.10	0.11 - 0.35 (0.23)	4 (100 %) LOD = 0.2425	Tröger et al., 2018
Sweden, lake	2015	0.54 ± 0.46	0.19 - 1.20 (0.39)	2 (100 %) LOD = 0.5425	Tröger et al., 2018
Swedish rivers	2013	9.5	0.03 - 19.00	40 (59 %) LOD = Not reported	Nguyen et al., 2017
Switzerland, Glatt river	2007	4.3	2.30 - 7.70 (2.80)	3 LOD = Not reported	Huset et al., 2008
UK, Thames River	2016	5.06	3.26 - 6.75	6 (100 %) LOD = 0.05	Pan et al., 2018
Xiaoqing River	2014	nd		LOD = Not reported	Heydebreck et al., 2015)
China, Chao Lake	2016	15.4	1.50 - 81.50	13 (100 %) LOD = 0.05	Pan et al., 2018
China, Huai River	2016	0.83	0.52 - 1.59	9 (100 %) LOD = 0.05	Pan et al., 2018
China, Liao River	2016	0.94	0.43 - 2.16	6 (100 %)	Pan et al., 2018

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min. – Max. (Median)	n (det. Freq), LOD	Reference
				LOD = 0.05	
China, Pearl River	2016	4.49	0.21 - 21.50	13 (100 %) LOD = 0.05	Pan et al., 2018
China, Shuangtaizi Estuary	2013	4.7	0.95 - 12.60 (4.45)	21 (100 %) LOD = 0.06-0.403	Shao et al., 2016
China, Tai Lake	2016	2.02	0.17 - 4.85	15 (100 %) LOD = 0.05	Pan et al., 2018
China, Taihu Lake	2012	3.76 ± 0.72		LOD = Not reported	Fang et al., 2014
China, Yangtze River	2016	1.84	0.22 - 4.68	35 (100 %) LOD = 0.05	Pan et al., 2018
China, Yellow River	2016	0.99	0.07 - 2.23	15 (100 %) LOD = 0.05	Pan et al., 2018
Hong Kong	2008-2010	6.83 ± 2.40	2.18 - 8.69 (7.30)	12 (100 %) LOD = Not reported	Loi et al., 2011
Korea, Han River	2016	2.27	1.34 - 3.17	6 (100 %) LOD = 0.05	Pan et al., 2018
South Korea, Shihwa Industrial zone	2004	2.5	4.27	LOD = 0.05	Rostkowski et al., 2006
Vietman	2013-2015		0.10 - 8.28 (4.19)	LOD = 0.03-0.52	Lam et al., 2017
Canada, urban	2012-2016	2.52 ± 1.04	1.40 - 4.00 (2.50)	5 (100 %) LOD = 0.05	D'Agostino and Mabury et al., 2017
Canada, rural	2012-2016	0.78 ± 0.90	0.00 - 1.70 (0.70)	4 (50 %) LOD = 0.05	D'Agostino and Mabury et al., 2017
NJ, Delaware River	2016	2.19	0.52 - 4.20	12 (100 %) LOD = 0.05	Pan et al., 2018

The PFBS concentrations measured in surface water vary within a wide range. In many cases the concentrations are at the low end, or even not detected. However, in other cases elevated levels are measured. The highest PFBS concentration reported is 1450 ng/L, measured in the Möhne river in Germany (Skutlarek et al., 2006). The high level was assumed to be related to the use of contaminated fertiliser in an agricultural area. However, the ubiquity of PFBS in water samples shows that it is highly mobile and well distributed in the freshwater environment, even apart from point sources.

PFBS concentrations found in surface water close to PFAS production facilities or other PFAS hot spots are presented in Table 11.

Table 11: Concentrations of PFBS in surface water near PFAS production facilities or known hotspots (ng/L).

Location	Date	Mean +/- SD	Min-Max. (Median)	n (det. Freq) LOD	Remark	Reference
Italy (Bormida, Tanaro)	2010-2013	6.8 ± 5.5	<LOD - 17.0 (5.9)	33 (80 %) LOD = 0.2-5	River basin survey	Valsecchi et al., 2015
Italy (Brenta, Fratta-Gorzzone)	2010-2013	878.0 ± 764.0	56.0 - 1,665.7 (894.2)	4 (100 %) LOD = 0.2-5	River basin survey	Valsecchi et al., 2015
Italy	2010-2013	11	0.0 - 66.0 (2.0)	(74 %) LOD = 0.2-5	Industrialised river	Castiglioni et al., 2015
Netherlands	2016	20.3 ± 3.7	12.0 - 27.0	18 (100 %) LOD = 0.04	River	Gebbink et al., 2017
Norway	2016		<7.5-1150	66 % LOD = Not reported	Stream	Lassen et al., 2017
China, Daling River	2014	1090	nd - 3,780.0 (668.0)	LOD = Not reported	River	Wang et al., 2016b
China, Fuxin	2009	320	7.8 - 445.0 (353.0)	LOD = Not reported	River	Bao et al., 2011

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min-Max. (Median)	n (det. Freq) LOD	Remark	Reference
China, Tangxun Lake	2013	4950	4,600.0 - 5,300.0	2 LOD = 0.5	WWTP effluent	Zhou et al., 2013
China, Tangxun	2013	3720	2,240.0 - 4,520.0	LOD = 0.5	Lake	Zhou et al., 2013
Taiwan	2013	0.3 ± 0.2		3 LOD = 0.1	Upstream WWTP	Lin et al., 2014
Taiwan	2013	329.0 ± 16.0		3 LOD = 0.1	WWTP	Lin et al., 2014
Taiwan	2013	392.0 ± 6.0		3 LOD = 0.1	Downstream WWTP	Lin et al., 2014
Taiwan	2013	282.0 ± 9.0		3 LOD = 0.1	Downstream WWTP	Lin et al., 2014
Taiwan	2013	262.0 ± 4.4		3 LOD = 0.1	Downstream WWTP	Lin et al., 2014
Canada	2012-2016	6.4 ± 4.5	0.8 - 13.0 (4.4)	14 (100 %) LOD = 0.05	AFFF source	D'Agostino and Mabury, 2017
Canada, lake	2010-2011	4.90 ± 1.00		5 LOD = 0.0091	local pollution from an airport	Lescord et al., 2015

The surface water concentrations of PFBS reported above were measured near PFAS production facilities in the Netherlands (Gebbinck et al., 2017) and China (Wang et al., 2016b; Bao et al., 2011; Zhou et al., 2013). PFBS was also detected in the influent and effluent from a waste-water treatment plant near an electronic production facility in Taiwan (Lin et al., 2014), in fire-fighting foam contaminated areas in Canada (D'Agostino and Mabury, 2017) and Norway (Lassen et al., 2017), as well as in a mixed industrial area that includes fluorochemical plants in Italy (Valsecchi et al., 2015; Castiglioni et al., 2015).

The PFBS concentrations reported from areas near point sources and industrial facilities are in many cases much higher compared to concentrations measured in surface waters without known PFBS sources. The highest concentrations of PFBS found, 5.3 µg/L, were found in Tangxu Lake in China, near a PFAS production facility. However, also in Europe high concentrations of PFBS have been found: 1.7 µg/L in water from the Fratta Garzone

river in Italy which receives water from textile industry and a fluorochemical plant, and 1.2 µg/L in a small stream in Norway near a production facility for fire-fighting foam.

3.2.5.2. PFBS in groundwater

Concentrations of PFBS measured in groundwater are presented in Table 12.

Table 12: Environmental concentrations of PFBS in groundwater (ng/L).

Location	Date	Mean +/- SD	Min-Max (Median)	n (det. Freq.) LOD	Remark	Reference
Europe - 26 countries	2008	0.3	25	164 (15 %) LOD = 0.3		Loos et al., 2010
France	2009		9	(4 %) LOD = 4		Boiteux et al., 2012
Netherlands Groundwater recharge area (several (PFAA) sources are present – former landfill, military base, small commercial/industrial area)	2011		OW 1: 1.6-104 OW2: <LOQ-104 PW:<LOQ LOQ= 0.01	LOD = Not reported	Observation wells (OW, n=2) Pumping wells (PW, n=5) – travel time through soil from landfill to pumping wells > 25 years	Eschauzier et al., 2013
Vietnam	2013-2015	<0.3		LOD = 0.03-0.52		Lam et al., 2017
China, Daling River	2014	375	865	4 LOD = 0.03		Wang et al., 2016b
China, Fuxin	2009	235	1.2 – 872 (34.4)	LOD = Not reported		Bao et al., 2011
China, Fuxin	2017	11 010	64-21 200	LOD = Not reported		Bao et al., 2019

A European survey of groundwater from 2008 reported PFBS in 15% of the samples, with a maximum PFBS concentration at 25 ng/L and mean at 0.3 ng/L (Loos et al., 2010). In a French study PFBS concentrations up to 9 ng/L were reported, but the substance was found in 4% of samples only (Boiteux et al., 2012). In the Netherlands a maximum PFBS concentration of 104 ng/L was found in a groundwater recharge area, in an observation well located downstream a landfill area.

Considerably higher PFBS levels were reported in groundwater collected near two PFAS production facilities in China, up to 21.2 µg/L (Wang et al., 2016b; Bao et al., 2011; 2019). A remarkable 24-fold increase in PFBS concentration from 0.87 to 21.2 µg/L in groundwater beneath one of the facilities was registered from 2009 to 2017 (Bao et al., 2019). Another study did not report PFBS in groundwater in Vietnam (Lam et al., 2017).

3.2.5.3. PFBS in drinking water

Drinking water samples, including finished water, bottled water, and water at different steps in the purification process, have been analysed for PFBS in areas near and far removed from PFAS source zones. Measured concentrations of PFBS in drinking water are listed in Table 13.

Table 13: Concentrations of PFBS in drinking water (ng/L)

Location	Date	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
Global	2015	0.1	1.6	38 (47 %) LOD = 0.02	Bottled water (Canada, Burkina Faso, Chile, Ivory Coast, France, Japan, Mexico, Norway, USA)	Kaboré et al., 2018
Global	2015	0.24	0.0 - 1.1 (0.2)	59 (88 %) LOD = 0.02	Tap water (Canada, Burkina Faso, Chile, Ivory Coast, France, Japan, Mexico, Norway, USA)	Kaboré et al., 2018
BE, DE, I, NL, NO, SE	2010		<0.09 - 18.8	(86 %) LOD = Not reported	Tap water samples in six European Countries (Sweden, Italy, Belgium, Netherlands, Norway, Germany) (n=7)	Ullah et al., 2011
France	2015	3.2	1.3 - 6.7 (2.9)	(32 %) LOD = 0.15-8.76	Bottled water	Schwanz et al., 2016
France	2015	6.8	2.0 - 15.0 (6.5)	(62 %) LOD = 0.15-8.76	Drinking water	Schwanz et al., 2016
Germany	2008-2009		<0.01 - 0.01(<0.01)	70 (3 %) LOD = 0.01	Drinking water	Wilhelm et al., 2010
Netherlands	2016	4.94	0.5 - 19.0	6 (100 %) LOD = 0.04	Source zone: Drinking water near a PFAS production facility	Gebbink et al., 2017
Germany Ruhr river	2006	12.2	<LOD - 26.0 (13.5)	37 LOD = 2	Drinking water	Skutlarek et al., 2006
Italy	2010-2012	6.7 ± 11.3	<0.5 - 45.0 (1.8)	46	Drinking water	IRSA-CNR, 2013

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
				(63 %) LOD = Not reported		
Netherlands	2008	24.0 ± 10.0	10.0 - 47.0	6 LOD = Not reported	Infiltrated river water	Eschauzier et al., 2010
Netherlands	2008	2.4 ± 4.9	0.5 - 18.0	6 LOD = Not reported	Infiltrated rainwater	Eschauzier et al., 2010
Spain	2009	4.5 ± 14.5	<0.07 - 69.4 (0.4)	LOD = 0.07	Drinking water	Ericson et al., 2009
Spain	2015	nd		LOD = 0.15-8.76	Bottled water	Schwanz et al., 2016
Spain	2015	11	2.8 - 24.0 (11.0)	(31 %) LOD = 0.15-8.76	Drinking water	Schwanz et al., 2016
Sweden	2015	0.8 ± 0.3	0.3 - 1.3 (0.7)	11 (92 %) LOD = 0.01	Various parts of DW treatment production	Tröger et al., 2018
China, Shuangtai zi Estuary	2013	5.42	1.0 - 13.8 (4.4)	21 (100 %) LOD = 0.06-0.403	Bottled water	Shao et al., 2016
China	2008-2010		0.0 - 18.0 (2.8)	70 (74 %) LOD = 0.04-1.6	Drinking water	Mak et al., 2009
New Jersey, US	2009-2010	nd	6	30	Raw water	Post et al., 2013

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
				(10 %) LOD = Not reported		
Brazil	2015	3.4	3.1 - 3.6 (3.5)	(50 %) LOD = 0.15-8.76	Bottled water	Schwanz et al., 2016
Brazil	2015	4.4	0.5 - 16.0 (1.3)	(90 %) LOD = 0.15-8.76	Drinking water	Schwanz et al., 2016
China, Daling River	2014	25.4	97.8	4 LOD = 0.03	Source zone: Drinking water	Wang et al., 2016b
China, Fuxin	2009	0.47	0.2 - 0.7 (0.5)	LOD = Not reported	Source zone: Raw water	Bao et al., 2011
China, Fuxin	2009	0.54	<0.1 - 0.6 (0.5)	LOD = Not reported	Source zone: Finished water	Bao et al., 2011
China, Fuxin	2009	0.54	<0.1 - 0.6 (0.5)	LOD = Not reported	Source zone: Drinking water	Bao et al., 2011

PFBS appears ubiquitously in drinking water, both from the tap and from bottles. Even in regions away from established PFAS source areas, substantial drinking water contamination can occur, with a France max of 15 ng/L (Schwanz et al., 2016), German max of 26 ng/L (Skutlarek et al., 2006), Italy max of 45 ng/L (IRSA-CNR, 2013), Spain max of 69 ng/L (Ericson et al., 2009), China max of 18 ng/L (Mak et al., 2009), and Brazil max of 16 ng/L (Schwanz et al., 2016). These concentrations even exceed those of some known source zones, such as 19.0 ng/L near a PFAS production facility in the Netherlands (Gebink et al., 2017), and 0.6 ng/L for drinking water near the PFAS production area of Fuxin, China (Bao et al., 2011). The highest concentration, however, was near a source zone, at 97.8 ng/L, along the Daling River, China (Wang et al., 2016b). This indicates that PFBS persists through drinking water production, and currently is ubiquitously present in global drinking water (Kaboré et al., 2018).

3.2.5.4. PFBS in effluents from wastewater treatment plants and landfills and occurrence in receiving waters

PFBS is commonly found in effluents from wastewater treatment plants (WWTP) and in landfill leachate. A list of PFBS concentrations measured is provided in Table 14.

Table 14: Concentrations of PFBS in WWTP effluent and in landfill leachate (ng/L).

Location	Date	Mean +/- SD	Min-Max (Median)	n (det. Freq) LOD	Remark	Reference
Faroe Islands	2004	0.2		LOD = 0.03	WWTP	Kallenborn et al., 2004
Finland	2004	64.6	<LOQ - 68.00	3 LOD = 0.03	Landfill	Kallenborn et al., 2004
Finland	2004	2.9	2.61 - 3.09	3 LOD = 0.03	WWTP	Kallenborn et al., 2004
Germany	2009		<0.39 - 1,356.00	20 LOD = 0.12	Landfill	Busch et al., 2010
Norway	2004	1.64	1.00 - 2.60	3 LOD = 0.03	WWTP	Kallenborn et al., 2004
Nordic countries	2017	5.9	0.92-13.1	100 % LOD = Not reported	WWTP	Kärroman et al., 2019
Sweden	2012- 2015	1.9	3.7	LOD = 0.07	WWTP	Eriksson et al., 2017
WWTP along River Elbe	2007	8.23	0.00 - 25.90 (5.10)	9 (100 %) LOD = 0.5	WWTP (9 wwtp, 18 samples)	Ahrens et al., 2009a
San Francisco	2014	2.70 ± 1.50		LOD = 1.9	WWTP	Houtz et al., 2016
USA (three temperate zones)	2013- 2014	231.00 ± 577.33	3.44 - 3,410.00 (41.05)	87 (100 %) LOD = 2.6	Landfill (Ref organises data based on climate zone and age)	Lang et al., 2017
Singapore	2015 – 2016	N/A	161.6- 1916.3 (752.6)	12/12 (100 %) LOD = 0.009	Landfill (raw leachate)	Yin et al., 2017
Ireland	June – November 2017	1100	<0.01 - 17,000 (79)	85 %	Landfill	Harrad et al., 2019

Concentrations of PFBS in landfill leachate are relatively high, with a max of 3410 ng/L measured in the US (Lang et al., 2017). High concentrations have also been found in Europe, with 1356 ng/L measured in untreated leachate at a landfill in Germany (Busch et al., 2010). The high level was explained by different uses of PFASs and different regulation and treatment processes on landfills in different countries. The effect of different treatment techniques was investigated in the study. Lowest PFAS concentrations were observed after membrane treatments (reverse osmosis and nanofiltration) and activated carbon, while the high PFAS levels were found after biological treatment and wet air oxidation (Busch et al., 2010).

Harrad et al. (2019) reported high concentrations of PFBS and other PFASs in samples of leachate from landfills in the Republic of Ireland. Leachate was collected from 40 municipal solid waste (MSW) landfill sites across the country. Samples from sites without high density polyethylene (HDPE) liners (i.e. unlined sites) were pumped from “boreholes” (pipes inserted into the landfill body for collection and sampling of leachate), while samples from newer state of the art sites (i.e. mixed and lined sites) were collected from on-site leachate storage tanks. PFBS (arithmetic mean = 1100 ng/L) was the predominant compound among the measured PFASs. The concentrations of PFBS were significantly higher (p<0.05) in leachate samples from newer, lined landfills than in samples from unlined landfills.

PFBS concentrations in WWTP effluent are typically lower, with a maximum in this study of 26 ng/L (Ahrens et al., 2009a). Evidently, landfills can be environmental hotspots for PFBS emissions which may result in a risk for the local environment. According to a Nordic screening study on PFAS by Kärrman et al. (2019), PFASs constituted between 11 % and 46 % of total PFASs (27 % on average) in WWTP effluent samples, and PFBS was the predominant compound detected in all samples with 37 % of the PFSA class on average.

3.2.5.5. PFBS in marine water

PFBS has been reported in marine water throughout the world both in remote regions and in populated regions. Concentrations of PFBS measured are given in Table 15.

Table 15: Environmental concentrations of PFBS in marine water samples (ng/L).

Location	Year	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
Mid-Atlantic	2009-2010	nd	nd-0.017	20 LOD = Not reported		Zhao et al., 2012
Baltic sea, Kattegat	2013	0.32	0.06 - 0.57	18 (65 %) LOD = 0.03		Nguyen et al., 2017
Baltic sea	2013-2014	0.10 ± 0.00		(100 %) LOD = 0.04		Gebbink et al., 2016a

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Year	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
North Sea, Baltic Sea and Norwegian Sea	2009		6.51	LOD = 0.105		Ahrens et al., 2010
Denmark	2004	0.8	0.30 - 1.09	4 LOD = 0.03		Kallenborn et al., 2004
Farao Islands	2004	0.1	0.05 - 0.20	4 LOD = 0.03		Kallenborn et al., 2004
Finland	2004	0.07	<LOQ - 0.93	3 LOD = 0.03		Kallenborn et al., 2004
Iceland	2004	0.06	0.05 - 0.08	4 LOD = 0.03		Kallenborn et al., 2004
Western Mediterranean Sea	2014	0.01 ± 0.01	nd - 0.02 (0.02)	0.012 (0 %) LOD = 0.01		Brumovský et al., 2016
China	2013	0.66	0.32 - 1.46 (0.49)	(100 %) LOD = 0.12	July	Chen et al., 2016
China	2013	nd	<MLQ - 0.24 (<MLQ)	(19 %) LOD = 0.12	November	Chen et al., 2016
Pacific Ocean	2007	0.02 ± 0.00		19 (16 %) LOD = Not reported		Wei et al., 2007
Pacific Ocean	2007	0.07 ± 0.01		LOD = Not reported		Wei et al., 2007
South China Sea	2012-2013	0.63 ± 0.43	0.25 - 1.48 (0.45)	17 (100 %) LOD = 0.01		Kwok et al., 2015

Concentrations of PFBS in marine water samples vary in the range not detected to 6.51 ng/L. The highest concentration was measured at the German coast. High concentrations were also detected in China, however at a considerably lower level, ca. 1.5 ng/L.

3.2.5.6. PFBS in soil and sediment

Concentrations of PFBS found in soil and sediments are compiled in Table 16.

Table 16: Environmental concentrations of PFBS in soil and sediment samples (ng/g dw).

Location	Date	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
Soil						
Norway (soil)	2015	<LOD	<LOD - 0.003 (<LOD)	10 (10 %) LOD = 5		Herzke et al., 2016
China, Fuxin (soil)	2017	11.9	<0.2-42 (ng/g fw)	LOD = Not reported	Garden soil near a fluorochemical industry park	Bao et al., 2019
Sediment						
Canada Resolute Lake, local point source (airport)	2003		nd - 0.1	LOD = 0.046	0-3 cm cores	Stock et al., 2007
Canada Char Lake, local point source (airport)	2003	<1.1		LOD = 1.0	0-3 cm cores	Stock et al., 2007
Canada Amituk Lake	2003		0.0 - 0.1	LOD = 0.029	0-3 cm cores	Stock et al., 2007
Baltic sea (sediment)	2013-2014	0.0002 ± 0.0001		(75 %) LOD = 0.0001		Gebbink et al., 2016a
China (sediment)	2013	0.06	0.0 - 0.2 (0.1)	(100 %) LOD = 0.04	Surface sediments (July)	Chen et al., 2016

Location	Date	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
China (sediment)	2013	0.06	0.1 - 0.1 (0.1)	(100 %) LOD = 0.04	Surface sediments (November)	Chen et al., 2016
Hong Kong (sediment)	2008-2010	<0.026		LOD = 0.03-0.1		Loi et al., 2011
Canada (sediment)	2012-2016	nd		LOD = 0.04	Urban	D'Agostino and Mabury et al., 2017

One study of PFBS in soil in Europe reported 10 % frequency of detection of PFBS and a max of 0.003 ng/g dw, which is relatively low (Herzke et al., 2016). In China, a study of PFBS in garden soil in the vicinity of a fluorochemical industrial park found high concentrations of PFBS with a maximum of 42 ng/g dw (Bao et al., 2019).

The range of PFBS in sediments is from < LOD to 0.2 ng/g dw, at the low end. Stock et al. (2007) was able to quantify PFBS in Arctic surface sediments, with a max concentration at 0.1 ng/g dw, however the contamination could be attributed to local emission from an airport.

In general, there is little data on PFBS in soil and sediments. However, PFBS is not primarily expected to be found in soil and sediments due to the preferred partitioning to water.

3.2.6. Summary and discussion of environmental distribution

The high complexity of the PFAS chemistry makes it difficult to predict the sorption of the substances from a single sorbent bulk property. Properties such as pH, identity of the soil/sediment and surface-bound cations interact to determine the binding of PFASs. The sulfonic acids tend to sorb more strongly than carboxylic acids, and the PFAS sorption tends to increase with increasing perfluorocarbon chain length. However, PFBS has been found to be a weakly sorbing substance with a high mobility in the environment. PFBS has a preference for distribution to the aqueous phase and is relatively readily transported when water flows through soil. Log K_{oc} for PFBS has been reported in the range 1.2 to 2.7.

PFBS is a strong acid which is fully deprotonated at environmentally relevant conditions. The corresponding sulfonate anion in solution, as well as dissolved PFBS salts do not volatilise. Preference for the water compartment has been confirmed by distribution modelling and by monitoring data, showing that PFBS is frequently found in surface water, groundwater, drinking water, marine water, effluents from waste water treatment plants and in landfill leachate. The concentrations vary from < LOD to 21200 ng/L in the proximity to known point sources (1666 ng/L as max in Europe). Near point sources, like a PFAS production facility in the Netherlands, PFBS was the dominating PFAS in both river samples (12-27 ng/L) and drinking water (0.5-19 ng/L), Gebbink et al. (2017). In China near a PFAS production facility at Tangxun Lake, the two most dominating PFAS in water were PFBA and PFBS, at means of 4770 and 3660 ng/L, respectively (Zhou et al., 2013). In groundwater a remarkable 24-fold increase in PFBS concentration was recorded in the

vicinity of a PFAS production plant in Fuxin, China, from 872 ng/L to 21 200 ng/L from 2009 to 2017 (Bao et al., 2019).

In many cases, PFBS is among the most dominating PFASs in environmental samples where PFBS has been observed to be the third most frequently detected PFAS in bottled water (47 %) and tap water (88%) in a global survey (Kaboré et al., 2018). The substance was also frequently detected with 27 % in drinking water samples from Brazil, Spain and France (Schwanz et al., 2016). Furthermore, in a recent survey of surface water in Northern Europe, with sampling in 2013, PFBS was found to be the dominating PFAS, contributing with 21% of the sum PFAS (Nguyen et al., 2017).

3.3. Removal from the environment, decontamination and purification

Methods for removal of PFBS from environmental media are important in order to have the possibility to purify contaminations and lower environmental and human exposure. As PFBS has a clear preference for the aqueous phase, purification techniques for water are most relevant. PFBS has been detected in surface water and ground water, but also in produced drinking water. Purification techniques suitable for drinking water treatment plants are therefore essential, while the purification of waste water is equally important.

Rahman et al. (2014) reviewed PFAS characteristics, their occurrence in surface water, and their fate in drinking water treatment processes. Occurrence data from full-scale drinking water treatment plants indicated that PFASs, if present in raw water, are not substantially removed by most drinking water treatment processes. Conventional coagulation, flocculation, and sedimentation cannot achieve more than 20% removal of PFAS, nor can rapid granular media filtration. Most PFASs are not substantially removed in oxidation or advanced oxidation processes, typical for drinking water treatment plants. Biodegradation of most PFASs under current drinking water treatment conditions is unlikely. Filtration through granular activated carbon (GAC) may be useful for removing PFASs from drinking water. Longer chain PFASs will sorb better than the shorter chain compounds. However, short-chain PFASs such as PFBA and PFBS may pass through or reach breakthrough very quickly. Ion exchange/non-ion exchange resins may be useful for removing PFASs. However, such equipment is not commonplace in drinking water treatment facilities. Nanofiltration and reverse osmosis membranes may be effective with a high rejection of most PFASs. However, low molecular weight PFASs may be less well rejected by some loose NF membranes. Besides, disposal of concentrate with elevated concentrations of PFASs, will need to be addressed.

Reducing the releases of PFBS in industrial emissions and wastewater also suffers from low efficiency due to the high mobility and low adsorption potential of PFBS. Reemtsma et al. (2016) point out that surface waters in densely populated areas are the recipients of effluents of wastewater treatment plants (WWTPs) as well as of runoff from urban surfaces and agricultural land. Wastewater treatment and subsurface barriers are in general ineffective for the removal of persistent and mobile substances, and therefore they may reach the raw waters used for drinking water production.

Ochoa-Herrera and Sierra-Alvarez (2008) investigated the removal of perfluorinated surfactants by sorption onto granular activated carbon, zeolite and sludge. They found that activated carbon adsorption is a promising treatment technique for the removal of PFOS from dilute aqueous streams, while the sorption was weaker for PFOA and PFBS. The estimated concentrations of PFBS sorbed onto granulated activated carbon (GAC) was 48 mg/g GAC for PFBS, compared to 57 mg/g GAC for PFOA and 182 mg/g GAC for PFOS. The adsorptive capacity decreased with increasing PFAS concentrations in the water.

Furthermore, the presence of precursor compounds may complicate the water treatment process as the precursors may have considerably different properties and behave differently through the purification steps compared to their degradation products. The precursors may even break down during or after purification which may lead to increased concentrations of the degradation products in the finished water (Rahman et al., 2014).

The behaviour of PFAAs from raw source water through a drinking water production chain to finished drinking water was examined in a study (Eschauzier et al., 2012). The raw source water was subjected to a series of purification steps, including coagulation, rapid sand filtration, dune passage, aeration, rapid sand filtration, ozonation, pellet softening, granular activated carbon (GAC) filtration, slow sand filtration. It was found that longer chain PFAA were readily removed by the GAC treatment step, while more hydrophilic shorter chain PFAA, especially PFBA and PFBS, were not removed by GAC and their concentrations remained constant through treatment.

In a study, Appleman et al. (2014) measured concentrations of PFASs in 18 raw drinking water sources and 2 treated wastewater effluents and evaluated 15 full-scale treatment systems for the attenuation of PFASs in water treatment utilities throughout the U.S. Water treatment techniques, such as coagulation followed by physical separation processes, and chemical oxidation, aeration and disinfection, were unable to remove PFASs. The levels of several PFASs were reduced by reverse osmosis (RO), granular activated carbon (GAC) and anion exchange (AIX). However, GAC and AIX were less effective at removing the shorter chain PFASs, whereas RO treatment was effective for even the smallest PFAS studied, PFBA. Despite RO's effectiveness, RO would likely be the most costly method for purification.

The low efficiency in most water treatment processes for PFBS is associated with the physicochemical properties of the substance. The high aqueous solubility and the low sorption potential results in a preferred distribution to the aqueous phase and low binding to soil and sediments, as well as adsorbents used in purification. Hence, most conventional purification techniques are ineffective in the removal of PFBS. Some more modern methods are promising and indicate a higher purification effect also for short-chain PFASs like PFBS. However, the equipment is not commonplace in water treatment facilities, and the treatment techniques may be expensive. In addition, it remains to be seen how the methods perform over time in practice with varying influent and the presence of other contaminants that affect the purification efficiency.

In summary, due to the high aqueous solubility and the low sorption potential, resulting in a preferred distribution to the aqueous phase, PFBS is not efficiently removed with conventional purification techniques. Both wastewater purification, drinking water production and removal of industrial emissions may suffer from the low purification efficiency for PFBS. The presence of PFBS precursors may complicate the water treatment process as the precursors may behave differently through the purification steps and may break down to PFBS either during or after purification. Efficient, practically and economically available methods do not exist today to remove PFBS contaminations and purify waste water and drinking water and lower environmental or human exposure. However, some more modern methods have promising removal efficiencies for PFBS and may be developed into practically available treatment techniques in the future.

3.4. Data indicating potential for long-range transport

3.4.1. Mobility and the potential for long-range transport

The hazards of short-chain perfluoroalkyl acids and other fluorinated alternatives to long-chain PFASs were assessed and compared to their long-chain homologues by Wang et al.

(2015). It was concluded that short-chain PFAAs, due to their higher solubility in water and lower sorption to solids, are more mobile in soil and sediment. This leads to higher mobility in the environment. In combination with the high environmental stability, the high mobility implies that the short-chain PFAAs have a high global contamination potential.

This is supported by the findings in Sections 3.2.1 and 3.2.2 on adsorption/desorption and Section 3.2.4 on the modelling of distribution, wherein physical-chemical properties and model estimates suggest that PFBS is a highly soluble, weakly sorbing substance, with a high mobility in the environment and a preference for distribution to the aqueous phase. PFBS maintains in the water phase and is subject to long-range transportation first of all via sea currents.

Wania (2003) looked at the partitioning properties that favour enrichment in the Arctic ecosystems and found that the long-term Arctic contamination potential of a perfectly persistent organic compound tends to be high for relatively volatile and water-soluble substances, and semivolatile and relatively hydrophobic substances with $\log K_{oa}$ 6.5 – 10 and $\log K_{AW} > -3$. Arp and Slinde (2018) refined the assessment and explained that for ionic substances like PFAS, one should use the D_{aw} rather than K_{aw} , and $K_{oc,a}$ rather than K_{oa} . Recalling from Table 9 the $\log K_{oc,a}$ values for PFBS (4.8) is very similar to PFHxS (5.0) and PFOS (4.7). The $\log D_{aw}$ values at pH 8 for PFBS (-14.5) are considered extremely low, similar to PFHxS (-13.8) and PFOS (-13.1), such that volatilisation from water to air for all three substances is negligible. This combination of physical-chemical properties implies that PFBS strongly will prefer the water phase, will be transported with sea currents, and will reach and accumulate in the Arctic if emissions are allowed to continue. A similar behaviour has been observed for PFOS and PFHxS.

A recent study of the partitioning behaviour of PFASs between the dissolved phase, surface sediment and suspended particulate matter around coastal areas of Bohai Bay, China, suggested that PFBS and PFHxS were entirely preserved in the water-dissolved phase, while PFDS tended to stay in the suspended particulate matter section (Liu et al., 2019). Ding et al. (2018) observed a somewhat different behaviour when similar measurements were performed in the Dalian Bay, China. PFOA, PFBA, and PFBS were the predominant PFASs in the water dissolved phase, while PFBS, PFOS and PFOA were the most prevalent compounds in suspended particulate matter. A $\log K_d$ for PFBS of 3.4 was reported, and it was concluded that PFASs (including PFBS) and the long-chain PFCAs were more inclined to prefer the suspended particulate matter phase. The differences between the two studies indicate that the tendency of PFBS to bind to particles in water may depend upon the specific properties of the particles.

A study of the role of sea spray aerosols in the long-range transport of perfluoroalkyl acids was performed by Johansson et al. (2018). It was measured from 200 to 62000 times higher concentrations of PFOS in the aerosols compared to the water concentrations, depending on the aerosol size. No specific values were given for PFBS, but the study indicated enrichment factors approximately one order of magnitude lower for PFBS than for PFOS. The results demonstrated that sea spray aerosols have the capacity to circulate significant amounts of PFAAs between the oceans and the atmosphere. A portion of the mass emitted from the oceans will deposit on land, thus re-entering the terrestrial system. This suggests that human exposure to PFAAs will continue even if strict global emission controls are implemented.

Reemtsma et al. (2016) pointed out that persistent and mobile organic compounds may be of concern for water quality because they are persistent in the environment and are not removed from water by sorption processes due to their high polarity and excellent water solubility. They may end up in drinking water, posing a potential risk to human health. However, there is no commonly accepted quantitative definition for a compound's mobility in regulations or elsewhere. Two possible quantifiers of mobility in water are water solubility and sorption tendency. Short-chain perfluoroalkyl carboxylates and sulfonates

(especially C2 and C3) were mentioned as a group of substances with properties of persistent and mobile compounds. Brendel et al. (2018) concluded that short-chain PFAAs, including PFBS, PFBA and PFHxA, have physicochemical properties that make them very mobile and prone to end up in the aqueous compartment. Due to the low adsorption potential, short-chain PFAAs will not bind to particles and stay mainly dissolved in the water phase. Furthermore, the short-chain PFAAs will have a higher potential for long-range transport compared to the long-chain homologues because of the high mobility and low adsorption potential.

As a mobile substance, there are no local or intermittent sinks for the pollution stock of PFBS, and therefore the substance has high potential for continuously increasing environmental concentrations and exposure of wildlife.

The ionic PFASs have high water solubility and low pKa values and are therefore almost completely dissociated at environmentally relevant pH values. Oceanwater is important as a sink and for transport of these compounds. The occurrence of high concentrations of PFASs in coastal waters could possibly be problematic, because the substances will be bioavailable and can accumulate in the marine food chain (Cai et al., 2012b).

3.4.2. Estimation of long-range transport potential

An atmospheric half-life of 76.4 days was estimated for PFBS with the Atmospheric Oxidation Program (AOPwin, v1.92) for EPISuite models, see Section 3.1.1.2. This value exceeds the threshold of two days which indicates a long-range transport potential for a substance.

The long-range transport potential of PFBS was modelled using the OECD P_{ov} & LRTP Screening Tool, a tool for estimation of the Long-Range Transport Potential (LRTP-Tool). Inputs to the model are half-lives in air, water and soil, as well as $\log K_{aw}$ and $\log K_{ow}$. The half-lives were estimated at $t_{1/2}(\text{air}) = 1830$ h, $t_{1/2}(\text{water}) = 4320$ h and $t_{1/2}(\text{soil}) = 8640$ h (Level III Fugacity Model, as present in OECD QSAR toolbox v4.2, February 2018). Air-water partitioning for an ionic substance like PFBS is pH-dependent. $\log K_{aw}$ represents the neutral form, while $\log D_{aw}$ varies with pH and shows the pH-dependence of the air-water partitioning. Values and ranges for $\log K_{aw}$ and $\log D_{aw}$ are presented in Table 5. In the present dossier, the estimate based on $\log K_{aw}$ for the neutral form ($\log K_{aw} = -2,59$, Table 5) is presented, as this allows for comparison with similar ionic substances for which estimates based on the neutral form exist. A substance with a lower $\log K_{aw}$ (e.g. PFBS at neutral pH) would be more influenced by the persistence in water, which for PFBS in this model is lower, compared to air. $\log K_{ow}$ is also pH-dependent and in Table 5 different values are reported for different pH-values. At pH 7-8, $\log K_{ow}$ is reported in the interval -8.0 to -3.0. The LRTP-Tool returns the same results for both extremes -8.0 and -3.0. Hence, $\log K_{ow} = -3.0$ was selected for the estimation.

The LRTP-Tool estimated for PFBS a characteristic travel distance, CTD = 17616 km and overall environmental persistence, $P_{ov} = 220$ days. CTD indicates the distance from a point source at which the chemical's concentration has dropped to 38% of its initial concentration. The results for PFBS indicate a high potential for long-range transport. The results are presented graphically in Figure 4. However, the present estimates represent screening estimates and are influenced by the physico-chemical input parameters. The results are in support of other evidence of transport of PFBS to remote areas.




Figure 4: Graphical output for the calculated CTD = 17616 km and POV = 220 days for PFBS (red dot). The lines in the figure represent the border for POP-like substances with high POV and high L RTP. The results indicate a high potential for long-range transport for PFBS.

According to Crookes and Fisk (2018), the overall CTD for substances with $\log K_{aw}$ down to around -3 is governed mainly by transport via the atmosphere, but at lower $\log K_{aw}$ values, the overall CTD is increasingly governed by transport via the water phase.

3.4.3. Occurrence of PFBS in remote areas

Butt et al. (2010) reviewed the levels and trends of PFASs in the Arctic environment and concluded that PFASs are ubiquitous in the Arctic. They pointed out that the marine food web has been well studied, particularly the top predators, while freshwater and terrestrial ecosystems have been poorly studied. PFOS was found to be most frequently detected in the Arctic wildlife, followed by PFNA and PFUnA. However, some whale species showed relatively high levels of perfluorooctane sulfonamide (PFOSA), and seabirds are typically characterised by high proportions of the C11–C15 PFCAs. PFOA is generally detected at low levels only in Arctic biota. PFBS is reported at low levels in Arctic biota, and this was explained by a low tendency to bioaccumulate due to the short biological half-life. PFBS was a more dominant contributor to the overall PFAS levels in Arctic water samples.

Furthermore, Butt et al. (2010) discussed the relative importance of the two dominant transport pathways of PFASs to the Arctic: Atmospheric transport of precursors and degradation, versus direct transport of PFASs via ocean currents. However, there was not enough evidence to draw any final conclusions.

Zhao et al. (2012) investigated the global distribution and long-range transport of PFASs using seawater samples collected from the Greenland Sea, East Atlantic Ocean and the Southern Ocean in 2009-2010. Relatively high concentrations of PFASs were detected near the European continent. In the Greenland Sea, the five most frequently detected compounds were PFOA, PFHxS, PFHxA, PFOS and PFBS.

In the following sections PFBS levels in different compartments in remote regions are compiled.

3.4.3.1. PFBS in remote areas – fresh water, air, snow and ice

Measurements of PFBS in samples of fresh water, air, snow and ice in remote regions are collected in Table 17.

Table 17: Environmental concentrations of PFBS in remote areas; fresh water, air, snow and ice samples (ng/L).

Location	Year	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
Arctic Ocean	2010	0.17 ± 0.42	<0.017 - 1.50 (0.02)	12 (50 %) LOD = 0.017	Snow/Ice	Cai et al., 2012a
Canada	2007 - 2008		0.01 - 0.02	11 LOD = Not reported	Lake	Veillette et al., 2012
Canada	2010 - 2011	0.07 ± 0.01		5 LOD = 0.0019	Lake	Lescord et al., 2015
Greenland	2015	0.0068		LOD = 0.00091	Ice cap (19 cm = 2015)	Pickard et al., 2018
Greenland	2015	0.0042		LOD = 0.00091	Ice cap (134 cm = 2013)	Pickard et al., 2018
Northern Sweden	2009	0.099	nd - 2.16	24 LOD = 0.021	Snow/Ice	Codling et al., 2014

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Year	Mean +/- SD	Min-Maks (Median)	n (det. Freq) LOD	Remark	Reference
Svalbard, Longyearbyen	2015	2.38		LOD = 0.003	Run-off water	Skaar et al., 2019
Svalbard, Longyearbyen	2015	2.41		LOD = 0.003	Waste water effluent	Skaar et al., 2019
Svalbard, Longyearbyen	2015	2.33		LOD = 0.003	Local firefighting training site (run-off)	Skaar et al., 2019
Svalbard, Ny-Ålesund	2015	22.99		LOD = 0.003	Freshwater, contaminated location	Skaar et al., 2019
Antarctica (Lake)	2011	0.038	<0.0083 - 0.05	4 LOD = 0.017	Lake (King George Island)	Cai et al., 2012
Antarctica	2011	0.017	<0.0083 - 0.02	4 LOD = 0.017	Snow/Ice (King George Island)	Cai et al., 2012
Alert, Greenland	2006-2015	0.1 ± 0.2	BDL - 1.5 (0.0)	169 (66 %) LOD = 0.008-1.8	Air, pg/m ³	Wong et al., 2018
Zeppelin, Svalbard	2006-2015	nd		383 LOD = 0.008-1.8	Air, pg/m ³	Wong et al., 2018

The PFBS levels measured in air, fresh water, snow and ice in remote regions vary between not detected and 2.16 ng/L when not including samples with known contamination. The concentrations are generally low, but the frequent detection shows that PFBS has spread via long-range transport to the Arctic, as well as the Antarctic. Waste water and runoff samples from Svalbard contained PFBS at concentrations in the range 2.33 – 22.99 ng/L.

3.4.3.2. PFBS in remote areas – marine water

PFBS has been reported in marine water throughout the world. Measurements from marine water collected in remote regions are shown in Table 18.

Table 18: Environmental concentrations of PFBS in marine water samples from remote regions (ng/L).

Location	Year	Mean +/- SD	Min-Max (Median)	n (det. Freq) LOD	Reference
Arctic Ocean	2010	0.03 ± 0.03	<0.017 - 0.08 (<0.017)	13 (38 %) LOD = 0.017	Cai et al., 2012
Central Arctic	2012	0.01 ± 0.01	0.04	LOD = 0.005	Yeung et al., 2017
Faroe Islands	2004		0.05 - 0.11	LOD = 0.03	Kallenborn et al., 2004
Iceland	2004		0.05 - 0.08	LOD = 0.03	Kallenborn et al., 2004
North Atlantic	2009-2010	nd	0.045	62 LOD = 0.0059-0.051	Zhao et al., 2012
Northwest Pacific Ocean	2010	0.02 ± 0.03	<0.017 - 0.10 (<0.017)	9 (33 %) LOD = 0.017	Cai et al., 2012
Russian Arctic Baydaratskaya Bay	2007	0.0132		LOD = Not reported	Saez et al., 2008
Antarctica	2007	0.0029		LOD = Not reported	Wei et al., 2007
South Atlantic	2009-2010	nd	nd-0.013	39 LOD = 0.0059-0.051	Zhao et al., 2012

Location	Year	Mean +/- SD	Min-Max (Median)	n (det. Freq) LOD	Reference
Mid-Atlantic	2009-2010	nd	nd-0.017	20 LOD = 0.0059-0.051	Zhao et al., 2012
Svalbard, Longyearbyen	2015	0.105		0.003	Skaar et al., 2019
Svalbard, Ny-Ålesund	2015	<0.001		0.001	Skaar et al., 2019

PFBS has been detected in marine water from remote regions in the range not detected to 0.11 ng/L in a sample from the Faroe Islands (Kallenborn et al., 2004). The ubiquitous presence of PFBS in marine waters over a very large area is an evidence of long-range transport of PFBS.

3.4.3.3. PFBS in remote areas – biota

According to Arp and Slinde (2018), prior to 2012, the majority of studies on Arctic marine biota did not report PFBS above the detection limits. In more recent studies, it is starting to appear more frequently. Routti et al. (2016) did not detect PFBS in any blood plasma samples in ringed seals from Svalbard in the period of 1990 to 2010. In a study by Gebbink et al. (2016) in East Greenland, PFBS was detected in all polar bear livers examined (8/8) at 0.032 ± 0.008 ng/g and in 67% of the killer whale livers at 0.0052 ± 0.0017 ng/g. However, PFBS was below detection limit in the ringed seal livers. Total PFSA in the three species was in the range 94-1825 ng/g, with PFOS being the predominant single substance. PFBS was measured in livers of both mothers and foetus of killer whale with a transfer rate of 2.3% (4.7% for PFOS). Routti et al. (2017) detected PFBS in polar bear plasma at Svalbard. In a study of temporal trends of perfluoroalkyl compounds in the period 1975-2011, Braune and Letcher (2013) detected PFBS in eggs of thick-billed murres (*Uria lomvia*) and northern fulmars (*Fulmarus glacialis*) from the Canadian Arctic in 2010 and 2011. Concentrations of PFBS measured in marine biota in the Arctic is listed in Table 19.

Table 19: Environmental concentrations of PFBS in Arctic biota (ng/g ww).

Location	Date	Mean +/- SD	Min – Max (Median)	n (df) LOD	Remark ^{a)}	Reference
Greenland	2012-2013	<0.002		10 LOD = 0.002	Ringed seal liver (<i>Pusa hispida</i>)	Gebbink et al., 2016
Greenland	2012-2013	0.03 ± 0.01		8 (100 %)	Polar bear liver (<i>Ursus maritimus</i>)	Gebbink et al., 2016

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min – Max (Median)	n (df) LOD	Remark ^{a)}	Reference
				LOD = 0.002		
Greenland	2012-2013	0.01 ± 0.00		6 (66 %) LOD = 0.002	Killer whale liver (<i>Orcinus orca</i>)	Gebbink et al., 2016
Canadian Arctic	1975-2011	0.57 ± 0.46		12 (4 pools of 3 eggs) detected in 2011 LOD = 0.1	Northern fulmars (<i>Fulmarus glacialis</i>)	Braune and Letcher (2013)
Canadian Arctic	1975 - 2011	0.04 ± 0.02 (2010) 0.07 ± 0.05 (2011)		15 (5 pools of 3 eggs) LOD = 0.1	Thick-billed murres (<i>Uria lomvia</i>)	Braune and Letcher (2013)
Nunavut, Canada	1972-2005	nd		184 LOD = Not reported	Ringed Seal liver	Butt et al., 2008
Svalbard	1990-2010	<0.07		LOD = Not reported	Ringed seal plasma	Routti et al., 2016
Svalbard	2000-2014	0.27	0.08 - 0.69 (0.27)	70 (100 %) LOD = Not reported	Polar bear plasma	Routti et al., 2017
Svalbard	2007	nd		LOD = 0.06-0.11	Black guillemots liver (<i>Cepphus grylle</i>)	Axelsson et al., 2014
Svalbard	2007	nd		LOD = 0.06-0.11	Glaucus gull liver (<i>Larus hyperboreus</i>)	Axelsson et al., 2014
Svalbard	2016	nd		LOD = Not reported	Ringed seal liver	Schlabach et al., 2018
Svalbard	2016	nd		LOD = Not reported	Polar bear whole blood	Schlabach et al., 2018
Svalbard	2017	<0.05		10	Common eider eggs	Schlabach et al., 2018

Location	Date	Mean +/- SD	Min – Max (Median)	n (df) LOD	Remark ^{a)}	Reference
				LOD = Not reported	(<i>Somateria mollissima</i>)	
Svalbard	2017	<0.07		5 LOD = Not reported	Black-legged Kittiwake eggs (<i>Rissa tridactyla</i>)	Schlabach et al., 2018
Svalbard	2017	<0.08		5 LOD = Not reported	Glaucus gull eggs	Schlabach et al., 2018
Svalbard	2017	0.04 ± 0.02	0.02 - 0.08 (0.04)	10 (60 %) LOD = Not reported	Polar bear plasma	Schlabach et al., 2018

The PFBS concentrations measured in Arctic biota vary between not detected and 0.69 ng/g ww. In most of the samples, concentrations were low. However, in a study of polar bear (*Ursus maritimus*) considerable levels (0.08 – 0.69 ng/g ww, median 0.27 ng/g ww) were found in plasma (Routti et al., 2017). Low levels were also found in killer whales (*Orcinus orca*) from Greenland in both mothers and the foetus (Gebbinck et al., 2016). PFBS is also found in eggs of northern fulmars and thick-billed murres in the Canadian Arctic.

3.4.4. Long-range transport of precursors and degradation to PFBS

In some cases, PFBS-related substances may also be efficiently transported over long distances or transported via other routes than PFBS itself, e.g. via atmospheric transport. Subsequent degradation with formation of PFBS will then contribute to the wide distribution of PFBS all over the globe. Due to the negative charge of the PFBS sulfonate anion, PFBS will have a low tendency to evaporate and for transport via air, while neutral precursors in many cases will have a higher vapour pressure and a stronger tendency to spread via air (Armitage et al., 2009). Another route of long-range transport is via distribution of PFBS-containing products. PFBS-related substances, including polymers, are in use in various products, like textiles, as a surface treatment agent (Chu and Letcher, 2014). Such products may be distributed world-wide for sale and use, and degradation of the precursors to PFBS during the use phase or in the waste stage will contribute to the increasing PFBS levels over a wide area.

A review of the microbial degradation of polyfluoroalkyl chemicals in the environment points out that perfluoroalkane sulfonamido derivatives may undergo aerobic biodegradation, via the relatively stable intermediate sulfonamides, to the sulfonic acids (e.g. PFBS) as the final degradation products (Liu and Avendaño, 2013). A comprehensive discussion of abiotic degradation of PFBS-precursors to PFBS may be found in the study by Nielsen (2017), and a summary in Section 1.3.3.

Monitoring data show that transport via air of neutral and volatile PFASs, like various substituted fluoroalkane sulfonamides, including the PFBS-precursors MeFBSE and MeFBSA, is actually happening (Lai et al., 2016). However, the dominant neutral PFAS-group detected in the study was the fluorotelomer alcohols (FTOHs). D'eon et al. (2006) showed that MeFBSE can degrade in reaction with OH radicals in the atmosphere to produce MeFBSA, in addition to PFBS and PFCAs as final degradation products. The atmospheric lifetime of MeFBSA was estimated at >20 days which suggests that the substance is long-lived enough to account for distribution throughout the North American troposphere, linking the production of MeFBSE to the ubiquity of this class of compounds in the remote environment (D'eon et al., 2006).

3.4.5. Summary of data indicating potential for long-range transport

Due to the high solubility in water and low adsorption to solids, PFBS has a preference for the aqueous compartment and is mobile in the environment. In combination with the high environmental stability, the high mobility implies that PFBS has a high global contamination potential. The long-range transport of PFBS is mainly via the water phase and sea currents, while sea spray aerosols have the capacity to circulate significant amounts of PFAAs between the oceans and the atmosphere. A portion of the mass emitted from the oceans will deposit on land, thus re-entering the terrestrial system.

An atmospheric half-life of 76.4 days was estimated for PFBS with the Atmospheric Oxidation Program, indicating a potential for long-range transport. This was confirmed by modelling with the OECD LRTP-Tool which concluded that PFBS has a characteristic travel distance of 17616 km and a high potential for long-range transport.

In addition to the long-range transport of PFBS itself, long-range transport of PFBS-related substances and degradation to PFBS may take place. The related substances have different properties than PFBS and may be transported via other routes, like atmospheric transport.

PFBS has been detected in samples of marine water, surface water, snow, ice, air and biota from remote areas, demonstrating that long-range transport has taken place. However, the measured concentrations of PFBS are generally low.

3.5. Bioaccumulation

REACH Annex XIII, Sections 1.1.2 and 3.2.2(a), defines the numerical criterion for bioaccumulation when assessing bioaccumulation data from fish or other aquatic species. However, due to their water solubility, short-chain PFASs are expected to be quickly excreted via gill permeation in water breathing animals. This reduces the bioaccumulation potential in such species.

Annex XIII (Section 3.2.2) further defines information which should be taken into account when the numerical criterion is not applicable, i.e. data on the bioaccumulation potential in terrestrial species and detection of elevated levels in biota, in particular in endangered species or in vulnerable populations, compared to levels in their surrounding environment. Annex XIII (Section 3.2.2 (b)) also allows taking data from human body fluids or tissues and the toxicokinetic behavior of a substance into account. In accordance with this, elimination half-lives have recently been used for certain PFAAs as a metric to estimate the bioaccumulation potential in air-breathing organisms. For example, the elimination half-life for PFOA in humans was important in the identification of PFOA (half-life 2 – 4 years) as a substance fulfilling the B criteria in Annex XIII (ECHA, 2013). PFHxS with a

half-life in humans of ca 7-8 years (or longer) was agreed to fulfil the vB criteria (ECHA, 2017).

3.5.1. Laboratory studies on bioaccumulation

3.5.1.1. Bioaccumulation in aquatic organisms

In a study by Martin et al. (2003a) in rainbow trout (*Oncorhynchus mykiss*), fish were exposed simultaneously to a homologous series of perfluoroalkyl carboxylates and sulfonates (including potassium PFBS) for 12 days, followed by 33 days of depuration in clean water in a flow-through system to determine compound-specific tissue distribution and bioconcentration parameters for perfluoroalkyl acids (PFAAs). In general, PFAAs accumulated to the greatest extent in blood > kidney > liver > gall bladder. PFBS was not detected in blood, liver and carcass at any sampling times, although as pointed out by the authors, a half-life of PFBS in liver of rainbow trout of 3.3 days is measured in a dietary accumulation study (see next paragraph). The study is well reported and regarded as reliable with a Klimisch reliability of 2 as the limit of detection in the different tissues is not reported.

In another study by Martin et al. (2003b) juvenile rainbow trout were exposed to a homologous series of perfluoroalkyl carboxylates and sulfonates (including potassium PFBS) in the diet for 34 days, followed by a 41 days depuration period. On six occasions during both the uptake period and the depuration period, fish were removed for analysis of carcass and liver concentrations. Concentrations were determined by LC-MS, and kinetic rates were calculated to determine compound-specific bioaccumulation parameters. Spiked food concentration was 0.32 µg/g for PFBS. PFBS was only detectable at the last three uptake sampling intervals and at the first sampling time of the depuration phase and was apparently only detected in liver. Limits of detection are not reported. A depuration half-life for PFBS for liver of 3.3 days is reported. The depuration half-lives for liver for PFHxS and PFOS were reported to be 13 days and 20 days, respectively. In general sulfonates bioaccumulated to a greater extent than carboxylates of equivalent perfluoroalkyl chain length. The recovery for PFBS in carcass and liver was low; around 60% or less. Otherwise, the study is regarded as reliable with a Klimisch reliability of 2. For PFBS a BAF could not be calculated. According to the authors, dietary exposure will not result in biomagnification of perfluoroalkyl acids in juvenile trout, but extrapolation to larger fish and homeothermic organisms should not be performed.

In a feeding study with adult rainbow trout the biomagnification potential and tissue-specific distribution of several perfluoroalkyl acids (PFAAs) were investigated (Goeritz et al., 2013). Tetrabutylammonium nonafluorobutane sulfonate at a chemical purity of sulfonic acid ≥ 98% was used. Biomagnification factors (BMFs) were determined based on a kinetic approach. Distribution factors were calculated for each test compound to illustrate the disposition of PFASs in rainbow trout after 28 d of exposure. Mean concentration of PFBS in the feed was 185 ± 26 µg/kg feed. BMF values of 0.02 for PFBS, 0.42 for PFOS and 0.18 for PFHxS were calculated. Liver, blood, kidney, and skin were identified as the main target tissues. Muscle tissue (37.6% of total body wt) comprised 41%, 41% and 26 % of the whole-body burden of PFBS, PFOS and PFHxS, respectively, on day 28 of the exposure period. The accumulation phase was followed by a 28-d depuration phase, in which the test animals were fed with non-spiked feed. The depuration half-life for whole fish was calculated to be 10.8 days for PFBS, 8.75 days for PFHxS and 16.1 days for PFOS. The authors pointed out that despite relatively low PFAS contamination in muscle and skin, the edible parts of the fish can significantly contribute to the whole-body burden.

In a follow up publication (Falk et al., 2015), based on data collected in the Goeritz study with rainbow trout described above, tissue specific uptake and elimination of the

perfluoroalkyl acids (PFAAs) were further investigated. Muscle, liver, kidneys, gills, blood, skin and carcass were examined individually. At the end of the accumulation phase 0.89% (PFBS), 3.79% (PFHxS) and 17.5% (PFOS) of the absolute, applied quantity of the PFSA was recovered in the whole fish. The main target organ was the liver with recovery rates between 0.11% (PFBS) and 4.01% (PFOS) of the total amount of PFSA supplied. For PFBS a maximum concentration of 110 µg/kg was found in the liver at the end of the accumulation phase (day 28), and 80% of the maximum was reached by day 7. Perfluoroalkyl sulfonic acids were taken up more readily and had longer estimated elimination half-lives than perfluoroalkyl carboxylic acids of the same chain length. The longest estimated tissue specific elimination half-life was found to be for PFOS and was 20.4 days in the liver. For PFBS the longest tissue specific elimination half-life was for liver and was 6.5 days.

KPFBS bioconcentration was examined in Bluegill sunfish (*Lepomis macrochirus*) in a test conducted under OECD 305 guidelines. Fish were exposed for 28 days with a 14-day depuration period. Average measured test substance concentrations were 0.53 mg/L and 5.2 mg/L. Steady-state BCFs (whole fish) was 0.3 and were attained in ≤ 7 days. Kinetic BCF values were 0.36 at 5.2 mg/L and 1.1 at 0.53 mg/L. The study was assigned a Klimisch reliability 1 (ECHA dissemination website). However, from the information provided the values reported do not seem to be growth corrected. → *Klimisch reliability 2-well documented Guideline study, acceptable for assessment.*

In a study by Chen et al. (2016a) adult female zebrafish (*Danio rerio*) were exposed for 24 days to an aqueous mixture of multiple PFASs (including free acid form of PFBS), followed by a 24 days depuration phase. The test was performed according to OECD Guideline 305 (Bioconcentration: Flow-through Fish Test). The water concentration of PFBS was 2.15 ± 0.28 µg/L in the low dose group and 17.8 ± 1.28 µg/L in the high dose group. PFBS was detected in all tissues except for muscle tissue of fish in the low-dose exposure group. Growth-corrected concentrations were calculated using the exponential growth model. Reported BCF values (whole fish) for PFBS were from 19.5 – 27.5. The body burden was calculated as the sum of tissue concentrations measured (plasma, liver, muscle, ovary) times the wet weight of the respective tissues, other tissues not included in estimating the whole-body burden. → *Klimisch reliability 2-well documented Guideline study, acceptable for assessment.*

Accumulation of PFAAs in the oligochaete *Lumbriculus variegatus* was investigated in a 28-days exposure study using sediments from contaminated areas (Lasier et al., 2011). The study is well documented, reliability 2 (ECHA dissemination website). Concentrations of PFAAs in oligochaete tissues revealed patterns similar to those observed in the respective sediments. Bioaccumulation of PFSA and PFCA by *L. variegatus* appeared to be primarily related to sediment concentrations. The tendency to bioaccumulate increased with PFAA chain length and the presence of the sulfonate moiety. With respect to PFBS a biota-sediment accumulation factor (BSAF) of 0.31 was determined that exceeded the BSAF values of PFOA (0.07), PFNA (0.20), PFDA (0.25), and PFUnDA (0.29). The authors concluded that sulfonates with four to seven carbons may be equally likely to bioaccumulate as PFOS in this species.

3.5.1.2. Bioaccumulation in earthworms

Earthworms (*Eisenia fetida*) were exposed to artificially contaminated soils with ten perfluoroalkyl substances (Zhao et al., 2013). A loamy surface soil (0 - 10 cm) without detectable PFASs was collected from a farm about 20 km southwest of Tianjin, China. The typical physicochemical properties of the soil were characterised as follows: pH 7.67; organic matter 4.88%; cation exchange capacity 38.47 cmol kg₋₁; moisture content 1.03% for airdried soil; clay 24%, silt 64%, and sand 12%. The soil was air-dried for 2

weeks, ground, and sifted through a 2 mm mesh. A small portion of pre-weighed soil was spiked with one mL of the mixed solution of 10 PFASs, dissolved in methanol and mixed thoroughly. The spiked soil was placed in a fume hood to allow the solvent to evaporate for 24 h. An aliquot of the untreated soil was added to the spiked soil and then mixed thoroughly. This step was repeated until all the pre-weighed soil was mixed. Three spiked soils were prepared and the concentration of each PFAS compound, including PFHxA, PFHpA, PFOA, PFNA, PFDA, PFUnDA, PFDoA, PFBS, PFHxS and PFOS, was at 100, 200, and 500 ng/g, respectively. The recoveries of PFASs were 76 - 105% for earthworms. The PFASs were not detected in the controls. The Method Detection Limits of PFASs were in the range of 0.5 - 1.0 ng/g dry weight (dw) for the target compounds. According to the authors, all the studied PFASs, including those with seven or less perfluorinated carbons, were bioaccumulated in the earthworms, and the biota-to-soil accumulation factors (BSAFs) increased with increasing perfluoroalkyl chain length and were greater for sulfonic acids than for the carboxylic acids of equal perfluoroalkyl chain length. The BSAFs were found to be dependent on the concentrations of PFASs in soil and decreased as the level of PFASs in soil increased. For PFBS, a BSAF value (kinetic) of 0.048 (0.768 for PFOS) was measured. The depuration half-life for PFBS was 9.6 days (18 days for PFOS). The BSAF measured based on wet weight of the organism, was 0.021 for PFBS at an exposure concentration of 100 ng/g (0.145 for PFOS).

In a study by Navarro et al. (2016) earthworms (*Eisenia andrei*) were exposed to sand soil spiked with a technical mixture of PFOS that contained PFBS. Worms were exposed for 21 days and allowed to depurate for 24 hours. PFBS was not detected in earthworms at the end of the depuration period although concentration in soil in some case (0.19 ng/g d.w.) reached levels similar to PFOS (0.21 ng/g d.w.). In another study by Navarro et al. (2017) the bioaccumulation behavior of perfluoroalkyl substances (PFASs) and halogenated flame retardants (HFRs) were examined in three horticultural crops and earthworms. Two species, spinach (*Spinacia oleracea*) and tomato (*Solanum lycopersicum* L.), were grown in field soil amended with a single application of biosolids (at agronomic rate for nitrogen), to represent the scenario using commercial biosolids as fertiliser. The third crop, corn (*Zea mays*), was grown in spiked soil with ~50 mg PFOS/kg soil (treatment 1), ~5 mg Deca-BDE/kg soil (treatment 2) and a mixture of both, ~50 mg PFOS and ~5 mg Deca-BDE/kg soil to represent a worst-case scenario (treatment 3). PFASs from wastes and soils (at the beginning $t=0$, and the end $t=final$; of the experiments) were extracted and purified. To examine the bioaccumulation in soil invertebrates, earthworms were exposed to the spiked soil after removal of the plants where corn had been grown. Worms were exposed for 28 days and allowed to depurate for 24 hours to avoid the presence of soil particulates that could interfere with the bioaccumulation study. Levels (treatment 1 and 3) found in earthworms were: PFBS 0.11 - 0.07 $\mu\text{g/g}$ d.w., PFHxS 3.18 - 2.93 $\mu\text{g/g}$ d.w. and PFOS 74.3 - 84.7 $\mu\text{g/g}$ d.w. BAFs (treatment 1 and 3) for individual substances, calculated based on levels in earthworms and levels in soil at $t=final$, were as follows: 3.89 - 6.09 (PFOS), 3.38 - 3.53 (PFHxS), 2.75 - 2.33 (PFBS).

3.5.1.3. Bioaccumulation in birds

In an acute and chronic effects (reproduction) study, adult Northern bobwhite quail (*Colinus virginianus*) was exposed to nominal dietary concentrations of 100, 300, or 900 mg PFBS/kg, ww feed for up to 21 weeks (Newsted et al., 2008). PFBS was present in liver and blood serum of adult quails in a dose-dependent manner. The ratio between mean PFBS concentrations in blood serum and liver for both adult male and female quail ranged from 3.4 to 5.1 with a mean value of approximately 4. Concentrations of PFBS in blood serum and liver were approximately 11- and 41-fold less than those in the diet, respectively, indicating that it is not bioaccumulating in the serum or liver of adult quails.

Concentrations of PFBS in eggs were directly proportional to dietary concentrations and were also dependent on when the eggs were laid during the study. Concentrations of PFBS in eggs laid during the seventh week were approximately 1.6-fold greater than that measured in eggs laid during the second week. Based on PFBS concentrations measured in adult females, the egg-to-serum ratio was approximately 1.0, while the egg-to-liver ratio was approximately 3.4 for all treatment groups. In 14-day-old offspring, PFBS concentrations in blood serum and liver were proportional but less than that measured in adult female quail or eggs from the same treatment group. PFBS concentrations in the tissues of offspring were at least 480-fold less than those measured in eggs from the seventh week for all treatment groups.

3.5.1.4. Bioaccumulation in mammals

A study by Numata et al. (2014) examined the transfer of a mixture of perfluoroalkyl acids (PFAAs) from contaminated feed into the edible tissues of 24 fattening pigs (*Sus scrofa domestica*). The feed, PFAA contaminated hay and barley, had been cultivated and harvested from a PFAA-contaminated region in Lower Saxony, Germany. The agricultural patch was polluted from the use of a soil improver provided by a recycling company that laced it with industrial waste containing high concentrations of PFAS. Mean concentration of PFBS in the feed was 132 ± 11 µg/kg on a dry weight basis. This study showed a tissue distribution of PFBS to blood plasma, liver, kidney, muscle tissue and fat. PFBS was eliminated through feces and urine. The geometric average elimination half-lives (up to 24 pigs) reflecting both plasma and edible tissue, was 43 days. The 95% variability elimination half-life intervals was 13-135 days for PFBS, taken from data analysis in ECHA, 2017. For comparison, the elimination half-lives for PFOS was 634 days (194-1970 days) and for PFOA 236 days (46-1074 days). Consequently, the individual variability was quite large, the highest values for PFBS are overlapping with the lower values for PFOA. BMF values reported for PFBS were 1.2 (whole pig), 0.8 (meat = mean of dorsal muscle, ventral muscle, and fat), 6.4 (liver), 14 (blood plasma), 2.2 (kidney) and 0.9 (fat). For comparison BMF values for PFOS were 17.9 (whole pig), 9.7 (meat), 503 (liver), 97 (blood plasma), 139 (kidney) and 8.3 (fat). The behavior of PFAAs in pigs is of interest because pigs are considered to be a biomedical model of human physiology (Numata et al., 2014).

In a study dairy cows (*Bos taurus*) were given feed that was contaminated with PFAAs, that is hay and grass silage which had been grown on a PFAA-contaminated farmland in Lower Saxony, Germany (Kowalczyk et al., 2013). Average concentration of PFBS was 68.4 ± 23.1 µg/kg in grass silage and 993.6 ± 224.4 µg/kg in hay. After the PFAA-feeding period of 28 days, three cows were slaughtered while the other 3 were fed PFAA-free feed for another 21 days (deuration period). The average daily intake of PFBS was 3.4 ± 0.7 µg/kg body weight, (for PFHxS, PFOS, and PFOA the average daily intake was 4.6 ± 1.0 , 7.6 ± 3.7 and 2.0 ± 1.2 µg/kg body weight (bw) respectively). During the PFAA-feeding period, PFBS concentrations above the detection limit (LOD of 0.1 µg/L) were detected only in milk samples (n = 11) of one cow with a mean concentration of 0.12 ± 0.02 µg/L and was not detected in milk during the PFAA-free feeding period. 0.005 and 0.003% of the ingested dose of PFBS were quantified in liver and kidney, respectively, of cows slaughtered directly after the PFAA-feeding period. The concentration of PFBS in liver after the feeding period was 0.3 ± 0.3 µg/kg, detectable in 2 of 3 cows (LOD = 0.2 µg/kg) and in kidney 1.0 ± 0.3 µg/kg. The concentration of PFBS in plasma was 1.8 ± 0.8 µg/L between the 1st and 28th days and PFBS was not detectable in plasma 4 days after the end of the PFAA-feeding period (day 32). For comparison, PFOA plasma concentrations were 8.6 ± 4.2 µg/L during the feeding of PFAA-contaminated feed and decreased rapidly after the end of the PFAA-feeding period, reaching the LOD (0.2 µg/L) on day 41 of the PFAA-free feeding period. Noticably, the authors concluded that overall, the kinetics of PFOA in cows were similar to those of PFBS and substantially differed from those of PFHxS and PFOS.

More information on mammalian toxicokinetics and elimination half-lives is provided in Sections 3.5.3.2 and 4.1.

3.5.2. Persistence and mobility in relation to bioaccumulation

The Peter Fisk Associates consultancy company evaluated the use of persistence and mobility of chemicals in the environment to fulfil the bioaccumulation criteria of the Stockholm Convention (Crookes and Fisk, 2018). They found that substances that have a low bioaccumulation potential could potentially reach similar levels in biota to substances that are known to bioaccumulate, provided that they are sufficiently persistent and mobile in the environment. For example, calculations in the study show that a substance with a half-life of 365 days and a BCF of 800 l/kg may reach comparable concentrations in a system as a substance with a half-life of 60 days and a BCF of 5000 l/kg, if time allows for steady-state to be reached (Figure 3 in reference).

In their report, Crookes and Fisk (2018) point out that substances with $\log K_{ow} < 5$ or $\log K_{oc} < 5$ and $\log K_{aw}$ in the range < -3 tend to be mobile in water. Furthermore, they found that the overall characteristic travel distance (CTD) for substances with $\log K_{aw}$ down to around -3 is governed mainly by transport via the atmosphere, but at lower $\log K_{aw}$ values the overall CTD is increasingly governed by transport via the water phase.

The Crookes and Fisk model was applied to PFBS, See Annex II. The calculations show that with an assumed half-life in water of 10 years and a reported BAF in crab of 110, the concentrations of PFBS in aquatic biota may be expected to exceed the biota concentrations for a persistent and bioaccumulative substance over time.

3.5.3. Field studies

In a study by Zhou et al. (2013) BAFs for short-chain PFASs in fish muscle from crucian carp (*Caracissius caracissius*) and fish sharpbelly (*Hemiculter leusisculus*), based on environmental measurements, were < 1 . The samples were collected from Tangxun lake which is located near a production base of the fluorochemical industry in Wuhan, China.

Naile et al. (2013) investigated the distribution and bioconcentration characteristics of PFASs in environmental samples collected from the west coast of Korea. During May of 2009, the concentrations of PFASs were determined in water ($n = 15$), sediment ($n = 12$), soil ($n = 13$), and biota ($n = 74$) from estuarine and coastal areas along the west coast of Korea. Water samples (1 L) were collected by dipping a clean, 1 L polypropylene (PP) bottle, which had been rinsed with methanol, just under the surface of the water. Samples of biota were collected by hand in coastal tidal pools and along the shore of inland bodies of water, and were transferred to and stored in clean PP bags. Duplicate samples and field blanks were collected daily, and were analyzed along with laboratory and procedural blanks. All samples of biota were pooled, homogenised, and freeze-dried. Compound-specific, bioaccumulation factors (BAFs) (reported as field-based bioconcentration factors (BCFs) by the authors) were calculated for aquatic organisms based on site-specific concentrations of PFASs in water samples. BAF for crab was reported as a mean for five different crab species: *Acanthogobius flavimanus*, *Sebastes schlegeli*, *Tridentiger obscurus*, *Hexagrammos otakii*, and *Mugil cephalus*.

Relatively high concentrations of PFBS were measured in crab, as compared to fish, gastropod and bivalve with a mean concentration of 0.25 ng/g ww in 44 samples. It was further found that in crab the greatest BAF values were measured for PFBS and PFOS, while in fish PFOS and PFDA had the greatest BCF values, and in gastropods and bivalves PFHxS. Whole body BAF as a mean for the five crab species was reported as: $\log BAF =$

2.04 ± 0.70 , corresponding to a BAF (crab¹) of 110. In comparison, the mean BAFs for PFHxS, PFOS and PFOA in crab were 575, 257 and 30, respectively. Mean BAFs were also reported for fish² (69), gastropod³ (107) and bivalve⁴ (12). Scatter plots of concentrations of PFASs in crab as a function of those in water, demonstrated a linear relationship. In general, the measured BAFs for PFASs were greater for fish and comparable for crab to those previously observed in Korea. The study would have benefitted from measuring of BAF for each individual crab species. However, BAF reported as a mean for five different crab species is also useful. The study is considered reliable with limitations, Klimisch score 2.

Similar results to the previous study were found by Hong et al. (2015) when they investigated the bioaccumulation characteristics of perfluoroalkyl acids (PFAAs) in coastal organisms from the west coast of South Korea. Twelve individual PFAAs in samples of water (n = 43) and biota (n = 59) were quantified by use of HPLC–MS/MS after solid phase extraction. Bioaccumulation of PFAAs in various organisms including fishes, bivalves, crabs, gastropods, shrimps, starfish, and polychaetes were determined. Samples of water and biota were collected from the same locations and according to the same method as in the previous study by Naile et al. (2013). The compositions of PFAAs accumulated in samples were slightly different among species. For example, PFOS was the predominant PFAA in fish and shrimp, while PFBS, PFPeA, and PFOA were dominant in bivalve, crab, and gastropod. The authors suggested that bioaccumulation of PFAAs in aquatic organisms is strongly dependent on the concentrations of PFAAs in water regardless of species. Field-based bioaccumulation factors (BAFs) of PFAAs in various aquatic organisms were calculated based on concentrations in water and biota (wet mass basis). For PFBS the following log BAFs were reported: 2.0 ± 0.46 (fish, n=2), 1.5 ± 0.53 (bivalve, n=8), 2.3 ± 0.57 (crab, n=13), 2.3 ± 0.79 (gastropod, n=14) and 1.6 ± 0.18 (shrimp, n=7). This corresponds to a BAF (crab) = 200, which is in the same range as the values reported by Naile et al. (2013). The study is considered a reliable, non-guideline study with some limitations in the documentation, Klimisch score 2.

Shi et al. (2018) investigated the processes governing uptake, distribution and elimination of PFASs in a study of the differential tissue distribution and bioaccumulation behavior of 25 PFASs in crucian carp from two field sites impacted by point sources. The highest median tissue-blood ratios for PFBS were observed in bile, gonad, heart, kidney and liver. However, when considering the relative body burdens, blood, gonads, and muscle together accounted for >90% of the amount in the organisms.

When comparing different PFASs and PFCAs the authors concluded that the trends in tissue blood ratios, relative body burdens and BAFs were most consistent with specific protein interactions as the governing mechanism for uptake and accumulation of PFASs although phospholipid partitioning may contribute to the accumulation of long-chain PFASs in specific tissues. Median tissue/blood ratios (TBRs) were consistently <1 for all PFASs and tissues except bile which displayed a distinct distribution pattern and enrichment of several perfluoroalkyl sulfonic acids.

¹ Mean concentrations including *Acanthogobius flavimanus*, *Sebastes schlegeli*, *Tridentiger obscurus*, *Hexagrammos otakii*, and *Mugil cephalus*.

² Mean concentrations including *Hemigrapsus sanguineus*, *Sesarma pictum*, *Hemigrapsus penicillatus*, *Helice tridens tridens*, and *Philyra pisum*

³ Mean concentrations including *Littorina brevicula*, *Monodonta labio*, *Umbonium thomasi*, *Glossaulax didyma*, and *Monodonta labio*

⁴ Mean concentrations including *Mytilus edulis*, *Macra veneriformis*, *Nuttallia olivacea*, and *Sinonovacula constricta*

Bioaccumulation of PFASs was quantified in eel (*Anguilla anguilla*) from several locations in the Netherlands (Kwadijk et al., 2010). Water and eels were collected at 21 different locations between May 2007 and August 2007. Fillets from 30 individual eels per location were randomly selected and homogenised prior to extraction and analysis. Historical eel tissue samples collected at three different locations from 1978 and onward for monitoring purposes, were also analysed for PFASs. Average BAF for PFBS in eel was 18 ± 4 (n=9). The BAF for PFBS was similar to that of L-PFOA (13) and PFOA (12), but lower than BAFs for the other measured PFASs.

In biota samples from Llobregat river in Catalonia, Spain, the highest BAF calculated for PFBS was 1736 (Campo et al., 2015). However, this BAF was calculated by the authors as highest fish concentration divided by highest water concentration (L/kg, fish not paired with water sample from sampling location). The biota sample size was low, and PFBS was detected in 2 of 14 water samples only. Three fish species were sampled from five different locations along the river, *Barbus graellsii*, *Cyprinus carpio* and *Micropterus salmoides*. PFBS was detected in 8/12 fish samples. → *Klimisch reliability 3 – not reliable, limitations in field-based estimations.*

3.5.3.1. Measured levels of PFBS in biota (other than remote arctic regions)

Measured levels of PFBS in biota in remote regions are reported in Section 3.4.3.3.

Marine biota

PFBS has been reported in several marine biota samples throughout the world. The largest concentration was found in shark liver (*Sphyrna tiburo* or *Rhizoprionodon terraenovae*) at a mean of 0.7 ng/g ww and max of 2.1 ng/g ww (Green et al., 2016). Van de Vijver et al. (2005) detected PFBS in the spleen of harbour seals from the Dutch Wadden Sea. Keller et al. (2012) detected PFBS in blood plasma from several species of turtles in the US, with a concentration in the herbivorous green turtle at many times higher than the concentrations in the other turtle species (Table 20). The concentration measured in green turtle plasma (*Chelonia mydas*), was at a mean of 0.09 and a max of 0.85 ng/g ww (Keller et al., 2012), however, the result was driven by one particular outlier. The green turtle is an endangered species according to International Union for Conservation of Nature (IUCN).

Lam et al. (2016) measured perfluoroalkane sulfonic acids (PFASs) and perfluoroalkyl carboxylic acids (PFCAs), in liver samples of Indo-Pacific humpback dolphins (*Sousa chinensis*) and finless porpoises (*Neophocaena phocaenoides*) from the South China Sea between 2002 and 2014. Levels of total perfluoroalkyl substances (PFASs) in samples ranged from 136–15,300 and 30.5–2,720 ng/g dw for dolphin and porpoise, respectively. PFBS concentrations in liver of dolphins and porpoises was from non-detect (<0.100 ng/g dw) to 12.3 ng/g dw and from non detect (<0.100 ng/g dw) to 1.76 ng/g dw, respectively. PFBS concentrations showed a significant increasing trend in the liver samples of dolphins from 2002 to 2014. The authors extrapolated, through regression analysis, that the levels of PFBS in dolphins were likely to increase 10-fold within 2024. Also, a significant increase in the ratio of PFBS to PFOS against time was found for the dolphin samples. No significant temporal trends of ΣPFASs appeared over the sampling period (Lam et al., 2016).

Table 20: Environmental concentrations of PFBS in marine biota from other regions than remote arctic areas (ng/g ww).

Location	Date	Mean +/- SD	Min – max (Median)	n (df) LOD	Remark	Reference
Baltic sea	2013-2014	<0.002		LOD = 0.002	Zooplankton	Gebbink et al., 2016a
Baltic sea	2013-2014	<0.01		LOD = 0.01	Herring whole fish	Gebbink et al., 2016a
Baltic sea	2013-2014	<0.06		LOD = 0.06	Sprat whole fish	Gebbink et al., 2016a
Baltic sea	2013-2014	<0.004		(100 %) LOD = 0.0004	Guillemot egg	Gebbink et al., 2016a
Black Sea	1997-1998	Nd		LOD = 1.4-3.2	Harbor porpoise organs	Van de Vijver et al., 2007
Wadden sea, Netherlands	2002	2.34 ± 0.68	1.74 – 3.28 (2.17)	4 LOD not available	Harbor seals (spleen)	Van de Vijver et al., 2005
Faroe Islands	1986-2013	<0.01		LOD = 0.01	Pilot whale muscle	Dassuncao et al., 2017
Norway	2015	<0.1	(0.10)	128 LOD = 0.1	Cod liver	Green et al., 2016
Spain	2009	ND		LOD = Not reported	Fish and shellfish (composite samples)	Domingo et al., 2012
Minnesota/ Mississippi R.	2007	ND		LOD = 0.01-1.89	Bluegill, Black crappie & Pumpkinseed fillet	Delinsky et al., 2010
USA, Georgia	2006	<0.3	0.3	LOD = Not reported	Shark soft tissue	Kumar et al., 2009
USA, Georgia	2006	0.7	<0.1 - 2.10	LOD = Not reported	Shark liver	Kumar et al., 2009
USA, Georgia	2006	0.4	<0.1 - 0.80	LOD = Not reported	Shark muscle	Kumar et al., 2009
USA	2007	ND		7 LOD = 0.02	Leatherback turtle plasma	Keller et al., 2012

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min – max (Median)	n (df) LOD	Remark	Reference
USA	2007	0.02 ± 0.04	<0.01 - 0.126 (0.0019)	15 (20 %) LOD = 0.02	Loggerhead turtle plasma	Keller et al., 2012
USA	2007	0.02 ± 0.02	<0.01 - 0.0558 (0.0177)	10 (30 %) LOD = 0.02	Kemp's ridley turtle plasma	Keller et al., 2012
USA	2007	0.04 ± 0.06	<0.02 - 0.139 (0.008)	5 (40 %) LOD = 0.02	Hawksbill turtle plasma	Keller et al., 2012
USA	2007	0.09 ± 0.27	<0.02 - 0.846 (0,0000547)	10 (20 %) LOD = 0.02	Green turtle plasma	Keller et al., 2012
South China Sea	2002 - 2014		< 0.100 – 12.3 (ng/g dw)	17 (71 %) LOD = 0.1	Indo-Pacific humpback dolphins' liver	Lam et al., 2016
South China Sea	2002 - 2014		<0.100 – 1.76 (ng/g dw)	50 (8%) LOD = 0.1	Finless porpoises' liver	Lam et al., 2016
Røst, Norway	1983	<0.009		5 LOD = 0.009	Herring Gull egg	Verreault et al., 2007
Røst, Norway	1993	<0.009		5 LOD = 0.009	Herring Gull egg	Verreault et al., 2007
Røst, Norway	2003	<0.009		5 LOD = 0.009	Herring Gull egg	Verreault et al., 2007
Røst, Norway	2017	<0.06		5 LOD = Not reported	European shag eggs (<i>Phalacrocorax aristotelis</i>)	Schlabach et al., 2018
Hornøya	1983	<0.009		5 LOD = 0.009	Herring Gull egg (<i>Larus argentatus</i>)	Verreault et al., 2007
Hornøya	1993	<0.009		5	Herring Gull egg	Verreault et al., 2007

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min – max (Median)	n (df) LOD	Remark	Reference
				LOD = 0.009		
Hornøya	2003	<0.009		5 LOD = 0.009	Herring Gull egg	Verreault et al., 2007
Tromsøya, Norway	2017	0.03 ± 0.03	0.01 - 0.07 (0.02)	5 (80 %) LOD = Not reported	Common gull egg (<i>Larus canus</i>)	Schlabach et al., 2018

Freshwater biota near point sources

Measurements in freshwater biota were primarily from Taihu Lake, China (Fang et al., 2014) and a national survey of river basins in Vietnam (Lam et al., 2017), see Table 21. The highest concentration in Taihu Lake was associated with zooplankton (1.68 ng/g ww) and in Vietnam with Tilapia liver (1.34 ng/g ww). In samples from Taihu Lake, PFBS was not detected in phytoplankton, invertebrates (mussels) and in 4/9 sampled fish species (muscle tissue), Fang et al. (2014). PFBS was among the PFASs detected in all aquatic biota species sampled (shrimp, fish and mallard (*Anas platyrhynchos*)) from a lake in China (Zhou et al., 2014).

Table 21: Environmental concentrations of PFBS in freshwater biota (ng/g ww).

Location	Date	Mean +/- SD	Min – max (median)	n LOD	Remark	Reference
China, Taihu Lake	2012	1.68 ± 0.04		3 LOD = 0.054	Zooplankton	Fang et al., 2014
China, Taihu Lake	2012	0.10 ± 0.00		3 LOD = 0.054	Crucian carp muscle (<i>Cyprinus carp</i>)	Fang et al., 2014
China, Taihu Lake	2012	0.08 ± 0.00		3 LOD = 0.054	Lake saury muscle (<i>Cololabis adocetus</i>)	Fang et al., 2014
China, Taihu Lake	2012	0.54 ± 0.01		20 LOD = 0.054	Mongolian culter muscle (<i>Culter mongolicus</i>)	Fang et al., 2014

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Date	Mean +/- SD	Min – max (median)	n LOD	Remark	Reference
China, Taihu Lake	2012	0.24 ± 0.01		10 LOD = 0.054	Mudfish, muscle (<i>Oriental weatherfish</i>)	Fang et al., 2014
China, Taihu Lake	2012	1.20 ± 0.03		3 LOD = 0.054	Whitebait, muscle (<i>Reganisalanx brachyrostralis</i>)	Fang et al., 2014
Vietnam	2013-2015	0.11 ± 0.54	1.34	149 LOD = 0.03-0.48	Fish and shellfish tissue (average of 5 fish species, 2 crustaceans (paddle crab, giant prawn), 1 gastropod (golden apple snail), 1 bivalve (golden freshwater clam))	Lam et al., 2017

Other biota samples

PFBS in the interval 0.07 - 0.33 ng/g ww were detected in the liver of mink (*Neovison vison*) from Norway (Schlabach et al., 2018). In a monitoring study on levels of PFAS in the terrestrial environment in Norway (Herzke et al., 2016) PFBS was detected in 5 out of 6 earthworms (*Lumbricidae*), levels ranged from 0.13 to 1.10 ng/g ww, see Table 22.

Table 22: Environmental concentrations in other biota samples (ng/g ww)

Location	Date	Mean +/- SD	Min – max. (median)	n LOD	Remark	Reference
Troms, Norway	2013-2014	0.19 ± 0.11	0.07 - 0.33 (0.19)	10 (100 %) LOD = Not reported	Mink liver (<i>Neovison vison</i>)	Schlabach et al., 2018
Norway, Oslo	2015	0.75	0.13 - 1.10 (0.94)	5/6 (83%) LOD = Not reported	Earthworms (<i>Lumbricidae</i>)	Herzke et al., 2016

3.5.3.2. Measured half-lives and levels in humans

Toxicokinetic data show that PFBS is found in many different tissues as it binds to the serum protein, albumin, and several other tissue proteins. The major route of elimination is through urine, although PFBS is also found in faeces and has been detected in milk of dairy cows.

The half-lives of PFBS in mice and rats are only a few hours, with a shorter half-life in female than male rodents. In monkeys, a half-life of up to a few days has been determined. Half-lives in humans have been measured in only 6 workers. The one female employee had the longest serum elimination half-life of 45.7 days, while the mean serum half-life of elimination in men was 24.1 days. A relatively large individual half-life variability for different other PFAAs was also measured using data from up to 66 individuals in a study by Zhang et al. (2013b). The half-life in pigs (43 days) seems to be similar to humans. See Section 4.1 and Table 27 for more details.

Comparing the measured half-lives of PFBS in rodents and humans shows that the half-life in humans is considerably longer than the half-life in rodents, although PFBS has a relatively short half-life in the human body compared to e.g. PFOA (2 – 4 years).

PFBS has been detected in human blood, lung, bone, kidney, urine and hair. Concentrations of PFBS in human blood have been measured in a number of studies from Europe, US and Asia. The PFBS level is often below the quantification limit, with a concentration frequently below 0.5 ng/mL. However, in a few studies concentrations up to 5 ng/mL were observed. In persons with known elevated exposure to PFBS, e.g. from drinking water or in occupational settings, also higher levels have been measured. Furthermore, PFBS has been detected in human cord blood with a mean concentration of 0.053 ng/mL, while the highest concentration was 0.46 ng/mL, see Section 4.3 and Table 28 for more information.

3.5.4. Summary of bioaccumulation

Due to being both hydrophobic and lipophobic, PFAAs do not bioaccumulate via distribution to fatty tissues like the legacy persistent pollutants (POPs) do. Due to their high water solubility, short-chain PFAAs are expected to be quickly excreted via gill permeation in water breathing animals. This reduces the bioaccumulation potential in such species.

Measured BCF values in fish are in the range 0.36 to 27.5. The highest fish BAF measured (field study) is 1736, however, the study had limitations; not reliable). In another field study a mean BAF for fish of 69 was reported while for crab a mean BAF of 110 was measured.

It has been estimated that substances with a low bioaccumulation potential could potentially reach similar levels in biota as substances that are known to bioaccumulate, provided that they are sufficiently persistent and mobile in the environment and when considering a longer time-scale.

PFBS has been detected in blood and different organs of animals and humans and is therefore bioavailable. PFBS has been shown to be able to bind to different proteins (transport proteins in blood, receptor proteins, enzymes and organic anion transporters), although the binding potential is lower compared to the longer chain PFAAs. According to Annex XIII also other information on the bioaccumulation potential can be considered in the bioaccumulation assessment, among others data from human body fluids or tissues, toxicokinetic data, results from bioaccumulation studies in terrestrial species and detection of elevated levels in biota, in particular in endangered species or in vulnerable populations.

PFBS has been detected in human blood, lung, bone, kidney, urine and hair. It has also been detected in human cord blood. Hence, bioavailability has been demonstrated. A serum elimination half-life of around one month (up to 46 days) has been measured in humans, which is considerably longer than the half-lives measured for rodents. The average measured half-life in pigs, 43 days, is similar to that in humans. The individual variability in the study with pigs was large and the higher half-life values for PFBS were overlapping with the lower values for PFOA. It should be remembered that data on human half-life is scarce and that data measuring half-lives of other PFAAs in humans has shown a great individual variability. No cut-off values for human elimination half-lives for fulfilling the B or vB criteria have been defined. Based on the lower half-life reported for PFBS in human blood compared to the half-lives of PFOA, 2 – 4 years (B), and PFHxS, 7 - 8 years (vB), it is concluded that PFBS is less bioaccumulative. It should be recognised that even though blood is used as a proxy for the entire body, the half-life in human blood is not necessarily identical with the whole body elimination half-life in humans (ECHA, 2017). BMF values > 1 (1.2 (whole pig), 0.8 (meat), 6.4 (liver), 14 (blood plasma), 2.2 (kidney) and 0.9 (fat)) have been reported in pigs.

PFBS has been detected in different species of wild life including green turtles and polar bears, both are endangered or vulnerable populations. PFBS has also been detected in other marine mammals, like dolphins and whales. In dolphins PFBS concentrations showed a significant increasing trend in the liver samples from 2002 to 2014. Also, a significant increase in the ratio of PFBS to PFOS against time was found for the dolphin samples. In the whale it was shown that PFBS transfers from mother to foetus. Additionally, PFBS was shown to transfer from mother to the eggs of birds. The findings of PFBS in sensitive life stages and in endangered species are of special concern.

3.6. Protein binding

PFAAs bind to proteins in the blood and tissues, and protein binding affects tissue distribution of PFAAs (Ng and Hungerbühler, 2014; Kerstner-Wood et al., 2003; Liu et al., 2017; Zhang et al., 2013a; Zhang et al., 2014). In blood, PFBS has been found to bind mainly to serum albumin, whereas PFOS, PFOA, and PFHxS have the potential to bind also to other human serum binding proteins, including plasma gamma-globulin, alpha-globulin, alpha-2-macroglobulin, transferrin, and beta-lipoproteins (Kerstner-Wood et al., 2003). The binding affinity to serum albumin seems to decrease in the order PFHxS > PFOS > PFBS (Bischel et al., 2011; Ng and Hungerbühler, 2014). A study by Liu et al. (2017) supports that the binding affinity to human serum albumin is lower for PFBS than for PFHxS and PFOS. PFBS binding did not cause apparent conformational changes to human serum albumin, but it induced functional changes as suggested by a decrease in esterase like activity of albumin at high concentrations (IC₅₀ 0.13 mM).

Perfluoroalkyl acids (PFAAs) are known to accumulate in the liver and induce hepatotoxicity in experimental animals. They resemble fatty acids in structure and concerns have been raised that PFAAs may disrupt fatty acid binding to transport proteins or receptors. Binding of PFAAs to L-FABP (liver-fatty acid binding protein) has been shown *in vitro* (Luebker et al., 2002; Zhang et al., 2013a). This binding may explain the high concentration in liver tissues or in kidneys where L-FABP is also expressed. Binding affinity of perfluoroalkyl acids to human L-FABP was reported to increase significantly with chain length from C4 to C11 and decrease with chain length above 11. PFBS was shown to bind human L-FABP, with the lowest affinity of the three tested PFASs: PFBS, PFHxS and PFOS (Zhang et al., 2013a). No binding was detected for the two fluorotelomer alcohols (6:2 FTOH and 8:2 FTOH).

Moreover, PFAAs may affect lipid metabolism through activation of peroxisome proliferator-activated receptors (PPARs). COS-1 cells were transfected with mouse or human PPAR α receptor-luciferase reporter plasmid to investigate the effects of different

PFAAs on PPAR α activation. PFNA (C20max mouse 5 μ M; human 11 μ M) and PFOA were predicted to have the highest PPAR α activating potencies of the tested PFAAs whereas PFBS had lower predicted PPAR α activation potencies (PFBS C20max mouse, 317 μ M; human, 206 μ M) (Wolf et al., 2008). As shown for PPAR α , PFAAs also bind to the human PPAR γ (Zhang et al., 2014). Similar to PPAR α , the binding affinity for the different PFASs to human PPAR γ *in vitro* increased with chain length from C4 to C11, and then decreased slightly with carbon chain length above 11 (Zhang et al., 2014). PFAS containing sulfonic acid groups bound more strongly than their carboxylic acid counterparts. Binding of PFBS to the ligand-binding domain of PPAR γ was not detected under the conditions used in this *in vitro* binding affinity assay. However, in the activity assay of the receptor, a small, but significant increase in activity was detected, albeit at higher concentrations of PFBS (150 μ M) compared to PFOS (10 μ M) (Zhang et al., 2014).

A recent publication by Rosenmai et al. (2018) found that cellular uptake of PFASs *in vitro* in HepG2-cells was in general low (below 1%). The sulfonates tested had lower uptake than the acids, although sufficient to activate PPAR α . Uptake varied also with chain length. For the sulfonates, the shorter chain PFBS showed higher intracellular uptake than PFOS (in contrast to the acids). PFBS showed also higher potency for PPAR α -activation at the lowest intracellular concentration. These data indicate that the potency of the short chain PFBS for activating PPAR α may be comparable to, or even higher than, its' longer counterparts (PFHxS and PFOS) (Rosenmai et al., 2018).

The capacity of several PFASs, including PFBS, to compete with T4 for binding to the human thyroid hormone transport protein transthyretin (TTR) was examined by Weiss et al., 2009. The binding potency decreased in the order: PFHxS > PFOS/PFOA > PFHpA (perfluoroheptanoic acid) > PFOSi (sodium perfluorooctane sulfinat) > PFNA (perfluorononanoic acid), with TTR binding potencies 12.5–50 times lower than the natural ligand T4. PFBS was determined to have more than 100 times lower potency than T4 (IC₅₀ of 19,460 nM that is approximately 20 times lower than the IC₅₀ determined for PFOS). Similar values for TTR binding potency for PFBS were found in a study by Ren et al. (2016).

Occurrence of perfluoroalkyl acids in animal protein feeds

In a recently published study by Li et al. (2019) the most prevalent animal protein supplement feeds (APF), which are blood meal, meat meal, feather meal, soybean meal and distillers dried grains with solubles (DDGS), were analysed for 16 PFAAs. The animal-derived APFs possessed higher Σ PFAAs with a mean of 10.9 ng/g dw compared to plant-derived APFs, mean 0.75 ng/g dw. The short chain PFAAs (PFBA, PFBS and PFHxS) were primarily found in blood meal, meat meal, soybean meal and DDGS, while the long-chain counterparts (PFOS) dominated in feather meal. In animal derived supplement feeds PFBS contributed with 17%, 13.3%, 9.2% of the total PFAAs in blood meal, feather meal, meat meal, respectively. In plant derived supplement feeds PFBS contributed with 30% in DDGS and 13% in soybean meal. The wide occurrence of short-chain PFAAs (PFBA, PFBS) in fish meal were found for the first time by Li et al. (2019b). In fish meal, which is the most important animal-derived feed in global husbandry, PFAAs were analysed and the average for Σ 16 PFAAs was 18.2 ng/g, (12% moisture). Short-chain PFAAs (PFBA, PFBS and PFHxS) were frequently detected and jointly contributed 15.7% of Σ 10 PFAAs in fish meal.

Increasing PFBS concentrations in the environment, its ability for protein binding and enrichment in plants results in increasing occurrence of PFBS in animal and plant derived protein supplement feeds. Since these supplemental feeds might be the main feed source for raised animals, human exposure through consumption of raised animals together with

PFBS-contaminated drinking water and plant-derived food, might lead to long-term continuous human exposure.

In summary, PFBS has been shown to bind to serum albumin and several other proteins, e.g. peroxisome proliferator-activated receptors (PPARs), liver-fatty acid binding protein (L-FABP) and thyroid hormone transport protein transthyretin (TTR). The ability to bind to proteins affects tissue distribution/accumulation and may be of toxicological significance. In general, the studies show a PFAA-protein binding affinity that depends on chain length. A carbon chain-length of 4 tends to have a lower binding affinity compared with a carbon chain length of 6-8. PFAAs containing the sulfonic acid group tend to bind more strongly to proteins than the corresponding carboxylic acids. Even though the binding affinity of PFBS to different transport proteins and receptors is lower than for its longer counterparts in vitro, the intracellular uptake and activation potencies may be different as indicated by Rosenmai et al. (2018).

The occurrence of PFAAs in animal protein supplement feeds (APFs) has been reported. In animal-derived supplement feeds PFBS contributed with 17%, 13.3% and 9.2% of the total PFAAs in blood meal, feather meal and meat meal, respectively. In plant derived supplement feeds PFBS contributed with 30% in distillers dried grains with solubles and 13% in soybean meal of the total PFAAs.

3.7. Enrichment in plants

Several studies investigated the uptake of PFBS and other PFASs from the surrounding environment into plants, Table 23. The studies were conducted under strictly controlled laboratory conditions, under semi-natural conditions, as well as field studies under environmental conditions, and monitoring data are compiled.

Krippner et al. (2015) examined PFAS accumulation in straw and grains of maize plants by looking at ten C4 to C10 substances containing the carboxylic (PFCAs) or sulfonic (PFSAs) functional groups, all individually spiked at 0.25 and 1 mg/kg soil. They confirmed the direct correlation between soil and plant tissue concentrations: the higher the soil PFAS content, the more the plants accumulate. They suggest that shorter-chain PFAS are more water soluble and therefore more mobile in soil and preferentially taken up by plants than longer-chain PFAS. Moreover, they observed lower translocation of long-chain PFAS within the plant than for short-chain PFAS, where long-chain PFAS largely remains in the roots and storage organs and short-chain PFAS transfers and accumulates in the above-ground plant parts. Müller et al. (2016) observed that tissue concentrations of PFAAs in hydroponic model plant system (*Arabidopsis thaliana*) during depuration rapidly declined in roots but remained constant in shoots, demonstrating irreversibility of the translocation process. Differences in accumulation levels of compounds in plants depends on the chain length, with the longer-chained compounds generally having lower accumulation rates, and on the associated functional groups, with PFSAs generally having lower accumulation rates than PFCAs (Scher et al., 2018).

Uptake mechanism for PFAS were studied by Wen et al. (2013) for PFOS and PFOA in corn. They suggested that PFOA and PFOS may have different uptake mechanisms in maize. For PFOA a potential active uptake and entry by anion channels for PFOA was suggested while for PFOS a passive entry by aquaporins (water channels) or anion channels (different than the ones used by PFOA) were suggested. Blaine et al. (2013) concluded that passive transport may be the primary mechanism for uptake and translocation due to the linearity of the plant uptake response to soil concentration of PFAS.

Ghisi et al. (2019) reviewed the accumulation of PFASs in agricultural plants. They emphasised two important sources to PFAS contamination of agricultural land: i) field

irrigation with contaminated surface or ground waters, and ii) the use of contaminated sewage sludge as a soil conditioner. They pointed out that from agricultural plants PFASs can transfer to humans through the food chain. The authors concluded that PFASs are absorbed by plants to different extents according to their concentrations, chain lengths, functional groups, plant species, variety and organs, growth media (hydroponics vs. soil), and soil and biosolid characteristics. In particular, the abundance and characteristics of soil organic matter are considered among the most important factors.

The C4–C6 compounds appear to accumulate preferably in leaves and fruits, whereas the compounds with longer chain lengths tend to be more concentrated in roots. The authors reported studies with PFBS concentrations up to 1843 µg/kg dw in maize straw, grown on soil with spiking concentration of 1 mg/kg PFAA mixture, and 205 µg/kg dw in lettuce leaves, grown on soil with industrially impacted biosolids.

It was pointed out by Ghisi et al. (2019) that agricultural soils close to airports, fire-fighting training locations, industrial sites and landfills are particularly at risk of PFAS pollution. However, the use of biosolids on agricultural soil must also be considered a potential source of agricultural plant contamination. Growing vegetables in hydroponics should also be discouraged in polluted areas, as it allows pollutants to interact directly with plants in the absence of soil that has the ability to adsorb pollutants.

In a study of whether PFASs in water used for yard and garden irrigation resulted in elevated concentrations of PFASs in soil and home-grown produce at homes with a history of PFAS-contaminated drinking water, it was found that short-chain PFASs have the highest potential to translocate to and bioaccumulate in edible plants (Scher et al., 2018). The findings were assumed to be globally relevant, as short-chain PFASs serve as commercial substitutes for longer-chain compounds and are increasingly detected in water due to their relatively high solubility and mobility. However, PFBS was not found in the groundwater in this particular area.

Blaine et al. (2013) investigated the accumulation of PFBS in lettuce and tomato grown on biosolid-amended soils and reported bioaccumulation factors (BAF) in lettuce up to 14.5 and PFBS concentrations in lettuce up to 205 µg/kg dw. Furthermore, a BAF of 0.42 was reported in the leaves and fruits from greenhouse tomatoes. PFBS enrichment was also demonstrated in other typical edible crop products, including radish, celery, tomato and peas with shoot concentration factors (SCF) of 3.4, 2.2, 3.7, 4.1, respectively (Blaine et al., 2013; 2014; 2014a) and SCF of up to 7.2 for thale cress (Müller et al., 2016). Felizeter et al. (2012; 2014) examined the uptake of PFASs into hydroponically grown lettuce, tomato, cabbage and zucchini. In the leaves the highest concentration factors were found, with LCF of 11-42, compared to other above-ground parts of the plants. In contrast, concentration factors for PFOS are much lower with 1.1 (Felizeter et al., 2012). A calculated mass distribution of different PFAAs showed that PFBS accumulates mainly in leaves (65%) and roots (21%), whereas PFOS accumulates mainly in roots (68-71%), Felizeter et al. (2014). The high potential for translocation in plants can be attributed to PFBS having a high solubility in water.

Krippner et al. (2014) demonstrated that shorter-chain PFASs are transferred predominantly and at higher concentrations to the shoot in maize plants with shoot/root ratio of 5.5 for PFBS. In contrast, long-chain PFAS accumulated at higher concentrations in the roots of maize plants with a shoot/root ratio of < 1 for PFOS. Krippner et al. (2015) calculated soil to plant transfer factors (TF) for PFBS in maize straw of 1.8-3.8, demonstrating uptake and storage of PFBS preferably in the leaves. However, Navarro et al. (2017) reported TF roots of 5.00 and TF leaves 4.00 for maize.

In wheat grown on biosolid amended soils, PFBS was not detected in grains but in roots and in shoots (Wen et al., 2014). Furthermore, Lan et al. (2018) reported that PFBS accumulates in wheat with lower BAF in root (2-5.6) than in shoot (9.1-10.5). The grass

plant *Bromus diandrus* was exposed to 0.5 and 1 µg/mL of PFAA mixture in nutrient solution for 20 days, and a mean TF in shoot of 3.2 for PFBS was determined (García-Valcárcel et al., 2014). In a mesocosm study with two aquatic macrophytes whole plant BCFs of 29.5 and 18.8 were reported for PFBS for one submerged and one free-floating aquatic macrophyte, respectively (Pi et al., 2017).

The leaching of PFAAs from field contaminated soil and their uptake into biota was investigated by Bräunig et al. (2019). Soil from firefighting training sites at two airports with historical usage of aqueous film-forming foam (AFFF) was sampled and a greenhouse study was carried out to investigate the uptake of PFAAs from soils into earthworms and wheat grass. The uptake of 12 different PFAAs was investigated. PFBS was found to have the third highest accumulation factor in wheat grass, after PFBA and PFPeA (but higher than PFHxS and PFOS). The authors concluded that the high accumulation of shorter chain PFAAs in grass has implications for their movement into the terrestrial food chain.

Navarro et al. (2017) further compared the uptake and distribution of PFASs in plant tissues with polybrominated diphenyl ethers (PBDEs). TF values higher for PFASs than for PBDEs were found in all crop plants: from 2 to 9-fold in spinach, 2 to 34-fold in tomato and 11 to 309-fold in corn. For PBDEs TFs for root and leaf were found in the intervals 0.32-0.50 and 0.01-0.03, respectively, as compared to 5.00 and 4.00 for PFBS. In another study Hurtado et al. (2016) determined the concentration of eight chemicals in leaf of lettuce and in the artificial soil (perlite and sand), in which they were grown in a greenhouse experiment. The tested substances were mostly neutral, except for ibuprofen and propranolol. Bisphenol-A, caffeine, propranolol and tonalide had calculated BAF values that ranged between 2.3 and 9.1 g_{dwt}/g_{dwt} . Ibuprofen and triclosan had a calculated maximum BAF value of 1.3 g_{dwt}/g_{dwt} , while sulfamethazine was not detected in plants. Carbamazepine had high BAF values ranging from 17 to 247 g_{dwt}/g_{dwt} , PFBS has higher or at least comparable BAFs as the substances bisphenol-A, caffeine, propranolol and tonalide, and higher BAFs than ibuprofen, triclosan and sulfamethazine.

Data on the uptake and concentrations of PFBS in plants is compiled in Table 23.

Table 23: Uptake and concentration of PFBS in plants

Method	Plant	Results	PFBS concentration (µg/kg dw)	Comments	Reference
Field and greenhouse study with soil and industrial impacted biosolids and municipal impacted biosolids PFAA mixture	Lettuce (<i>Lactuca sativa</i>) Tomato (<i>Lycopersicon lycopersicum</i>)	Greenhouse, Municipal soil L. sativa: BAF: 14.5 Field trial L. sativa BAF: 2.02 L. lycopersicum industrially impacted soil BAF: 0.42 µg/kg dw	Lettuce leaves 205 Tomato fruits 19.4	Greenhouse and field	Blaine et al., 2013
Greenhouse, Plants grown in soils amended with industrial biosolids, loading 49 ng	Radish (<i>Raphanus sativus</i>), celery (<i>Apium graveolens var. dulce</i>), tomato	Shoot concentration factor (SCF) Radish= 3.4 Celery=2.2 Tomato= 3.7 Pea= 4.1	Celery 107 Pea fruits 16 Radish root	Greenhouse	Blaine et al., 2014

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Method	Plant	Results	PFBS concentration (µg/kg dw)	Comments	Reference
PFBS/g	<i>(Lycopersicon lycopersicum)</i> , and sugar snap pea (<i>Pisum sativum</i> var. <i>macrocarpon</i>)		62		
Greenhouse hydroponic system (PFAA-spiked nutrient solution with nominal concentration of 10 ng/L to 10 µg/L)	Lettuce (<i>Lactuca sativa</i>)	Root Conc. Factors (RCF)= 0.002-0.008 Foliage Conc. Factors (RCF)= 0.002-0.008		Greenhouse	Felizeter et al., 2012
Greenhouse – hydroponic system (PFAA-spiked nutrient solution with nominal concentration of 10 ng/L to 10 µg/L)	Tomato (<i>Solanum Lycopersicum</i> var. <i>moneymaker</i>), Cabbage (<i>Brassica oleracea</i> convar. <i>capitata</i> var. <i>alba</i>) Zucchini (<i>Cucurbita pepo</i> var. <i>Black Beauty</i>)	Root concentration factor (RCF): <i>cabbage</i> : PFBS > 10 <i>zucchini</i> : PFBS > 10 <i>tomato</i> : PFBS > 10 - Stem concentration factor (SCF): <i>cabbage</i> : PFBS≈ 0.3 <i>zucchini</i> : PFBS≈ 1.5 <i>tomato</i> : PFBS≈ 2.1 -Leaf concentration factor (LCF) <i>cabbage</i> : PFBS≈ 11 <i>zucchini</i> : PFBS≈ 18 <i>tomato</i> : PFBS≈ 42 Edible part concentration factor (ECF) <i>cabbage</i> : PFBS≈ 0.8 <i>zucchini</i> : PFBS≈ 0.6 <i>tomato</i> : PFBS≈ 0.4 - Edible part/leaf transfer factor:		Greenhouse	Felizeter et al., 2014

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Method	Plant	Results	PFBS concentration (µg/kg dw)	Comments	Reference
		All factors were < 1, which indicates that leafy crops with open leaves (spinach or some lettuce) accumulate higher amounts in the edible part than fruit-bearing crops. Leafy crops pose a higher risk for human exposure			
Hydroponically grown grass plants. With 0.5 and 1 µg/mL PFAA mixture	Grass (<i>Bromus diandrus</i>)	TF shoot= 3.3 (0.5 µg/mL) TF shoot= 3.1 (1 µg/mL)		Lab	García-Valcárcel et al., 2014
Corn grown in pots with nutrient containing 0.1 mg/L PFBS at pH 5,6,7. Plants harvested after 5 d, shoots and roots analysed	Corn (<i>Zea mays</i>)	Shoot/root ratio= 5.5		Lab	Krippner et al., 2014
Pot experiment (soil was spiked with an aqueous solution of 0.25 mg individual PFAA/kg soil and 1 mg individual PFAA/kg soil) After 128 days straw and kernels were harvested PFAA mixture, ten compounds	Corn (<i>Zea mays</i>)	Transfer factor soil/ straw (TF straw): PFBS= 3.85 (0.25 mg/kg treatment) PFBS=1.84 (1.0 mg/kg treatment)	Treatment 1 mg/kg Maize straw:1843 kernels:5.4	Lab test All concentrations are measured and based on dry weight.	Krippner et al., 2015
Pot	Wheat	BAF root:		Greenhouse	Lan et al.,

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Method	Plant	Results	PFBS concentration (µg/kg dw)	Comments	Reference
experiment soil spiked with solution of 0.2 and 2 mg/kg PFAA. Harvested after 4 weeks.	<i>(Triticum aestivum L.)</i>	PFBS= 2.0 (0.2 mg/kg) PFBS= 5.6 (2 mg/kg) BAF shoot PFBS= 9.1 (0.2 mg/kg) PFBS= 10.5 (2 mg/kg)			2018
Hydroponic system, plants exposed to 2 µg/L in growth media for 10 or 14 days.	Thale cress (<i>Arabidopsis thaliana</i>)	Root Conc. Factors (RCF)12.6 Shoot Conc. Factors (SCF)= 7.2		Lab test	Müller et al., 2016
Field soil amended with biosolids	Corn (<i>Zea mays</i>)	Transfer factor (TF root): PFBS= 5.00 (0.2 mg/kg) (TF leaves): PFBS= 4.00 (0.2 mg/kg)	Maize: roots 150 leaves 120	Field test	Navarro et al., 2017
Semi-static mesocosm study. Uptake phase 15 days. Water concentration (nominal and measured): 20 µg/L	Submerged and free-floating aquatic macrophytes (<i>E. horemanii</i> / <i>E. crassipes</i>)	Leaf BCF _{ss} : 33.0 / 21.2 Root BCF _{ss} : 12.9 / 14.9 Whole-plant BCF _{ss} : 29.5 / 18.8			Pi et al., 2017
Field soil amended with biosolid	Wheat (<i>triticum aestivum</i>)		Wheat roots 22- 60 Wheat shoots 22 Grains, husks < MDL	Field test	Wen et al., 2014

MDL: method- detection limit

Data on the uptake of PFBS in terrestrial plants near hotspot pollution sites are presented in Table 24.

Table 24: Uptake factors and concentrations of PFBS in terrestrial plants near hot spots

Monitoring data				
Method	Plant	Result	Location	Reference
Grass silage and hay cultivated on PFAA contaminated farmland.	Grass	Mean: Grass silage: 68.4 ± 23.1 µg/kg dw	Europe, Germany	Kowalczyk et al., 2013

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

		Hay: 933± 224 µg/kg dw		
Leaves and bark, soil, water etc collected in 2012 from an area surrounding a large fluorochemical industry park (Jiangsu-Hi) in China. Leaves not washed before analysis.	Tree camphor	Leaf/ accumulation factors LSAF leaves= 2.0 LSAF bark= 0.33 Mean values used to calculate LSAF factors	China	Shan et al., 2014
Leaf and soil samples obtained from a campus of Dalian University, China.	Pine needels (<i>Pinus massoniana</i>) Cypress (<i>Platycladus orientalis</i>) Ginkgo (<i>Ginkgo biloba</i>) Popular, willow, sophora, plane-tree	Leaf/ accumulation factors LSAF leaves=17 pine LSAF leaves=0.59 cypress LSAF leaves=0.12 ginkgo LSAF leaves=0.38 popular, willow, sophora and plane tree.	China	Zhang et al., 2015
Vegetables from five gardens (0.2-1.5 km away from fluorochemical industrial park	Tomato (<i>Solanum lycopersicum</i>), Cucumber (<i>Cucumis sativus</i>), Eggplant (<i>Solanum melongena</i>), Pepper (<i>Capsicum annum</i>), Chinese cabbage (<i>Brassica rapa ssp. Pekinensis</i>),	Vegetables from home gardens (µg/kg fw) 0.2 km 11 0.4 km 11 0.6 km 5.7 1 km <0.2 1.5 km <0.2	China, Fuxin	Bao et al., 2019

3.7.1. Measured levels of PFBS in plants

PFBS has shown a high propensity to transfer and enrich in plants, due to its high mobility in plant xylem (Arp and Slinde, 2018).

In Europe, Kowalczyk et al. (2013) investigated PFBS, PFHxS, PFOS, and PFOA in contaminated feed and the absorption, distribution, and milk secretion of the substances in dairy cows, see Table 24. Hay and grass silage were obtained from an area with a pollution incident in Germany in 2006. The mean content of PFBS in grass silage was $68.4 \pm 23.1 \mu\text{g}/\text{kg dw}$ and in hay $993.6 \pm 224.4 \mu\text{g}/\text{kg dm}$, showing that PFBS is taken up in plants used in agriculture.

In Asia, Shan et al. (2014) collected i.a. leaves and bark of camphor trees and soil from the area surrounding a large fluorochemical industry park (Jiangsu-Hi) in China. The authors compared with previous studies on PFOA and hypothesised that PFASs predominantly were transported from the industry sites via atmospheric transportation on particulate matter to the nearby environment and deposited on trees and plants. Hence, tree samples were used to monitor airborne PFASs in the environment. In leaves and bark PFBS was detected with a mean of $0.42 \mu\text{g}/\text{kg dw}$ and $0.07 \mu\text{g}/\text{kg dw}$, respectively. However, compared with other studies in this dossier, it may seem that direct uptake in plants and trees from soil via water may have been somewhat underestimated in this study. However, airborne transport of PFASs on particulate matter may be a significant means of local transportation of PFASs from industrial emission sites to the nearby environment. In any case, the sum of PFASs in the tree leaves decreased significantly as the distance from the facilities increased. For comparison, the leaf/soil and bark/soil accumulation factors were calculated from the mean concentration of PFBS in soil of $0.21 \mu\text{g}/\text{kg dw}$, 2 and 0.3 respectively, see Table 24.

In another study several PFAAs were monitored in the leaves of coniferous and deciduous forests grown in the urban areas of Dalian, China (Zhang et al., 2015). The results showed that coniferous tree leaves take up more PFAAs than broad-leaved tree leaves. For PFBS calculated leaf/soil accumulation factors (LSAF) were highest for the pine needles of cypress 17, followed by 0.6 for cypress, 0.12 for ginkgo and 0.04 for poplar, willow, sophora and plane tree, see Table 24.

A recently published study investigated whether PFAS-contaminated groundwater around a fluorochemical industrial park (FIP) in Fuxin, China, could be introduced into home-produced vegetables in local residences via the application of groundwater for the irrigation or feeding purposes (Bao et al., 2019), see Table 24. Samples of garden soil and vegetables (tomato, cucumber, eggplant, pepper and chinese cabbage) were obtained from five home gardens in the distance of 0.2-1.5 km from the FIP. After PFBA, PFBS was the most dominant contaminant discovered in vegetables with a concentration range between <0.2 and $11 \mu\text{g}/\text{kg fw}$. This case is described in detail in Section 3.9. The authors conclude that contamination of vegetables with PFBS might be attributed to irrigation with contaminated groundwater, contaminated soil or atmospheric depositions.

Li et al. (2019) analysed the most prevalent animal protein supplement feeds (APF) for 16 PFAAs, for more details see Section 3.6 In plant derived supplement feeds PFBS contributed with 30% in distillers dried grains with solubles and 13% in soybean meal of the total PFAAs.

In summary, several studies and field data have shown that PFBS can be taken up in plants especially in its edible parts like leaves, vegetables and fruits. Uptake factors in shoot ranged from 2 to 4 for several crops like radish, celery, tomato and pea (Blaine et al., 2014) until up to 14 in lettuce leaves (Blaine et al., 2013). Leaf concentration factors of up to 42 in tomato were determined (Felizeter et al., 2014). The highest plant concentrations of PFBS were reported in maize straw $1843 \mu\text{g}/\text{kg dw}$ and in vegetables like lettuce with $205 \mu\text{g}/\text{kg dw}$ (Ghisi et al., 2019). Furthermore, monitoring data from crops grown 0.4 km from a PFAS point source showed concentrations of up to $11 \mu\text{g}/\text{kg fw}$ PFBS, which might be attributed to contamination via groundwater or soil. Occurrence of PFBS in plant derived supplemental feeds has been reported by Li et al. (2019).

Kowalczyk et al. (2013) demonstrated transfer of PFBS from contaminated feed into meat and milk of dairy cows, see Section 3.5.1.4.

3.8. Temporal trends and comparison with other PFASs

In the present section the data from the field studies above are put into context and compared with other PFASs or with the development over time.

PFBS and precursor substances were largely introduced into the market as a replacement chemical for PFOS-substances (Olsen et al., 2009b). It should be expected that as PFOS use has declined, the PFBS emissions have increased. Corresponding to this, several studies have reported that PFBS environmental concentrations are increasing with time, and other recent studies have reported PFBS to be the dominating PFAS present in monitoring studies.

Water. The emergence of PFBS in the environment is particularly noticeable in water monitoring studies. In a global survey of surface water from 2016, both PFBS and PFOS were detected in all samples. However, PFBS was significantly higher in concentration, with the median PFBS/PFOS ratio being 3.9 (Pan et al., 2018). A recent survey of surface water in Northern Europe, with sampling in 2013, found PFBS to be the dominating PFAS, contributing 21% of the sum PFAS (Nguyen et al., 2017). Similarly, a survey of European and Chinese rivers from 2013-2014 found PFBS to be the dominating PFAS (average 15.6 ng/L), followed by PFOA (4.8 ng/L) and PFPeA (4.7 ng/L) (Heydebreck et al., 2015). In the South China Sea, PFBS was one of the major PFASs analysed, contributing 21% of the total PFAS concentrations, together with PFOA (~26%) and PFOS (~20%). In the Bohai Sea, PFBS was the third most abundant PFAS, after PFOA and PFHxA (Chen et al., 2016).

Drinking water. A study of drinking water from Brazil, France and Spain (Schwanz et al., 2016) reported that PFBS was one of the most frequently detected PFAS, found in 27.2% of the samples, along with PFOS (100%), PFHpA (51.3%) and PFOA (23.0%). A global survey found similar results, with PFBS being the third most frequently detected PFAS in bottled water (47%) and tap water (88%) (Kaboré et al., 2018).

Source zones. Increasing concentrations of PFBS are noticed in source areas as well. Near a PFAS production facility in the Netherlands, PFBS was the dominating PFAS in both river samples (12-27 ng/L) and drinking water (0.5-19 ng/L) (Gebbinck et al., 2017). Near the PFAS production facility at Tangxun Lake, China, the two most dominating PFAS in water were PFBA and PFBS, at mean concentrations of 4770 and 3660 ng/L, respectively (Zhou et al., 2013). A remarkable 24-fold increase in PFBS concentration in groundwater was recorded in the vicinity of a PFAS production plant from 872 ng/L to 21 200 ng/L from 2009 to 2017 (Bao et al., 2019).

WWTP and Landfills. In a recent survey of WWTP in Sweden, PFBS, PFOS, and PFHxS had similar concentrations (1.9 ng/L, 1.9 ng/L, and 1.5 ng/L, respectively) (Eriksson et al., 2017). In San Francisco in 2014, PFBS was not the most dominating, but was emitted on average at 2.7 ng/L, which is a considerable level (Houtz et al., 2016).

Human blood. Glynn et al. (2012) reported that in the period 1996-2010, PFOS in blood decreased, while replacement substances like PFBS and PFHxS increased by 11% and 8.3% per year, respectively. The source of the PFBS exposure was documented to be a local source, see Section 4.3.

Biota. Prior to 2012, PFBS was rarely reported in Arctic marine biota, but it is more common in more recent samples, particularly in polar bears (Gebbinck et al., 2016; Routti et al., 2017; Schlabach et al., 2018). Further indication that PFBS is rising in biota while

PFOS levels are sinking, is found in a time series observed in dolphin livers in the South China Sea (Lam et al., 2016). The ratio of PFBS/PFOS concentrations in the liver was found to have increased from 2002 to 2014. Hence, despite that the measured concentrations of PFBS in biota are generally low, some studies report a clear tendency of an increase of these concentrations.

In summary, recent studies indicate that PFBS is one of the PFASs that show increasing concentrations in various environmental media, including in water and in biota. If PFBS emissions continue to increase, it follows that concentrations in environmental media would increase even further.

3.9. Case study: Environmental distribution of PFBS – Fuxin, China

In this section a case study for the environmental distribution of PFBS in the vicinity of a point source in China is presented. The case study is based on a publication by Bao et al. (2019). In the surroundings of the Fuxin fluorochemical industrial park (FIP), northwestern China, the extent of PFAS contamination in several home gardens, groundwater from private wells and garden soils was measured. PFAS contaminations were also characterised in both home-produced vegetables and eggs from the same residences. The study is included to illustrate the environmental fate for PFBS when released from a point source. The study represents high PFBS concentrations but may nevertheless be relevant if the PFBS concentrations in Europe are allowed to reach similar levels.

The Fuxin FIP was built in 2004 and contains several fluorochemical production plants, mainly for the manufacture of PFBS and polytetrafluoroethylene (PTFE). High levels of PFAS contaminations have been detected in surface water and groundwater around the FIP in Fuxin over the past few years (Bao et al., 2011). In a previous study, PFBS contamination was determined in river and groundwater samples from the environment surrounding the FIP, with maximum measured concentrations at 0.445 µg/L and 0.872 µg/L, respectively.

In the study by Bao et al. from 2019, ten PFASs were measured to investigate the extent of PFAS contamination in the groundwater, soil and home-produced vegetable and egg samples near Fuxin. Sampling was performed in the home gardens of five residences close to the local fluorochemical plant, Table 25. The distances from the FIP site were 0.2 (H1), 0.4 (H2), 0.6 (H3), 1.0 (H4) and 1.5 (H5) km. Duplicate groundwater samples from shallow aquifers at depths of 5-7 m from five private wells in all home gardens were collected. Duplicate samples of surface soil at a depth of 2 cm from the gardens were also collected. Additionally, five common home-produced vegetable samples (tomato, cucumber, eggplant, pepper and Chinese cabbage) were obtained from all the five gardens. Home produced eggs were acquired from three residences only. Samples were collected of public groundwater normally used for both irrigation of home-produced vegetables and feeding of domestic poultry by the residents around the FIP.

PFBS and PFOA were the predominant PFAS contaminants in the groundwater beneath the FIP with maximum concentrations of 21.2 and 2.51 µg/l, respectively. A 24-fold elevation in PFBS contamination from 0.872 to 21.2 µg/L was detected in the local groundwater from 2009 to 2017. In the soil samples from the three sites closest to the FIP, PFBS was the main contaminant, contributing between 62% and 72% of all the target analytes. The concentrations of PFBS in soil showed a declining trend with increasing distance from the FIP site, ranging from 42.1 to < 0.20 ng/g dw.

Table 25: PFBS concentrations in environmental and dietary samples

Sampling site (distance from FIP)	Groundwater ($\mu\text{g/L}$)	Soil (ng/g dw)	Vegetables (ng/g fw)	Whole egg (ng/g fw)	Egg white (ng/g fw)	Egg yolk (ng/g fw)
H1 (0.2 km)	21.2	42	11	32	22	43
H2 (0.4 km)	20.6	11	11	n.a.	n.a.	n.a.
H3 (0.6 km)	13.1	5.6	5.7	24	19	24
H4 (1.0 km)	0.117	0.75	< 0.20	20	16	39
H5 (1.5 km)	0.064	< 0.20	< 0.20	n.a.	n.a.	n.a.

The contamination of vegetables and soils in residential gardens is likely due to irrigation with groundwater from the local public water system. However, the authors could not rule out a possible contribution from atmospheric depositions of PFAS contaminations in the vicinity of the release points.

Table 26: Estimated daily intakes of PFBS (ng/kg bw/day)⁵

Location	Eggs	Vegetables	Groundwater*
H1	24	34	707
H2	n.a	33	687
H3	19	17	436
H4	15	0.42	3.9
H5	n.a	0.42	2.1

*Possible drinking of groundwater from private wells

Residents around the Fuxin FIP prefer to consume drinking water from the municipal public water system (PWSs), rather than groundwater from their own private wells (PWs), Bao et al. (2011). The estimated daily intake of PFBS through ingestion of groundwater, home-produced vegetables and eggs was 765 ng/kg bw per day for residents living in closest proximity to the FIP (H1), Table 26. The levels detected in groundwater near the FIP (H1, H2, and H3) were exceeding the health advisories for PFBS in drinking water, 7 $\mu\text{g/L}$, as issued by the Minnesota Department of Health (2011), Table 25.

Contaminations from point sources to the groundwater could be a threat to drinking water quality in many areas around the world. Previous studies have shown that groundwater in shallow aquifers could suffer from migration of PFAS contaminations through the release of AFFF (Moody et al., 2003), discharge from fluorochemical facilities (Hoffman et al., 2011), or the application of biosolids (Lindstrom et al., 2011). In addition to the Fuxin case, there was a large-scale contamination of drinking water with short-chain PFASs in

⁵ Adapted from Bao et al. (2019).

Uppsala, Sweden, and of arable land in Rastatt, Germany. The contamination in Sweden was mainly caused by the use of AFFF at a firefighting training site at a military airport. The Rastatt case, however, concerned contamination of former arable land with short-chain PFASs and precursors due to the use of compost mixed with sludge from paper production. This resulted in severe contamination of drinking water and crops. Considering the dramatic 24-fold increase in PFBS concentration in water between 2009 and 2017 at Fuxin FIP, the levels are likely to increase further.

3.10. Summary and discussion of environmental fate properties

PFASs are among the most stable organic compounds due to the high bond energy of the C-F bond, which makes the substances resistant to acids, bases, oxidation and reduction, even at high temperatures. PFBS has been found to be stable towards hydrolysis, oxidation and photodegradation in the atmosphere, as well as resistant to degradation in conventional biodegradation tests. Its hydrolytic half-life in water is more than 1 year.

PFBS has been found to be a weakly sorbing substance with a high mobility in the environment. PFBS has a preference for distribution to the aqueous phase and is relatively readily transported when water flows through soil. PFBS is a strong acid which is fully deprotonated at environmentally relevant conditions. The corresponding sulfonate anion in solution, as well as dissolved PFBS salts, do not volatilise.

Preference for the water compartment has been confirmed by distribution modelling and by monitoring data, showing that PFBS is frequently found in surface water, groundwater, drinking water, marine water, effluents from waste water treatment plants and in landfill leachate. The concentrations vary from low to very high in proximity to known point sources. There is evidence that PFBS has had an increasing trend over the last years in water samples, as well as in biota. A similar trend is also seen for human blood, see Section 4.3.

Due to the high aqueous solubility and the low sorption potential, resulting in a preferred distribution to the aqueous phase, PFBS is not readily removed with conventional purification techniques. Both wastewater purification, drinking water production and removal of industrial emissions may suffer from the low purification efficiency for PFBS.

PFBS has a high global contamination potential due to its high environmental stability and high mobility. Long-range transport of PFBS is mainly via the water and sea currents, while sea spray aerosols may also contribute. Long-range transport of PFBS-related substances and degradation to PFBS may take place, often via different routes of transportations than PFBS, like atmospheric transport.

Long-range transport has been demonstrated when PFBS has been found in samples of marine water, surface water, snow, ice, air and biota from remote areas. However, the measured concentrations of PFBS are generally low.

An increasing trend for PFBS in the environment has been indicated, which can be explained by a combination of increasing annual production and a continuous accumulation of a persistent substance. However, the detection of PFBS in remote areas, like the Arctic, is taken as evidence of the persistence and mobility of PFBS.

As a short-chain, ionic perfluorinated substance, PFBS is a highly mobile substance, and there are no local or intermittent sinks for the pollution stock. Oceanwater is an important compartment for storage and transport of the compound. In coastal water PFBS is bioavailable and can accumulate in the marine food chain, and therefore there is a high potential for further increase in the exposure of wildlife to the substance.

PFBS is bioavailable and has been detected in blood and different organs of animals and humans. Bioaccumulation and protein binding is discussed in detail in Sections 3.5 and 3.6, respectively, while the levels of PFBS in humans are presented in Section 4.3. PFBS has been found to bind to serum albumin, and thereby the blood can distribute PFBS within the body resulting in a potential to enrich particularly in blood rich tissues. PFBS has also been shown to bind to different other proteins (peroxisome proliferator-activated receptors (PPAR α , PPAR γ), human thyroid hormone transport protein transthyretin (TTR), and organic anion transporting polypeptides (OATPs)), although the binding potential is generally lower compared to the longer chain PFAAs. However, tissue distribution and accumulation of PFBS may be affected by its binding to proteins and may be of toxicological significance.

PFBS has been detected in human blood, lung, bone, kidney, urine and hair. It has also been detected in human cord blood. A serum elimination half-life of around one month (up to 46 days) has been measured in humans, which is considerably longer than the half-lives measured for rodents. However, PFBS has a relatively short half-life in the human body compared to the longer chain counterparts. The half-life of PFBS in pigs was reported to be similar to that in humans, and BMF values > 1 were reported for pigs. PFBS has been detected in different species of wild life including green turtles and polar bears, both are threatened species. PFBS has also been detected in marine mammals like dolphins and whales. In the whale it was shown that PFBS transfers to the fetus. Additionally, PFBS was shown to be transferred to the eggs of birds. The findings of PFBS in sensitive life stages and in threatened species are of special concern.

Substances with a low bioaccumulation potential have been estimated to potentially reach similar levels in biota to substances that are known to bioaccumulate, provided that they are sufficiently persistent and mobile in the environment. Calculations even show that with an assumed half-life in water of 10 years and a reported BAF in crab of 110, the concentrations of PFBS in aquatic biota may be expected to exceed the biota concentrations for a persistent and bioaccumulative substance over time.

The occurrence of PFAAs in animal protein supplement feeds (APFs) has been reported. In animal derived supplement feeds PFBS contributed with 17%, 13.3%, 9.2% to the total amount of PFAAs in blood meal, feather meal, meat meal respectively. In plant derived supplement feeds PFBS contributed with 30% in distillers dried grains with solubles and 13% in soybean meal to total PFAAs.

Due to its high water solubility, PFBS is bioavailable to plants. Enrichment in plants has been demonstrated by studies and field data. PFBS can be found in high concentrations in the edible parts of plants like leaves, vegetables and fruits, and long-term exposure of humans and animals can not be excluded.

4. Human health hazard assessment

The studies referred to in this chapter are according to standard tests, guidelines or GLP if stated so. Otherwise they are studies published in scientific peer-reviewed journals or public reports.

4.1. Toxicokinetics (absorption, metabolism, distribution and elimination)

4.1.1. Non-human information

One study investigated the toxicokinetics of PFBS and perfluorohexanoic acid (PFHxA) in Sprague–Dawley rats and cynomolgus monkeys. A single intravenous dose of 10 mg/kg bw was given to both species followed by measuring the serum concentration of PFBS and PFHxA. The serum clearance (see Table 27) was more rapid in female than male rats for both substances, while no apparent gender difference was seen for monkeys (Chengelis et al., 2009).

In a pharmacokinetic study by Olsen et al. (2009b) in rats and monkeys, PFBS was administered orally or by a single intravenous (i.v.) dose in rats (30 mg K+PFBS/kg body weight and monkeys of 10 mg KPFBS/kg body weight), a half-life of 4 hours in female rats, 4.8 hours in male rats and 3.5 days in female monkeys and 4 days in male monkeys were measured. The volume of distribution (V_{dss}) was estimated and shown to be similar between sexes, and the V_{dss} suggested that distribution was primarily extracellular regardless of sex according to the authors. Urine appeared to be a major route of elimination (Olsen et al., 2009b).

In a toxicokinetic evaluation, male Sprague–Dawley rats received a single 30 mg/kg bw dose of a PFBS-precursor, Me-FBSA (ECHA dissemination website, CAS 34454-97-2, <https://echa.europa.eu/registration-dossier/-/registered-dossier/5299/7/2/2>) in propylene glycol by oral gavage at a volume of 5 mL/kg body weight. The vehicle control group rats received a single dose of ethylene glycol at a volume of 5 mL/kg. Necropsies were performed at 4 hours, 28 hours and 100 hours post dose. Animals that received the test article at 30 mg/kg bw had average total fluorine concentrations in serum of 10.8 ppm (4 hours post dose), 23.1 ppm (28 hours post dose) and 1.0 ppm (100 hours after dose). The half-life elimination times for the test compound in the serum was 17 hours. The average liver total fluorine concentrations were 14.1, 18.3 and 2.2 ppm after 4 hours, 28 hours or 100 hours post dose, respectively. The apparent total fluorine elimination rates were similar in both serum and liver for animals treated with the test compound.

Toxicokinetic parameters were reported in a recent study in parallel to the recent NTP-study (NTP, 2019) on three fluorinated compounds (PFBS, PFHxS and PFOS) (Huang et al., 2019a). A single intravenous dose (4 mg/kg bw as 4mL/kg) or gavage administration (three dose levels (4, 20 and 100 mg/kg bw as 5 mL/kg) of PFBS were given to male and female Hsd:Sprague-Dawley rats. Concentrations of these PFAS were measured in the plasma, liver, kidney, and brain. A two-compartment model was the best fit for male rats (intravenous and gavage) and female rats (intravenous) while a one-compartment model was a better fit for the female gavage exposure data. Plasma half-life increased with longer chain length after gavage administration: PFBS- males averaged 3.3 h, females 1.3 h; PFHxS- males averaged 16.3 days, females 2.1 days; PFOS- males and females averaged ~ 20 days. There were dose-dependent changes in clearance and systemic exposure for all administered chemicals. Overall, concentrations of PFBS were higher in the liver than in the kidney in both sexes. PFBS concentrations in all tissues decreased slightly over time, with females having a faster decrease than males. In males, liver:plasma ratios were

generally above 1, dropping below 1.0 at 12 h, while kidney:plasma ratios remained around 0.29 to 0.38. In female rats, the liver:plasma ratios were lower and kidney:plasma ratios were higher than those in males. Brain:plasma ratios ranged from 0.01 to 0.02 in males and was 0.02 in females at 0.5 h. Sex differences in plasma half-life and tissue distribution were observed for PFBS and PFHxS, but not PFOS.

The tissue distribution of PFBS in mice was characterized (Bogdanska et al., 2014) in the same manner as for the eight-carbon homologue perfluorooctane sulfonate (PFOS) to allow direct comparisons. Following dietary exposure of adult male C57/BL6 mice for 1, 3 or 5 days to 16 mg (35)S-PFBS/kg bw/day, both scintillation counting and whole-body autoradiography (WBA) revealed the presence of PFBS in all of the 20 different tissues examined, demonstrating its ability to leave the bloodstream and enter tissues. After 5 days of treatment, the highest levels were detected in liver, gastrointestinal tract, blood, kidney, cartilage, whole bone, lungs and thyroid gland. WBA revealed relatively high levels of PFBS in male genital organs as well, except for the testis. The tissue levels increased from 1 to 3 days of exposure but appeared thereafter to level-off in most cases. The estimated major body compartments were whole bone, liver, blood, skin and muscle. This exposure to PFBS resulted in 5-40-fold lower tissue levels compared to PFOS. PFBS also showed different pattern of tissue distribution compared to PFOS, including lower levels in liver and lungs relative to blood. In, general, the level in tissues relative to blood were lower for PFBS than for PFOS, with the difference being most pronounced for liver and lungs (Bogdanska et al., 2014).

In another study male and female CD-1 mice were given a single dose with 0, 30, or 300 mg/kg bw PFBS via gavage. Trunk blood was collected at 0.5, 1, 2, 4, 8, 16, 24, and 48 hours and urine at 24 hours after dosing. The half-life of PFBS was estimated to be 4.6 hours in the male mice and 2.5 hours in the females. Within 24 hours, more than 95% of the serum PFBS was excreted into urine (Rumpler et al., 2016). Thus, the half-life of PFBS in mice is short, similar to that of rats (see Table 27 below).

Another study examined the transfer of a mixture of perfluoroalkyl acids (PFAAs) from contaminated feed into the edible tissues of 24 fattening pigs and this study showed a tissue distribution of PFBS to blood plasma, liver, kidney, muscle tissue and fat. PFBS was eliminated through feces and urine (Numata et al., 2014). The geometric average half-life was estimated to be 43 days, the individual variability was large and the higher half-life values for PFBS were overlapping with the lower values for PFOA. PFBS have also been detected in milk from dairy cows, although at very low concentrations compared to other PFASs (Kowalczyk et al., 2013).

The enterohepatic circulation of PFASs likely contributes to their extended elimination half-lives in humans. It was demonstrated that PFBS, PFHxS, and PFOS were transported into hepatocytes both in a sodium-dependent and a sodium-independent manner by Na⁺/taurocholate cotransporting polypeptide (NTCP). PFBS, PFHxS, PFOS and PFCAs with 7-10 carbons are substrates of organic anion transporting polypeptides (OATPs). Chinese Hamster Ovary (CHO) and Human Embryonic Kidney 293 (HEK293) cells were used to demonstrate that human OATP1B1, OATP1B3, and OATP2B1 can transport PFBS, PFHxS, PFOS and the 2 PFCAs (C8 and C9). In addition, it was shown that rat OATP1A1, OATP1A5, OATP1B2, and OATP2B1 transport all 3 sulfonates. This study suggests that besides NTCP and the human apical sodium-dependent bile salt transporter (ASBT), OATPs also are capable of contributing to the enterohepatic circulation and extended human serum elimination half-lives of PFBS and other PFASs (Zhao et al., 2017).

4.1.2. Human information

Olsen and co-workers (Olsen et al., 2009a) also performed an elimination assessment in humans where 6 employees (1 woman and 5 men) donated total of 10 blood samples to obtain serum samples over a 6 months period when the employees were not working in the production line. Spot urine samples were also collected at three time points. Determination of the serum elimination half-lives for the six subjects included all serum PFBS concentrations measured above the LOQ of 5 ng/mL (0.005 µg/mL). The mean serum elimination half-life for PFBS was 27.7 days (95% confidence interval 16.1–39.3). The geometric mean serum elimination half-life was 25.8 days (95% CI 16.6–40.2). The female employee had the longest serum elimination half-life of 45.7 days (see Table 27), while the mean serum half-life of elimination in men was 24.1 days (95% CI 13.1–32.5). Urine seems to be the major route of elimination. Due to the low number of subjects (only one female) and variations in the half-life obtained for one male subject who was tested twice, the elimination half-life obtained should be used with caution. However, a relatively large individual half-life variability for different other PFAAs was also observed using data from up to 66 individuals in a study by Zhang et al. (2013b).

4.1.3. Summary

PFBS is transported through the body and is found in many different tissues as it binds to the serum protein, albumin, and several other proteins. The major route of elimination is through urine, although PFBS is also found in feces and has been detected in milk of dairy cows. The half-life of PFBS in mice and rats are similar with shorter half-life in female than male rodents. PFBS has a longer half-life in monkeys compared to rodents, while the half-life in pigs seem to be similar to humans. PFBS may also bind to several different transporter polypeptides (NTCP, ASBT and OATPs) which are all capable of contributing to the enterohepatic circulation and thus extended human serum elimination half-lives of PFBS and other PFAAs. A serum elimination half-life of around one month (up to 46 days) has been measured in humans. Comparing the measured half-lives (Table 27) shows that the half-life of PFBS in humans is considerably longer than the measured half-lives in rodents. Based on the half-life reported for PFBS in human blood compared to the half-lives of PFOA, 2 – 4 years, (B) and PFHxS, 7 - 8 years, (vB), it is concluded that PFBS is less bioaccumulative. However, no final conclusion on the bioaccumulation potential can yet be reached, as there is yet no approach to interpret the human and mammalian data. However, the available information indicates a moderate bioaccumulation potential.

Table 27: Serum t_{1/2} of PFBS in different species.

Serum t _{1/2} of PFBS in different species		Female	Male	Reference
Sprague-Dawley rat <i>PFBS</i>	half-life for the terminal elimination phase serum T _{1/2}	0.64 hours	2.1 hours	Chengelis et al., 2009
Sprague-Dawley rat <i>Potassium PFBS</i>	Terminal mean half-life of serum elimination Two-compartmental model and first-order elimination	4.0 hours	4.8 hours	Olsen et al., 2009b

Serum t_{1/2} of PFBS in different species		Female	Male	Reference
Sprague-Dawley rat <i>Potassium PFBS</i>	Mean half-life was measured in plasma.	1.3 hours	3.3 hours	Huang et al., 2019a
D-1 mice <i>Potassium PFBS</i>	Mean serum elimination half-life	2.5 hours	4.6 hours	Rumpler et al., 2016 (abstract)
Cynomolgus monkeys <i>PFBS</i>	Half-life for the terminal elimination phase	0.34 days	0.63 days	Chengelis et al., 2009
Cynomolgus monkeys <i>Potassium PFBS</i>	Mean serum elimination half-life	3.5 days	4 days	Olsen et al., 2009b
Pigs <i>PFBS</i>	Geometric average values reflect both plasma and edible tissue elimination half-lives	43 days	43 days	Numata et al., 2014
Humans <i>Potassium PFBS</i>	Mean serum elimination half-life one-compartment model	45.7 days (1 female)	24.1 days (5 male)	Olsen et al., 2009a

4.2. Human exposure routes

Exposure via the environment

The presence of PFBS in drinking water was presented in Section 3.2.5.3, while enrichment of PFBS in plants and exposure through the consumption of vegetables was discussed in Section 3.7.

Consumer exposure - household dust

The presence of PFBS in house dust and exposure via the indoor environment has been documented in the scientific literature. Several studies report findings of PFBS in dust collected in European homes (D'Hollander et al., 2010; Haug et al., 2011; Jogsten et al., 2012; Bohlin-Nizzetto et al., 2015; Eriksson and Kärrman, 2015; Karásková et al., 2016; Torre et al., 2019). The European PFBS-concentrations in house dust range between 0.01 ng/g (Bohlin-Nizzetto et al., 2015) and 72.8 ng/g (Eriksson and Kärrman, 2015) with a detection frequency ranging between 21 (derived from Haug et al., 2011) and 60% (Jogsten et al., 2012).

A study of world-wide indoor exposure to polyfluoroalkyl phosphate esters (PAPs) and other PFASs in household dust by Ericsson and Kärrman (2015) reports that PFBS was detected more frequently (approximately 50%) than in previous studies of dust sampled in 2008/2009 from Norway (Haug et al., 2011) and the U.S. (Fraser et al., 2013; Knobeloch et al., 2012), but at a similar level.

In a recent study by Torre et al. (2019) the concentrations of PFASs were quantified in 65 house dust samples obtained from Belgium, Italy and Spain. The authors report an increase in perfluorobutane sulfonate (PFBS) concentrations in Belgian house dust in comparison to previously published data. The levels of PFBS seem to have increased in the last ten years (0 to 0.40 ng/g; D'Hollander et al., 2010; Torre et al., 2019). No correlations between PFOS and PFHxS with PFBS ($p > 0.05$) were detected. The authors conclude that this suggests different application areas for PFBS. The potassium salt of PFBS is marketed as flame retardants for polycarbonate resins used in electronics (OECD, 2013). The study did not find any significant correlations between PFBS content in dust and information on the presence and use of electric and electronic devices in the homes.

4.3. Levels of PFBS in humans

Human blood has been the most frequently used matrix for determining the internal dose of PFBS in humans as described in detail below. However, PFBS has also been detected in other human tissues, see Table 28. In a study on autopsy tissues from 20 subjects who had been living in Tarragona, Catalonia, Spain, PFBS was detected in bone, lung and kidney, but for bone and kidney only in one sample each (Perez et al., 2013). The concentrations varied between <LOD (14.4 ng/g) and 17.8 ng/g for bone, <LOD (3.4 ng/g) and 80.4 ng/g for kidney and <LOD (2.10 ng/g) to 9.7 ng/g for lung. In a study from Belgium on 30 human hair samples, PFBS was detected in all samples, and the concentrations ranged from 34 to 120 pg/g (Alves et al., 2015). PFBS was also determined in urine ($n=24$, LOD 0.74 ng/mL) and hair samples ($n=30$ LOD 0.09 ng/g) from Barcelona, Spain, collected in 2010-2011 and 2009, respectively (Perez et al., 2012). Only one urine sample and two hair samples had detectable concentrations of PFBS, and the concentrations were 1.3 ng/mL in urine, and 0.8 and 10.7 ng/g in hair, respectively.

In human blood concentrations of PFBS have been measured (Table 28) in both the general population and populations which have experienced known elevated PFBS exposure, including workers.

In a study comprising human blood plasma samples from the German environmental specimen bank collected between 1982 and 2010 ($n=258$, age 20-29 years, both genders), PFBS was not found in concentrations above the LOQ of 0.5 ng/mL in any samples (Schröter-Kermani et al., 2013).

In another study from Germany, PFBS was determined in samples ($n=396$) collected between 2009 and 2016. However, only 1% of the samples had detectable concentrations of PFBS (LOQ 0.4 ng/mL), and the concentrations in these samples were below the LOQ of 0.4 ng/mL (Fromme et al., 2017).

A Norwegian study comprising pooled serum samples of men ($n= 24$ pools, each comprising >20 samples, age 40-50 years) collected between 1977 and 2006, reported PFBS concentrations above LOQ (0.05 ng/mL) in 16 of 24 samples. The concentrations varied between 0.065 and 0.18 ng/mL, but no time trend was observed (Haug et al., 2009).

In a more recent study from Norway, paired serum, plasma and whole blood samples were collected from 61 adults (men and women) in 2013-2014 (Poothong et al., 2017). PFBS

was detected in 51, 81 and 100% of the serum, whole blood and plasma samples, respectively, and the concentrations ranged from below LOQ to 0.22 ng/mL. The LOQs for serum, whole blood and plasma were 0.009 ng/mL, 0.009 ng/mL and 0.018 ng/mL, respectively.

Serum samples of nulliparous women from the Aarhus Birth Cohort Biobank, Denmark (n=1533, median age 29 years) were collected in the period 2008-2013 (Bjerregaard-Olesen et al., 2016). PFBS was detected only in 1.2% of the samples (LOD 0.02 ng/mL).

During 2001 to 2004 men and women turning 70 years (from Uppsala, Sweden) were invited to donate blood samples (n=1006), to study their PFBS concentrations (Salihovic et al., 2015). Detectable levels of PFBS were found in 14% of the samples from women and in 11 % of the samples from men. The levels ranged from 0.035 to 0.086 ng/mL (LOD 0.01-0.17 ng/mL).

46 blood samples from adults (age 19-53 years, males and females) living in Barcelona, Spain, were investigated, and PFBS was found in 16 of 46 samples, in concentrations up to 0.43 ng/mL (LOD 0.04 ng/mL) (Gómez-Canela et al., 2015). The samples were collected in 2009-2010.

In the US, PFBS was determined in two cross sectional studies comprising blood donors that gave samples in 2006 or 2010 (Olsen et al., 2012). In both sample sets the 75 percentile was below the LOQ (0.025 ng/mL), and the 95 percentile was 0.3 ng/mL. In a follow up study on samples collected from blood donors in 2015 (n=616) only 8.4% of the samples had quantifiable concentrations of PFBS, in the range from below LOQ to 4.2 ng/mL (Olsen et al., 2017).

In a French study on women and their newborns (n=100 pairs, age 20-46 years, samples collected in 2010-2013), PFBS was detected (LOD 0.05-0.2 ng/mL) in less than 1% of the paired maternal and cord sera samples (Cariou et al., 2015). The quantified levels were always lower than 0.40 ng/mL. Samples of breast milk from the same mothers were collected and analysed, but no samples had detectable concentrations of PFBS (LODs: 0.01 to 0.04 ng/mL). In a Spanish cohort of mother and children, the concentration of PFBS was assessed in paired samples of maternal and cord blood (N=66 pairs, mean maternal age 32 years) collected in 2003–2006, but no samples had levels above LOQ (0.10 ng/mL) (Manzano-Salgado et al., 2015).

Concentrations of PFBS were assessed in 687 samples of cord blood collected in Shanghai, China between 2011 and 2012 (Wang et al., 2016a). The mean concentration of 0.053 ng/mL, while the highest concentration was 0.46 ng/mL. Concentrations above LOQ (0.009 ng/mL) were found in 97% of the samples. Hence it is demonstrated that fetuses are exposed to PFBS through trans-placental transfer.

In a study of primiparous women from Uppsala, Sweden (n=36 pools, age 19–41 years) the serum concentrations of PFBS increased with 11% per year in the period from 1996 to 2010 (Glynn et al., 2012). The highest concentration was found in a pool from 2010 with a concentration of 0.1 ng/mL (LOD 0.013 ng/mL). The increasing trend of PFBS was later confirmed and linked to contamination of the drinking water in the city of Uppsala (Gyllenhammar et al., 2015). Concentrations of PFBS in serum of first-time mothers (n=297, age 20-41 years) sampled in the Uppsala county in two different time periods were determined (Gyllenhammar et al., 2015). Statistically significantly higher concentrations of PFBS (LOD 0,01 ng/g) were found in the samples collected between 2008–2011 (<LOD-0.80 ng/mL) compared to those collected between 1996–1999 (<LOD-0.21 ng/mL), with median concentrations of 0.019 and 0.027 ng/mL in 1996-99 and 2008-2011, respectively. The highest levels were observed among women living in areas with the most elevated drinking water levels, in both time periods. This support the suggestion that drinking water intake has been the major pathway of exposure for PFBS in Uppsala

over the studied period. However, in another study on primiparous women living in Uppsala County, Sweden (n= 30 pools, each comprising 9-10 individuals, age 19 to 42 years) no significant change in concentration was seen in the serum samples for PFBS over the time period 1997 to 2012 (Gebbink et al., 2015). PFBS was observed above LOQ (0.009 ng/mL) in around 60% of the samples, in concentrations in the range 0.011 to 0.058 ng/mL.

In a study from the Veneto region in Italy where the drinking water had been contaminated with PFASs, including PFBS, elevated serum concentrations were observed among people living in the exposed area (<LOQ – 4.26 ng/mL) compared to the people living in the non-exposed area (<LOQ-0.36 ng/mL), Ingelido et al. (2018).

The level of PFBS was explored in a study on serum samples collected in 2011 from the general population of Henan, China (n=133, age 0-88 years, male and females) (Fu et al., 2014). In around 30% of the samples PFBS was observed, and the concentrations ranged from 0.03 to 2.2 ng/mL (LOD 0.05 ng/mL).

In a study on 103 males from Guangzhou city, Guangdong province, China, collected between 2012 and 2013, PFBS was detected (LOD 0.013 ng/mL) in all samples with a mean blood concentration of 0.18 ng /mL (95 percentile; 0.43 ng/mL), Song et al. (2018).

PFBS was determined in a cohort of women (n=335) from Zhejiang, China, included in a case control study on endometriosis-related infertility (Wang et al., 2017). Samples were collected in 2014-2015. As many as 98.5% of the plasma samples had detectable (LOD 0.009 ng/mL) concentrations of PFBS, and the median (IQR) concentrations were 0.091 ng/mL (0.088-0.097 ng/mL) and 0.089 ng/mL (IQR 0.085-0.095 ng/mL) for cases and controls, respectively.

In 2004, a fluorochemical industrial park was built in Fuxin, China, for the production of PFBS and other PFAAs. Serum samples from non-occupationally exposed residents from Fuxin, China (n=120, age 0.1 to 87 years, males and females) were collected in 2009 (Bao et al., 2011). PFBS was found above LOQ (0.07 ng/mL) in 93% samples, with a median concentration of 0.12 ng/mL while the max concentration was 1.3 ng/mL. In another study on non-occupationally exposed healthy residents from Fuxin, China (n=100, 50 males and 50 females, age 1-87 years), with samples collected in 2015, PFBS was detected in all serum samples (LOQ: 0.05 ng/mL), Bao et al. (2017). The median (0.14 ng/mL) and max concentrations (1.5 ng/mL), were very similar to those observed in the samples collected in 2009. In a later study from the same area it was demonstrated that the area is considerably contaminated with PFBS, see Section 3.9, Bao et al. (2019).

In a Chinese study serum samples from fishery employees and their families, living close to the Tangxun Lake with elevated PFBS concentrations were compared to serum samples from people in a reference area in the Hubei Province (Zhou et al., 2014). PFBS was detected (LOQ 0.10 ng/mL) in 72% and 29% of the serum samples of the fishery employees and fishery families, respectively, whereas it was detected only in one serum sample from the reference area. The concentrations in the fishery employees (detected in 28 of 39 samples) were highest with a median concentration of 11.3 ng/mL, followed by the fishery families (detected in 2 of 7 samples) with a median concentration of 2.34 ng/mL. The PFBS concentration measured in the sample from the reference area (detected in 1 of 9 samples) was 0.12 ng/mL.

Concentrations of PFBS in the range < LOD (0.02 ng/mL) to 0.09 ng/mL have been reported in blood serum from Australian fire fighters (n=149, age 17-66 years), Rotander et al. (2015). In a study on blood levels in employees of the 3M Company (n=18, including six researchers, six production workers, six corporate employees, age 29-79 years), PFBS was measured (Ehresman et al., 2007). Most samples were below LOQ, but this was rather

high, 5 ng/mL. However, in the samples from some of production workers PFBS concentrations in the range 7-32 ng/mL were observed.

Table 28: Concentrations of PFBS in human tissues (ng/g ww) and blood (ng/mL)

Location	Year	Mean +/- SD ng/mL	Min-Maks (Median) ng/mL	n (det. Freq)	Remark	Reference
Germany, Münster	1982-2010	nd		258 (0%)	Plasma	Schröter-Kermani et al., 2013
Germany	2009 - 2016	<0.4		396 (1%)	Plasma	Fromme et al., 2017
Norway	1997 - 2006		0.065 - 0.18	24 pools (67%)	Serum (men 40 – 50 years)	Haug et al., 2009
Norway	2013-2014	0,05; 0,13; 0,04 in serum, whole blood, plasma respectively	< LOQ - 0.22	61 (51, 81, 100% of serum, whole blood, plasma respectively)	Paired serum, plasma and whole blood (adult men and women)	Poothong et al., 2017
Denmark	2008-2013			1533 (1.2%)	Serum (nulliparous women, median age 29)	Bjerregaard-Olesen et al., 2016
Italy, Veneto	2018	<LOQ	<LOQ - 4.26	257 (31%)	Blood (exposed areas)	Ingelido et al., 2018
Italy, Veneto	2018	<LOQ	<LOQ - 0.36	250 (18%)	Blood (non-exposed areas)	Ingelido et al., 2018
France	2010 - 2013		LOD (0.05-0.2) - 0.4 (max)	< 1% of paired samples	Paired maternal and cord sera (women and their newborns)	Cariou et al., 2015
Spain, Tarragona	2008	0.9*	1.5 (max)		Liver	Pérez et al., 2013
Spain, Tarragona	2008	3.2*	17.8 (max)		Bone	Pérez et al., 2013
Spain, Tarragona	2008	<LOD			Brain	Pérez et al., 2013
Spain, Tarragona	2008	17.8*	9.7 (max)		Lung	Pérez et al., 2013

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Year	Mean +/- SD ng/mL	Min-Maks (Median) ng/mL	n (det. Freq)	Remark	Reference
Spain, Tarragona	2008	8*	80.4 (max)		Kidney	Pérez et al., 2013
Spain, Barcelona	2009 - 2010	0.20 (0.11)	< LOQ - 0.43	46 (35%)	Blood adults (age 19-53, males and females)	Gómez-Canela et al., 2015
Spain	2003 - 2006		< LOQ	66	Paired samples maternal and cord blood (mean maternal age 32 years)	Manzano-Salgado et al., 2015
Sweden, Uppsala	1996- 1999	0.03 ± 0.00	<MDL - 0.21 (0.02)	132 (77 %)	Serum pregnant mothers (1996-1999)	Gyllenhammar et al., 2015
Sweden, Uppsala	2008- 2011	0.06 ± 0.01	<MDL - 0.80 (0.03)	134 (86 %)	Serum pregnant mothers (2008-2011)	Gyllenhammar et al., 2015
Sweden, Uppsala	1996- 2010		<LOD - 0.10 (0.09)	36 pools (72 %)	Serum primiparous mothers	Glynn et al., 2012
Sweden, Uppsala	1997- 2012		<LOD (0,009) - 0,058	30 pools (60%)	Serum primiparous mothers	Gebbink et al., 2015
Sweden, Uppsala	2001 - 2004	0,068(0.05) for women and 0.055 (0,03) for men	0.035 - 0.086	1006 (14% women and 11 % of men)	Blood (Men and women age 70)	Salihovic et al., 2015
USA	2006 and 2010				Plasma, blood donors	Olsen et al., 2012
USA	2015		<LOQ-4.2	616 (8,4%)	Plasma, blood donors	Olsen et al., 2017
USA			< LOQ (5)- 32		Serum, plasma and blood (Fluorochemic al workers)	Ehresman et al., 2007
Australia			<LOD (0.02) - 0.09	149	Serum (Fire fighters)	Rotander et al., 2015

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Location	Year	Mean +/- SD ng/mL	Min-Maks (Median) ng/mL	n (det. Freq)	Remark	Reference
China, Henan	2011		0.03 - 2.2	133 (30%)	Serum (0-88 years, male and females)	Fu et al., 2014
China, Guangzhou city	2012-2013	0.18	(95 percentile 0.43)	103 (100%)	Blood	Song et al., 2018
China, Zhejiang	2014-2015		Case: median 0.091 (IQR: 0.088-0.097) Controls: median 0.089 (IQR: 0.085-0.095)	335 (98.5%)	Plasma, women (case study) endometriosis-related infertility	Wang et al., 2017
China, Shanghai	2011-2012	0.053	LOD - 0.46 (0.05)	687 (97%)	Cord blood (ng/mL)	Wang et al., 2016a
China, Tangxun Lake			Employee: 11.3 (median) fishery family: 2.34 (median) ref. area: 0.12 (median)	39 employees (72%) 7 family members (29%) 9 background exposed (11%)	Serum, fishery employees, fishery family, reference area population	Zhou et al., 2014
China, Fuxin	2009	0.19 ± 0.22	0.01 - 1.30 (0.12)	120 (93%)	Serum	Bao et al., 2011
China, Fuxin	2015	0.23 ± 0.27	0,05-1.5 (0,14)	100 (100%)	Serum	Bao et al. 2017

* ng/g

4.3.1. Summary

PFBS has been detected in human blood, lung, bone, kidney, urine and hair. Concentrations of PFBS in human blood have been analysed in a number of studies from Europe, US and Asia. The PFBS level is often below the quantification limit, with a concentration frequently below 0.5 ng/mL. However, in a few studies concentrations up to 5 ng/mL were observed. In persons with known elevated exposure to PFBS, e.g. from drinking water or in occupational settings, also higher levels have been measured. Further,

PFBS has been detected in human cord blood with a mean concentration of 0.053 ng/mL, while the highest concentration was 0.46 ng/mL.

Studies from Uppsala, Sweden, as well as a study from the Veneto region in Italy, in both cases with known elevated PFBS exposure from drinking water, have shown that persons experiencing elevated exposure to PFBS have elevated PFBS concentrations in their blood compared to people from the same region experiencing only the general background exposure. Furthermore, elevated PFBS blood concentrations were observed in fishery employees and their families working and living nearby a lake in China with elevated PFBS concentrations. High blood PFBS concentrations were also observed in employees from a company manufacturing PFBS. Hence, it has been demonstrated that elevated exposure to PFBS leads to elevated blood concentrations.

4.4. Repeated dose toxicity

4.4.1. Non-human information

4.4.1.1. In vitro data

An in vitro study showed that PFBS treatment for 6 days extensively promoted the differentiation of 3T3-L1 preadipocytes to adipocytes, resulting in significantly increased triglyceride levels in cells (Qi et al., 2018). In particular, the treatments with PFBS (10, 50, 100 and 200 µM) at the early adipogenic differentiation period (day 0–2) were positively correlated with increased triglyceride accumulation on day 6. Protein and messenger-RNA (mRNA) levels of the transcription factors in adipocyte differentiation; CCAAT/enhancer-binding protein α (C/EBPα) and peroxisome proliferator-activated receptor gamma (PPARγ), along with acetyl-CoA carboxylase (ACC) and fatty acid synthase (FAS), the key proteins in lipogenesis, increased after PFBS treatment. PFBS significantly activated the phosphorylation of extracellular signal-regulated kinase1/2 (ERK1/2) after 4-h treatment, and the effect of PFBS on triglyceride was abolished by U0126, a specific MAPK/ERK kinase (MEK) inhibitor. In conclusion, PFBS increased the adipogenesis of 3T3-L1 adipocytes, in part, via MEK/ERK-dependent pathway.

4.4.1.2. In vivo data: oral

Data available on the toxicity of PFBS in animals have identified the liver, kidneys, stomach, and hematological systems as targets of toxicity. However, the data are limited. A statistically significant dose dependent decrease in hemoglobin and hematocrit levels were observed in male rats administered orally 200 mg/kg bw/day and 600 mg/kg bw/day PFBS for 90 days (Lieder et al., 2009a). A significant decrease in erythrocyte levels were observed at 600 mg/kg bw/day. The NOAEL was determined to be 60 mg/kg bw/day in male rats (based on haematological parameters) and 600 mg/kg bw/day in females (based on kidney effects described below) (Lieder et al., 2009a). No histological changes or abnormalities in bone marrow of male rats were observed at 600 mg/kg bw /day. →Klimisch reliability 1 - well documented, guideline study. A study by 3M reported no significant hematological alterations in rats dosed with up to 900 mg/kg bw /day PFBS by gavage for 28 days (3M, 2001) →Klimisch reliability 1 - guideline study.

In a repeated oral dose toxicity study for 28-days performed by NTP, a significant decrease in serum phosphorus and potassium levels in male rats treated at 300 mg/kg bw/day and 900 mg/kg bw/day was observed, giving NOAEL values of 100 mg/kg bw/day for males and 300 mg/kg bw/day for females (NTP, 2019). The decrease for potassium was approximately 20% of the control values at a level of significance of $p < 0.05$ in both

treatment groups. The decrease for phosphorus was approximately 16% of control values at a level of $p < 0.05$ in both treatment groups.

Administration of 600 mg/kg bw/day for 90 days resulted in tubular and ductal papillary epithelial hyperplasia in the kidneys and necrosis and hyperplasia/hyperkerosis in the forestomach of rats (Lieder et al., 2009a). Increases in absolute and relative liver weight were reported in male rats administered 900 mg/kg bw/day for 28 days (3M, 2001). At 300 mg/kg bw/day treatment-related microscopic changes were observed in the kidney and liver for males and kidneys for females, and at 1,000 mg/kg bw/day hepatocellular hypertrophy was observed in a 2-generation study (Lieder et al., 2009b). → Klimisch reliability 1 - well documented, guideline study.

Treatment of male rats with 900 mg/kg bw/day PFBS by gavage for 28 days induced a significant increase in absolute and relative liver weight (25–30%) relative to controls, which was no longer detected following a 14-day recovery period (3M, 2001). Clinical chemistry tests of liver function were unremarkable, and there were no chemical-related microscopic alterations. The NOAEL for liver weight effects was 300 mg/kg bw/day. Treatment of female rats with 900 mg/kg bw/day PFBS by gavage for 28 days caused a significant increase (9–11%) in absolute and relative kidney weight, but no significant alterations in the microscopic appearance of the kidneys was observed (3M, 2001). The weight of the kidneys returned to control levels following a recovery period of approximately 14 days. The NOAEL for kidney weight effects was 300 mg/kg bw/day PFBS.

A mice study with PFBS administered daily to male mice (APOE*3-Leiden.E3L CETP) at 30 mg/kg bw/day for 4-6 weeks modestly reduced plasma triglycerides and cholesterol, and increased clearance of triolein (a triglyceride) (Bijland et al., 2011).

4.4.1.3. In vivo data: inhalation

In a subchronic study, Crl:CD SD rats were exposed (whole body) to vapours of perfluorobutane sulfonyl fluoride (PBSF) for 6 hr/day, 5 days/week for 4 weeks (ECHA dissemination website⁶). The mean analysed exposure concentrations of the chemical were 47, 162 and 459 ppm. Clinical signs observed immediately post exposure included vocalising and agitation when handled, walking on toes (abnormal gait) and hyperactivity, consistent with a neurological effect. These signs were transient and generally resolved the following day with no evidence of sustained neurotoxicity. There were no effects on haematological or blood chemistry parameters. Histopathological examination of the respiratory tract revealed no treatment-related findings (ECHA dissemination website⁶).

4.4.2. Human information

One low-confidence study (Zeng et al., 2015) used the controls from the case-control study of asthma described below (Dong et al., 2013) and examined the association between PFBS exposure and serum lipids. The models were adjusted for age, gender, BMI, parental education level, exercise and ETS (environmental smoking) exposure. There was a statistically significant increase in total cholesterol ($\beta = 19.3$ mg/dL increase per 1 $\mu\text{g}/\text{l}$

⁶ <https://echa.europa.eu/registration-dossier/-/registered-dossier/10698/7/6/3>

increase in PFBS, 95% CI = 0.6–38.0, $p < 0.05$) but when PFBS exposure was analysed in quartiles, no exposure-response gradient was observed.

10 different PFASs were analysed in cord plasma samples (N=687) collected in Shanghai between 2011 and 2012, a region widely polluted with PFASs in China. The PFAS levels were compared with different maternal factors, including BMI, and it was found an inverse association with PFUnDA and PFDODA but a positive association with PFHxS and PFBS (Wang et al., 2016a).

4.4.3. Summary and discussion of repeated dose toxicity

The 28-day and the 90-day oral rat study show effect on liver, kidneys, stomach and hematological systems as PFBS-targets of toxicity. A dose dependent decrease in hemoglobin and hematocrit levels were observed in male rats with NOAEL determined to be 60 mg/kg bw/day (based on haematological parameters). In the 90-day study epithelial hyperplasia was observed in the kidney, and necrosis and hyperplasia/hyperkerosis was observed in the forestomach, both in male and female rats at 600 mg/kg bw/day. The NOAEL is 200 mg/kg bw/day in this study based on these microscopic effects seen in kidney and forestomach. 3M reported a NOAEL of 300 mg/kg bw/day for increased kidney weight. Furthermore, an investigation on mice indicated modest changes in lipid metabolism after 4-6 weeks daily exposure to PFBS (30 mg/kg bw/day). As supporting evidence for lipid metabolism distortion, an in vitro study showed that PFBS increased the adipogenesis of 3T3-L1 adipocytes significantly increasing the triglyceride level, in part, via MEK/ERK-dependent pathway. In addition, a Zebra fish study showed that PFBS can disrupt pancreatic organogenesis and perturb expression of genes involved in lipid metabolism (described in Chapter 5). These data together with human data linking PFAS to an increased BMI or changes in blood cholesterol or triglycerides, points towards a possible metabolic effect related to PFBS-exposure. However, other data in the literature may be less clear. In addition, a sub-chronic rat inhalation study with PFBS reported a transient effect on neurological parameters.

4.5. Mutagenicity

4.5.1. Non-human information

4.5.1.1. In vitro data

According to NICNAS (2005) the potassium salt of PFBS, KPFBS, was not found to be mutagenic to bacteria or clastogenic to Chinese Hamster Ovary-W-B1 cells under relevant test conditions.

A study by Eriksen et al. (2010) examined possible PFAS-induced reactive oxygen species (ROS) generation and oxidative DNA damage in HepG2 cells (a liver cell line). PFBS did not induce any ROS or oxidative DNA-damage which is seen for PFOS and PFOA.

Hence, the available information does not indicate that PFBS has in vitro mutagenic potential under testing conditions.

4.6. Carcinogenicity

4.6.1. Non-human information

4.6.1.1. In vivo data- Carcinogenicity: oral

In a repeated dose toxicity 90-day study, kidney hyperplasia was reported at 600 mg/kg bw/day. However, histopathological evaluation of the kidney by an independent expert reported that no consistent changes were seen in the kidneys (NICNAS, 2005; Lieder et al., 2009a). Increased incidence of necrosis in epithelial cells in the forestomach of male and female rats at 600 mg/kg bw/day was observed, including a thickening of mucosa due to hyperplasia and hyperkeratosis. In the two-generation study of Lieder et al. (2009b) treatment related microscopic changes were observed in the kidney of F1-generation female rats in the 300- and 1000 mg/kg bw/day dosage groups. The changes were minimal to moderate severity of hyperplasia in the epithelium of the inner medulla at the dose groups mentioned, respectively. The effects were seen in a two-generation study and a 2-year cancer study is not available.

4.7. Toxicity for reproduction

4.7.1. Effects on fertility

4.7.1.1. Non-human information

From the 2-generation study conducted by Lieder et al. (2009b) parental-generation (P) rats were dosed orally by gavage with 0, 30, 100, 300 and 1000mg KPFBS/kg bw/day for 10 weeks prior to and through mating (males and females), as well as during gestation and lactation (females only). First generation (F1) pups were dosed similarly, beginning at weaning. Second generation (F2) pups were not directly dosed but potentially exposed to PFBS through placental transfer and nursing, and the study was terminated 3 weeks after their birth, in which rats were exposed to gavage doses of potassium PFBS as high as 1,000 mg/kg bw/day. In this study, no alterations in fertility, estrus cycling, or histological alterations in female reproductive tissues were observed. However, exposure to 100 but not 300 or 1000 mg/kg bw/day through gestation showed a significant increase in diestrus cycling (>6 consecutive days). There were no treatment-related microscopic changes in sex organs of males although a decreased number of spermatids per gram testes and increased incidence of abnormal sperm were noted at the highest dose, although this was not seen in the F1-generation and was within historical variations. Mating and fertility parameters were seemingly unaffected in P and F1 generation (Lieder et al., 2009b). Furthermore, administration of up to 900 mg/kg bw/day PFBS to rats by gavage for 28 days did not cause any significant gross or microscopic alterations in primary or secondary sex organs from males or females (3M, 2001).

Caenorhabditis elegans (*C. elegans*) has been proposed as an alternative model to screen for reproductive toxicity in mammals (Bressers et al., 2018). It is however not officially accepted by EU regulatory bodies or validated by ECVAM. A recent study by Chen et al. (2018) shows that egg production and brood number of *C. elegans* decreased markedly following exposure to PFBS, although at a much higher exposure concentration than what was seen for PFOS. However, the internal concentrations of PFBS and PFOS were similar. The results indicate that PFBS exerts reproductive toxicity in *C. elegans* via germ-cell apoptosis resulting from elevated levels of reactive oxygen species. However, since the relevance to mammalian toxicity has not been evaluated properly the data must be interpreted with care.

4.7.1.2. Human information

In China and Taiwan, two studies examined different reproductive outcomes in women and men (Song et al., 2018; Zhou et al., 2016). Biomonitoring PFAAs in blood and semen samples from 103 male participants from Guangdong province in China was performed cross-sectionally to investigate potential link between PFAAs exposure and semen mobility in China. Negative correlations were significantly observed between sperm motility and PFBS concentrations in semen ($p < 0.01$), as well as for other PFASs in semen. (Song et al., 2018). Another study, (Zhou et al., 2016) in 225 Taiwanese adolescents, looked at associations between PFBS levels and reproductive hormones and they reported no clear correlations in males or females. However, both of these studies are considered of low-confidence and low sensitivity due to low exposure levels of PFBS and low participant numbers, and thus there is a lack of clear associations.

4.8. Developmental toxicity

4.8.1. Non-human information

In a two-generation reproductive toxicity study in rats administered 0, 30, 100, 300 and 1000 mg/kg bw/day of potassium PFBS (as reported above in more detail), there was no clear evidence of adverse effects on reproduction, fertility and lactation at the highest dose tested of 1000 mg/kg bw/day (Lieder et al., 2009b). The NOAEL for reproductive toxicity in males and female P (parent) generation is 1000 mg/kg bw/day. The P generation male and female rats LOAEL toxicity was 300 mg/kg bw/day based on treatment-related microscopic changes in the kidney and liver for P-generation males and kidneys for P-generation females, respectively. The mean number of liveborn F1 pups was statistically significantly decreased in the 30 mg/kg bw/day group, but this change was not dose-dependent. The viability index in F1 pups and the lactation index in F1 and F2 pups showed statistically significant changes at various doses but were not dose-dependent (Lieder et al., 2009b). Similarly, no clear effects were observed in delivery and litter parameters (e.g., implantations, litter sizes, live fetuses, corpora lutea, and early resorptions) following prenatal exposure from GDs 6 to 20 (Lieder et al., 2009b). F1 litter developmental NOAEL was 300 mg/kg bw/day based on body weight effects for males. The F1 generation male and female rats LOAEL was 300 mg/kg bw/day based on treatment-related microscopic changes in the kidney and liver for males and kidneys for females. No adverse effects were observed in second generation (F2) pups in doses as high as 1000 mg/kg bw/day. → *Klimisch reliability 1- well documented, guideline study.*

A recent publication, reporting a non guideline GLP-study performed by Feng et al. (2017), found that offspring mice, orally exposed before pregnancy to potassium PFBS (50, 200 and 500 mg/kg bw/day) from gestational day (GD)1-GD20, showed a delay in perinatal growth, vaginal opening and first estrus cycle, and diestrus was prolonged from 200 mg/kg bw/day. The study was conducted on twelve week old ICR Mice (China), 30 dams in each dose group were randomly signed to one of 3 experimental groups examining; 1. survival, growth, pubertal onset and ovarian and uterine development (50 offspring/10 dams), 2. Thyroid- and gonadal hormone measurements in offspring: PND1(n=30), PND30 (n=10), PND60 (n=10) and 3. plasma concentration. → *Klimisch reliability 2- well documented, non-guideline study.*

Ovarian and uterine size, as well as follicle and corpus luteum numbers were reduced in adult PFBS exposed offspring (PND 60). They exhibited fewer primordial follicles, primary follicles, secondary follicles, early antral follicles, antral follicles, and preovulatory follicles, as well as fewer corpora lutea at both 200 and 500 mg/kg bw/day compared to control (Feng et al., 2017). Importantly, no effects on the health (e.g., weight gain) of the exposed dams were observed at any dose (Feng et al., 2017). Furthermore, pubertal and adult

PFBS-offspring exhibited decreases in serum estrogen (E2) and progesterone (P4) levels with the elevation of luteinising hormone levels. These results indicate that prenatal PFBS exposure (≥ 200 mg/kg bw/day) causes delay in pubertal onset and reproductive organ development in female offspring mice. In addition, PFBS-dams exhibited significant decreases in total T4 and T3 levels and free T4 levels and increases in TSH levels, but no changes in E2 and P4 levels (described in Section 4.11.2). Serum-PFBS level were measured to be 74, 332 and 720 ng/mL at GD20 in dams exposed to 50, 200 and 500 mg/kg bw/day respectively. These results together with the effects mentioned above (under fertility), indicate that prenatal PFBS exposure (≥ 200 mg/kg/day) causes permanent hypothyroxinemia accompanied by deficits in perinatal growth, pubertal onset, and reproductive organ development in female mice. However, the study did not correlate the observed effects for the statistically significant decrease in pup body weight at the relevant dose levels. → *Klimisch reliability 2- well documented, not a guideline study.*

4.8.2. Human information

A recent study reported that prenatal exposure to PFBS was positively associated with the risk of preeclampsia and overall hypertensive disorders of pregnancy, HDP (Huang et al., 2019). PFASs were measured by liquid chromatography system coupled with tandem mass spectrometry in 687 umbilical cord plasma samples collected between 2011 and 2012 in Shanghai, China. Information on HDP including gestational hypertension and preeclampsia was collected from medical records. In this fully adjusted statistical model, PFBS from cord blood plasma had an increased odds of preeclampsia [adjusted odds ratio (AOR): 1.81, 95% confidence interval (CI): 1.03–3.17], and overall HDP (AOR: 1.64, 95% CI: 1.09–2.47). However, this study found that preeclampsia was not associated with PFOA and PFOS, which were different from that found in the US-based C8 Health Project which reported a probable link between PFOA and pregnancy-induced hypertension. In addition, PFBS was measured in cord plasma and not maternal plasma which may indicate a reverse causation since preeclampsia might affect liver and kidney functions leading to less secretion of PFBS and more accumulation in the body. Due to the short carbon-chain of PFBS high transmission through the placenta might be expected and cord plasma level may well reflect maternal level. In a recent human mother-child cohort study (Gao et al., 2019) similar concentrations of PFBS were detected in both cord blood and maternal blood, and in 70% of the cases PFBS was detected in both the cord and respective maternal sample.

Chen et al. (2019b) investigated the association between prenatal exposure to PFASs, including PFBS, and childhood adiposity at 5 years of age in a birth cohort study in Shanghai. The study involved 1140 pregnant women from 2012 to 2017. The median concentration of PFAS in the cord plasma ranged from 0.05 (PFBS) to 6.74 ng/mL (PFOA). The findings indicated that in girls, PFBS had a significant positive association with waist circumference and waist to height ratio in a multivariable linear regression model. Girls in the highest tertile of PFBS concentrations had more fat mass, as well as higher body fat percentage, waist circumference, and waist to height ratio compared to those in the lowest tertile. The authors concluded that high prenatal exposure to PFBS may be of concern for childhood adiposity.

4.8.3. Summary and discussion of reproductive toxicity

The two-generation reproduction study with PFBS in rats demonstrate that PFBS has a low potential to produce effects on reproduction or development in rats under testing conditions used. No clear adverse effects on functional aspects of reproduction were observed in either generation. In both the P- and F1-generation males, minimal to mild

hepatocellular hypertrophy in males and histomorphological changes in the kidney in both males and females was observed microscopically in the 300 and 1000 mg/kg bw/day KPFBS dose groups. Prenatal exposure to PFBS in mice indicated reproductive toxic effects on several parameters such as delay in perinatal growth and pubertal onset in addition to changes in reproductive organ development from 200 mg/kg bw/day.

4.9. Other effects: Neurotoxicity

4.9.1. Non-human information

4.9.1.1. In vitro data

In vitro studies exposing transformed neuronal PC12 cells (a standard in vitro model for neuronal development) with different fluorinated compounds showed that all compounds tested had an adverse effect on neuronal development of these cells, seemingly through different mechanisms (Slotkin et al., 2008). The rank order of adverse effects on PC12 cells was PFOSA>PFOS>PFBS≈PFOA. In the case of PFBS, PC12 cell differentiation was suppressed for two different neurotransmitter phenotypes. PFBS, unlike other fluorinated chemicals, induced a concentration dependent reduction of ChAT (choline acetyl transferase) and TH (tyrosine hydroxylase) activity needed for differentiation into two distinct neurotransmitter phenotypes, i.e. acetylcholine (Ach) and dopamine (DA), respectively.

A study by Liao et al. (2009) shows that PFASs exhibit adverse effects on cultured primary rat hippocampal neurons to various extents, which was generally dependent on the carbon chain length and functional group attached to the fully fluorinated alkyl chain. PFBS had a significant effect on calcium homeostasis by increasing voltage dependent calcium influx through calcium channels after exposure to high concentration (100 µM). In addition, exposure to 50 µM PFBS disturbed neuronal developmental processes by significantly reducing the total length of neurites and changing the pattern of the number of bifurcated neurites.

4.9.1.2. In vivo data

The 90-day oral rat study (test guideline OECD 408) of potassium PFBS included a FOB (Functional Observational Battery) and motor activity assessment but not a peripheral neuropathy assessment per se. The authors did not find any significant alterations in motor activity or performance on functional observation tests at doses up to 600 mg/kg bw/day (Lieder et al., 2009a).

However, neurological alterations were reported in an oral 28-day study in rats (NTP, 2019), where treated males differed from control males in a decrease in tail flick, rotorod latency and foot splay, but did not exhibit a dose-response at the doses tested. In contrast, treated females exhibited an increase in rotorod latency.

A recent study examined long-term potentiation (LTP) in rats after exposure to PFOS and its alternatives, such as PFBS, aiming to provide some evidence about the potential to affect cognitive ability. Different dosages of PFOS, PFHxS, PFBS and chlorinated polyfluorinated ether sulfonate (Cl-PFAES), were dosed to rats via acute intracerebroventricular injection. The field excitatory postsynaptic potential (fEPSP) amplitude of the input/output functions, paired-pulse facilitations, and LTP in vivo were recorded. PFOS and its alternatives including PFBS inhibited LTP in varying degrees without significant effects on the normal synaptic transmission. There was a higher potency of

PFHxS and PFOS than PFBS to inhibit LTP and points to the possibly higher neurotoxicity potential of the long carbon chain perfluoroalkyl compounds. The results suggest that acute exposure to PFOS and its alternatives, including PFBS, impaired the synaptic plasticity by a postsynaptic rather than a presynaptic mechanism (Zhang et al., 2016).

4.9.2. Summary and discussion of neurotoxicity

In summary, there are only very few studies providing some evidence that the nervous system is a sensitive target for PFBS. Clinical observations and morphological examinations of the nervous system in the 28-day study reported a decrease in tail flick latency test in males after PFBS exposure and acute PFBS exposure of rats had an inhibiting effect on LTP (although at a lower potency than PFOS) which indicate a potential to affect cognitive ability. In addition, PFBS has shown to disturb neuronal development in vitro in two separate studies. However, more studies are needed to verify the significance of these results.

4.10. Other effects: Immunotoxicity

4.10.1. Non-human information

4.10.1.1. In vitro

The release of the pro-inflammatory cytokines IL-6 and TNF- α was evaluated in liposaccharide stimulated primary human lymphocytes and in the THP-1 cell line, while the release of IL-10 and IFN- γ was evaluated in PAH stimulated human peripheral blood lymphocytes (Corsini et al., 2012). After exposure to several PFAS, TNF- α production was suppressed, and PFBS showed to inhibit IL-10 release. Mechanistic investigations demonstrated that inhibition of TNF- α release in THP-1 cells occurred at the transcriptional level. All PFAS, including PFBS, decreased LPS-induced NF- κ B activation. None of the PFAS tested, except PFOA, were able to activate PPAR α driven transcription in transiently transfected THP-1 cells, excluding a role for PPAR α in the immunomodulation as observed. PFBS and PFDA prevented LPS-induced I- κ B degradation. Overall, this study suggests that PFAS, including PFBS, affect NF- κ B activation, which directly suppresses cytokine secretion by immune cells.

The US EPA ToxCast program has tested potassium PFBS (PFBS-K). A positive activation of the G-protein coupled receptor (GPCR) LTB4r (Leukotriene receptor B4) was noted for PFBS-K. This GPCR NovaScreen assay reported in ToxCast, assess binding to the guinea pig LTB4r. Several other PFASs (i.e. PFOS, PFHxS, PFOA, PFHxA) show binding to this receptor in ToxCast as well. The functional relevance of this findings is still unclear, however it is known that this receptor is involved in pro-inflammatory responses and inhibition of this receptor is used in asthma prescribed drugs.

4.10.1.2. Respiratory effects in rats

Administration of PFBS at gavage doses of up to 900 mg/kg bw/day for 28 days or 600 mg/kg bw/day for 90 days had no significant effect on the gross or microscopic morphology of the lungs and trachea in rats (3M, 2001; Lieder et al., 2009a); no increases in nasal lesions were observed in the 90 day study (Lieder et al., 2009a).

4.10.2. Human information

Associations between different PFAS and various infectious diseases (i.e. common cold, lower respiratory tract infections, otitis media, pneumonia, respiratory syncytial virus infection and/or varicella) in children have been reported (Granum et al., 2013; Pennings et al., 2016; Dalsager et al., 2016; Goudarzi et al., 2017; Impinen et al., 2018). However, PFBS was not among the PFAS in these studies.

The relationship between PFAS and asthma- and allergy related outcomes has been investigated in several studies. The findings were not consistent between the different studies and PFBS was not a part of most of the investigations (Granum et al., 2013; Impinen et al., 2018; Smit et al., 2015). However, a case-control study on 231 asthmatic children and 225 non-asthmatic control children (aged 10–15 years) from Northern Taiwan showed that asthmatics had significantly higher serum PFAAs concentrations compared to the healthy controls (Dong et al., 2013). The adjusted odds ratios (OR) for asthma among those with the highest versus lowest quartile of PFAS exposure ranged from 1.81 (95% CI: 1.02, 3.23) for PFDoA, 1.90 (95% CI 1.08–3.37) for PFBS to 4.05 (95% CI: 2.21, 7.42) for PFOA. When stratified by gender, a greater number of significant associations between PFAAs and asthma outcomes were found in males than in females. Among males, adjusted odds ratios for asthma among those with the highest versus lowest quartile of PFAAs exposure ranged from 2.59 (95% CI: 1.14, 5.87) for the PFBS to 4.38 (95% CI: 2.02, 9.50) for PFOS.

To evaluate the association between serum PFAA concentrations and T-lymphocyte-related immunological markers of asthma in children, and to assess whether gender modified this association, serum concentrations of ten PFAAs and serum levels of TH1 [interferon (IFN)- γ , interleukin (IL)-2] and TH2 (IL-4 and IL-5) cytokines were measured (Dong et al., 2013; Zhu et al., 2016). The results showed that serum PFAAs, including PFBS, were associated positively with TH2 cytokines and inversely with TH1 cytokines among male asthmatics. In addition, a 16-kDa club cell secretory protein (Clara) (CC16), which plays a key role in anti-inflammatory and antioxidant functions in epithelial cells and a prominent biomarker for asthma, was measured in a later study (Zhou et al., 2017). A negative association between several PFAAs, except PFBS, and CC16 was observed. When effects of multiplicative interactions between asthma and individual PFAAs on the CC16 outcome were estimated among children, a marginally significant interaction effect ($p < 0.15$) on CC16 levels was found between asthma and PFOS, PFOA, PFBS and PFBS in all participants.

4.10.3. Summary and discussion of immunotoxicity

Taken together, *in vitro* studies as well as epidemiological studies indicate an association between PFAS and different immunological outcomes, with lower potency for the shorter chain PFBS. Only very few studies have analysed PFBS among the PFAAs and results show a weak relationship between PFBS and asthma-related outcomes. However, more studies are needed to verify these associations.

4.11. Other effects: Endocrine disruption

4.11.1. *In vitro*

An *in vitro* PFAS-exposure study of human cell lines such as MCF-7, H295R, LNCaP and MDA-kb2 was set up in order to study the underlying mechanisms of the reproductive toxic effects and in particular study the endocrine properties of different PFASs including PFBS (Behr et al., 2018). The obtained *in vitro* data suggested that the PFASs examined in this study displayed either a weak or no (anti)-estrogenic and no (anti)-androgenic activity in human cells and had only weak impact on steroid hormone secretion as well as on

steroidogenic gene expression. The steroidogenesis assay in H295R-cells was conducted following OECD TG 456. Effects were predominantly observed for PFOA and PFOS and not for their substitutes. Moreover, these effects were only observed at high concentrations, at least three to four orders of magnitude above human blood serum concentrations which are in the range of about 10 nM. PFBS was not found to have (anti)-estrogenic or (anti)-androgenic activity or effects on steroid hormone secretion or steroidogenic gene expression at concentrations up to 10 µM. The authors concluded that PFASs do not affect estrogen or androgen receptor signalling or steroid hormone biosynthesis at concentrations relevant to human exposure.

PFASs have been shown *in vitro* to disturb steroidogenesis in human placental choriocarcinoma cell line (JEG-3) by inhibiting CYP19 aromatase gene expression, increasing several membrane lipids (Gorrochategui et al., 2014). The study highlights the ability of the PFAS mixture to alter cellular lipid pattern at concentrations well below those that generate toxicity, and the potential for PFBS to significantly inhibit aromatase activity in placental cells (IC₅₀ of 68 µM).

The US EPA ToxCast program has tested potassium PFBS (KPFBS) for its ability to disrupt androgen, estrogen, thyroid or steroidogenesis pathways using a screening battery of endocrine bioactivity assays. ToxCast summary showed no significant activity in the tests conducted on these parameters.

In vitro studies are useful as supporting information and may suggest that the tested chemical act via a particular mode of action. However, *in vitro* assays do not reflect the true complex hormonal axis as present in a living organisms and results must be interpreted with care. In addition, several important endocrine mechanisms, e.g. additional mechanisms for disruption of thyroid hormone signaling, are currently not covered in the ToxCast assays.

4.11.2. *In vivo*

In the 28-day rat study by 3M in 2001 (3M, 2001) the thyroid weight and histology was reported to be normal after PFBS exposure. However, no thyroid hormones were measured. In the 90-day and two-generation reproductive rat study with PFBS-K by Lieder et al. (2009a and b) (reported in Sections 4.4 and 4.8, respectively), no levels of thyroid hormones were measured either. They reported however that no effects of PFBS was seen on thyroid histology in the 90-day rat study. Thyroid weight was not performed in this study nor in the 2-generation study which in fact was recommended in the guide line. Additional requirements on thyroid measurements have been added to the updated OECD guideline for the 90-day study from 2018. One statistical significant finding in the 2-generation study that may relate to ED disruption was a transient increase in number of pups showing ≥6 Days of diestrus cycle at the 100 mg/kg bw/day. → *Klimisch reliability 1- well documented, guideline studies according to the guidelines used.*

In the 28-day PFBS exposure study by NTP (2019), statistically significant dose-dependent decreases in total T3, total T4, and free T4 levels were reported in both male and female rats at all doses tested (≥62.6 mg/kg bw/day). The reported reductions in total T3 were up to -57% and -43% in male and female rats, -86% and -77% in free T4, and -97% and -71% in total T4, respectively. Thyroid gland weight, thyroid histopathology, and TSH levels were not changed after 28 days of PFBS exposure in male or female rats at up to 1,000 mg/kg bw/day (NTP, 2019). → *Klimisch reliability 1- well documented, guideline study.*

The study by Feng et al. (2017), described in more detail in Section 4.8.1, showed that when KPFBS (200 and 500 mg/kg bw/day) was orally administered to pregnant mice (10

dams in each group) on gestational days 1–20, their female offspring exhibited a statistically significant decrease in perinatal body weight gain, delay in eye opening, significant delay of vaginal opening and prolonged first estrus and diestrus. The ovarian and uterine size, as well as follicle and corpus luteum numbers were reduced in adult offspring. Furthermore, pubertal and adult offspring exhibited decreases in serum estrogen (E2) and progesterone (P4) levels with the elevation of luteinising hormone levels. Notably, decreases in serum total thyroxine (T4) and 3,3', 5-triiodothyronine (T3) levels were observed in fetal, pubertal, and adult offspring in conjunction with slight increases in thyroid-stimulating hormone (TSH) and thyrotropin-releasing hormone levels. In addition, dams exhibited a significant decrease in total T4 and T3 levels and free T4 levels and increases in TSH levels, but no changes in E2 and P4 levels. The BWs of GD20 PFBS-dams were not different from those of control dams, irrespective of dosage. These results may indicate that prenatal PFBS exposure (≥ 200 mg/kg bw/day) may cause hypothyroxinemia accompanied by deficits in perinatal growth, pubertal onset, and reproductive organ development in female mice. → *Klimisch reliability 2- well documented, non-guideline study.*

In support for this, both PFHxS (Ramshøj et al., 2016) and PFOS (Yu et al., 2009; Lau et al., 2003) have been found to reduce total T3 and/or total T4 level(s) in rodents leading to hypothyroxinemia in pups. PFHxS dose-dependently decreased dam T4 level on GD15 and pup T4 and T3 levels on PND 16/17 from 5 mg PFHxS/kg/day while gestational exposure to PFOS (3.2 mg PFOS/kg) alone significantly decreased T4 levels in pups on PNDs 21 and 35. Postnatal exposure to PFOS alone also induced T4 depression on PNDs 21 and 35. Thus, there seems to be a decrease at least in thyroxine (T4) levels in both offspring and dams exposed to either PFOS (Yu et al., 2009), PFHxS (Ramshøj et al., 2016) or PFBS (Feng et al., 2017) with decreasing potency from PFOS to PFBS.

Endocrine disturbances upon exposure to PFBS have also been observed in ecotoxicity studies, see Section 5.6.

4.11.3. Summary

In vitro studies as well as ToxCast screening show little or no support that the PFBS-K substance acts via the endocrine pathway with the limited tests available. However, *in vivo* studies give supporting evidence for possible endocrine disruptive properties of PFBS.

In mice a decrease in both T3 and T4 was seen after prenatal PFBS-exposure, together with deficits in perinatal growth, pubertal onset and reproductive organ development at concentrations ≥ 200 mg/kg bw/day (Feng et al., 2017). Similar patterns of decrease in total T3, total T4, and free T4 were observed in adult female and adult male rats from a 28-day study at doses ≥ 62.6 mg/kg bw/day. These decreases were of a concerning magnitude in both dams and offspring and they were shown to persist at least 60 days after gestation. Unfortunately, levels of thyroid hormones were not available in the two-generation reproductive rat study by Lieder et al. (2009b).

Taken together, there is sufficient evidence from different *in vivo* studies that the substance has the potential to perturb the hypothalamic–pituitary–thyroid axis as the findings seem to be consistent. *In vitro* studies are useful as supporting information and may suggest mode of action analyses but do not reflect the true complex hormonal axis as present in a living organism. In addition, several important endocrine mechanisms, e.g. for thyroid disruption, are currently not covered in the *in vitro* ToxCast assays. No information concerning these endpoints in humans is available, however, animal models are considered informative for evaluating the potential for thyroid effects of chemicals in humans (Zoeller et al., 2007). The thyroid seems therefore to be a potential target organ for PFBS toxicity in animals as well as humans.

4.12. Summary and discussion of human exposure and health hazard assessment

In blood PFBS binds to albumin, is transported through the body and is found in many different tissues. PFBS has also been detected in human cord blood. PFBS binds to a number of other proteins, including receptors. A chain-length of C4 is generally reported to have a low binding affinity to albumin and other proteins, however, PFAAs containing the sulfonic acid group may bind stronger to proteins than their carboxylic acid counterpart. PFBS may also bind to several different transporter polypeptides (NTCP, ASBT and OATPs) which are all capable of contributing to the enterohepatic circulation and thus extended human serum elimination half-lives of PFBS and other PFAAs.

Since PFBS has a relatively high aqueous solubility, it can be found in different crops and in drinking water, and it can be taken up in the food chain, which may result in increased exposure. Household dust has also a relevant source of exposure for humans to PFBS.

Concentrations of PFBS in human blood have been determined in a number of studies. In the studies many samples were below the quantification limit, and in most studies the maximum concentration was below 0.5 ng/mL. However, in some studies concentrations up to around 5 ng/mL were observed. Studies of people with known elevated exposure to PFBS, for example via drinking water or in an occupational setting, have demonstrated that increased exposure leads to higher PFBS concentrations in the blood, even though the half-life of PFBS is relatively short (around 1 month) compared to for instance PFOA (2 – 4 years).

Table 29 shows an overview of health effects caused by PFBS.

Table 29: Overview of PFBS-induced health effects

Target organ	Type of effect	NOAEL/LOAEL	Reference	Klimisch score
Liver	Weight increase	LOAEL: 600 mg/kg bw/day NOAEL: 300 mg/kg bw/day	3M, 2001 Lieder et al., 2009a	1
	Modest changes in lipid metabolism	LOAEL: 30 mg/kg bw/day	Bijland et al., 2011	3
Kidney	Absolute and relative kidney weight, hyperplasia was observed in the kidney, and necrosis and hyperplasia/hyperkerosis was observed in the forestomach, both in male and female rats	LOAEL: 600 mg/kg bw/day NOAEL: 200 mg/kg bw/day	Lieder et al., 2009a	1
Haematological system	Decrease in hemoglobin and hematocrit levels	LOAEL: 200 mg/kg bw/day NOAEL (male): 60 mg/kg bw/day	Lieder et al., 2009a	1
Repro/Developmental Tox	Perinatal growth, pubertal onset and reproductive organ development	LOAEL: 200 mg/kg bw/day NOAEL: 50 mg/kg bw/day	Feng et al., 2017	2
	Delay in perinatal growth, vaginal opening and first estrus cycle, and diestrus was prolonged Significant increase in diestrus cycling	LOAEL: 100 mg/kg bw/day NOAEL: 30 mg/kg bw/day (no effects were seen at 300 or 1000 mg/kg bw/day)	Lieder et al., 2009b	1
Endocrine effects	Disturbed thyroid axis:	LOAEL: 62.6 mg/kg bw/day in rats	NTP 2019	1
	Decreases in total T4 and T3 levels and free T4 levels in rats and mouse and, in addition for the mice study, increases in TSH levels (Feng et al., 2017)	LOAEL: 200 mg/kg bw/day NOAEL: 50 mg/kg bw/day (74 ng/mL in plasma) in mice	Feng et al., 2017	2

Different studies in rodents, supported by in vitro studies or ecotoxicological studies, provide evidence for adverse health effects. Repeated dose toxicity studies in rat have shown PFBS-mediated effects on liver, kidneys, and hematological systems, although at relatively high doses. A dose dependent decrease in hemoglobin and hematocrit levels was observed in male rats with a NOAEL determined to be 60 mg/kg bw/ day (90 days) as the most sensitive.

A study on mice indicated modest changes in lipid metabolism after 4-6 weeks daily exposure to PFBS (30 mg/ kg bw/day). Only a few studies have, as of yet, seen similar effects on lipid metabolism. Together with human data indicating that PFBS, as well as other PFAAs, have been associated to increased BMI or changes in blood cholesterol and triglycerides, these data may point towards an effect of PFBS on metabolic disorder or disturbance of the lipid metabolism.

Evidence for genotoxic or mutagenic effects of PFBS has not been found. The current knowledge of carcinogenic effects of PFBS is insufficient. However, some indications for carcinogenicity were found in a study showing minimal to moderate effects on the kidneys. Indications for reproductive toxic effects of PFBS were found when prenatal exposure to PFBS was linked to a delay in pubertal onset and reproductive organ development in female offspring mice.

Other effects of PFBS, such as neurotoxicity and immunotoxicity, have also been reported. However, there are very few studies providing evidence that the nervous system or the immune system are sensitive targets for PFBS.

Endocrine disturbances, such as decrease in T3 and T4 levels, have been shown after prenatal PFBS-exposure, both in mother and their offspring, together with deficits in perinatal growth, pubertal onset and reproductive organ development in mice at doses ≥ 200 mg/kg bw/day. These effects are supported by data from ecotoxicological studies. Hence, PFBS exposure seems to disturb the levels of thyroid hormone in intact organisms. In addition, PFBS exposure of frogs resulted in increased expression of the estrogen and androgen receptor, as well as providing effects on sexual development.

Taken together, toxicological data obtained for PFBS point towards similar health effects as seen for other PFAAs, although the potency seems to be lower for PFBS than for the C6 or C8 counterparts for several of the endpoints. This was seen in a report on relative toxic potency to liver for 20 different PFASs, investigated by Zeilmaker et al. (2018). PFBS was found to have lower liver toxicity compared to C6 and C8 PFASs and C4 to C18 PFCAs. However, in general, the different endpoints have not necessarily been adjusted for the internal PFBS-concentration in the specific target organ. Due to the extreme persistency of PFBS in the environment and documented uptake and distribution in the human body, continuous exposure through drinking water, food as well as dust may cause higher levels in the blood, leading to unforeseen and unwanted health challenges. In addition, the combinatorial effect when exposed to a mixture of PFASs is of concern. The co-exposure of PFBS and other very persistent fluorochemicals present in the environment may lead to a combination of effects on human health.

5. Environmental hazard assessment

5.1. Aquatic compartment (including sediment)

5.1.1. Fish

5.1.1.1. Short-term toxicity to fish

Summaries of two standard acute toxicity tests are available in the registration dossier for KPFBS. Both tests followed the OECD TG 203 guideline. Fathead minnows were exposed to nominal concentrations (0, 204 (actual 220), g/L, 408 (actual 437) mg/L, 816 (actual 888) mg/L, 1632 (actual 1655) mg/L and 3263 (actual 3341) mg/L) of PFBS for 96 hours in a static system. Two negative control groups were kept in blank test medium. No mortality was observed at concentrations up to 888 mg/L, however 30% cumulative mortality was observed at 1655 mg/L and 100 % cumulative mortality was observed at 3341 mg/L. The 96-hour LC₅₀ of KPFBS to fathead minnow was 1938 mg/L. → *Klimisch reliability 1- well documented, guideline study.*

Furthermore, bluegill sunfish were exposed to nominal concentrations between 612 (actual 629) mg/L to 9790 (actual 9433) mg/L of KPFBS in a static test. Control fish were kept in blank test medium. The NOEC for bluegill sunfish was 2715 mg/L. At 5252 mg/L 15% cumulative mortality was observed, and 100% mortality was observed at the highest exposure concentration. Reported LC₅₀ value for bluegill sunfish was 6452 mg/L. → *Klimisch reliability 1- well documented, guideline study.*

Hagenaars et al. (2011): A zebrafish study assessed the structure activity relationship of PFBS and three other PFASs following a prolonged zebrafish early life stage test according to OECD 236 guideline Fish embryo toxicity test (FET) The test procedure is previously described by Nagel (2002). Normal fertilised zebrafish eggs were exposed to nominal concentrations (50, 100, 250, 500, 1000 and 3000 mg/L) of PFBS potassium salt. Groups of twenty normally shaped fertilised eggs per exposure concentration were divided over a 24-well plate and each egg was placed individually in 2 mL of the test solution. Remaining four wells were filled with clean water and used for control eggs. 3,4-dichloroaniline (3.7 mg/L) was used as a positive control and resulted in mortality between 72% and 92%. Two replicate plates were used for each exposure concentration (i.e. 40 embryos per exposure concentration). The eggs were exposed until 120 hours post fertilisation (hpf). The embryos were checked for mortality at 8 hpf, and for mortality and hatching every 12 h until 120 hpf to include each developmental stage. Heart rate was recorded at 48 hpf (non-hatched) and 72 hpf (hatched). Malformations of the head were apparent in exposed larvae starting from 96 hpf. Another effect was uninflated swim bladder which resulted in abnormal swimming. The results also demonstrated significant altered heart rates in embryos exposed to PFBS. At 3000 mg/L PFBS significantly increased heart rates were detected after 48 hpf, however the heart rates significantly decreased after 72 hpf. The EC₅₀ value of PFBS was 1529.32 mg/L and was a regression estimate of the concentrations at which 50 % of the embryos showed an effect (mortality or malformation). There was a low mortality rate of PFBS, reported LC₅₀ was >3000 mg/L. NOEC at 96 hpf and 120 hpf were 500 mg/L and 250 mg/L respectively (Table 30). *Klimisch reliability 2- modified guideline study.*

Ulhaq et al. (2013a): Zebrafish embryos were exposed to nominal (actual) concentrations (blank, 10 mg/L, 30 mg/L, 100 mg/L, 1000 mg/L and 3000 mg/L) of PFBS and six other PFASs following OECD TG 236 with an extended exposure period and additional sublethal endpoint observations. Normally developed fertilised eggs were distributed on a 48 well plate along with 750 µL of the exposure medium (PFASs and reconstituted water). A total of 24 embryos were tested per PFAS concentration, as well as 24 embryos in the water control group. The embryos were exposed until 144 hpf.

Observations of lethal and sublethal endpoints were made after 24, 48, 120 and 144 hpf. Heart rate was recorded at 48 hpf. Effects observed in the literature following PFBS exposure were affected heart rate and pericardial oedema, which is a sign of compromised cardiac output. A combined EC₅₀ (450 mg/L) was determined for lethal and non-lethal effects due to inadequate statistical power. NOEC (heart rate) for PFBS was 300 mg/L. Similar sublethal effects in zebrafish embryos following exposure to other PFASs, such as PFOA and PFOS, have been observed in the studies by Zheng et al. (2012) and Hagenaaers et al. (2011). *Klimisch reliability 2- modified guideline study.*

Ulhaq et al. (2013b): Locomotor behavior of zebrafish larvae was also assessed on the same exposed individuals described in the previous section. Sublethal behavioural endpoints were recorded at 144 hpf using an automated video tracking system. The effect on larvae activity was determined both in light and darkness. The reported EC₅₀ (450 mg/L) in this study was based on the combined sublethal and lethal embryotoxicity effect data from the study described in the previous paragraph (Ulhaq et al., 2013a; Ulhaq et al., 2013b). Overall activity was reduced at 1000 mg/L PFBS, but active swimming speed was increased to that of the control group. Compared to the control group, there was a reduction in overall activity in the group exposed to the highest concentration of PFBS (3000 mg/L). This effect is considered as a disturbance in the behavioural response pattern in zebrafish larvae. In this study, PFBS was the compound that exhibited the highest impact on the tested behavioural endpoints. Exposure to PFBS was positively correlated with active swimming speed and negatively correlated to all the other endpoints. This indicates that PFBS can cause disturbances in zebrafish behaviour that might imply ecological consequences for wild fish. *Klimisch reliability 2- modified guideline study.*

Sant et al., 2019: Embryos were exposed to PFBS (0, 4.8 mg/L and 9.6 mg/L) daily, beginning at 1-day post fertilisation (dpf) until 4 and 7 dpf. They were examined for viability following the OECD Fish Embryo Acute Toxicity Test Guidelines No. 236 (OECD, 2013), and examined under microscope for hatching at 3 dpf and swim bladder inflation at 4 dpf. Incidences of fish with significantly stunted growth and truncated exocrine pancreas length was significantly increased after PFBS exposure, although these two effects occurred independently. Pancreatic islet morphology revealed an increased incidence of hypomorphic islets (areas lower than the 1st percentile of controls) and an elevated occurrence of fragmented islets. RNA Seq data (4 dpf) also identified disruptions in regulation of lipid homeostasis. The findings demonstrate that PFBS can disrupt pancreatic organogenesis and perturb expression of genes involved in lipid metabolism. The overall impact is not as severe as when compared to findings with embryonic PFOS exposures, as reported by Sant et al. (2017). *Klimisch reliability 2- modified guideline study.*

Table 30: Overview of short-term effects on fish

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
Fathead minnows	96 h	Mortality	LC ₅₀	1938 mg/L	OECD 203	ECHA dissemination website; NICNAS, 2005
Bluegill sunfish	96 h	Mortality	LC ₅₀	6452 mg/L	OECD 203	ECHA dissemination website; NICNAS, 2005

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
Zebrafish larvae	96h/120 hpf	Lethal, sublethal and developmental	LC ₅₀ EC ₅₀ NOEC	>3000 mg/L 1900.78/ 1529.32 mg/L 500/ 250 mg/L	OECD 236	Hagenaars et al., 2011
Zebrafish larvae	144 hpf	Developmental Heart rate	EC ₅₀ LC ₅₀ NOEC	450 mg/L 1500 mg/L 300 mg/L	OECD 236	Ulhaq et al., 2013a
Zebrafish larvae	144 hpf	Locomotor behaviour	EC ₅₀	450 mg/L		Ulhaq et al., 2013b
Zebrafish	1dpf, 4dpf and 7dpf (day post fertilisation)	Viability and organ development	LC ₅₀ (est.)	393.13 mg/L (1310 µM)	OECD 236	Sant et al., 2019

In summary, exposure to high concentrations of PFBS resulted in disturbances in heart rate, morphological malformation and altered swimming behavior in zebra fish. In the presented acute and long-term studies, an overall low mortality rate was observed in exposed fish. In the currently available literature, there is a large range in observed effect concentrations for zebra fish.

5.1.1.2. Long-term toxicity to fish

Chen et al. (2018a): F0 marine medaka (*Oryzias melastigma*) eggs were exposed to PFBS (0, 1, 3, and 10 µg/L, nominal) until sexual maturity in a semi static system with daily water renewal to maintain the appropriate PFBS concentrations. Actual measured concentrations of PFBS were 1.0 ± 0.1 µg/L, 2.9 ± 0.1 µg/L, and 9.5 ± 0.3 µg/L, respectively. DMSO (<0.001% v/v) was used as a carrier solvent. Nominal (1, 3, 10 µg/L) and measured (1±0.1, 2.9±0.1 and 9.5±0.3 µg/L) concentrations did not deviate more than 10% during the full duration of exposure. The medaka juvenile and adults were fed twice daily with flake food and newly hatched *Artemia nauplii*. *O. melastigma* is not a widely used OECD model fish but has been used in some ecotox studies as a model teleost in marine toxicology, probably because it is closely related to Japanese medaka (*Oryzias latipes*) which is a well described test model. Exposure concentrations used in the study are environmentally realistic considering the high occurrence of PFBS in leachate of landfill sites in Singapore and Ireland, as well as near point sources in China (Bao et al., 2019; Harrad et al., 2019; Yin et al., 2017).

O. melastigma exposure was conducted in 100 mL exposure medium for the first 30 dpf, then 4 L medium until 2 months, and then 20 L medium for the last 4 months of exposure (until sexual maturity at 6 months). Three replicates of 150 embryos were used for each exposure concentration. F0 fish were then dissected for the tissues. F1 eggs were collected by pairing adult F0 medaka and cultured in clean seawater until the F1 fish were 6 months old, when the F1 fish were dissected. Prior to dissection, F1 fish were paired to produce F2 egg which were collected and cultured in clean seawater until 15 dpf to evaluate impaired early development. The effects on thyroïdal axis, TH levels, TBG contents, expression of genes along the HPT axis, body weight and length were investigated to elucidate multigenerational thyroid disrupting potency of PFBS across three generations (F0, F1 and F2).

Body weight and length of sexually mature female and male fish of the F0 generation were significantly decreased in 2.9 and 9.5 µg/L PFBS exposure groups when compared to the control. Reduced body length was observed in male medaka exposed to 2.9 and 9.5 µg/L PFBS when compared to the control. Parental exposure (F0) to 9.5 µg/L PFBS caused decreased body weight in F1 adult male and female medaka, when compared to the control. F1 fish from exposed parents also showed decreased body length. A significant increase in brain somatic index (BSI) levels for the F1 generation after exposure to 1.0 (males and females) and 9.5 (females) compared to control was observed (supplementary data).

Hatching of F1 larvae was significantly delayed following parental PFBS exposure (at 1.0 µg/L exposure group, $P < 0.05$). Body weight of both sexes and body length of male medaka were significantly ($P < 0.05$) decreased in 2.9 µg/L and 9.5 µg/L exposure groups relative to the control group, NOEC 1.0 µg/L (Chen et al., 2018a). The observed effects in this study indicate that PFBS can impair development in fish. See Section 5.6 for further discussions. *Klimisch score 2 – reliable, non-guideline study.*

Chen et al. (2018b): The objective of the second study by Chen et al. (2018b) was to investigate the toxic effects of PFBS on vision in marine medaka. Culture and life-cycle exposure were performed according to the study described in the previous section (Chen et al., 2018a). The effects measured in the eyes were PFBS accumulation, contents of various neurotransmitters, enzyme activity in eyes, AChE activity, MAO activity and protein expression. Eyes from five fish of the same sex were pooled as a replicate. PFBS exposure resulted in concentration dependent accumulation of PFBS in the eye tissue, and impaired both visual development and function in a sex-dependent manner. The visual systems of male medakas appeared to be more severely impacted by PFBS compared to the females. In exposed females, eye weight was significantly decreased, and water content in eyes increased. Concentrations of PFBS in the eyes increased in a dose dependent manner. Multiple neural signalling processes were perturbed by PFBS life-cycle exposure, including cholinergic, glutamergic, GABAergic and monoaminergic signals. Levels of norepinephrine and epinephrine in the eyes were increased, which may adaptively decrease the intraocular hypertension. PFBS exposure also adversely affected expressions of eye proteins that were related to visual functions and motor activity, and thus the muscular control of the eyes of the fish. These results highlight the eye as a sensitive organ to environmental pollutants. Impaired visual functions perception is likely to reduce survival of wildlife. The findings in this study indicate a direct effect of PFBS on visual development and function through life span of marine medaka, Chen et al. (2018b). Thyroid hormones tightly regulate the development and differentiation of visual systems in vertebrates (Bohnsack and Kahana, 2013; Dong et al., 2014), see Section 5.6 for further discussions. *Klimisch score 2 – reliable, non-guideline study.*

Chen et al. (2018c): Furthermore, Chen et al. performed a study on the effects of PFBS exposure on the gut microbiota of marine medaka (methods are described in previous sections). It was found that even at environmentally realistic concentrations, PFBS caused dysbiosis of gut microbiota which was durable in parents and persisted in the offspring.

Decreased relative abundances of *Cetobacterium* were observed in intestines of F1 males derived from parents exposed to 1.0 µg/L PFBS. Fluctuations in abundances of *Cetobacterium* might affect the supply of vitamin B12 in the intestine, which could result in deficiencies and mental disorders. *Klimisch score 2 – reliable, non-guideline study.*

Chen et al. (2019a): Furthermore, Chen et al. investigated the effects on the reproductive system and the underlying mechanisms related to endocrine disruption and transgenerational toxicities after a life-cycle exposure of parent marine medaka to environmentally realistic concentrations of PFBS, as previously described by Chen et al. (2018a). Exposure of the F0 generation to PFBS started at the embryonic stage and continued until sexual maturity (6 months). F0 adult medaka were dissected at 6 months. Some F0 adults were depurated for 2 months in clean seawater to monitor the PFBS elimination rate. The sex ratio of medaka fish was determined based on the development of secondary sex characteristics. To calculate egg production of the F0 generation, spawn was collected daily during the final 2 weeks of the exposure period.

Hardness of the egg envelope was significantly increased in eggs from parents exposed to 2.9 (P<0.01) and 9.5 (P<0.05) µg/L PFBS. Eggs spawn of the F1 generation in the 9.5 µg/L exposure group exhibited a significant (P<0.05) decrease in weight compared with that of the control group. F1 larvae mortality increased significantly after parental exposure to 2.9 (P<0.001) and 9.5 (P<0.5) µg/L PFBS. Furthermore, 0.2, 0.9 and 2.9 ng/g dw PFBS was transferred from exposed parents to F1 eggs in the 1.0, 2.9 and 9.5 µg/L exposure groups. However, no residual PFBS was detected in the F1 fish after depuration in clean seawater until sexual maturity. PFBS exposure was also shown to result in a skewed sex ratio in fish toward male dominance (LOEC 9.5 µg/L, NOEC 2.9 µg/L). Furthermore, the gonado somatic index and the oogenesis of female fish was lowered (LOEC 9.5 µg/L, NOEC 2.9 µg/L) and decrease in egg production (LOEC 2.9 µg/L, NOEC 1.0 µg/L). Additionally, endocrine disruption through the hypothalamus-pituitary-gonad axis was induced by PFBS exposure. Some observed effects were skewed sex ratio toward male dominance, increased larval mortality and impaired reproductive functions of female fish, see Section 5.6. *Klimisch score 2 – reliable, non-guideline study.*

Table 31: Summary of long-term effects on fish

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
Marine medaka	Life cycle	Body weight	LOEC NOEC	2.9 µg/L 1.0 µg/L	No test guideline. Exposure of F0 from egg until sexual maturity	Chen et al., 2018a
Marine medaka (male)	Life cycle	Body length (male)	LOEC NOEC	2.9 µg/L 1.0 µg/L	No test guideline. Exposure of F0 from egg until sexual maturity	Chen et al., 2018a
Marine medaka	Life cycle	Eye effects	NOEC	Males: 1.0 µg/L: Water percentage per eye, ACh content in eyes Females: 1.0 µg/L: ACh	No test guideline. Exposure of F0 from egg until sexual maturity	Chen et al., 2018b

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
				content in eyes 2.9 µg/L: choline, glutamate and GABA content in eyes of female fish, MAO activity in eyes of female fish		
Marine medaka	Life cycle	Eye weight	NOEC	2.9 µg/L	No test guideline. Exposure of F0 from egg until sexual maturity	Chen et al., 2018b
Marine medaka	Life cycle	Thyroid disruption	LOEC	1.0 µg/L	No test guideline. Exposure of F0 from egg until sexual maturity	Chen et al., 2018a
Marine medaka	Life cycle	Larvae mortality	LOEC NOEC	2.9 µg/L 1.0 µg/L	No test guideline. Exposure of F0 from embryo until sexual maturity (6 months)	Chen et al., 2019a
Marine medaka	Life cycle	Dysbiosis of gut microbiota	LOEC	1.0 µg/L	No test guideline. Exposure of F0 from egg until sexual maturity	Chen et al., 2018c

In summary, it has been demonstrated that long term exposure to PFBS may lead to a range of different effects on marine medaka. Exposure of marine medaka from eggs to sexual maturity lead to significantly increased F1 larvae mortality, decreased body weight of both sexes and decreased body length of male medaka, developmental and functional defects in eyes, and dysbiosis of gut microbiota durable in parents and persisting in the offspring. The long-term no-observed effect concentration (NOEC) for marine medaka is 1.0 µg/L based on reduced body weight and length observed in a life cycle test. The NOEC based on anomalies in visual function was 1.0 µg/L.

5.1.2. Aquatic invertebrates

5.1.2.1. Short-term toxicity to aquatic invertebrates

A 96 hour acute toxicity test performed on mysid shrimp (*Americamysis bahia*, former scientific name *Mysidopsis bahia*) is described in the registration dossier for KPFBS and reviewed by NICNAS (2005). The test was conducted following EPA OPPTS 850.1035 (Mysid Acute Toxicity Test) in 2000 mL polyethylene buckets containing 1500 mL of test or control medium and nominal concentrations ranged from 31 to 1000 mg/L KPFBS. Negative controls were kept in blank test medium. The mysids were subjected to a photoperiod of 16 h light and 8 h dark, temperature was held at 25 ± 2 °C, and measurements of pH ranged from 8.0 to 8.3. Observations were made at 4, 24, 48, 72 and 96 hours after test initiation. The 96 h LC₅₀ of mysid shrimp was 372 mg/L. → *Klimisch reliability 1- well documented, guideline study.*

A 48 hour immobilisation test was performed on *Daphnia magna* according to OECD TG 202 (ECHA dissemination website; NICNAS, 2005). Analytical exposure concentrations were < LOQ, 234 mg/l, 470 mg/L, 886 mg/L, 1707 mg/L and 3767 mg/L. Negative controls were kept in blank test medium. The daphnids were subjected to a photoperiod of 16 h light and 8 h dark, the temperature was held at 20 ± 1 °C, and measurements of pH ranged from 7.9 to 8.5. Observations were made after 4, 24 and 48 hours after initiation to determine the numbers of mortalities and immobile organisms. The EC₅₀ (48 h) was 2183 mg/L based on immobilisation of *Daphnia magna*. → *Klimisch reliability 1- well documented, guideline study.*

Table 32: Summary of short-term effects on aquatic invertebrates

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
Mysid shrimp	96 h	Lethality	LC ₅₀	372 mg/L	EPA OPPTS 80.1035	ECHA dissemination website; NICNAS, 2005
<i>Daphnia magna</i>	48 h	Immobilisation	EC ₅₀	2183 mg/L	OECD 202	ECHA dissemination website; NICNAS, 2005

5.1.2.2. Long-term toxicity to aquatic invertebrates

The chronic toxicity of KPFBS was determined in a 21 day *Daphnia magna* life-cycle chronic toxicity test equivalent to OECD guideline TG 211. The study is described in the registration dossier for KPFBS. The daphnids were exposed to six different concentrations of KPFBS (nominal/actual concentrations were 63/60, 125/121, 250/247, 500/502, 1000/995 and 2000/1876 mg/L). Negative controls were kept in blank test medium. Temperature was held at 20 ± 1 °C except for two occasions when the temperature was 18 °C. The daphnids were subjected to a photoperiod of 16 h light and 8 h dark, temperature was held at 20 ± 1 °C, and measurements of pH ranged from 7.9 to 8.5. The daphnid neonates were less than 24 hours at test initiation. The first generation was observed daily for survival, onset of reproduction, number of neonates and signs of toxicity. Length and dry weight of surviving first generation daphnids was determined. The 21 day NOEC was 502 mg/L, while LOEC was 995 mg/L. The study showed that the chemical was practically non-toxic

to daphnids after chronic exposure. → *Klimisch reliability 1- well documented, guideline study.*

Table 33: Summary of long-term effects in aquatic invertebrates

Species	Duration	Endpoint	Effect level	Conc.	Comment	Reference
<i>Daphnia magna</i>	21 days	Several endpoints, see text	NOEC LOEC	502 mg/L 995 mg/L	OECD TG 211	ECHA dissemination website; NICNAS, 2005

5.1.3. Algae and aquatic plants

A 96-h EbC₅₀ (algal biomass) and ErC₅₀ (growth) of 2347 mg/L and 5733 mg/L study, following OECD TG 201, has been reported for *Raphidocelis subcapitata* (formerly formerly known by the scientific names *Selenastrum capricornutum* and *Pseudokirchneriella subcapitata*) on the ECHA dissemination website and is also reviewed by NICNAS (2005). The EbC₅₀ and ErC₅₀ values are the concentration at which 50% reduction of biomass or growth rate is observed, respectively. The test was carried out by exposing *Selenastrum capricornutum* to nominal concentrations of KPFBS (nominal/ actual concentrations; 313/285, 625/563, 1250/1077, 2500/2216, 5000/4561 and 10000/9478 mg/L) for 96 h. Negative controls were kept in blank test medium. Temperature was held at 24 ± 2 °C and the algae were held under continuous cool-white fluorescent lighting. The effect of PFBS exposure on cell count, cell morphology, cell densities, area under growth curves, growth rates and percent inhibition values were determined (NICNAS, 2005). Based on the observed results in this study, the PFBS potassium salt was practically non-toxic to algae. → *Klimisch reliability 1- well documented, guideline study.*

Rosal et al. (2010): The toxicity of PFBS towards *P. subcapitata* was also examined. Chronic toxicity was determined following an algal growth inhibition test according to OECD TG 201 *P. subcapitata* open system, using 96-well microplate in which the algae were cultured in a total volume of 200 µL. Low toxicity was observed, with 37 % growth inhibition at 20250 mg/L. → *Klimisch reliability 2 - comparable to guideline study with acceptable restrictions*

Table 34: Summary of effects on algae

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
<i>Raphidocelis subcapitata</i>	96 h	Biomass	EbC ₅₀	2347 mg/L	OECD 201	ECHA dissemination website; NICNAS, 2005
		Growth	ErC ₅₀	5733 mg/L		
<i>Raphidocelis subcapitata</i>	72 h	Growth	EC ₅₀	>20 250 mg/L	37 % growth inhibition	Rosal et al., 2010

5.1.4. Sediment organisms

Stefani et al. (2014): A multigeneration toxicity test on *Chironomus riparius* aimed to investigate the evolutionary consequences of exposure to PFBS and two other perfluoroalkyl substances. The test was performed according to OECD TG 233, but extended to 10 generations exposed at a low concentration (nominal concentration of 10 µg/L). Native *C. riparius* L3 larvae were collected and bred until emergence and egg deposition. Six-hundred larvae were bred per treatment and per generation until emergence and egg deposition under a nominal concentration of 10 µg/L of contaminants. New-born larvae were used to start the next generation. Total exposure time was 300 days and the study followed ten generations until emergence of adults in generation ten. The test temperature was 20±1 °C and measured pH was 7.8-8.2. Exposure to PFBS resulted in an increased mutation rate based on five microsatellite loci, which indicated a stronger genetic variability in the PFBS exposed population compared that of controls. This may indicate a potential risk of mutational load caused by exposure to PFBS. The study provides indication of long-term relevant population effects on *C. riparius* caused by PFBS. → *Klimisch reliability 2 - comparable to guideline study with acceptable restrictions. Test extended to 10 generations at low concentration.*

Table 35: Summary of effects on sediment organisms

Species	Duration	Endpoint	Effect level	Exposure concentration	Comment	Reference
<i>Chironomus riparius</i>	10 generations	Population effect	Not assessed	10 µg/L (exposure concentration)	Extended OECD 233	Stefani et al., 2014

5.1.5. Other aquatic organisms

The toxicity of PFBS was examined by using two bioluminescence inhibition assays on the marine bacterium *Vibrio fischeri* and the cyanobacterial recombinant strain *Anabeana CPB4337* (Rosal et al., 2010). Bioassays with the photo-luminescent bacteria *Vibrio fischeri* were performed according to ISO 11348-3 standard protocol (International Organization for Standardization, 2007). The incubation period for this assay was 15 min. This bioassay measures the decrease in bioluminescence induced in the cell metabolism due to the presence of a toxic substance. A third bioassay was performed using the recombinant bioluminescent cyanobacterium *Anabeana CPB4337* and was based on the inhibition of constitutive luminescence caused by the presence of any toxic substance (Rodea-Palomares et al., 2009). Low toxicity was observed in *V. fischeri* and *Anabeana*, with effect concentrations (EC₅₀) of 17520 mg/L and 8386 mg/L respectively (Rosal et al., 2010). → *Klimisch reliability 2 - comparable to guideline study with acceptable restrictions*

Table 36: Summary of effects on marine microorganisms

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
<i>Vibrio fischeri</i>	15 min	Bioluminescence inhibition	EC ₅₀	17520 mg/L	ISO 11348-3	Rosal et al., 2010
<i>Anabeana CPB4337</i>	15 min	Bioluminescence inhibition	EC ₅₀	8386 mg/L	ISO 11348-3	Rosal et al., 2010

Lou et al. (2013): A study on *Xenopus laevis* aimed to investigate comparatively adverse effects of longterm exposure to PFOS and PFBS on the growth and sexual development of amphibians using *Xenopus laevis*, which is the most used model amphibian species in toxicological studies (Cevasco et al., 2008). Adult *X. laevis* female and male (3 years old) were raised separately in glass tanks and fed with chopped pork liver and commercial amphibian diet three times a week. Breeding was induced in a pair of *X. laevis* by injecting human chorionic gonadotropin. Fertilized eggs were incubated. On the fifth day postfertilization, tadpoles were exposed to PFOS (0.1; 1; 100; 1,000 µg/l), PFBS (0.1; 1; 100; 1,000 µg/l), 100 ng/l 17-beta-estradiol (E2), and 100 ng/l 5 alpha-androstan-17-beta-ol-3-one (DHT), respectively until 2 months postmetamorphosis, with water renewal every other day to maintain the appropriate exposure concentrations. DMSO (0.001% v/v) was used as a carrier solvent. Stock solutions were diluted by charcoalfiltered tap water to prepare test water. The tadpoles were fed three times daily with *Artemia*.

X. laevis exposure was conducted in 18L water tanks. Three replicates of 25 tadpoles were used for each exposure concentration. At the end of exposure (exact test duration not detailed), the survival rate of the frogs was recorded. The frogs were then anaesthetised and dissected for the tissues. Effects on gonads, livers, and sex-related genes were investigated. It was found that PFBS did not affect growth or survival but had the potential to promote expression of estrogen and androgen receptor activity. However, aromatase activity in the brain was not altered by PFBS. The authors indicated that the increase in expression of ER and AR suggests an increase in the responsiveness to the corresponding sex hormone and potential adverse effects on sexual development, see Section 5.6. Furthermore, exposure to PFOS and PFBS resulted in similar adverse effects on hepatohistology at >100 µg/L ($P < 0.05$). PFBS induced hepatocyte degeneration in both male and female frogs. Liver tissues also exhibited hepatocyte hypertrophy in both sexes following exposure to PFBS at the highest concentration (1000 µg/L, $P < 0.05$). Hepatohistological impairment caused by PFBS exhibited no sex-difference. In conclusion, these results show that PFBS have adverse effects on hepato-histology and sexual development on *X. laevis*. *Klimisch score 2 – well documented, non-guideline study.*

5.2. Terrestrial compartment

5.2.1. Toxicity to soil micro-organisms

Wójcik et al. (2018): The effects of PFBS and other water soluble perfluorinated pollutants on phospholipids in model soil decomposer membranes were investigated. PFASs are toxic to a wide range of soil bacteria and this biocide activity is related with their membrane activity (Chen et al., 2013; Pasquini et al., 2013). Such effects can lead to lowered degradation of dead organic matter by decomposer organisms.

Phospholipid Langmuir monolayers were used as a simplified model of bacterial membranes and their interactions with selected environmentally relevant PFASs, such as PFOS, PFOA and PFBS were studied (Wójcik et al., 2018). The effects of PFASs on the texture of model membranes were studied by Brewster angle microscopy, while the influence on molecular packing in the 2D bacterial crystal lattice was searched by the Grazing Incidence X-ray diffraction technique. PM-IRRAS spectroscopy was used to study the effect of PFASs on the phospholipid polar head group conformation, while penetration tests were used to monitor the effectiveness of PFASs incorporation into the model membrane. Results showed that Gram negative bacteria are more susceptible to PFASs than Gram positive species.

PFBS caused significant changes in the model membranes. Both PFOS and PFBS changed the orientation of the phospholipids within the 2D crystal lattice inducing the tilt of the hydrocarbon chains from the monolayer normal. PFBS turned out to be more structure

breaking than PFOS. The tilt of the hydrophobic chains induced by the PFASs presence in the subphase was greater for PFBS than for other PFASs. Results also showed that PFBS was not built into the hydrophobic regions of the membranes, but rather interacted with the polar headgroups of the monolayer. Tests proved that PFBS and PFOS behave similarly in that they can be incorporated to the monolayer at low surface pressure. The authors concluded that the switch from eight-carbon atom PFASs to shorter chained homologues does not necessarily lower their environmental toxicity.

5.2.2. Toxicity to terrestrial plants

Lan et al. (2018): The adverse effects of six perfluoralkyl acids PFBA, PFHxA, PFOA and PFBS, PFHxS, PFOS on the growth of wheat seedling were studied at two different spiking levels (200 and 2000 µg/kg soil). Biomass and chlorophyll levels were analyzed, and plant uptake and translocation were calculated. Pot experiments with pre-germinated seedlings of wheat (*Triticum aestivum*) in spiked soils were conducted in greenhouse for four weeks, with four replicates of each treatment. Fresh weight of shoot and roots were determined per pot and the chlorophyll contents (chlorophyll a and b) were measured at two different wavelengths (663 nm and 645 nm) with a spectrophotometer.

The exposure to PFBA/PFBS reduced the synthesis of Chlorophyll *a* in wheat by 35% at spiking levels of 2000 µg/kg, indicating their phytotoxicity. In contrast, PFOA/PFOS can promote the wheat chlorophyll content. Shoot biomass was significantly reduced (22.5% for PFBA/PFBS) at spiking levels of 2000 µg/kg PFAAs and the reduction increased with the carbon chain length of PFAAs. However, wheat root biomass showed a promotion at both spiking levels, possibly due to the enhancement in permeability of nutrients.

Qu et al. (2010): In another study by PFOS stimulated the growth of wheat seedlings and induce the synthesis of chlorophyll and soluble protein in wheat seedlings at concentrations less than 10 mg/L. However, PFOS concentrations >10 mg/L treatment could exert inhibition to the elongation and biomass of roots and leaves, and lead to damage to chlorophyll accumulation and soluble protein synthesis. When the concentration of PFOS was raised up to 200 mg/L, the activity of superoxide dismutase (SOD) and peroxidase (POD), two enzymes controlling oxidative stress in plants, decreased significantly with 12.6% and 33.7% inhibition for roots respectively, indicating that the antioxidative defensive system in wheat seedlings might be damaged by PFOS. Oxidative stress in *Arabidopsis thaliana* caused by PFOA had also been reported by (Yang et al., 2015).

The exposure of wheat seedlings to PFBS/PFBA led to a reduced chlorophyll a content of 35% and reduced shoot biomass of 22.5%, even though the test concentrations were much higher than concentrations detected in monitoring studies. High concentration of PFCAs and PFSAs can lead to a decrease in biomass in wheat and can cause oxidative stress, indicating their potential for phytotoxicity.

5.3. Microbiological activity in sewage treatment systems

Inhibition of microbial activity by KPFBS was tested on sewage microorganisms according to OECD TG 209 (NICNAS, 2005; Echa dissemination website). The nominal concentrations were 100 mg/L, 1000 mg/L and 10000 mg/L. Two negative controls were included in the study. 3,5- dichlorophenol was used as positive control. The exposure period was 3 hours. There were two inoculum controls. No clear dose-response was observed, and no toxic effects on respiration were observed in the tested inoculum as compared to controls. The results from this study indicated that KPFBS was not inhibitory to sewage microorganisms (EC₅₀ > 1000 mg/L). → *Klimisch reliability 1- well documented, guideline study.*

5.4. Toxicity to birds

Newsted et al. (2008): One acute dietary study on two bird species and one reproduction study on bobwhite quail have been performed by Newsted et al. (2018). The acute dietary study was performed according to OECD 205 and the species tested were bobwhite quail and mallard. Nominal test concentrations of potassium PFBS were 0 (control), 1000, 1780, 3160, 5620 and 10 000 mg PFBS/kg feed. Control birds were fed a plain diet. The bobwhite quails and mallards were exposed through their feed for 5 days at 39 °C and 31 °C respectively, with a 16 hours light and 8 hours dark period. Few treatment related effects were observed. There was a statistically significant reduction in body weight gain in quail in the 5620 and 1000 mg/kg group during day 1-8. Statistically significant reductions in body weight gain in the mallards was observed in the 10000 mg/kg group only. Hence the dietary LC50 values were determined to be >10000 mg/kg for both species. The NOEC was determined to be 3160 mg/kg for bobwhite quail and 5620 mg/kg for mallard.

For the bobwhite quail the NOEC was 3160 mg/kg due to reduction in body weight gain in the 5620 and 10000 mg/kg dose groups. The NOEC for the mallard was 5620 mg/kg due to reduction in body weight gain only in the 10000 mg/kg dose group. Regarding the reproduction study, the estimated NOEC for northern bobwhite quail exposed to PFBS in diet was 900 mg/kg, the highest concentration tested. Chronic dietary exposure of adult quail also led to concentration-dependent increase in PFBS accumulation in offspring eggs and juveniles, indicating maternal transfer to the progeny. → *Klimisch reliability 1- well documented, guideline study.*

The reproduction study was conducted according to OECD 206. Eighteen-week-old northern bobwhite quail were exposed to potassium PFBS. The nominal dietary concentrations were 100, 300 and 900 mg/kg, and the exposure period lasted 21 weeks. Each group consisted of sixteen pairs consisting of one female and one male bird per pen. Sixteen breeding pairs were included as negative controls and were fed an untreated diet. Recorded effects were adult health, body weight gain and feed consumption. Eggs were studied for egg shell thickness, egg shell cracks/abnormalities, infertility or embryo mortality. Cracked and abnormal eggs were discarded. Remaining eggs were incubated and allowed to hatch on day 21 of incubation. Evaluated reproductive endpoints were egg production, embryo viability, hatchability and hatchling health and survival. Hatchlings were housed in brooding pens and fed an untreated diet. Blood samples were collected from one 16-week old chick from each brooding pen. Adult birds and chicks were then euthanised and subjected to gross necropsy. There were no treatment related mortalities in any of the treatment groups. No PFBS related effects were noted on any lethal or nonlethal endpoints measured in the study and the dietary NOEC was determined to be 900 mg/kg ww feed. The authors concluded that PFBS exposure does not pose a significant risk to avian species. → *Klimisch reliability 1- well documented, guideline study.*

Two chronic studies with NOEC values of 200 (pilot study, Klimisch score 2) and 900 mg/kg feed with birds have also been reported on ECHAs dissemination website.

Table 37: Summary of effects on birds

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
Northern bobwhite quail (<i>Colinus virginianus</i>)	5 days	Acute toxicity	LC ₅₀ NOEC	>10000 mg/kg 3160 mg/kg	OECD TG 205	Newsted et al., 2008

Species	Duration	Endpoint	Effect level	Concentration	Comment	Reference
Northern bobwhite quail	21 weeks	Reproduction	NOEC	900 mg/kg	OECD TG 206	Newsted et al., 2008
Mallard (<i>Anas platyrhynchos</i>)	5 days	Acute toxicity	LC ₅₀ NOEC	>10000 mg/kg 5620 mg/kg	OECD TG 205	Newsted et al., 2008

5.5. Mammalian wildlife

The hazards described in Chapter 4 are also relevant for mammalian wildlife, especially to wildlife species with low reproductive output, because any negative effect on development or reproduction has a high likelihood of leading to serious effects at the population level for such species. Effects of PFBS on hormone levels or/and development/reproduction are of particular concern. In mice (Feng et al., 2017) and rats (NTP, 2019) decreases in total T4 and T3 levels and free T4 levels, and in addition for the mice study, increases in TSH levels. In the mice study (Feng et al., 2017) PFBS exposure caused a delay in perinatal growth, vaginal opening and first estrus cycle, and diestrus was prolonged (see Chapter 4).

PFBS has been shown to enrich in the edible parts of plants (see Section 3.7). For herbivorous mammalian wildlife species eating plants as the sole energy source this may be of particular concern. A quantitative risk characterisation is, however, not part of the SVHC assessment.

5.6. Other effects: Endocrine disruption

There are currently no T parameters validated for investigation of the disturbances of the thyroid pathway in fish. Thus, the findings from these studies should be treated with caution.

Chen et al. (2018a): The study aimed to investigate the multigenerational endocrine disrupting potential of PFBS by exposing F0 marine medaka (*Oryzias melastigma*) eggs to PFBS at different environmentally relevant concentrations (0, 1.0, 2.9, and 9.5 µg/L), until sexual maturity (see Section 5.1.1.2) for further description of the study design). In addition to the decreased body weight and body length discussed in Section 5.1.1.2, the hatching success of F1 larvae was significantly delayed at 1.0 µg/L exposure group following parental PFBS exposure. However, PFBS exposure of F0 eggs did not influence the hatching percentage at 15 dpf. This study demonstrated that PFBS has the potential ability to disturb the thyroid endocrine system across three generations following exposure to F0 marine medakas.

PFBS exposure resulted in a reduction in plasma T3 levels in F0 female exposed to 9.5 µg/L PFBS, while it increased the levels of T4 in the brains of the 1.0 and 9.5 µg/L exposure groups and subsequently decreased the levels of *Dio2*. As a characteristic of hyperthyroidism, increased T3 content suppress the transcription and activity of 5' - deiodinase (*Dio2*). *Dio2* is an enzyme involved in the activation and deactivation of thyroid

hormones, and it is important in the regulation of active T3 from the T4 prohormone (Orozco et al., 2005). Exposure to 2.9 and 9.5 µg/L PFBS resulted in a significant down-regulation of *Dio2* in male brains. Transgenerational disturbances in the thyroid endocrine system were noted in F1 and F2 generations at different life stages, implying the heritability of parental alterations caused by PFBS. The decreased levels of T3 in F0 female blood following exposure to 9.5 µg/L PFBS, was transferred to F1 eggs. Delayed hatching was coupled with elevated T3 levels in F1 larvae. F1 adults showed similar symptoms of thyroidal disruption as F0 adults. A slight recovery was noted in the F2 generation, although F2 larvae still exhibited thyroid disruption and synthesised excessive T4 suggesting that thyroidal disturbances may occur without direct exposure. The thyroid system plays a key role in several physiological processes, such as growth, development and reproduction (Power et al., 2001). Disruption of the thyroid hormone axis due to exposure to endocrine disrupting chemicals have been shown to affect larval development, growth and viability (Shi et al., 2009; Spaan et al., 2019). Furthermore, proper regulation of thyroid hormones is necessary for eye development and differentiation of visual systems in vertebrates (Bohnsack et al., 2013; Dong et al., 2014; Baumann et al., 2016). Thus, alterations in thyroid hormone signaling may lead to ocular developmental abnormalities, see Section 5.1.1.

Table 38: Overview of transcriptional effect in F1 and F2 offspring (larvae). Modified from Table S4 in the supplementary material. Significant upregulation is shown by upwards arrow and significant downregulation is shown by downwards arrow. A dash indicates non-significant results.

Genes	F1 offspring (15 dpf)			F2 offspring (15 dpf)		
	1.0µg/L	2.9 µg/L	9.5 µg/L	1.0 µg/L	2.9µg/L	9.5µg/L
<i>CRH</i>	↓	-	↓	-	↓	-
<i>CRHR</i>	-	-	-	-	-	-
<i>TSHβ</i>	-	-	-	-	-	-
<i>TSHR</i>	↓	↓	-	-	-	-
<i>NIS</i>	↓	↓	-	-	-	-
<i>TPO</i>	↓	↓	↓	↓	↓	-
<i>TG</i>	↓	↓	↓	-	↓	-
<i>TBG</i>	↓	↓	↓	-	↓	↑
<i>TRα</i>	-	-	↓	-	-	-
<i>TRβ</i>	↓	-	-	-	↓	-
<i>Dio 1</i>	↓	↓	↓	-	↓	↑
<i>Dio 2</i>	-	-	-	-	-	-
<i>Dio 3</i>	↓	↓	↓	-	↓	-
<i>UGT1</i>	↓	↓	↓	-	-	↑
<i>SULT</i>	↓	↓	↓	-	↓	-
<i>MCT8</i>	-	-	-	-	-	-
<i>SLCO1c1</i>	↓	↑	↓	-	↓	-
<i>pax8</i>	-	-	-	-	↓	-
<i>hhx</i>	↓	↓	↓	-	-	-

Chen et al. (2019): A recent article by Chen et al. (2019) exposing F0 marine medaka (*Oryzias melastigma*) eggs to PFBS at different environmentally relevant concentrations (0, 1.0, 2.9, and 9.5 µg/L), until sexual maturity (see section 5.1.1.2) demonstrates that PFBS significantly ($P < 0.05$) skewed the sex ratio in fish toward male dominance at 9.5 µg/L.

Only secondary sexual characteristics were used to assess the sex. The sex ratio in controls were quite high 83% and no mention of what the normal sex ratio of marine medaka is was mentioned. The acceptable sex ratio in other species (freshwater) is 30-70% males or females. A significant increase in number of males was observed in the highest test concentration, indicating that PFBS has implications for sexual development. The gonado somatic index (GSI) values were significantly lowered ($P < 0.001$) in the F0 females exposed to 9.5 µg/L. This correspond to the observed shift in F0 oocyte distribution where a significant increase in previtellogenic and significant decrease in vitellogenic and postvitellogenic oocytes were observed. This delay in oogenesis could in turn result in the observed significant decrease in F0 fecundity. The systemic effects on the gonads shown by GSI, delayed oogenesis and reduced fecundity are cohesive and indicate an effect of PFBS on reproduction. The observed significant decrease in hormonal level of E2, 11KT and E2/T in F0 females exposed to 9.5 µg/L are indicative of an anti-estrogenic effect.

Vongphachan et al. (2011): A gene expression assay was performed on avian embryonic neuronal cells of domestic chicken (*Gallus domesticus*) and herring gull (*Larus argentatus*) investigating key genes in the thyroid hormone pathway. Primary cultures of chicken embryonic neuronal (CEN) and herring gull embryonic neuronal (HGEN) cells were prepared for an in vitro screening method to determine the effects of PFAS exposure on the mRNA expression of TH-responsive genes. CEN and HGEN cells were prepared from the cerebral cortices of day 11 and 14 embryos, respectively. The cells were treated with PFBS at different concentrations 0.01, 0.1, 1, 3, 10, and 50 µM, the latter concentration was administered to the CEN cells only. Due to a decrease in cell viability following PFHxA treatment at 30 µM and 50 µM, 10 µM was the highest PFAS concentration used for subsequent mRNA expression in CEN and HGEN cells. The cells were incubated for 24 h. Cell viability was estimated using the Calcein-AM assay. RNA was extracted and

complementary DNA (cDNA) was synthesised using reverse transcriptase (RT) prior to PCR assay. To investigate gene expression mRNA levels were quantified using real-time PCR. T3 was used as a positive control for gene expression analyses. PFBS significantly altered the mRNA of TH-responsive genes in primary cultures of CEN and HGEN cells. Exposure to PFBS altered expression of TH-responsive genes (type III iodothyronine 5'-deiodinase (D3) at 3 µM and 10 µM, and neurogranin (RC3) at 10 µM) in chicken embryonic cells to a greater extent than that of longer chained PFASs. These effects could affect TH-dependent processes, while changes in RC3 expression could have consequences in synaptic plasticity, associative with learning and memory (Iniguez et al., 1993; Iniguez et al., 1996). In herring gull embryonic cells there was an increase in Octamer motif-binding factor 1 (Oct-1) mRNA following PFBS treatment at 10 µM. This study provide evidence that PFBS and other PFASs have the potential to alter TH homeostasis in birds.

Lou et al. (2013): A study on *Xenopus laevis* aimed to investigate comparatively adverse effects of longterm exposure to PFOS and PFBS on the growth and sexual development of amphibians using *Xenopus laevis*, see Section 5.1.5. Effects on gonads, livers, and sex-related genes were investigated. It was found that PFBS did not affect growth or survival but had the potential to promote expression of estrogen and androgen receptor activity. However, aromatase activity in the brain was not altered by PFBS. The authors indicated that the increase in expression of ER and AR suggests an increase in the responsiveness to the corresponding sex hormone and potential adverse effects on sexual development.

In summary, endocrine disruptive effects of PFBS are observed in several species. However, effects are seen at concentrations lower than those of longer chained PFASs such as PFOS. Exposure of F0 marine medaka to PFBS, resulted in multigenerational effects on the thyroid hormone system. These effects persisted in offspring that were reared in PFBS-free seawater. PFBS exposure also caused a direct effect on visual development and function across generations in the same fish. An in vitro study of primary cultures of avian neuronal cells reported effects on mRNA expression levels associated with the TH pathway. Considering these results, the thyroid seems to be a potential target organ for PFBS toxicity in fish.

Table 39: Summary of endocrine disruptive effects

Species	Duration	Endpoint	Observed effect concentration	Comment	Reference
Marine medaka	Life cycle	Thyroid disruption	LOEC: 1.0 µg/L (T3 contents and TBG contents in F1 larvae) 9.5 µg/L (T4 contents in F2 larvae)	Non-guideline	Chen et al., 2018a
		Delayed hatching	LOEC: 1.0 µg/L		
		Hatching percentage (F1 larvae)	LOEC: 1.0 µg/L		
		Skewed sex ratio (bias towards male)	LOEC: 9.5 µg/L NOEC: 2.9 µg/L		Chen et al., 2019

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

Species	Duration	Endpoint	Observed effect concentration	Comment	Reference
		F0 GSI values	LOEC: 9.5 µg/L NOEC: 2.9 µg/L		
		F0 fecundity (eggs/female/day)	LOEC: 2.9 µg/L NOEC: 1.0 µg/L		
		Oocyte development	LOEC: 9.5 µg/L NOEC: 2.9 µg/L		
		F1 larval mortality	LOEC: 2.9 µg/L NOEC: 1.0 µg/L		
Domestic chicken and herring gull embryonic neuronal cells	24 h	Effects on TH-responsive genes (fold change)	≥3 µM (D3) ≥10 µM (RC3)	Non-guideline	Vongphachan, 2011
<i>Xenopus laevis</i>	Life cycle	Upregulation of estrogen receptor (ER)	LOEC (brain): 0.1 µg/L (female and male) LOEC (liver): 0.1 µg/L (female and male)	Non-guideline	Lou et al., 2013
		Upregulation androgen receptor (AR)	LOEC (brain): 0.1 µg/L (male) NOEC (brain): 1000 µg/L (female) LOEC (liver): 1000 µg/L (female and male) NOEC (liver): 100 µg/L (female and male)		

5.7. Summary and discussion of the environmental hazard assessment

Ecotoxicity data are available for PFBS and the potassium salt of PFBS, KPFBS, which is the form of PFBS most often used in laboratory testing. With regards to ecotoxicity of the potassium salt, several proprietary GLP studies have been included in the registration

dossier. Ecotoxicological studies, supported by in vivo studies in rodents and in vitro studies, provide some evidence for adverse health effects. A full acute base set is available showing low acute toxicity of PFBS.

The acute toxicity (LC₅₀) to fish was reported to be 1938 mg/L (fathead minnow). A few long-term toxicity studies on fish are available. PFBS exposure caused similar effects as PFOS in zebrafish and marine medakas, although at higher concentrations (Hagenaars et al., 2011; Ulhaq et al., 2013a; Ulhaq et al., 2013b). Exposure to PFBS of zebrafish larvae resulted in malformations, altered heart rates, oedemas and distinct changes in behavioral patterns, such as abnormal swimming behavior.

Multigenerational disruption of the thyroid endocrine system by PFBS has been described for marine medaka. A life-cycle exposure of F0 marine medaka to environmentally realistic concentrations of PFBS led to marked and persistent disturbances in the thyroid endocrine system, such as hormone levels, transport and signalling, across three generations. A significant delay in hatching of F1 larvae following parental PFBS exposure was observed at 1.0 µg/L. Life-cycle exposure of F0 medaka from eggs to sexual maturity led to significantly decreased body weight of male and female fish (at 2.9 µg/L and 9.5 µg/L PFBS) and decreased body length of male medaka (at 2.9 µg/L and 9.5 µg/L PFBS), NOEC 1.0 µg/L. Furthermore, egg production significantly decreased upon exposure to PFBS (at 2.9 µg/L and 9.5 µg/L). Although offspring were reared in PFBS-free seawater, thyroid disruptive effects were observed in both F1 and F2 generations. Similar thyroid disruptive effects have also been reported in rat and mice studies, see Section 4.1.1.2. In the same marine medakas, PFBS exposure was also shown to result in a skewed sex ratio in fish toward male dominance (NOEC 2.9 µg/L) and egg production was significantly decreased (NOEC 1.0 µg/L). Furthermore, PFBS accumulated in eye tissue in a concentration dependent manner. Significant and sex-dependent effects were seen on visual proteins and neurotransmitter signalling which are essential for proper visual function. In addition, endocrine disturbances caused by PFBS have been shown in tadpoles and in avian neuronal cells (Lou et al., 2013; Vongphachan et al., 2011).

Acute algal toxicity tests (96 hour duration) with the freshwater algae *Pseudokirchneriella subcapitata* indicate EC₅₀ values > 5000 mg/L. While bioluminescence inhibition assays on bacteria and cyanobacteria indicated similar EC₅₀ values, exposure to PFBS caused significant changes in model bacteria membranes. Data from acute studies with the freshwater crustacean *Daphnia magna* were similar with 48 hour EC₅₀ value of >2000 mg/L. A 96 hour acute toxicity test with the marine crustacean *Mysidopsis bahia* is reported with an EC₅₀ value of 372 mg/L. A chronic 28 day daphnid reproduction test resulted in a NOEC value of 502 mg/L. Long-term relevant population effects were reported in *C. riparius* following exposure to 10 µg/L PFBS. In addition, two acute toxicity studies with birds were reported by Newsted et al. (2008) with LC₅₀ values >10000 mg/kg feed.

Exposure of wheat seedlings to PFBS/PFBA led to a reduced chlorophyll a content of 35% and reduced shoot biomass of 22.5%. PFCAs and PFSAs can cause a decrease in biomass in wheat and oxidative stress, indicating their potential for phytotoxicity.

Based on the available ecotoxicity data showing that PFBS exposure may lead to effects on development in fish, including reduced body weight and length (effects seen at 2.9 µg/L), delay in hatching (effects seen at 1.0 µg/L), reduced egg production (effect seen at 2.9 µg/L) and a skewed sex ratio (effects seen at 9.5 µg/L), it is concluded that PFBS fulfils the T criteria for the environment of Annex XIII of REACH (i.e. NOEC or EC₁₀ for marine or freshwater organisms less than 10 µg/L). The conclusion is supported by demonstrated effects on the visual system of marine medaka with a NOEC 1.0 µg/L, as well as dysbiosis of gut microbiota in the same fish following PFBS exposure. There are indications that PFBS may have endocrine disrupting properties.

6. Conclusions on the SVHC Properties

6.1. CMR assessment

Not relevant for the identification of the substance as SVHC in accordance with Article 57 (f) of the REACH Regulation.

6.2. PBT and vPvB assessment

Not relevant for the identification of the substance as SVHC in accordance with Article 57 (f) of the REACH Regulation.

6.3. Assessment under Article 57 (f)

In the present SVHC dossier it is evaluated whether perfluorobutane sulfonic acid (PFBS) and its salts should be regarded as “substances for which there is scientific evidence of probable serious effects to human health or the environment which give rise to an equivalent level of concern to those of other substances listed in Article 57 points (a) to (e) of the REACH Regulation”.

The assessment follows a case-by-case approach and is based on available information in a weight of evidence evaluation.. All studies and publications referenced in this dossier are considered relevant for the equivalent level of concern consideration for PFBS and its salts, and are used in this weight of evidence assessment. Despite the fact that some studies are not conducted according to OECD guidelines, or may not be considered highly relevant as standalone studies, there is no reason to discard any study on the basis of reliability.

As a strong acid, PFBS is readily deprotonated and forms sulfonate salts with bases, e.g. sodium, potassium and ammonium salts. In aqueous solution the salt forms and sulfonate ion will exist in equilibrium with PFBS itself. In the human body and in the environment PFBS will quickly be deprotonated by available bases and will be present dissolved on its anionic form. The concentrations reported in environmental and human monitoring studies will always be a sum of the acid PFBS, its anionic sulfonate form and its salt forms, see Chapter 1.

Several different precursors to PFBS exist and are in use in mixtures and in products, including side-chain fluorinated polymers. Such related substances may degrade to PFBS during the use-phase of a product, in the waste stage or in the environment and may be used in higher overall volumes compared to PFBS itself or its salts. However, the concern in this evaluation assessment is first and foremost related to the ultimate degradation product PFBS that the precursor substances eventually will turn into, see Section 1.3.3.

6.3.1. Summary of the data on the hazardous properties

6.3.1.1. Persistency/degradation

The perfluoroalkyl chain is persistent due to the high stability of the C-F-bond, while the sulfonic acid group is at its highest oxidation state and cannot be oxidised further under environmentally relevant conditions. There is experimental evidence that the shorter chain trifluoromethane sulfonic acid is non-degradable, while the structurally similar longer chain

sulfonic acids PFHxS and PFOS have already been identified as (very) persistent fulfilling the persistency criteria of REACH Annex XIII (see Section 3.1.4).

No abiotic or biotic degradation of PFBS has been shown at relevant environmental conditions, see Section 3.1. The hydrolytic half-life of PFBS in water was demonstrated to be more than 1 year and PFBS is considered to by far fulfil the P and vP criteria in REACH Annex XIII. This is supported by read-across to trifluoromethane sulfonic acid and PFOS. PFOS was found to be hydrolytically stable under environmental conditions with a hydrolytic half-life of 41 years (UNEP, 2006). QSAR predictions and screening tests confirmed that PFBS is meeting the screening criteria for "potentially persistent/very persistent" according to ECHA Guidance on PBT/vPvB assessment (ECHA, 2017a). For PFOS no biodegradation has been demonstrated (UNEP, 2006). High persistency is also demonstrated by the presence of PFBS in remote areas where there is no indication of a possible direct emission source.

Based on experimental data and read-across with relevant perfluoroalkane sulfonic acids like PFOS and trifluoromethane sulfonic acid, PFBS should be identified as a very persistent (vP) substance (hydrolytic half-life, $t_{1/2}$ at 25 °C of more than 1 year) that by far fulfils the P and vP criteria in REACH Annex XIII.

6.3.1.2. Partitioning in the environment and mobility

PFBS is a highly soluble, weakly sorbing substance with a high mobility in the environment. PFBS has a preference for distribution to the aqueous phase which is confirmed by distribution modelling and by field data demonstrating that PFBS is found in marine and surface water, and in ground water and drinking water samples throughout the world. Furthermore, PFBS is readily transported when water flows through soil. Being a strong acid, PFBS will be fully deprotonated at environmentally relevant conditions, and volatilisation of PFBS will be negligible, see Section 3.2.

Near fluorochemical industry facilities high PFBS concentrations are often found in surface water and groundwater. At an industrial site in China a 24-fold increase of the PFBS-concentration in the local groundwater was registered from 2009 to 2017.

6.3.1.3. Decontamination and release reduction for removal of PFBS from the environment and from drinking water

Due to the preference for the aqueous phase in the environment, the most important compartment for PFBS decontamination is water. PFBS is not readily removed with conventional water purification techniques. The same properties that make PFBS a highly mobile substance in the environment are also the reason why removal of PFBS usually fails. Due to the high aqueous solubility and the low sorption potential, PFBS will only to a low degree bind to adsorption materials and will rather remain in the water phase through the purification process. Drinking water production, as well as purification of wastewater industrial emissions may suffer from the low purification efficiency for PFBS, see Section 3.3.

The presence of PFBS precursors may complicate the water treatment process even further as the precursors may behave differently through the purification steps and may break down to PFBS either during or after purification.

The methods available today to remove PFBS contaminations from drinking water and lower the human exposure to PFBS are expensive and not commonplace. In a recent global

survey on drinking water, PFBS was the third most frequently detected PFAS in bottled water (47% of the samples) and tap water (88% of the samples).

The Swedish National Food Agency has recommended limits for drinking water based on the presence of 11 PFASs (PFBS, PFHxS, PFOS, 6:2 FTSA, PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFNA and PFDA)⁷. If the sum of these 11 PFASs occurs at concentrations greater than 0.09 µg/l, the Agency recommends that measures are taken as soon as possible to reduce the pollution. That PFBS is specified as one of the 11 PFASs covered by the limit value, demonstrates that there is specific concern associated with this particular compound.

6.3.1.4. Potential for long-range transport

The long environmental lifetime and the high mobility of PFBS results in a high global contamination potential. A characteristic travel distance of 17616 km and an overall environmental persistence, P_{ov} , of 220 days have been estimated. Long-range transport of PFBS is mainly via water and sea currents. Long-range transport of PFBS-related substances and degradation to PFBS may take place via other routes of transportation than those of PFBS, such as atmospheric transport. This also contributes to the concentrations of PFBS in the environment, see Section 3.4.

Sea spray aerosols contribute to the global transport of PFAAs, including PFBS, through the capacity to circulate significant amounts of PFAAs between the oceans and the atmosphere. A portion of the mass emitted from the oceans will deposit on land, thus re-entering the terrestrial system. Hence, human exposure to PFAAs like PFBS may continue even if emissions of the substances are stopped.

The global contamination potential has been demonstrated by findings of PFBS in samples of surface water, snow, ice, air and marine water in remote areas such as the Arctic, as well as in the Antarctic. PFBS has also been detected in Arctic biota, including polar bears and killer whales.

6.3.1.5. Bioaccumulation

Measured BCF values in fish are in the range 0.36 to 27.5. The highest fish BAF measured (field study) is 1736, however we considered this study unreliable with a Klimisch score 3. In another field study a mean BAF for fish of 69 was reported while for crab a mean BAF of 110 was measured. Compared to the numeric criteria in Annex XIII it is concluded that PFBS is not bioaccumulative in aquatic organisms (BCF < 2000).

However, PFBS is bioavailable and has been shown to enrich in blood and internal organs of animals and humans. Unlike the enrichment of contaminants in adipose tissue, the accumulation in specific organs has a higher potential to cause adverse effects, see Section 3.6.

PFBS has been detected in different species of wildlife, including green turtles and polar bears which are both listed on the IUCN red list of threatened species. PFBS was also

⁷ <https://www.livsmedelsverket.se/en/food-and-content/oonskade-amnen/miljogifter/pfas-in-drinking-water-fish-risk-management?AspxAutoDetectCookieSupport=1>

detected in birds and in marine mammals like dolphins and killer whales. In the killer whale the substance was shown to transfer from mother to foetus. Likewise, PFBS has been found to transfer from mother to eggs in birds. Hence, birds and animals, including endangered species, are exposed to PFBS at sensitive life stages.

In humans PFBS has been detected in blood, lungs, bone, kidneys, urine, hair and cord blood. A serum elimination half-life of around one month (up to 46 days) has been measured in humans. The half-life in pigs was reported to be similar to that in humans, and BMF values > 1 have been reported for pigs (1.2 (whole pig), 0.8 (meat), 6.4 (liver), 14 (blood plasma), 2.2 (kidney) and 0.9 (fat), indicating that PFBS might be bioaccumulative in pigs. No cut-off values for human elimination half-lives for fulfilling the B or vB criteria have been defined. Based on the lower half-life reported for PFBS in human blood compared to the half-lives of PFOA, 2 – 4 years, (B) and PFHxS, 7 - 8 years, (vB), it is concluded that PFBS is less bioaccumulative. However, it should be remembered that the data on human half-life is scarce and that a great individual variability in the data on half-lives of other PFAAs in humans has been observed. Furthermore, even though blood is used as a proxy for the entire body, the half-life in blood is not necessarily identical with the whole-body elimination half-life in humans. The presence of PFBS in human cord blood is of particular concern since the foetus is highly vulnerable to exposure to toxic substances.

No final conclusion on the bioaccumulation potential can yet be reached, as there is yet no approach to interpret the human and mammalian data. However, the available information indicates moderate bioaccumulation potential.

6.3.1.6. Enrichment in plants

Field irrigation with contaminated surface or ground waters and the use of contaminated sewage sludge as a soil conditioner are sources to PFAS in agricultural plants. From the plants PFASs can transfer to humans through the food chain. Several studies and field data have demonstrated the uptake of PFBS in plants. PFBS and other short-chain PFASs tend to enrich significantly in edible parts like leaves, vegetables and fruits. Uptake factors in shoot are in the range 2 – 4 for several crops like radish, celery, tomato and pea (Blaine et al., 2014), while uptake factors of up to 14 in lettuce leaves (Blaine et al., 2013) and 42 in tomato leaves have been measured (Felizeter et al., 2014). Considerable enrichment of PFBS has also been demonstrated for cabbage, zucchini and maize.

In plant derived supplement feeds PFBS contributed with 30% of the total amount of PFAAs in distillers dried grains with solubles and with 13% in soybean meal (Li et al., 2019b). Furthermore, in an area with a pollution incident in Germany in 2006, PFBS was determined in grass silage and hay as feed for dairy cows at $68.4 \pm 23.1 \mu\text{g}/\text{kg dm}$ and $993.6 \pm 224.4 \mu\text{g}/\text{kg dm}$, respectively. This shows that PFBS in this particular case was actually taken up in plants used as feed in agriculture, see Section 3.7.

6.3.1.7. Protein binding

PFBS can bind to serum albumin, and thereby the blood can distribute PFBS within the body, resulting in a potential to enrich particularly in blood rich tissues. Furthermore, PFBS acts as a substrate of organic anion transporting polypeptides, which function as an aid for uptake, elimination or reabsorption from urine to blood. It has also been demonstrated that PFBS can be transported into hepatocytes by the bile acid transporters, see Sections 3.6 and 4.1.1.

PFAAs resemble fatty acids in structure and concerns have been raised that PFAAs may disrupt fatty acid binding to transporter proteins or receptors and thus affect lipid metabolism. For example, PFBS was shown to bind to the liver fatty acid binding protein (L-FABP), which partly may explain the concentrations found in liver tissues or in kidneys where L-FABP is expressed. PFBS has also shown a potential to bind to and/or to activate the peroxisome proliferator-activated receptors (PPAR α , PPAR γ).

PFBS was found to compete with T4 for binding to the human thyroid hormone transport protein transthyretin (TTR), although the binding potency of PFBS was found to be considerably lower compared to T4, as well as compared to PFOS.

In animal derived supplement feeds PFBS contributed with 17%, 13.3%, 9.2% to total PFAAs in blood meal, feather meal and meat meal, respectively.

Altogether, PFBS has been shown to bind to different transport proteins and receptor proteins, although the binding affinities and activating potentials in general are lower for PFBS than for the longer chain counterparts.

6.3.1.8. Effects on human health

Repeated dose toxicity studies in rats have documented PFBS-mediated effects on liver, kidneys and the hematological system. A dose dependent decrease in hemoglobin and hematocrit levels was observed in male rats as the most sensitive effect with a NOAEL at 60 mg/kg/day. Evidence for genotoxic or mutagenic effects of PFBS has not been reported, but the current knowledge of possible carcinogenic effects is still limited and insufficient, see Chapter 4, and summary in Section 4.12.

A developmental study in mice has documented endocrine disturbances, such as decrease in T3 and T4 following prenatal PFBS-exposure, together with deficits in perinatal growth, pubertal onset and reproductive organ development at doses ≥ 200 mg/kg/day. Effects on thyroid hormone levels were also observed in a 28-day rat study. Support for effects of PFBS on the thyroid hormone levels is found in data from ecotoxicological studies as well, see Section 5.6.

Other effects of PFBS, including effects on lipid metabolism, neurotoxicity and immunotoxicity have been reported, but data are insufficient.

In general, toxicological data point towards similar health effects for PFBS as documented for some long-chain PFAAs, although the potency seems to be lower.

6.3.1.9. Effects in the environment

The currently available ecotoxicity studies have documented low acute toxicity following exposure to PFBS. However, long-term effects are reported in fish, see Section 5.1.1. PFBS-mediated effects in zebrafish are related to altered heart rate, head morphology and distinct changes in behavioural patterns, such as swimming speed. The lowest EC₅₀ for zebrafish was 450 mg/L. A life-cycle exposure of F0 marine medaka from eggs to sexual maturity led to marked disturbances in the thyroid endocrine system (e.g. hormone levels, transport and signalling) across three generations (F0, F1 and F2), although offspring were cultured in PFBS-free seawater. A significant delay in hatching of F1 larvae (at 1.0 $\mu\text{g/L}$) following parental PFBS exposure was observed, as well as a significantly decreased body weight of male and female fish and decreased body length of male medaka (NOEC 1.0 $\mu\text{g/L}$). Furthermore, exposure to PFBS was shown to result in a skewed sex ratio in marine

medakas toward male dominance (NOEC 2.9 µg/L) and egg production was significantly decreased (NOEC 1.0 µg/L). These adverse effects were observed at environmentally relevant exposure concentrations, 2.9 µg/L and 9.5 µg/L PFBS. Such concentrations have been measured near pollution hot-spot sites, see Section 3.2.5.1 and Table 11. In the same fish, PFBS was shown to accumulate in eye tissue in a concentration dependent manner and cause significant effects on visual proteins that are essential for proper visual signalling, see Section 5.6. PFBS was also found to cause dysbiosis of gut microbiota durable in parents and persisted in the offspring in marine medaka.

The ecotoxicological data showing developmental effects in fish (reduced body weight and length (NOEC of 1.0 µg/L); delay in hatching (LOEC 1.0 µg/L); reduced egg production (NOEC 1.0 µg/L); and a skewed sex ratio (NOEC of 2.9 µg/L)) are considered to fulfil the T criteria for the environment of Annex XIII of REACH. This conclusion is supported by effects of PFBS on the visual system of marine medakas with a NOEC < 1.0 µg/L.

In addition, available toxicological data indicate that PFBS can affect wildlife by interfering with endocrine processes, such as the thyroid hormone system, see Section 5.6. This has been demonstrated in fish and rodents. In frogs PFBS was found to promote expression of estrogen and androgen receptor activity in tadpoles.

Exposure of wheat seedlings to PFBS/PFBA led to a reduced chlorophyll a content of 35% and reduced shoot biomass of 22.5%. PFCAs and PFSAs can cause a decrease in biomass in wheat and oxidative stress, indicating their potential for phytotoxicity, see Section 5.2.2.

6.3.2. Concerns arising from the substance properties

The Guidance on the preparation of an Annex XV dossier for the identification of substances of very high concern, v2.1 (2018)⁸ provides general instructions to support the drafting. On the interpretation of the equivalent level of concern (ELoC), as laid down in Article 57 (f) of REACH, further support has been developed to assist all interested parties in the assessment of substances that may meet this category of concern. In 2015, ECHA, the Commission, and Member States prepared a discussion paper on the identification of substances as SVHC under Article 57 (f) with sensitisers as an example.⁹ Some years later the Joint Research Centre (JRC) repeated this exercise for the example of neurotoxic and immunotoxic substances, testing to what extent the same assessment elements as for sensitisers could be used on those health concerns.¹⁰ It was concluded by JRC and discussed among Member States, ECHA, and the Commission that this same set of relatively generic assessment elements could be applied to assist the more detailed case-by-case assessment of an equivalent level of concern for these two endpoints. Since then, several substances were included into the Candidate List on the basis of an Equivalent Level of Concern for endocrine disrupting effects in the environment, for endocrine disrupting effects on human health and on effects on specific target organs after repeated exposure, i.e. kidney and bone. Assessment of these different cases suggests several additional elements that could be considered, for example in the context of evaluating an equivalent level of concern for the environment like e.g. the possibility to adversely impact

⁸ https://echa.europa.eu/documents/10162/23036412/svhc_en.pdf/8faef33c-b46e-4186-8b7c-8cfbeccd0812

⁹ https://echa.europa.eu/documents/10162/13657/svhc_art_57f_sensitisers_en.pdf

¹⁰ Identification of Substances of Very High Concern (SVHC) under the 'equivalent level of concern' route (REACH Article 57(f)) – neurotoxicants and immunotoxicants as examples; <http://publications.jrc.ec.europa.eu/repository/bitstream/JRC96572/jrc96572-identification%20svhc%20reach%20article%2057f.pdf>

future generations and the possibility to impact regions or environmental organisms that are spatially remote from the point of environmental release of the substance.

In the ELoC assessment for PFBS, hazard related arguments and arguments related to environmental abundance and fate play an important role in concluding on the weight of evidence for an equivalent level of concern. This is in line with a recent judgement of the Court (Case C-323/15 P) where ELoC provision "*encompasses the possibility of taking into consideration, for the purposes of comparison, material going beyond merely the hazards arising from the intrinsic properties of the substances concerned*" (paragraph 34).

6.3.2.1. Concern for an irreversible and increasing presence in the environment

The available information on the physicochemical properties of PFBS, and its degradation and environmental presence, gives rise to concern that once the substance enters the environment, its presence will be irreversible. PFBS is stable towards both abiotic and biotic environmental degradation and has been shown to by far exceed the trigger values of being a very persistent (vP) substance, see Section 3.1. The hydrolytic half-life of PFBS in water was demonstrated to be more than 1 year. However, it may be much more than 1 year, and it is likely that PFBS will persist in the environment and expose future generations. The P/vP assessment is supported by read-across with relevant perfluoroalkane sulfonic acids like PFOS and trifluoromethane sulfonic acid. PFBS tends to distribute to the aqueous phase and is readily transported when water flows through soil. There are no natural barriers to prevent PFBS from being distributed to surface waters, groundwater and oceans once PFBS has entered the environment

Hence, PFBS may remain in the environment for such long times that it becomes increasingly difficult to predict possible exposures. As long as PFBS-releases to the environment continue, including the degradation of precursor compounds, its presence and concentrations may continue to increase. Consequently, slowly increasing environmental concentrations may be unavoidable and irreversible.

Cousins et al. (2019) point out that since actions to reduce emissions of PFAAs started in 2000–2002, there has been no evidence of consistent declines of any PFAA in the environment or biota. This illustrates the irreversibility of releases of such persistent chemicals as PFAAs.

There is a wide, dispersive use of PFBS and its precursors, see Section 1.3.3. The substances are used in common products, e.g. as a flame retardant in polycarbonate plastic, as catalyst or additive in manufacture, in metal plating or as water and stain repellent protection for leather, textiles and for hard porous surfaces, see Part II and Chapter 8.

6.3.2.2. Intrinsic substance properties result in irreversible and increasing contamination of surface water, marine water and groundwater

Building on the concern laid down in Section 6.3.2.1, the information presented in Section 3.2 gives rise to the additional concern that once PFBS enters the environment, the substance is free to move with the water phase and may thus lead to an irreversible contamination. The combination of high persistence and mobility allows PFBS to distribute on a great spatial scale in the environment even far from the release sources.

The high solubility (52.6 g/L at 22.5-24 °C), low adsorption potential (Log K_{oc} 1.2-2.7) and low volatility (Henry's Law constant: 0.040 Pa * m³/mol) of the ionised form ensure

that PFBS will remain in the water phase. PFBS will only to a low degree be “filtered” out when water moves through different layers of sediment, see Section 3.2. Based on today’s knowledge, there is no natural “sink” that will remove PFBS from the water phase, and therefore PFBS has the potential to reach remote and pristine areas, as well as ground water and drinking water sources.

Currently available methods to remove PFBS from water are expensive and not commonplace, see Section 3.3. Consequently, it is difficult to control emissions of PFBS from industry, purify wastewater or remove PFBS in drinking water production facilities. Suitable purification techniques may potentially become available in the future. However, their practical and economical applicability is as of yet uncertain. In light of this information and remaining uncertainties, emissions to the environment are considered irreversible, leading to an irreversible exposure of the environment, as well as of man via the environment.

Monitoring data confirm the uncontrollable spread of PFBS in the worldwide aqueous environment.

As such, there is high concern that the irreversible presence of PFBS in the environment may lead to worldwide unavoidable contamination of the environment and of drinking water sources.

6.3.2.3. PFBS contamination worldwide, including in remote regions

The concerns expressed in Sections 6.3.2.1 and 6.3.2.2 lead to the concern for a worldwide spread for PFBS. The presence of PFBS at background concentrations in the low ng/L range in the global environment is already being reported and supports this concern. Due to the global water cycle and the fact that the aqueous compartments are all well connected, the very high persistency and the high mobility of PFBS lead to long distance transport processes in the environment (estimated CTD: 17616 km, see Section 3.4.2).

PFBS will with time gradually spread to the entire aquatic environment, mainly via sea currents and sea spray aerosols, see Section 3.4. Contamination of remote areas has been demonstrated by field analyses. PFBS has been detected in surface water, snow, ice, air and marine water from remote areas in the Arctic and the Antarctic, as well as in biota such as polar bears, killer whales, northern fulmars and thick-billed murre.

Monitoring data indicate an increasing trend for PFBS in the environment, which can be explained by a combination of increasing annual production and a continuous accumulation of a persistent substance. However, the detection of PFBS in remote areas, like the Arctic, is taken as evidence of the persistence and mobility of PFBS, see Section 3.4.3.

Furthermore, long-range transport of volatile PFBS-related substances may take place via atmospheric transport. Subsequent degradation will release PFBS and contribute to the overall environmental concentrations of the substance. Thus, PFBS may affect humans and the environment far away from its points of emission into the environment and vulnerable populations and ecosystems in remote regions are affected.

6.3.2.4. Continuous presence in water results in continuous bioavailability

The intrinsic properties of PFBS lead to the concern for a continuous bioavailability of the substance worldwide. Information available on PFBS in different aqueous compartments demonstrate that PFBS is ubiquously present in water samples at background

concentrations, or higher in the vicinity of release sources, see Section 3.2.5. Section 3.5.3 shows that PFBS is found in marine biota, in fresh water biota and in other biota, while Section 4.1.2 shows that PFBS is taken up in humans. Biota levels are higher near point sources where the external concentrations are higher. PFBS is also enriched in plants, see Section 3.7. These monitoring data support that PFBS can be bioavailable and that exposure may take place via food and drinking water. Because PFBS stays in the water phase, the substance will be continuously bioavailable to organisms that live in water, organisms that drink water, plants that extract water from soil, organisms that eat plants or water-living organisms and eventually humans who will be exposed e.g. through food and via drinking water. In the absence of any natural degradation processes or sink, PFBS will remain bioavailable over multiple generations and hence can lead to inter-generational effects. A similar concern is also described for highly soluble and highly polar substances by Reemtsma et al. (2016).

It is of particular concern that PFBS has been detected in the human body - in blood, lung, bone, kidney, urine, hair and cord blood. Concentrations are generally low, but in a few studies concentrations up to 5 ng/mL have been observed and in persons with known elevated exposure even higher levels have been measured, see Section 4.3.

The bioavailability, in combination with the concerns expressed in Sections 6.3.2.1, 6.3.2.2 and 6.3.2.3, lead to concern for increasing internal exposures in wildlife and humans over time, which can be expected to trigger effects.

6.3.2.5. PFBS may enter biota and humans via several routes

PFBS enters the biosphere as a result of manufacturing and industrial emissions, and through the use of PFBS or precursor compounds in mixtures and articles. PFBS and its precursors have a wide dispersive use, including in consumer products, e.g. for the surface treatment of textiles and porous hard surfaces, see Part II and Section 8.

Humans and animals are exposed to PFBS in water and plants through the diet, see Sections 3.2.5 and 3.7. Near point sources, local drinking water, homegrown vegetables and eggs from domestic hens may have elevated PFBS levels, see Section 3.9. It has been shown that PFBS enriches in the edible parts of agricultural products and feed. PFBS is frequently found in surface water, tap water, bottled water and produced drinking water, with the highest concentrations found in areas with high PFBS contamination. Humans are also exposed to PFBS through the indoor environment and household dust, see Section 4.2.

Occurrence of PFBS in animal- and plant-derived protein supplement feed may increase as a result of increasing PFBS concentrations in the environment and the compound's ability to bind to proteins and enrich in plants. Since these supplemental feeds might be the main feed source for raised animals, human exposure through consumption of raised animals together with PFBS contaminated drinking water and plant derived food, might lead to long-term continuous elevated human exposure.

Continuous exposure is accompanied by a continuous bioavailability and constantly elevated levels of PFBS within the body. If releases of PFBS to the environment are not minimised, concentrations in vegetables and drinking water relevant for human nutrition will increase further.

Lam et al. (2016) observed a significant increasing temporal shifting trend from PFOS to PFBS in dolphin liver from the South China Sea in the period 2002 – 2014. This pattern was assigned to the substitution of PFOS by PFBS as an alternative in various applications.

The observation supports that increased use of PFBS and its precursors may lead to increased levels of PFBS in biota.

6.3.2.6. Weight of evidence evaluation of concern for adverse effects on human health and the environment

Chapters 4 and 5 and their summaries in Section 6.3.1.8 and 6.3.1.9, respectively, suggest that PFBS has the potential to cause serious effects to human health and the environment. PFBS or its salts do not currently have a harmonised classification under 1272/2008. However, self-classifications do exist for e.g. PFBS and KPFBS, see Part II and Chapter 7.

Human health

PFBS bioaccumulates differently compared to the classic lipophilic substances, due to the binding to serum albumin and other proteins in blood and tissues, see Sections 3.5, 3.6 and 4.1. The protein binding and atypical distribution influence the distribution in the human body. Moreover, due to the differences in half-lives observed for PFBS in humans, around one month (up to 46 days), compared to rats and mice (less than 1 day), the toxicity observed in animal studies may underestimate the toxicity in humans. In particular, long-term effects cannot be sufficiently assessed, and a precautionary approach should be taken when considering the risks related to the substance.

PFBS is bioavailable and has been detected in human blood, lung, bone, kidney, urine and hair. It has also been detected in human cord blood which is of particular concern since the foetus is highly vulnerable to exposure to toxic substances.

Toxicological data obtained for PFBS point towards similar health effects as documented for some long-chain PFAAs, although the potency seems to be lower. Effects on the liver, kidneys and the hematological system have been observed in rats, see Chapter 4 and summary in Section 6.3.1.8.

It is of particular concern that studies have found that PFBS may affect the thyroid system together with deficits in growth and development, see Chapters 4 and 5. PFBS thereby has a potential for causing adverse effects in both humans and wildlife. Effect on thyroid hormones has been shown for different species (rats, mice and fish). Furthermore, endocrine disturbances of PFBS have been shown in tadpoles, see Section 5.6.

The high persistency, mobility and bioavailability of PFBS leads to continuous exposure, and this adds to the concern for the observed effects, see Sections 6.3.2.1 to 6.3.2.5. It is not possible to reverse the exposure, and effects on human health and the environment are to be considered irreversible. With a continuous and irreversible exposure even effects that would be reversible in standard toxicity tests upon removal of exposure, would be irreversible if the environmental concentrations are allowed to reach levels that could trigger effects. Furthermore, there is a concern for currently unknown effects of PFBS in humans that may not be observed in standard toxicity tests or may only develop after life-long exposure.

Environment

PFBS has been detected in different species of wild-life, including in endangered or vulnerable populations and in sensitive life stages, see Sections 3.4.3.3 and 3.5.3.1.

PFBS affects several different functions in the fish marine medaka. Life-cycle exposure to PFBS was followed by marked disturbances in the thyroid endocrine system across three generations, although offspring were cultured in PFBS-free seawater. A significant delay in hatching of larvae was observed, together with decreased body weight of male and female fish and decreased body length of male medaka. Furthermore, exposure to PFBS was shown to result in a skewed sex ratio in marine medakas toward male dominance, and egg production was significantly decreased. It was also shown that PFBS accumulated in eye tissue in a concentration dependent manner and caused significant effects on visual proteins that are essential for proper visual signalling, see Section 5.6. The effects observed may have serious implications on a population level. PFBS was also found to cause dysbiosis of gut microbiota durable in parents and persisted in the offspring in marine medaka. Based on the effects observed for marine medaka (NOEC values down to 1.0 µg/L), it is considered that the T criteria for the environment according to Annex XIII of REACH are fulfilled for PFBS.

It is particularly concerning that the severe ecotoxicological effects demonstrated for PFBS are disturbances on fish, while the substance has physicochemical properties that make it persist in the aqueous environment for ages, available to fish.

In addition, there is a concern for currently unknown effects of PFBS on the environment that may not be observed in standard toxicity tests or may only develop after life-long exposure. This concern originates from the fact that PFBS will remain in the environment for so long that there is a high probability that effects on the environment that are not known today may be discovered in coming years, which is similar to what is described in the context of unknown effects for human health.

Concern for sensitive life stages, vulnerable species and future generations

PFBS has been measured in different species of wildlife, including green turtles and polar bears, which are both listed on the IUCN red list of threatened species. PFBS was also detected in marine mammals like dolphins and killer whales. In the whale the substance was shown to transfer from mother to foetus. Likewise, PFBS has been found to transfer from mother to eggs in birds, see Section 3.4.3.3.

It is of particular concern that PFBS is transferred from mother to offspring in humans, whales and birds. The foetus is highly vulnerable and sensitive to exposure to toxic substances. Future generations are exposed to PFBS at this sensitive development stage, including endangered species.

The weight-of-evidence for potential effects on human health and the environment can be characterised by two types of effects:

1. Observed adverse effects: concern for serious effects observed in the available studies that can be assessed based on the available data with (high) probability for human health and the environment, see paragraphs on human health and the environment earlier in this section.
2. Yet unknown effects: there is concern for possible effects that may *only become apparent after life-long exposure that are normally not tested for in standard toxicity tests. This concern is inherent to the high persistency of the substance and its continuous bioavailability.*

Unexpected effects from life-long exposure to PFBS represent a particular concern when taken together with the findings that PFBS transfers from mother to foetus in humans and in animals and may affect sensitive life stages and future generations.

Possibility to remedy effects

Following the concerns expressed in Sections 6.3.2.1 to 6.3.2.5, exposures may occur with a delay, as measured from the moment of emission, which complicates management or prevention of effects. Methods to remove PFBS from water are expensive and not commonplace, see Section 3.3. Control of emissions of PFBS from industry, purification of wastewater or removal of PFBS during drinking water production may therefore be challenging. Suitable purification techniques may potentially become available in the future. However, their practical and economical applicability is highly uncertain, and such remediation techniques may come with high societal costs. Degradation of precursor compounds with formation of PFBS represents a future source of PFBS, even if the emissions are stopped immediately. Consequently, once effects become apparent, it may be too late to take measures to protect us from exposure.

Brendel et al. (2018) point out that the lack of appropriate water treatment technologies results in everlasting background concentrations in the environment, and thus, organisms are permanently and poorly reversibly exposed. Considering such permanent exposure, it is very difficult to estimate long-term adverse effects in organisms.

Concern related to co-exposure

The concerns brought forward in Sections 6.3.2.1 to 6.3.2.5 lead to a concern for co-exposure with other contaminants with similar health effects. One possible large group of contaminants with roughly similar health effects (with different potencies) are other PFASs. PFBS is a member of the large group of PFASs, which currently counts about 4730 different substances, see Section 1.4. We are exposed to an increasing number of new and unidentified fluorinated products, some of which may share common mechanisms of actions and have additive effects. Often more than one PFAS can be identified in environmental samples, suggesting that PFASs are likely to co-occur in soil and the aqueous environment. The relative toxic potency of individual PFASs to liver was examined in a study. However, in general the current knowledge about combination effects of PFASs is limited. Co-exposure may last for a very long time.

Summary of the overall assessment of effects on human health and the environment

Toxicological data obtained for PFBS point towards similar health effects as documented for some long-chain PFAAs, although the potency seems to be lower. Due to the differences in half-lives observed for PFBS in humans, around one month (up to 46 days), compared to rats and mice (less than 1 day), the toxic effects observed in animal studies may underestimate the toxicity in humans. PFBS has been found to affect thyroid hormones in mice together with deficits in perinatal growth, pubertal onset and reproductive organ development. Effects on liver, kidneys and the hematological system has been observed in rats.

PFBS has been shown to cause severe toxic effects on the fish marine medaka in different ways. The thyroid endocrine system is disturbed across generations. PFBS causes significant delay in hatching of larvae, decreased body weight of male and female fish and

decreased body length of male medaka. Furthermore, exposure to PFBS was shown to result in a skewed sex ratio toward male dominance, and egg production was significantly decreased. PFBS also accumulates in eye tissue and causes significant effects on visual proteins that are essential for proper visual signalling. The mentioned effects may have serious implications on a population level.

Based on the effects on the marine medaka, it is considered that the T criteria for the environment according to Annex XIII of REACH are fulfilled for PFBS.

PFBS has been detected in human blood, cord blood, lung, bone, kidney, urine and hair. Furthermore, PFBS has been found to transfer from mother to foetus in humans and in animals and may affect sensitive life stages and future generations.

6.3.2.7. Derivation of limit values is highly uncertain

Due to the inherent properties of PFBS (e.g. very persistent and highly mobile in the environment), it is difficult, maybe even impossible, to reliably estimate current and future environmental concentrations and thereby exposure to animals and humans, see Section 6.3.2.1. Potential adverse effects due to life-long exposure are largely unknown. Considering the irreversibility of the environmental contaminations and the continuously increasing background levels, estimation of acceptable releases is not possible. Hence, the only way to adequately control the risk is through minimisation of emissions.

Furthermore, the toxicity observed in animal studies may underestimate the toxicity in humans due to the differences in half-lives observed for PFBS in humans, around one month (up to 46 days), compared to rats and mice (less than 1 day).

Estimation of future environmental concentrations and exposure via the environment is further complicated by the use and emissions of precursor compounds that may degrade to PFBS. There is considerable use of such precursor compounds, see Part II and Chapter 8. The degradation time for the precursors depends on their specific identity, see Section 1.3.3.

PFBS is only one member of the large group of PFASs that humans and the environment are exposed to, see Section 1.4. The number of PFASs in products encountered in daily life has expanded, and in the last count by OECD, 4730 PFAS-related CAS-numbers were found. We are exposed to an increasing number of new and unidentified fluorinated products, some of which may share common mechanisms of actions and have additive effects. Often more than one PFAS can be identified in environmental samples, suggesting that PFASs are likely to co-occur in soil and the aqueous environment. The relative toxic potency of individual PFASs to liver was examined in a study. However, in general the current knowledge about combinatorial effects of PFASs is limited.

Long-term exposure cannot be reliably estimated. Hence, the concerns outlined in this dossier cannot be readily quantified with a sufficient level of certainty using the currently available information. Previous emissions that already have happened would need to be accounted for together with current and future releases. In addition, the degradation of precursors with formation of PFBS would constitute a secondary source of PFBS, and the PFBS levels in the environment can continue to increase even if emissions are stopped. Consequently, the establishment of a 'safe concentration' for PFBS in the environment is not advisable.

6.3.2.8. Comparison of above concerns to concerns of PBT/vPvB substances

The ECHA *Guidance for PBT/vPvB assessment (Chapter R.11) (ECHA (2017a) states:*

“Experience with PBT/vPvB substances has shown that they can give rise to specific concerns that may arise due to their potential to accumulate in parts of the environment and

- that the effects of such accumulation are unpredictable in the long-term;*
- such accumulation is in practice difficult to reverse as cessation of emission will not necessarily result in a reduction in substance concentration.”*

“Furthermore, PBT or vPvB substances may have the potential to contaminate remote areas that should be protected from further contamination by hazardous substances resulting from human activity because the intrinsic value of pristine environments should be protected” (ECHA Guidance R.11).”

The concerns expressed in Sections 6.3.2.1 to 6.3.2.7 are the same as these three key concerns for PBT/vPvB substances. PFBS with its high persistence, high mobility and long-range transport potential matches very well with the guidance description above. Cousins et al. (2019) reflect on this type of concern in a more general manner for compounds that are highly persistent and poorly adsorb to organic matter and sediments. They even demonstrate that if a chemical is highly persistent, its continuous release will lead to continuously increasing contamination irrespective of the chemical's other physical-chemical properties. It is argued that the increasing concentrations will result in increasing probabilities of the occurrence of known and unknown effects and that, once adverse effects are identified, it will take decades, centuries or even longer to reverse contamination and therefore effects.

Section 4.0.1 of REACH Annex I explains that a hazard assessment addressing all the long-term effects and the estimation of the long-term exposure of humans and the environment cannot be carried out with sufficient reliability for PBT/vPvB-substances. This is also the case for PFBS and the concerns are of similar nature. Long-term exposure estimations for PFBS will be highly uncertain because of the very high persistence, the irreversibility of contaminations and the formation of PFBS from precursor compounds. Effects of PFBS on human health and the environment have been documented, and this is accompanied by the uncertainty of effects arising at longer time scales.

The environmental concentrations of PFBS will increase if the use and emissions continue. Because of its high long-range transport potential, due to the high persistency and high mobility in water, remote areas and pristine environments will also experience increasing PFBS levels. The environmental abundance of PFBS is considered impossible to reverse as there are no practically available methods for the removal of PFBS from the environment. Degradation of precursor substances to PFBS will result in continuous increase in the environmental concentrations and will further complicate the environmental distribution and potential clean-up. Hence, exposure of humans and the environment will continue for generations even after cessation of use. See also Section 6.3.2.1, Cousins et al. (2019).

Continued manufacture, use and release of PFBS will lead to higher levels of PFBS in the environment as there is no natural sink or way of removal for PFBS. Hence, all the mass of PFBS that gets emitted will be bioavailable. The exposure will increase, together with the levels of PFBS in humans and biota. How the concentrations will compare to PBT/vPvB-substances depends on many factors and is difficult to predict. However, common for both PFBS and classic PBT/vPvB-substances is that once harmful levels in the environment have been reached, it is impossible to remove the exposure and thereby reverse the harm.

Cousins et al. (2019) used model calculations to show the importance of persistence on environmental concentrations. The results indicated that concentrations of very persistent chemicals (e.g. with a degradation half-life in the order of 2000 days) may continue to increase for a certain time even after the emissions start to reduce. After the stop of emissions, the decreasing concentrations show a long tail that extends for many years.

A further similarity between PFBS and PBT/vPvB substances is the concern that irreversible or poorly reversible internal exposures to a substance can potentially lead to toxic effects that are not already known, as it appeared in several historical PBT/vPvB cases. This same concern is expressed for PFBS in Section 6.3.2.4. Both highly persistent and bioavailable substances, as well as PBT/vPvB substances, lead to unpredictable and uncontrollable internal exposures in the organism. PFBS which is less readily adsorbed to soil, sediment and suspended matter compared to the classic PBT/vPvB substances due to its properties, will likely have a higher proportion of the emitted substance bioavailable for organisms.

Regarding vPvB substances, ECHA guidance (ECHA 2017a) states:

"In the case of vPvB substances, there is concern that even if no toxicity is demonstrated in laboratory testing, long-term effects might be possible since high but unpredictable levels may be reached in man or the environment over extended time periods"

PFBS as a very persistent and highly mobile contaminant behaves differently in the environment as compared to a classic PBT/vPvB-substance that has a higher bioaccumulation potential and a lower mobility due to higher adsorption potential and lower aqueous solubility. The mobility of PFBS in soil and along waterways and the potential for long-range transport with sea currents will be higher. On the other hand, a low bioaccumulation potential may lead to less accumulation in the food-chain. If the half-life of excretion is lower, the levels of the substance in the body will decrease faster if the exposure is removed. On the other hand, calculations show that with an assumed half-life for PFBS in water of 10 years and a reported BAF in crab of 110 (Naile et al., 2013), the concentrations of PFBS in aquatic biota may be expected to exceed the biota concentrations for a persistent and bioaccumulative substance over time, see Section 3.5.2. Persistent and bioavailable substances, if either mobile or bioaccumulative, would seem to share the same concern for the development of high internal concentrations which may trigger effects. The continuous exposure, high mobility in water, no removal of PFBS from the environment and the enrichment in plants may ensure a continuous exposure via drinking water and diet.

6.3.3. Overall concern and assessment of the level of concern

PFBS is considered to be of equivalent level of concern to the very high concern for substances meeting the criteria laid down in Article 57 (a) to (e) according to Article 57 (f) of the REACH Regulation ((EC) 1907/2006) because of the overall concern arising from the concern elements described in Section 6.3.2. The elements which are used in this case for assessing the level of concern and description of how PFBS compares to those elements are listed in Table 40.

PFBS has due to its persistence, mobility, potential for long-range transport, observed effects to human health and environment, contamination of water and accumulation in plants, a very high potential to cause effects in wildlife and in man via environment. The very high persistence together with low adsorption potential (and therefore difficulty for end-of-pipe treatment) and high mobility imply a very high potential for increasing pollution stock. Irreversible, bioavailable, increasing exposures of both wildlife and man via environment will maintain as long as emissions continue, and even if emissions are stopped, the concentrations will not start to decrease immediately. The consequence is a

high potential for irreversible effects once effect levels have been reached, and an increasing seriousness of effects while exposures keep increasing. The substance has due to its intrinsic properties also a high potential to cause widespread exposures. Due to the challenging decontamination of drinking water sources, as well as the large variety of exposure routes via drinking water and food intake, there are no possibilities to avoid the continuous and increasing exposure in any human populations. Neither will any wildlife populations be protected from the whole released mass of the substance. It follows that both environment and humans are susceptible to impairment at large.

Table 40: Overview of the qualitative components relevant for assessing the level of concern

Irreversibility of the exposure of wildlife and man via the environment	<p>PFBS has high <i>potential to cause irreversible exposures</i>. Based on experimental results from screening tests on KPFBS, read-across to similar measurements for the corresponding C1 and C8 sulfonic acids, as well as supportive evidence from BIOWIN predictions, it can be concluded that the degradation potential of PFBS in all environmental compartments is very low or negligible. Its high persistency implies that PFBS will remain in the environment much longer than most other substances that are identified as exhibiting P or vP properties. PFBS by far exceeds the criteria for P and vP as laid down in Annex XIII of the REACH regulation.</p> <p>Exposures are not expected to decrease upon cessation of releases because of the <i>high persistence of the substance</i>. Degradation of precursors with formation of PFBS constitutes a secondary source of PFBS, and the PFBS levels in the environment can continue to increase even if emissions of PFBS are stopped.</p> <p>The high potential to cause very long-term exposures causes a difficulty to quantify exposures with sufficient certainty.</p>
Potential for rapid and wide geographic scale contamination	<p>Due to the global water cycle and the fact that the aqueous compartments are all well connected, the high persistency and the high mobility of PFBS lead to long distance transport processes in the environment. PFBS has already been detected in samples of surface water, snow, ice, air, marine water and biota from remote areas such as the Arctic and the Antarctic. Wide dispersive use of PFBS and its precursors in various applications further facilitates the global distribution of PFBS.</p>
Potential to continuous increase of exposures	<p>PFBS has a very high potential to cause an increasing pollution stock due to the combination of high persistence and difficulty of using <i>end-of-pipe emission reduction measures (as a result of low adsorption potential, negligible degradation and high water solubility)</i>. Furthermore, as a mobile substance, there are no local or intermittent sinks for the pollution stock and therefore the substance has high potential to cause continuous increase of exposure of wildlife.</p> <p>Additionally, due to the inefficiency and costs of</p>

decontamination and remediation techniques, PFBS has a high potential to cause continuously increasing exposure of humans via the environment. It is very difficult to remove PFBS from water. Adsorption of PFBS to soil, sediment and organic matter is poor. Available techniques suggest that remediation of PFBS-containing water comes with high costs for society and the generation of serious amounts of waste from the water purification process. Removing PFBS from the water compartment of the environment is impossible in practice due to the widespread presence in the aqueous environment and the high mobility of the substance.

Potential for causing serious effects although those would not be observed in standard tests

The continuously increasing pollution stock (see above), will lead to increasing internal exposures and a high likelihood of reaching levels which would cause effects. The whole released mass of the substance will be continuously bioavailable to organisms that live in water, organisms that drink water, plants that extract water from soil, animals that drink water and eat plants or water living organisms and humans who will be exposed e.g. through food and via drinking water. With time serious effects even for endpoints where the substance would show a low or moderate intrinsic toxicity based on standard tests, may be expected. Wildlife feeding on e.g. water and plants which enrich PFBS may be susceptible to reaching effect levels.

Potential for causing serious effects on human health (known and unknown), and the environment (including the potential for irreversible effects)

In the environment, severe effects have been observed for fish (marine medaka). A significant delay in hatching of larvae following parental PFBS exposure has been observed. Life-cycle exposure of F0 medaka from eggs to sexual maturity leads to significantly decreased body weight of male and female fish and decreased body length of male medaka. Furthermore, exposure to PFBS was shown to result in a skewed sex ratio in marine medakas toward male dominance, and egg production was significantly decreased. The effects were observed with NOECs less than 10 µg/L, consequently the T criteria (ENV) for PBT substances are fulfilled. PFBS was also found to accumulate in eye tissue of marine medaka in a concentration dependent manner and cause effects on visual proteins that are essential for proper visual signalling.

For human health, the concerns relate to thyroid hormone disturbances observed in mice, together with developmental effects. Effects have also been observed on the liver, kidney, haematological system and thyroid hormones in rats.

The continuous long-term and increasing exposure could eventually trigger effects that could be considered irreversible, even if they are normally reversible in standard toxicity studies upon the removal of exposure.

The highly mobile character of PFBS together with its

Delay of effects	<p>high persistence <i>mean that exposures may occur with a delay, as measured from the moment of emission. It can be argued for PFBS that it is equally difficult to manage or prevent effects when exposure could occur with a delay, as it is difficult to manage or prevent effects that could occur after a short-time exposure. Degradation of precursor substances with formation of PFBS represents a long-term source of PFBS to the environment that will contribute long after emissions are ceased.</i></p> <p>PFBS is expected to contribute to cumulative exposure with several other PFASs. Co-exposure may eventually occur and may last for a very long time <i>as degradation of PFBS and other PFASs in the environment is extremely slow or negligible.</i></p>
Potential to cause combined effects (co-exposure)	<p>The increasing and irreversible exposures lead to a potential of PFBS to cause yet unknown effects for human health and the environment. With time, effects may be discovered that may lead to more stringent safety levels for PFBS. The derivation of safe exposure levels may therefore be possible in principle but is considered not to be sufficiently reliable.</p>
Uncertainties in deriving safe concentration limits	
Possibility to remedy effects	<p>Reversing effects may hardly be possible due to the difficulty to reduce exposure. Consequently, for PFBS prevention of emissions is preferred over remediation of releases and treatment of effects. Monitoring data have demonstrated the global transport of PFBS as the substance has been detected in samples of surface water, snow, ice, air and marine water from remote areas such as the Arctic and the Antarctic.</p>
Uncertainties in quantifying exposures with sufficient certainty	<p>Due to its high persistency and mobility, and the degradation of precursors as a long-term source of PFBS, quantifications of future exposure to PFBS are highly uncertain. No currently available exposure estimation tools would reliably predict exposures from such substances.</p>
Potential to impair humans and the environment at large	<p>There are no natural barriers or environmental sinks that may reduce environmental concentrations of <i>PFBS, and hence exposure. There is a concern that the high potential for wide geographical scale contamination, in combination with its high persistence and adverse effects (as described above), may eventually lead to an unpredictable and uncontrollable risk for human health or the environment.</i></p> <p>PFBS has already been detected in human blood in studies from Europe, US and Asia. Furthermore, PFBS has been found in biota such as dolphins and whales, as well as green turtles and polar bears, which are both threatened species.</p>

Inter-generational effects PFBS persists in the environment for a very long time, and effects of current emissions may be observed or only become apparent in future generations. PFBS has been detected in human cord blood, and it has been demonstrated to be transferred from mother to foetus in killer whales and to eggs in birds.

Societal concern Art. 7.3 of the Water Framework Directive (2000/60/EC) stipulates that "Member States shall ensure the necessary protection for the bodies of water identified with the aim of avoiding deterioration in their quality of water to reduce the level of purification treatment required in the production of drinking water."

PFBS is frequently found in surface water, tap water, bottled water and produced drinking water. Highest concentrations are found in areas with high PFBS contamination. *Decontamination can only be achieved at high societal costs, if at all.*

Consequently, there is societal concern for the presence of PFBS in drinking water that requires immediate action.

6.3.4. Conclusion on the hazard properties and equivalent level of concern assessment

Perfluorobutane sulfonic acid (PFBS) and its salts are proposed to be identified as substances of very high concern in accordance with Article 57(f) of Regulation (EC) 1907/2006 (REACH) as there is scientific evidence of probable serious effects to the environment and human health which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of REACH.

PFBS-salts will be fully indistinguishable from PFBS in the environment as the salts will exist in their dissociated anionic sulfonate form, just like PFBS itself, and they will all be a part of the same equilibrium. Hence, all conclusions on different end-points apply to any and all salt forms, as well as PFBS itself.

Several concerns are caused by the intrinsic properties of PFBS. Elements of concern are triggered by individual intrinsic properties or by different combinations of the properties.

PFBS is persistent in the environment. No abiotic or biotic degradation of PFBS at relevant environmental conditions has been shown. This is supported by read-across to perfluoroalkane sulfonic acids with both shorter and longer chain lengths, which also have a very high environmental stability. PFBS shows a preference for distribution to the aqueous phase due to its high solubility in water (fully miscible with water at 20 °C) and low sorption potential ($\log K_{oc}$ 1.2 to 2.7) and is considered very mobile in the environment.

Overall, PFBS has a very high potential to cause widespread effects in wildlife and in humans exposed via environment, due to its persistence, mobility, potential for long-range transport (CTD 17616 km, $P_{ov} = 220$ days), observed adverse effects that may be relevant for human health and the environment, and exposure via drinking water and food. It is considered that the T criteria for ecotoxicity in Annex XIII of REACH are fulfilled for PFBS based on its documented effects on marine medaka that include significant reduction in body length and weight (NOEC 1.0 µg/L), reduced egg production (NOEC 1.0 µg/L), delayed hatching (LOEC 1.0 µg/L), a skewed sex ratio toward male dominance (NOEC 2.9 µg/L). Disturbances of the visual system (NOEC 1.0 µg/L) of marine medaka represent a supporting argument in the ecotoxicity assessment.

It is particularly concerning that PFBS prefers and persists in the aqueous environment, while severe effects of PFBS have been demonstrated for fish, which belong to this particular environmental compartment.

The very high persistence, together with low adsorption potential and high mobility, imply a high potential for increasing environmental concentrations and irreversible exposures of wildlife and of humans via the environment. Hence, PFBS is bioavailable via the aqueous environment. Furthermore, the low adsorption potential and high water solubility leads to difficulties and high costs in removing PFBS using end-of-pipe treatment. Together, these elements of concern lead to a high potential for irreversible effects once effect levels have been reached.

The high global transport potential is demonstrated by detection of PFBS in samples of surface water, snow, ice, air and marine water from remote areas such as the Arctic and the Antarctic. PFBS has also been found in biota like dolphins and whales, as well as green turtles and polar bears, which are both threatened species. This shows that PFBS is bioavailable, and that exposure may occur throughout the food chain and via drinking water and that this is already taking place worldwide.

In pigs BMF values > 1 have been reported (BMF 1.2, whole pig), indicating that PFBS might be bioaccumulative in pigs. A serum elimination half-life of around one month (up

to 46 days) has been measured in humans, which is considerably longer than the half-lives measured for rodents (less than 1 day).

The continuous and increasing exposure in human populations cannot be avoided if releases are not minimised. Similarly, wildlife populations cannot be protected from the total quantity of the substance released.

The level of concern is considered very high due to the combination of:

- high potential for irreversible exposure due to very high persistence and mobility and, in the case of human exposure via the environment, difficulties and costs to decontaminate drinking water,
- high potential for increasing contamination and increasing, bioavailable exposures as the intrinsic properties cause a difficulty to remove the substance after release,
- high potential for wide geographic scale contamination,
- ecotoxic effects fulfilling the T criteria in Annex XIII of REACH based on several different effects on marine medaka, including reduced egg production, delayed hatching of larvae, significant reduction in body length and weight, and a skewed sex ratio toward male dominance, supported by disturbances in the visual system,
- thyroid hormonal disturbances observed in rodents and fish and effects on expression of sex hormone receptor activity in tadpoles,
- effects on thyroid hormones together with deficits in perinatal growth, pubertal onset and reproductive organ development in mice,
- observed effects in experimental toxicity studies are of such nature that in combination with the above aspects, they lead to a high potential for serious effects on humans and the environment on a global scale,
- potential for causing serious effects that are not observed in standard tests or effects caused by life-long exposure or inter-generational effects following transfer from mother to offspring in several species,
- high societal concern.

PFBS has been detected in humans worldwide and in different species of wild life, including in endangered species and in remote areas. The substance has been found to transfer from mother to offspring in humans, whales and in birds and may disturb development at sensitive life stages and in vulnerable populations. It may be difficult in practice to manage exposures due to the high mobility of PFBS and the fact that exposures may take place at a different location than where releases occurred and at a different moment in time. The high persistence and high mobility of PFBS together lead to a concern for co-exposure with other contaminants with similar effects on human health and the environment. Co-exposure may last for a very long time, because natural degradation processes for these substances are slow or negligible. This is brought into the weight-of-evidence as supportive information.

Limitations of the available remediation techniques raise a concern that the removal of PFBS from drinking water may only be possible with high societal costs. Remediation of environmental pollution may even be practically impossible due to the high mobility of the substance. Furthermore, PFBS will quickly diffuse from contaminated sites.

Taken together in a weight-of-evidence consideration, the above arguments demonstrate that there is scientific evidence of probable serious adverse effects of PFBS and its salts to the human health and the environment, which gives rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of REACH Regulation.

REFERENCES FOR PART I

- 3M (2001). A 28-day oral (gavage) toxicity study of T-7485 in sprague-dawley rats. https://hero.epa.gov/hero/index.cfm/reference/details/reference_id/4241246
- Ahrens, L., Felizeter, S. & Ebinghaus, R. (2009): Spatial distribution of polyfluoroalkyl compounds in seawater of the German Bight. *Chemosphere*, 76, 179-184.
- Ahrens, L., Felizeter, S., Sturm, R., Xie, Z. & Ebinghaus, R. (2009a): Polyfluorinated compounds in waste water treatment plant effluents and surface waters along the River Elbe, Germany. *Marine Pollution Bulletin*, 58, 1326-1333.
- Ahrens, L., Gerwinski, W., Theobald, N. & Ebinghaus, R. (2010): Sources of polyfluoroalkyl compounds in the North Sea, Baltic Sea and Norwegian Sea: Evidence from their spatial distribution in surface water. *Marine Pollution Bulletin*, 60, 255-260.
- Alves, A., Jacobs, G., Vanermen, G., Covaci, A. & Voorspoels, S. (2015): New approach for assessing human perfluoroalkyl exposure via hair. *Talanta*, 144, 574-583.
- Appleman, T. D., Higgins, C. P., Quinones, O., Vanderford, B. J., Kolstad, C., Zeigler-Holady, J. C. & Dickenson, E. R. V. (2014): Treatment of poly- and perfluoroalkyl substances in U.S. full-scale water treatment systems. *Water Research*, 51, 246-255.
- Armitage, J. M., Macleod, M. & Cousins, I. T. (2009): Comparative Assessment of the Global Fate and Transport Pathways of Long-Chain Perfluorocarboxylic Acids (PFCAs) and Perfluorocarboxylates (PFCs) Emitted from Direct Sources. *Environmental Science & Technology*, 43, 5830-5836.
- Arp, H. P. & Slinde, G. A., Norwegian Geotechnical Institute (NGI) (2018). PFBS in the Environment: Monitoring and Physical-Chemical Data Related to the Environmental Distribution of Perfluorobutanesulfonic Acid (M-1122). <http://www.miljodirektoratet.no/Documents/publikasjoner/M1122/M1122.pdf>
- Ateia, M., Maroli, A., Tharayil, N. & Karanfil, T. (2019): The overlooked short- and ultrashort-chain poly- and perfluorinated substances: A review. *Chemosphere*, 220, 866-882.
- Australian Government, NICNAS (2005). Existing chemical hazard assessment report: Potassium perfluorobutane sulfonate. https://www.nicnas.gov.au/_data/assets/pdf_file/0004/4927/Potassium_Perfluorobutane_Sulfonate_PDF.pdf
- Axelsson, S. (2014). Perfluoroalkyl substances in Arctic birds - A comparison between glaucous gulls and black guillemots from Svalbard, Swedish University of Agricultural Sciences. *Bachelor of Science*. https://stud.epsilon.slu.se/6902/11/axelsson_s_140625.pdf
- Bao, J., Liu, W., Liu, L., Jin, Y., Dai, J., Ran, X., Zhang, Z. & Tsuda, S. (2011): Perfluorinated Compounds in the Environment and the Blood of Residents Living near Fluorochemical Plants in Fuxin, China. *Environmental Science & Technology*, 45, 8075-8080.

- Bao, J., Liu, L., Wang, X., Jin, Y.-H. & Dong, G.-H. (2017): Human exposure to perfluoroalkyl substances near a fluorochemical industrial park in China. Environmental Science and Pollution Research, 24, 9194–9201.
- Bao, J., Yu, W.-J., Liu, Y., Wang, X., Jin, Y.-H. & Dong, G.-H. (2019): Perfluoroalkyl substances in groundwater and home-produced vegetables and eggs around a fluorochemical industrial park in China. Ecotoxicology and Environmental Safety, 171, 199–205.
- Barzen-Hanson, K. A., Davis, S. E., Kleber, M. & Field, J. A. (2017): Sorption of Fluorotelomer Sulfonates, Fluorotelomer Sulfonamido Betaines, and a Fluorotelomer Sulfonamido Amine in National Foam Aqueous Film-Forming Foam to Soil. Environmental Science & Technology, 51, 12394-12404.
- Baumann, L., Ros, A., Rehberger, K., Neuhauss, S. C. F. & Segner, H. (2016): Thyroid disruption in zebrafish (*Danio rerio*) larvae: Differentmolecular response patterns lead to impaired eye development andvisual functions. Aquatic Toxicity, 172, 44-55.
- Behr, A.-C., Lichtenstein, D., Braeuning, A., Lampen, A. & Buhke, T. (2018): Perfluoroalkylated substances (PFAS) affect neither estrogen and androgen receptor activity nor steroidogenesis in human cells in vitro. Toxicology Letters, 291, 51–60.
- Bijland, S., Rensen, P. C. N., Pieterman, E. J., Maas, A. C. E., Hoorn, J. W. v. d., Erk, M. J. v., Havekes, L. M., Dijk, K. W. v., Chang, S.-C., Ehresman, D. J., Butenhoff, J. L. & Princen, H. M. G. (2011): Perfluoroalkyl Sulfonates Cause Alkyl Chain Length-Dependent Hepatic Steatosis and Hypolipidemia Mainly by Impairing Lipoprotein Production in APOE*3-Leiden CETP Mice. Toxicological Sciences, 123, 290–303.
- Bischel, H. N., MacManus-Spencer, L. A., Zhang, C. & Luthy, R. G. (2011): Strong associations of short-chain perfluoroalkyl acids with serum albumin and investigation of binding mechanisms. Environmental Toxicology and Chemistry, 30, 2423–2430.
- Bjerregaard-Olesen, C., Bach, C. C., Long, M., Ghisari, M., Bossi, R., Bech, B. H., Nohr, E. A., Henriksen, T. B., Olsen, J. & Bonefeld-Jørgensen, E. C. (2016): Time trends of perfluorinated alkyl acids in serum from Danish pregnant women 2008–2013. Environment International, 91, 14-21.
- Blaine, A. C., Rich, C. D., Hundal, L. S., Lau, C., Mills, M. A., Harris, K. M. & Higgins, C. P. (2013): Uptake of Perfluoroalkyl Acids into Edible Crops via Land Applied Biosolids: Field and Greenhouse Studies. Environmental Science & Technology, 47, 14062-14069.
- Blaine, A. C., Rich, C. D., Sedlacko, E. M., Hundal, L. S., Kumar, K., Lau, C., Mills, M. A., Harris, K. M. & Higgins, C. P. (2014): Perfluoroalkyl Acid Distribution in Various Plant Compartments of Edible Crops Grown in Biosolids-Amended soils. Environmental Science & Technology, 48, 7858–7865.
- Blaine, A. C., Rich, C. D., Sedlacko, E. M., Hyland, K. C., Stushnoff, C., Dickenson, E. R. V. & Higgins, C. P. (2014a): Perfluoroalkyl Acid Uptake in Lettuce (*Lactuca sativa*) and Strawberry (*Fragaria ananassa*) Irrigated with Reclaimed Water. Environmental Science & Technology, 48, 14361–14368.
- Bogdanska, J., Sundström, M., Bergström, U., Borg, D., Abedi-Valugerdi, M., Bergmanb, Å., DePierre, J. & Nobel, S. (2014): Tissue distribution of 35S-labelled

- perfluorobutanesulfonic acid in adult mice following dietary exposure for 1–5 days. Chemosphere, 98, 28-36.
- Bohlin-Nizzetto, P., Hanssen, L. & Herzke, D., Norwegian Environment Agency (2015). PFASs in house dust (M-430). <https://www.miljodirektoratet.no/globalassets/publikasjoner/m430/m430.pdf>
- Bohnsack, B. L. & Kahana, A. (2013): Thyroid hormone and retinoic acid interact to regulate zebrafish craniofacial neural crest development. Developmental Biology, 373, 300-309.
- Boiteux, V., Dauchy, X., Rosin, C. & Munoz, J.-F. o. (2012): National Screening Study on 10 Perfluorinated Compounds in Raw and Treated Tap Water in France. Archives of Environmental Contamination and Toxicology, 63, 1-12.
- Brady, N. C. & Weil, R. R. (2010). Elements of the Nature and Properties of Soils, 3rd Ed. Pearson Education Limited, Harlow, UK.
- Braune, B. M. & Letcher, R. J. (2013): Perfluorinated Sulfonate and Carboxylate Compounds in Eggs of Seabirds Breeding in the Canadian Arctic: Temporal Trends (1975–2011) and Interspecies Comparison. Environmental Science & Technology, 47, 616–624.
- Bräunig, J., Baduel, C., Barnes, C. M. & Mueller, J. F. (2019): Leaching and bioavailability of selected perfluoroalkyl acids (PFAAs) from soil contaminated by firefighting activities. Science of the Total Environment, 646, 471-479.
- Brendel, S., Fetter, E., Staude, C., Vierke, L. & Biegel-Engler, A. (2018): Short-chain perfluoroalkyl acids: environmental concerns and a regulatory strategy under REACH. Environmental Sciences Europe, 30, 1-11.
- Bressers, S., Tobor-Kaplon, M. & Beekhuijzen, M. (2018): Optimization of the use of *Caenorhabditis elegans* (C. elegans) in reproductive toxicology testing: describing the most optimal test conditions. Toxicology Letters, 295 Supplement 1, S82.
- Brumovský, M., Karaskova, P., Borghini, M. & Nizzetto, L. (2016): Per- and polyfluoroalkyl substances in the Western Mediterranean Sea waters. Chemosphere, 159, 308-316.
- Busch, J., Ahrens, L., Sturm, R. & Ebinghaus, R. (2010): Polyfluoroalkyl compounds in landfill leachates. Environmental Pollution, 158, 1467–1471.
- Butt, C. M., Berger, U., Bossi, R. & Tomy, G. T. (2010): Levels and trends of poly- and perfluorinated compounds in the arctic environment. Science of the Total Environment, 408, 2936-2965.
- Cai, M., Yang, H., Xie, Z., Zhao, Z., Wang, F., Lu, Z., Sturm, R. & Ebinghaus, R. (2012): Per- and polyfluoroalkyl substances in snow, lake, surface runoff water and coastal seawater in Fildes Peninsula, King George Island, Antarctica. Journal of Hazardous Materials, 209-210, 335– 342.
- Cai, M., Zhao, Z., Yin, Z., Ahrens, L., Huang, P., Cai, M., Yang, H., He, J., Sturm, R., Ebinghaus, R. & Xie, Z. (2012a): Occurrence of Perfluoroalkyl Compounds in Surface Waters from the North Pacific to the Arctic Ocean. Environmental Science & Technology, 46, 661–668.

- Cai, M., Zhao, Z., Yang, H., Yin, Z., Hong, Q., Sturm, R., Ebinghaus, R., Ahrens, L., Cai, M., He, J. & Xie, Z. (2012b): Spatial distribution of per- and polyfluoroalkyl compounds in coastal waters from the East to South China Sea. Environmental Pollution, 161, 162-169.
- Campbell, T. Y., Vecitis, C. D., Mader, B. T. & Hoffmann, M. R. (2009): Perfluorinated Surfactant Chain-Length Effects on Sonochemical Kinetics. Journal of Physical Chemistry A, 113, 9834-9842.
- Campo, J., Pérez, F., Masiá, A., Picó, Y., Farré, M. I. & Barceló, D. (2015): Perfluoroalkyl substance contamination of the Llobregat River ecosystem (Mediterranean area, NE Spain). Science of the Total Environment, 503-504, 48-57.
- Cariou, R., Veyrand, B., Yamada, A., Berrebi, A., Zalkod, D., Durand, S., Pollono, C., Marchand, P., Leblanc, J.-C., Antignac, J.-P. & Bizec, B. L. (2015): Perfluoroalkyl acid (PFAA) levels and profiles in breast milk, maternal and cord serum of French women and their newborns. Environment International, 84, 71-81.
- Castiglioni, S., Valsecchi, S., Polesello, S., Rusconi, M., Melis, M., Palmiotto, M., Manenti, A., Davoli, E. & Zuccato, E. (2015): Sources and fate of perfluorinated compounds in the aqueous environment and in drinking water of a highly urbanized and industrialized area in Italy. Journal of Hazardous Materials, 282, 51-60.
- Cevasco, A., Urbatzka, R., Bottero, S., Massari, A., Pedemonte, F., Kloas, W. & Mandich, A. (2008): Endocrine disrupting chemicals (EDC) with (anti)estrogenic and (anti)androgenic modes of action affecting reproductive biology of *Xenopus laevis*: II. Effects on gonad histomorphology. Comparative Biochemistry and Physiology, Part C, 147, 241-251.
- Chen, H., Yao, J., Wang, F., Cai, M. & Liu, H. (2013): Toxicity of perfluorooctanoic acid to *Pseudomonas putida* in the aquatic environment. Journal of Hazardous Materials, 262, 726- 731.
- Chen, H., Sun, R., Zhang, C., Han, J., Wang, X., Han, G. & He, X. (2016): Occurrence, spatial and temporal distributions of perfluoroalkyl substances in wastewater, seawater and sediment from Bohai Sea, China. Environmental Pollution, 219, 389-398.
- Chen, F., Gong, Z. & Kelly, B. C. (2016a): Bioavailability and bioconcentration potential of perfluoroalkyl-phosphinic and -phosphonic acids in zebrafish (*Danio rerio*): Comparison to perfluorocarboxylates and perfluorosulfonates. Science of the Total Environment, 568, 33-41.
- Chen, F., Wei, C., Chen, Q., Zhang, J., Wang, L., Zhou, Z., Chen, M. & Liang, Y. (2018): Internal concentrations of perfluorobutane sulfonate (PFBS) comparable to those of perfluorooctane sulfonate (PFOS) induce reproductive toxicity in *Caenorhabditis elegans*. Ecotoxicology and Environmental Safety, 158, 223-229.
- Chen, L., Hu, C., Tsui, M. M. P., Wan, T., Peterson, D. R., Shi, Q., Lam, P. K. S., Au, D. W. T., Lam, J. C. W. & Zhou, B. (2018a): Multigenerational Disruption of the Thyroid Endocrine System in Marine Medaka after a Life-Cycle Exposure to Perfluorobutanesulfonate. Environmental Science & Technology, 52, 4432-4439.
- Chen, L., Tsui, M. M. P., Shi, Q., Hu, C., Wang, Q., Zhou, B., Lam, P. K. S. & Lam, J. C. W. (2018b): Accumulation of perfluorobutane sulfonate (PFBS) and impairment of visual function in the eyes of marine medaka after a life-cycle exposure. Aquatic Toxicology, 201, 1-10.

- Chen, L., Lam, J. C. W., Hu, C., Tsui, M. M. P., Wang, Q., Giesy, J. P. & Lam, P. K. S. (2018c): Perfluorobutanesulfonate Exposure Causes Durable and Transgenerational Dysbiosis of Gut Microbiota in Marine Medaka. Environmental Science & Technology Letters, 5, 731–738.
- Chen, L., Lam, J. C. W., Hu, C., Tsui, M. M. P., Lam, P. K. S. & Zhou, B. (2019a): Perfluorobutanesulfonate Exposure Skews Sex Ratio in Fish and Transgenerationally Impairs Reproduction. Environmental Science & Technology, 53, 8389–8397.
- Chen, Q., Zhang, X., Zhao, Y., Lu, W., Wu, J., Zhao, S., Zhang, J. & Huang, L. (2019b): Prenatal exposure to perfluorobutanesulfonic acid and childhood adiposity: A prospective birth cohort study in Shanghai, China. Chemosphere, 226, 17-23.
- Chengelis, C. P., Kirkpatrick, J. B., Myers, N. R., Shinohara, M., Stetson, P. L. & Sved, D. W. (2009): Comparison of the toxicokinetic behavior of perfluorohexanoic acid (PFHxA) and nonafluorobutane-1-sulfonic acid (PFBS) in cynomolgus monkeys and rats. Reproductive Toxicology, 27, 400-406.
- Chu, S. & Letcher, R. J. (2014): In vitro metabolic formation of perfluoroalkyl sulfonamides from copolymer surfactants of pre- and post-2002 Scotchgard fabric protector products. Environmental Science & Technology, 48, 6184-6191.
- Codling, G., Halsall, C., Ahrens, L., Vento, S. D., Wiberg, K., Bergknut, M., Laudon, H. & Ebinghaus, R. (2014): The fate of per- and polyfluoroalkyl substances within a melting snowpack of a boreal forest. Environmental Pollution, 191, 190-198.
- Corsini, E., Sangiovanni, E., Avogadro, A., Galbiati, V., Viviani, B., Marinovich, M., Galli, C. L., Dell'Agli, M. & Germolec, D. R. (2012): In vitro characterization of the immunotoxic potential of several perfluorinated compounds (PFCs). Toxicology and Applied Pharmacology, 258, 248–255.
- Cousins, I. T., Vestergren, R., Wang, Z., Scheringer, M. & McLachlan, M. S. (2016): The precautionary principle and chemicals management: The example of perfluoroalkyl acids in groundwater. Environment International, 94, 331-340.
- Cousins, I. T., Ng, C. A., Wang, Z. & Scheringer, M. (2019): Why is High Persistence Alone a Major Cause of Concern? Environmental Science Processes & Impacts, 21, 781-792.
- Crookes, M. J. & Fisk, P., Peter Fisk Associates (2018). Evaluation of using mobility of chemicals in the environment to fulfil bioaccumulation criteria of the Stockholm Convention. <http://files.chemicalwatch.com/26%20-%20CA%2017%202019%20Assessment%20mobility%20criteria%20POPs.docx>
- D'Agostino, L. A. & Mabury, S. A. (2017): Certain Perfluoroalkyl and Polyfluoroalkyl Substances Associated with Aqueous Film Forming Foam Are Widespread in Canadian Surface Waters. Environmental Science & Technology, 51, 13603–13613.
- Dalsager, L., Christensen, N., Husby, S., Kyhl, H., Nielsen, F., Høst, A., Grandjean, P. & Jensen, T. K. (2016): Association between prenatal exposure to perfluorinated compounds and symptoms of infections at age 1–4 years among 359 children in the Odense Child Cohort. Environment International, 96, 58–64.
- Dassuncao, C., Hu, X. C., Zhang, X., Bossi, R., Dam, M., Mikkelsen, B. & Sunderland, E. M. (2017): Temporal Shifts in Poly- and Perfluoroalkyl Substances (PFASs) in North

- Atlantic Pilot Whales Indicate Large Contribution of Atmospheric Precursors. Environmental Science & Technology, *51*, 4512–4521.
- Delinsky, A. D., Strynar, M. J., McCann, P. J., Varns, J. L., McMillan, L., Nakayama, S. F. & Lindstrom, A. B. (2010): Geographical Distribution of Perfluorinated Compounds in Fish from Minnesota Lakes and Rivers. Environmental Science & Technology, *44*, 2549–2554.
- D'eon, J. C., Hurley, M. D., Wallington, T. J. & Mabury, S. A. (2006): Atmospheric Chemistry of N-methyl Perfluorobutane Sulfonamidoethanol, C₄F₉SO₂N(CH₃)CH₂CH₂OH: Kinetics and Mechanism of Reaction with OH. Environmental Science & Technology, *40*, 1862–1868.
- D'Hollander, W., Roosens, L., Covaci, A., Cornelis, C., Reynders, H., Campenhout, K. V., Voogt, P. d. & Bervoets, L. (2010): Brominated flame retardants and perfluorinated compounds in indoor dust from homes and offices in Flanders, Belgium. Chemosphere, *81*, 478–487.
- Ding, G., Xue, H., Yao, Z., Wang, Y., Ge, L., Zhang, J. & Cui, F. (2018): Occurrence and distribution of perfluoroalkyl substances (PFASs) in the water dissolved phase and suspended particulate matter of the Dalian Bay, China. Chemosphere, *200*, 116–123.
- Domingo, J. L., Ericson-Jogsten, I., Perelló, G., Nadal, M., Bavel, B. V. & Kärrman, A. (2012): Human Exposure to Perfluorinated Compounds in Catalonia, Spain: Contribution of Drinking Water and Fish and Shellfish. Journal of Agricultural and Food Chemistry, *60*, 4408–4415.
- Dong, G.-H., Tung, K.-Y., Tsai, C.-H., Liu, M.-M., Wang, D., Liu, W., Jin, Y.-H., Hsieh, W.-S., Lee, Y. L. & Chen, P.-C. (2013): Serum Polyfluoroalkyl Concentrations, Asthma Outcomes, and Immunological Markers in a Case-Control Study of Taiwanese Children. Environmental Health Perspectives, *121*, 507–513.
- Dong, W., Macaulay, L. J., Kwok, K. W. H., Hinton, D. E., Ferguson, P. L. & Stapleton, H. M. (2014): The PBDE metabolite 6-OH-BDE 47 affects melanin pigmentation and THRb mRNA expression in the eye of zebrafish embryos. Endocrine Disruptors, *2*, 1–15.
- Droge, S. T. J. & Goss, K.-U. (2013): Development and Evaluation of a New Sorption Model for Organic Cations in Soil: Contributions from Organic Matter and Clay Minerals. Environmental Science & Technology, *47*, 14233–14241.
- Droge, S. T. J. & Goss, K.-U. (2013a): Sorption of Organic Cations to Phyllosilicate Clay Minerals: CEC Normalization, Salt Dependency, and the Role of Electrostatic and Hydrophobic Effects. Environmental Science & Technology, *47*, 14224–14232.
- ECHA (2013). Support document for the identification of PFOA as a substance of very high concern. <https://echa.europa.eu/documents/10162/8059e342-1092-410f-bd85-80118a5526f5>
- ECHA (2015). Support document for the identification of PFNA as a substance of very high concern. <https://www.echa.europa.eu/documents/10162/48ae5fe3-9436-4a10-a533-ed642b92ce47>
- ECHA (2017). Member State Committee Support Document for the Identification of Perfluorohexane-1-sulphonic Acid and its Salts as Substances of Very High Concern

Because of Their vPvB (Article 57e) Properties. <https://echa.europa.eu/candidate-list-table/-/dislist/details/0b0236e18184a0e1>

ECHA (2017a). Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT/vPvB assessment. https://echa.europa.eu/documents/10162/13632/information_requirements_r11_en.pdf

Ehresman, D. J., Froehlich, J. W., Olsen, G. W., Chang, S.-C. & Butenhoff, J. L. (2007): Comparison of human whole blood, plasma, and serum matrices for the determination of perfluorooctanesulfonate (PFOS), perfluorooctanoate (PFOA), and other fluorochemicals. *Environmental Research*, 103, 176–184.

Elvers, B. (2011). *Ullmann's Encyclopedia of Industrial Chemistry*, 7th ed. Wiley-VCH Verlag GmbH, New York, USA.

Ericson, I., Domingo, J. L., Nadal, M., Bigas, E., Llebaria, X., Bavel, B. v. & Lindström, G. (2009): Levels of Perfluorinated Chemicals in Municipal Drinking Water from Catalonia, Spain: Public Health Implications. *Archives of Environmental Contamination and Toxicology*, 57, 631-638.

Eriksen, K. T., Raaschou-Nielsen, O., Sørensen, M., Roursgaard, M., Loft, S. & Møller, P. (2010): Genotoxic potential of the perfluorinated chemicals PFOA, PFOS, PFBS, PFNA and PFHxA in human HepG2 cells. *Mutation Research*, 700, 39–43.

Eriksson, U. & Kärrman, A. (2015): World-Wide Indoor Exposure to Polyfluoroalkyl Phosphate Esters (PAPs) and other PFASs in Household Dust. *Environmental Science & Technology*, 49, 14503–14511.

Eriksson, U., Haglund, P. & Kärrman, A. (2017): Contribution of precursor compounds to the release of per- and polyfluoroalkyl substances (PFASs) from waste water treatment plants (WWTPs). *Journal of Environmental Sciences*, 61, 80-90.

Eschauzier, C., Beerendonk, E., Scholte-Veenendaal, P. & Voogt, P. D. (2012): Impact of Treatment Processes on the Removal of Perfluoroalkyl Acids from the Drinking Water Production Chain. *Environmental Science & Technology*, 46, 1708-1715.

Eschauzier, C., Haftka, J., Stuyfzand, P. J. & Voogt, P. d. (2010): Perfluorinated Compounds in Infiltrated River Rhine Water and Infiltrated Rainwater in Coastal Dunes. *Environmental Science & Technology*, 44, 7450–7455.

Eschauzier, C., Raat, K. J., Stuyfzand, P. J. & Voogt, P. D. (2013): Perfluorinated alkylated acids in groundwater and drinking water: Identification, origin and mobility. *Science of the Total Environment*, 458–460, 477–485.

Falk, S., Failing, K., Georgii, S., Brunn, H. & Stahl, T. (2015): Tissue specific uptake and elimination of perfluoroalkyl acids (PFAAs) in adult rainbow trout (*Oncorhynchus mykiss*) after dietary exposure. *Chemosphere*, 129, 150–156.

Fang, S., Chen, X., Zhao, S., Zhang, Y., Jiang, W., Yang, L. & Zhu, L. (2014): Trophic Magnification and Isomer Fractionation of Perfluoroalkyl Substances in the Food Web of Taihu Lake, China. *Environmental Science & Technology*, 48, 2173–2182.

Felizeter, S., McLachlan, M. S. & Voogt, P. d. (2012): Uptake of Perfluorinated Alkyl Acids by Hydroponically Grown Lettuce (*Lactuca sativa*). *Environmental Science & Technology*, 46, 11735–11743.

- Felizeter, S., McLachlan, M. S. & Voogt, P. D. (2014): Root Uptake and Translocation of Perfluorinated Alkyl Acids by Three Hydroponically Grown Crops. Journal of Agricultural and Food Chemistry, 62, 3334–3342.
- Feng, X., Cao, X., Zhao, S., Wang, X., Hua, X., Chen, L. & Chen, L. (2017): Exposure of Pregnant Mice to Perfluorobutanesulfonate Causes Hypothyroxinemia and Developmental Abnormalities in Female Offspring. Toxicological Sciences, 155, 409-419.
- Fraser, A. J., Webster, T. F., Watkins, D. J., Strynar, M. J., Kato, K., Calafat, A. M., Vieira, V. M. & McClean, M. D. (2013): Polyfluorinated compounds in dust from homes, offices, and vehicles as predictors of concentrations in office workers' serum. Environment International, 60, 128-136.
- Fromme, H., Wöckner, M., Roscher, E. & Völkel, W. (2017): ADONA and perfluoroalkylated substances in plasma samples of German blood donors living in South Germany. International Journal of Hygiene and Environmental Health, 220, 455–460.
- Fu, Y., Wang, T., Wang, P., Fu, Q. & Lu, Y. (2014): Effects of age, gender and region on serum concentrations of perfluorinated compounds in general population of Henan, China. Chemosphere, 110, 104-110.
- Gao, K., Zhuang, T., Liu, X., Fu, J., Zhang, J., Fu, J., Wang, L., Zhang, A., Liang, Y., Song, M. & Jiang, G. (2019): Prenatal Exposure to Per- and Polyfluoroalkyl Substances (PFASs) and Association between the Placental Transfer Efficiencies and Dissociation Constant of Serum Proteins–PFAS Complexes. Environmental Science & Technology, 53, 6529–6538.
- García-Valcárcel, A. I., Moleroa, E., Escorial, M. C., Chueca, M. C. & Tadeo, J. L. (2014): Uptake of perfluorinated compounds by plants grown in nutrient solution. Science of the Total Environment, 472, 20-26.
- Gebbink, W. A., Asseldonk, L. v. & Leeuwen, S. P. J. v. (2017): Presence of Emerging Per- and Polyfluoroalkyl Substances (PFASs) in River and Drinking Water near a Fluorochemical Production Plant in the Netherlands. Environmental Science & Technology, 51, 11057-11065.
- Gebbink, W. A., Bossi, R., Rigét, F. F., Rosing-Asvid, A., Sonne, C. & Dietz, R. (2016): Observation of emerging per- and polyfluoroalkyl substances (PFASs) in Greenland marine mammals. 144, 2384-2391.
- Gebbink, W. A., Bignert, A. & Berger, U. (2016a): Perfluoroalkyl Acids (PFAAs) and Selected Precursors in the Baltic Sea Environment: Do Precursors Play a Role in Food Web Accumulation of PFAAs? Environmental Science & Technology, 50, 6354–6362.
- Gebbink, W. A., Glynn, A. & Berger, U. (2015): Temporal changes (1997-2012) of perfluoroalkyl acids and selected precursors (including isomers) in Swedish human serum. Environmental Pollution, 199, 166-173.
- Ghisi, R., Vamerali, T. & Manzetti, S. (2019): Accumulation of perfluorinated alkyl substances (PFAS) in agricultural plants: A review. Environmental Research, 169, 326–341.
- Glynn, A., Berger, U., Bignert, A., Ullah, S., Aune, M., Lignell, S. & Darnerud, P. O. (2012): Perfluorinated Alkyl Acids in Blood Serum from Primiparous Women in Sweden:

Serial Sampling during Pregnancy and Nursing, And Temporal Trends 1996–2010. Environmental Science & Technology, 46, 9071–9079.

- Goeritz, I., Falk, S., Stahl, T., Schäfers, C. & Schlechtriem, C. (2013): Biomagnification and tissue distribution of perfluoroalkyl substances (PFASs) in market-size rainbow trout (*Oncorhynchus mykiss*). Environmental Toxicology and Chemistry, 32, 2078–2088.
- Gómez-Canela, C., Fernández-Sanjuan, M., Farrés, M. & Lacorte, S. (2015): Factors affecting the accumulation of perfluoroalkyl substances in human blood. Environmental Science and Pollution Research, 22, 1480–1486.
- Gorochategui, E., Pérez-Albaladejo, E., Casas, J., Lacorte, S. & Porte, C. (2014): Perfluorinated chemicals: Differential toxicity, inhibition of aromatase activity and alteration of cellular lipids in human placental cells. Toxicology and Applied Pharmacology, 277, 124–130.
- Goudarzi, H., Miyashita, C., Okada, E., Kashino, I., Chen, C.-J., Ito, S., Araki, A., Kobayashi, S., Matsuura, H. & Kishi, R. (2017): Prenatal exposure to perfluoroalkyl acids and prevalence of infectious diseases up to 4 years of age. Environment International, 104, 132–138.
- Granum, B., Haug, L. S., Namork, E., Stølevik, S. B., Thomsen, C., Aaberge, I. S., Loveren, H. v., Løvik, M. & Nygaard, U. C. (2013): Pre-natal exposure to perfluoroalkyl substances may be associated with altered vaccine antibody levels and immune-related health outcomes in early childhood. Journal of Immunotoxicology, 10, 373–379.
- Grayson, M. (1978). Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Edition. John Wiley & Sons, New York, USA.
- Green, N. W., Schøyen, M., Øxnevad, S., Ruus, A., Allan, I., Hjermann, D. & Severinsen, G., Norwegian Environment Agency (2016). Contaminants in coastal waters of Norway 2015 (M-618). <http://www.miljodirektoratet.no/Documents/publikasjoner/M618/M618.pdf>
- Guelfo, J. L. & Higgins, C. P. (2013): Subsurface Transport Potential of Perfluoroalkyl Acids at Aqueous Film-Forming Foam (AFFF)-Impacted Sites. Environmental Science & Technology, 47, 4164–4171.
- Gyllenhammar, I., Berger, U., Sundström, M., McCleaf, P., Eurén, K., Eriksson, S., Ahlgren, S., Lignell, S., Aune, M., Kotova, N. & Glynn, A. (2015): Influence of contaminated drinking water on perfluoroalkyl acid levels in human serum – A case study from Uppsala, Sweden. Environmental Research, 140, 673–683.
- Hagenaars, A., Vergauwen, L., Coen, W. D. & Knapen, D. (2011): Structure–activity relationship assessment of four perfluorinated chemicals using a prolonged zebrafish early life stage test. Chemosphere, 82, 764–772.
- Harrad, S., Drage, D. S., Sharkey, M. & Berresheim, H. (2019): Brominated flame retardants and perfluoroalkyl substances in landfill leachate from Ireland. Science of the Total Environment, 695, 1–7.
- Haug, L. S., Thomsen, C. & Becher, G. (2009): Time Trends and the Influence of Age and Gender on Serum Concentrations of Perfluorinated Compounds in Archived Human Samples. Environmental Science & Technology, 43, 2131–2136.

- Haug, L. S., Huber, S., Schlabach, M., Becher, G. & Thomsen, C. (2011): Investigation on Per- and Polyfluorinated Compounds in Paired Samples of House Dust and Indoor Air from Norwegian Homes. Environmental Science & Technology, 45, 7991–7998.
- Hellsing, M. S., Josefsson, S., Hughes, A. V. & Ahrens, L. (2016): Sorption of perfluoroalkyl substances to two types of minerals. Chemosphere, 159, 385-391.
- Herzke, D., Nygård, T. & Heimstad, E. S., Norwegian Environment Agency (2016). Environmental pollutants in the terrestrial and urban environment 2015 (M-570). <https://www.miljodirektoratet.no/Documents/publikasjoner/M570/M570.pdf>
- Heydebreck, F., Tan, J., Xie, Z. & Ebinghaus, R. (2015): Alternative and Legacy Perfluoroalkyl Substances: Differences between European and Chinese River/Estuary Systems. Environmental Science & Technology, 49, 8386-8395.
- Higgins, C. P. & Luthy, R. G. (2006): Sorption of Perfluorinated Surfactants on Sediments. Environmental Science & Technology, 40, 7251-7256.
- Hoffman, K., Webster, T. F., Bartell, S. M., Weisskopf, M. G., Fletcher, T. & Vieira, V. M. (2011): Private Drinking Water Wells as a Source of Exposure to Perfluorooctanoic Acid (PFOA) in Communities Surrounding a Fluoropolymer Production Facility. Environmental Health Perspectives, 119, 92-97.
- Hong, S., Khim, J. S., Wang, T., Naile, J. E., Park, J., Kwon, B.-O., Song, S. J., Ryu, J., Codling, G., Jones, P. D., Lu, Y. & Giesy, J. P. (2015): Bioaccumulation characteristics of perfluoroalkyl acids (PFAAs) in coastal organisms from the west coast of South Korea. Chemosphere, 129, 157-163.
- Houtz, E. F., Sutton, R., Park, J.-S. & Sedlak, M. (2016): Poly- and perfluoroalkyl substances in wastewater: Significance of unknown precursors, manufacturing shifts, and likely AFFF impacts. Water Research, 95, 142-149.
- Huang, R., Chen, Q., Zhang, L., Luo, K., Chen, L., Zhao, S., Feng, L. & Zhang, J. (2019): Prenatal exposure to perfluoroalkyl and polyfluoroalkyl substances and the risk of hypertensive disorders of pregnancy. Environmental Health, 18, 1-10.
- Huang, M. C., Dzierlenga, A. L., Robinson, V. G., Waidyanatha, S., DeVito, M. J., Eifrid, M. A., Granville, C. A., Gibbs, S. T. & Blystone, C. R. (2019a): Toxicokinetics of perfluorobutane sulfonate (PFBS), perfluorohexane-1-sulphonic acid (PFHxS), and perfluorooctane sulfonic acid (PFOS) in male and female Hsd:Sprague Dawley SD rats after intravenous and gavage administration. Toxicology Reports, 6, 645-655.
- Hurtado, C., Domínguez, C., Pérez-Babace, L., Cañameras, N., Comas, J. & Bayona, J. M. (2016): Estimate of uptake and translocation of emerging organic contaminants from irrigation water concentration in lettuce grown under controlled conditions. Journal of Hazardous Materials, 305, 139–148.
- Huset, C. A., Chiaia, A. C., Barofsky, D. F., Jonkers, N., Kohler, H.-P. E., Ort, C., Giger, W. & Field, J. A. (2008): Occurrence and Mass Flows of Fluorochemicals in the Glatt Valley Watershed, Switzerland. Environmental Science & Technology, 42, 6369–6377.
- Impinen, A., Nygaard, U. C., Carlsen, K. C. L., Mowinckel, P., Carlsen, K. H., Haug, L. S. & Granum, B. (2018): Prenatal exposure to perfluoroalkyl substances (PFASs)

associated with respiratory tract infections but not allergy- and asthma-related health outcomes in childhood. Environmental Research, 160, 518–523.

- Ingelido, A. M., Abballe, A., Gemma, S., Dellatte, E., Iacovella, N., Angelis, G. D., Zampaglioni, F., Marra, V., Miniero, R., Valentini, S., Russo, F., Vazzoler, M., Testai, E. & Felip, E. D. (2018): Biomonitoring of perfluorinated compounds in adults exposed to contaminated drinking water in the Veneto Region, Italy. Environment International, 110, 149-159.
- Iniguez, M. A., Lecea, L. D., Guadano-Ferraz, A., Morte, B., Gerendasy, D., Sutcliffe, J. G. & Bernal, J. (1996): Cell-specific effects of thyroid hormone on RC3/neurogranin expression in rat brain. Endocrinology, 137, 1032–1041.
- Iñiguez, M. A., Rodriguez-Peña, A., Ibarrola, N., Aguilera, M., Muñoz, A. & Bernal, J. (1993): Thyroid hormone regulation of RC3, a brain-specific gene encoding a protein kinase-C substrate. Endocrinology, 133, 467–473.
- International Organization for Standardization (2007). ISO 11348-3:2007 Water quality -Determination of the inhibitory effect of water samples on the light emission of *Vibrio fischeri* (Luminescent bacteria test). <https://www.iso.org/standard/40518.html>
- IRSA-CNR, Ministero dell’Ambiente e della Tutela del Territorio e del Mare, Italy (2013). Realizzazione di uno studio di valutazione del Rischio Ambientale e Sanitario associato alla contaminazione da sostanze perfluoroalchiliche (PFAS) nel Bacino del Po e nei principali bacini fluviali italiani. http://www.minambiente.it/sites/default/files/archivio/allegati/reach/progetto_PFAS_sintesi.pdf
- Jogsten, I. E., Nadal, M., Bavel, B. v., Lindström, G. & Domingo, J. L. (2012): Per- and polyfluorinated compounds (PFCs) in house dust and indoor air in Catalonia, Spain: Implications for human exposure. Environment International, 39, 172-180.
- Johansson, J., Salter, M., Navarro, J. A., Leck, C., Nilsson, E. D. & Cousins, I. T. (2018): Global transport of perfluoroalkyl acids via sea spray aerosol. ChemRxivPreprint.
- Kaboré, H. A., Duy, S. V., Munoz, G., Méité, L., Desrosiers, M., Liu, J., Sory, T. K. & Sauv e, S. (2018): Worldwide drinking water occurrence and levels of newly-identified perfluoroalkyl and polyfluoroalkyl substances. Science of the Total Environment, 616-617, 1089-1100.
- Kallenborn, R., Berger, U. & J rnberg, U., Nordic Council of Ministers (2004). Perfluorinated alkylated substances (PFAS) in the Nordic environment. https://www.miljodirektoratet.no/globalassets/publikasjoner/klif2/nyheter/dokumenter/pfas_nmr2004.pdf
- Kar skov , P., Venier, M., Melymuk, L., Be anov , J., Vojta, Œ., ProkeŒ, R., Diamond, M. L. & Kl nov , J. (2016): Perfluorinated alkyl substances (PFASs) in household dust in Central Europe and North America. Environment International, 94, 315–324.
- Keller, J. M., Ngai, L., McNeill, J. B., Wood, L. D., Stewart, K. R., O’Connell, S. G. & Kucklick, J. R. (2012): Perfluoroalkyl contaminants in plasma of five sea turtle species: Comparisons in concentration and potential health risks. Environmental Toxicology and Chemistry, 31, 1223–1230.
- Kerstner-Wood, C., Coward, L., Gorman, G. (2003): Protein binding of perfluorohexane sulfonate, perfluorooctane sulfonate and perfluorooctanoate to plasma (human,

rat, and monkey), and various human-derived plasma protein fractions. Southern Research Institute, submitted to the U.S. Environmental Protection Agency's Administrative Record. AR226-1354.

- Kissa, E. (2001). Fluorinated Surfactants and Repellents. Marcel Dekker, Inc., USA.
- Knobeloch, L., Imm, P. & Anderson, H. (2012): Perfluoroalkyl chemicals in vacuum cleaner dust from 39 Wisconsin homes. Chemosphere, 88, 779–783.
- Kotthoff, M. & Bücking, M. (2018): Four Chemical Trends Will Shape the Next Decade's Directions in Perfluoroalkyl and Polyfluoroalkyl Substances Research. Frontiers in Chemistry, 6, 1-6.
- Kowalczyk, J., Ehlers, S., Oberhausen, A., Tischer, M., Fürst, P., Schafft, H. & Lahrssen-Wiederholt, M. (2013): Absorption, Distribution, and Milk Secretion of the Perfluoroalkyl Acids PFBS, PFHxS, PFOS, and PFOA by Dairy Cows Fed Naturally Contaminated Feed. Journal of Agricultural and Food Chemistry, 61, 2903–2912.
- Krippner, J., Brunn, H., Falk, S., Georgii, S., Schubert, S. & Stahl, T. (2014): Effects of chain length and pH on the uptake and distribution of perfluoroalkyl substances in maize (*Zea mays*). Chemosphere, 94, 85–90.
- Krippner, J., Falk, S., Brunn, H., Georgii, S., Schubert, S. & Stahl, T. (2015): Accumulation Potentials of Perfluoroalkyl Carboxylic Acids (PFCAs) and Perfluoroalkyl Sulfonic Acids (PFASs) in Maize (*Zea mays*). Journal of Agricultural and Food Chemistry, 63, 3646–3653.
- Kumar, K. S., Zushi, Y., Masunaga, S., Gilligan, M., Pride, C. & Sajwan, K. S. (2009): Perfluorinated organic contaminants in sediment and aquatic wildlife, including sharks, from Georgia, USA. Marine Pollution Bulletin, 58, 601–634.
- Kwadijk, C. J. A. F., Korytár, P. & Koelmans, A. A. (2010): Distribution of Perfluorinated Compounds in Aquatic Systems in The Netherlands. Environmental Science & Technology, 44, 3746–3751.
- Kwok, K. Y., Wang, X.-H., Ya, M., Li, Y., Zhang, X.-H., Yamashita, N., Lam, J. C. W. & Lam, P. K. S. (2015): Occurrence and distribution of conventional and new classes of per and polyfluoroalkyl substances (PFASs) in the South China Sea. Journal of Hazardous Materials, 285, 389–397.
- Kärrman, A., Wang, T. & Kallenborn, R., Nordic Council of Ministers (2019). Screening of Poly- and Perfluoroalkyl Substances (PFASs) and Extractable Organic Fluorine (EOF) in the Nordic Environment (*in press*).
- Lai, S., Song, J., Song, T., Huang, Z., Zhang, Y., Zhao, Y., Liu, G., Zheng, J., Mi, W., Tang, J., Zou, S., Ebinghaus, R. & Xie, Z. (2016): Neutral polyfluoroalkyl substances in the atmosphere over the northern South China Sea. Environmental Pollution, 214, 449-455.
- Lam, J. C. W., Lyu, J., Kwok, K. Y. & Lam, P. K. S. (2016): Perfluoroalkyl Substances (PFASs) in Marine Mammals from the South China Sea and Their Temporal Changes 2002–2014: Concern for Alternatives of PFOS? Environmental Science & Technology, 50, 6728–6736.
- Lam, N. H., Cho, C.-R., Kannan, K. & Cho, H.-S. (2017): A nationwide survey of perfluorinated alkyl substances in waters, sediment and biota collected from

- aquatic environment in Vietnam: Distributions and bioconcentration profiles. Journal of Hazardous Materials, 323, 116-127.
- Lan, Z., Zhou, M., Yao, Y. & Sun, H. (2018): Plant uptake and translocation of perfluoroalkyl acids in a wheat–soil system. Environmental Science and Pollution Research, 25, 30907-30916.
- Lang, J. R., Allred, B. M., Field, J. A., Levis, J. W. & Barlaz, M. A. (2017): National Estimate of Per- and Polyfluoroalkyl Substance (PFAS) Release to U.S. Municipal Landfill Leachate. Environmental Science & Technology, 51, 2197–2205.
- Lasier, P. J., Washington, J. W., Hassan, S. M. & Jenkins, T. M. (2011): Perfluorinated chemicals in surface waters and sediments from northwest Georgia, USA, and their bioaccumulation in *Lumbricus variegatus*. Environmental Toxicology and Chemistry, 30, 2194–2201.
- Lassen, C., Brinch, A. & Jensen, A. A., COWI A/S (2017). Investigation of sources to PFBS in the environment (M-759). <http://www.miljodirektoratet.no/Documents/publikasjoner/M759/M759.pdf>
- Lath, S., Navarro, D. A., Losic, D., Kumar, A. & McLaughlin, M. J. (2018): Sorptive remediation of perfluorooctanoic acid (PFOA) using mixed mineral and graphene/carbon-based materials. Environmental Chemistry, 15, 472-480.
- Lau, C., Thibodeaux, J. R., Hanson, R. G., Rogers, J. M., Grey, B. E., Stanton, M. E., Butenhoff, J. L. & Stevenson, L. A. (2003): Exposure to Perfluorooctane Sulfonate during Pregnancy in Rat and Mouse. II: Postnatal Evaluation. Toxicological Sciences, 74, 382–392.
- Lescord, G. L., Kidd, K. A., Silva, A. O. D., Williamson, M., Spencer, C., Wang, X. & Muir, D. C. G. (2015): Perfluorinated and Polyfluorinated Compounds in Lake Food Webs from the Canadian High Arctic. Environmental Science & Technology, 49, 2694–2702.
- Li, Y., Oliver, D. P. & Kookana, R. S. (2018): A critical analysis of published data to discern the role of soil and sediment properties in determining sorption of per and polyfluoroalkyl substances (PFASs). Science of the Total Environment, 628-629, 110-120.
- Li, X., Gao, K., Dong, S., Liu, X., Fu, K., Wang, P., Zhang, A., Su, X. & Fu, J. (2019): Length-specific occurrence and profile of perfluoroalkyl acids (PFAAs) in animal protein feeds. Journal of Hazardous Materials, 373, 224-231.
- Li, F., Fang, X., Zhou, Z., Liao, X., Zou, J., Yuan, B. & Sun, W. (2019a): Adsorption of perfluorinated acids onto soils: Kinetics, isotherms, and influences of soil properties. Science of the Total Environment, 649, 504–514.
- Liao, C., Wang, T., Cui, L., Zhou, Q., Duan, S. & Jiang, G. (2009): Changes in Synaptic Transmission, Calcium Current, and Neurite Growth by Perfluorinated Compounds Are Dependent on the Chain Length and Functional Group. Environmental Science & Technology, 43, 2099–2104.
- Lieder, P. H., Chang, S.-C., York, R. G. & Butenhoff, J. L. (2009a): Toxicological evaluation of potassium perfluorobutanesulfonate in a 90-day oral gavage study with Sprague–Dawley rats. Toxicology, 255, 45–52.

- Lieder, P. H., York, R. G., Hakes, D. C., Chang, S.-C. & Butenhoff, J. L. (2009b): A two-generation oral gavage reproduction study with potassium perfluorobutanesulfonate (K+PFBS) in Sprague Dawley rats. Toxicology, 259, 33–45.
- Lin, A. Y.-C., Panchangam, S. C., Tsai, Y.-T. & Yu, T.-H. (2014): Occurrence of perfluorinated compounds in the aquatic environment as found in science park effluent, river water, rainwater, sediments, and biotissues. Environmental Monitoring and Assessment, 186, 3265–3275.
- Lindstrom, A. B., Strynar, M. J., Delinsky, A. D., Nakayama, S. F., McMillan, L., Libelo, E. L., Neill, M. & Thomas, L. (2011): Application of WWTP Biosolids and Resulting Perfluorinated Compound Contamination of Surface and Well Water in Decatur, Alabama, USA. Environmental Science & Technology, 45, 8015–8021.
- Liu, J. & Avendaño, S. M. (2013): Microbial degradation of polyfluoroalkyl chemicals in the environment: A review. Environment International, 61, 98–114.
- Liu, Y., Cao, Z., Zong, W. & Liu, R. (2017): Interaction rule and mechanism of perfluoroalkyl sulfonates containing different carbon chains with human serum albumin. RSC Advances, 7, 24781–24788.
- Liu, Y., Zhang, Y., Li, J., Wu, N., Li, W. & Niu, Z. (2019): Distribution, partitioning behavior and positive matrix factorization based source analysis of legacy and emerging polyfluorinated alkyl substances in the dissolved phase, surface sediment and suspended particulate matter around coastal areas of Bohai Bay, China. Environmental Pollution, 246, 34–44.
- Loi, E. I. H., Yeung, L. W. Y., Taniyasu, S., Lam, P. K. S., Kannan, K. & Yamashita, N. (2011): Trophic Magnification of Poly- and Perfluorinated Compounds in a Subtropical Food Web. Environmental Science & Technology, 45, 5506–5513.
- Loos, R., Locoro, G., Comero, S., Contini, S., Schwesig, D., Werres, F., Balsaa, P., Gans, O., Weiss, S., Blaha, L., Bolchi, M. & Gawlik, B. M. (2010): Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. Water Research, 44, 4115–4126.
- Lou, Q.-Q., Ren, D.-K., Zhang, Y.-F., Xu, H.-M., Zhou, Z., Shi, Y.-L., Zhao, Y.-X., Ge, Y.-N., Wei, W.-J. & Qin, Z.-F. (2013): Effects of perfluorooctanesulfonate and perfluorobutanesulfonate on the growth and sexual development of *Xenopus laevis*. Ecotoxicology, 22, 1133–1144.
- Luebker, D. J., Hansen, K. J., Butenhoff, J. L., Bass, N. M. & Seacat, A. M. (2002): Interactions of fluorochemicals with rat liver fatty acid-binding protein. Toxicology, 176, 175–185.
- Mak, Y. L., Taniyasu, S., Yeung, L. W. Y., Lu, G., Jin, L., Yang, Y., Lam, P. K. S., Kannan, K. & Yamashita, N. (2009): Perfluorinated Compounds in Tap Water from China and Several Other Countries. Environmental Science & Technology, 43, 4824–4829.
- Manzano-Salgado, C. B., Casas, M., Lopez-Espinosa, M.-J., Ballester, F., Basterrechea, M., Grimalt, J. O., Jiménez, A.-M., Kraus, T., Schettgen, T., Sunyer, J. & Vrijheid, M. (2015): Transfer of perfluoroalkyl substances from mother to fetus in a Spanish birth cohort. Environmental Research, 142, 471–478.

- Martin, J. W., Mabury, S. A., Solomon, K. R. & Muir, D. C. G. (2003a): Bioconcentration and tissue distribution of perfluorinated acids in rainbow trout (*Oncorhynchus mykiss*). Environmental Toxicology and Chemistry, 22, 196–204.
- Martin, J. W., Mabury, S. A., Solomon, K. R. & Muir, D. C. G. (2003b): Dietary accumulation of perfluorinated acids in juvenile rainbow trout (*Oncorhynchus mykiss*) Environmental Toxicology and Chemistry, 22, 189-195.
- Meyer, T. & Wania, F. (2007): What environmental fate processes have the strongest influence on a completely persistent organic chemical's accumulation in the Arctic? Atmospheric Environment, 41, 2757–2767.
- Milinovic, J., Lacorte, S., Vidal, M. & Rigol, A. (2015): Sorption behaviour of perfluoroalkyl substances in soils. Science of the Total Environment, 511, 63-71.
- Minnesota Department of Health (2011). Perfluorobutane sulfonate. <http://www.health.state.mn.us/divs/eh/risk/guidance/gw/pfbs.pdf>
- Moody, C. A., Hebert, G. N., Strauss, S. H. & Field, J. A. (2003): Occurrence and persistence of perfluorooctanesulfonate and other perfluorinated surfactants in groundwater at a fire-training area at Wurtsmith Air Force Base, Michigan, USA. Journal of Environmental Monitoring, 5, 341-345.
- Moody, C. A. & Field, J. A. (2000): Perfluorinated Surfactants and the Environmental Implications of Their Use in Fire-Fighting Foams. Environmental Science & Technology, 34, 3864-3870.
- Munoz, G., Giraudel, J.-L., Botta, F., Lestremau, F., Dévier, M.-H., Budzinski, H. & Labadie, P. (2015): Spatial distribution and partitioning behavior of selected poly- and perfluoroalkyl substances in freshwater ecosystems: A French nationwide survey. Science of the Total Environment, 517, 48-56.
- Müller, C. E., LeFevre, G. H., Timofte, A. E., Hussain, F. A., Sattely, E. S. & Luthy, R. G. (2016): Competing mechanisms for perfluoroalkyl acid accumulation in plants revealed using an Arabidopsis model system. Environmental Toxicology and Chemistry, 35, 1138-1147.
- Möller, A., Ahrens, L., Surm, R., Westerveld, J., Wielen, F. v. d., Ebinghaus, R. & Voogt, P. d. (2010): Distribution and sources of polyfluoroalkyl substances (PFAS) in the River Rhine watershed. Environmental Pollution, 158, 3243-3250.
- Nagel, R. (2002): DarT: The Embryo Test with the Zebrafish *Danio rerio* - a General Model in Ecotoxicology and Toxicology. ALTEX, 19, 38-48.
- Naile, J. E., Khim, J. S., Hong, S., Park, J., Kwon, B.-O., Ryu, J. S., Hwang, J. H., Jones, P. D. & Giesy, J. P. (2013): Distributions and bioconcentration characteristics of perfluorinated compounds in environmental samples collected from the west coast of Korea. Chemosphere, 90, 387–394.
- Navarro, I., Torre, A. d. I., Sanz, P., Porcel, M. Á., Pro, J., Carbonell, G. & Martínez, M. d. I. Á. (2017): Uptake of perfluoroalkyl substances and halogenated flame retardants by crop plants grown in biosolids-amended soils. Environmental Research, 152, 199-206.
- Navarro, I., Torre, A. d. I., Sanz, P., Pro, J., Carbonell, G. & Martínez, M. d. I. Á. (2016): Bioaccumulation of emerging organic compounds (perfluoroalkyl substances and halogenated flame retardants) by earthworm in biosolid amended soils. Environmental Research, 149, 32-39.

- Newsted, J. L., Beach, S. A., Gallagher, S. P. & Giesy, J. P. (2008): Acute and Chronic Effects of Perfluorobutane Sulfonate (PFBS) on the Mallard and Northern Bobwhite Quail. Archives of Environmental Contamination and Toxicology, 54, 535–545.
- Ng, C. A. & Hungerbühler, K. (2014): Bioaccumulation of Perfluorinated Alkyl Acids: Observations and Models. Environmental Science & Technology, 48, 4637–4648.
- Nguyen, M. A., Wiberg, K., Ribeli, E., Josefsson, S., Futter, M., Gustavsson, J. & Ahrens, L. (2017): Spatial distribution and source tracing of per- and polyfluoroalkyl substances (PFASs) in surface water in Northern Europe. Environmental Pollution, 220, 1438-1446.
- Nielsen, C. J., University of Oslo (2017). Potential PFBS and PFHxS Precursors - Literature study and theoretical assessment of abiotic degradation pathways leading to PFBS and PFHxS (M-792). <http://www.miljodirektoratet.no/Documents/publikasjoner/M792/M792.pdf>
- Norwegian Environment Agency (2018). Risk Management Option Analysis Conclusion Document: Perfluorobutane sulfonic acid (PFBS), its salts and related substances. <https://echa.europa.eu/documents/10162/1cb252bc-3952-3d5e-48b3-e78d5f5f0255>
- National Toxicology Program, NTP (2019). The toxicity studies of perfluoroalkyl sulfonates (perfluorobutane sulfonic acid, perfluorohexane sulfonate potassium salt, and perfluorooctane sulfonic acid) administered by gavage to Sprague Dawley (Hsd:Sprague Dawley SD) rats. https://ntp.niehs.nih.gov/ntp/htdocs/st_rpts/tox096_508.pdf
- Numata, J., Kowalczyk, J., Adolphs, J., Ehlers, S., Schafft, H., Fuerst, P., Müller-Graf, C., Lahrssen-Wiederholt, M. & Greiner, M. (2014): Toxicokinetics of Seven Perfluoroalkyl Sulfonic and Carboxylic Acids in Pigs Fed a Contaminated Diet. Journal of Agricultural and Food Chemistry, 62, 6861–6870.
- Ochoa-Herrera, V. & Sierra-Alvarez, R. (2008): Removal of perfluorinated surfactants by sorption onto granular activated carbon, zeolite and sludge. Chemosphere, 72, 1588–1593.
- OECD (2002). Hazard assessment of perfluorooctane sulfonate (PFOS) and its salts. <http://www.oecd.org/env/ehs/risk-assessment/2382880.pdf>
- OECD/UNEP Global PFC Group (2013). Synthesis paper on per- and polyfluorinated chemicals (PFCs). <http://www.oecd.org/chemicalsafety/risk-management/synthesis-paper-on-per-and-polyfluorinated-chemicals.htm>
- OECD/UNEP, the Global PFC Group (2018) "Toward a new comprehensive Global Database of Per- And Polyfluoroalkyl Substances (PFASs)." Available at <http://www.oecd.org/chemicalsafety/portal-perfluorinated-chemicals/> (accessed on 2019-01-31).
- Olsen, G. W., Butenhoff, J. L. & Zobel, L. R. (2009a): Perfluoroalkyl chemicals and human fetal development: An epidemiologic review with clinical and toxicological perspectives. Reproductive Toxicology, 27, 212-230.
- Olsen, G. W., Chang, S.-C., Noker, P. E., Gorman, G. S., Ehresman, D. J., Lieder, P. H. & Butenhoff, J. L. (2009b): A comparison of the pharmacokinetics of

- perfluorobutanesulfonate (PFBS) in rats, monkeys, and humans. Toxicology, 256, 65-74.
- Olsen, G. W., Lange, C. C., Ellefson, M. E., Mair, D. C., Church, T. R., Goldberg, C. L., Herron, R. M., Medhdizadehkashi, Z., Nobiletti, J. B., Rios, J. A., Reagen, W. K. & Zobel, L. R. (2012): Temporal Trends of Perfluoroalkyl Concentrations in American Red Cross Adult Blood Donors, 2000–2010. Environmental Science & Technology, 46, 6330–6338.
- Olsen, G. W., Mair, D. C., Lange, C. C., Harrington, L. M., Church, T. R., Goldberg, C. L., Herron, R. M., Hanna, H., Nobiletti, J. B., Rios, J. A., Reagen, W. K. & Ley, C. A. (2017): Per- and polyfluoroalkyl substances (PFAS) in American Red Cross adult blood donors, 2000-2015. Environmental Research, 157, 87-95.
- Orozco, A. & Valverde-R, C. (2005): Thyroid Hormone Deiodination in Fish. Thyroid, 15, 799-813.
- Pan, Y., Zhang, H., Cui, Q., Sheng, N., Yeung, L. W. Y., Sun, Y., Guo, Y. & Dai, J. (2018): Worldwide Distribution of Novel Perfluoroether Carboxylic and Sulfonic Acids in Surface Water. Environmental Science & Technology, 52, 7621–7629.
- Parsons, J. R., Sáez, M., Dolfin, J. & Voogt, P. d. (2008): Biodegradation of Perfluorinated Compounds. Reviews of Environmental Contamination and Toxicology, 196, 53-71.
- Pasquini, L., Merlin, C., Hassenboehler, L., Munoz, J.-F., Pons, M.-N. & Görner, T. (2013): Impact of certain household micropollutants on bacterial behavior. Toxicity tests/study of extracellular polymeric substances in sludge. Science of the Total Environment, 463–464, 355–365.
- Pennings, J. L. A., Jennen, D. G. J., Nygaard, U. C., Namork, E., Haug, L. S., Loveren, H. v. & Granum, B. (2016): Cord blood gene expression supports that prenatal exposure to perfluoroalkyl substances causes depressed immune functionality in early childhood. Journal of Immunotoxicology, 13, 173-180.
- Pereira, H. C., Ullberg, M., Kleja, D. B., Gustafsson, J. P. & Ahrens, L. (2018): Sorption of perfluoroalkyl substances (PFASs) to an organic soil horizon - Effect of cation composition and pH. Chemosphere, 207, 183-191.
- Perez, F., Llorca, M., Farré, M. & Barceló, D. (2012): Automated analysis of perfluorinated compounds in human hair and urine samples by turbulent flow chromatography coupled to tandem mass spectrometry. Analytical and Bioanalytical Chemistry, 402, 2369–2378.
- Pérez, F., Nadal, M., Navarro-Ortega, A., Fàbrega, F., Domingo, J. L., Barceló, D. & Farré, M. (2013): Accumulation of perfluoroalkyl substances in human tissues. Environment International, 59, 354-362.
- Pi, N., Ng, J. Z. & Kelly, B. C. (2017): Uptake and elimination kinetics of perfluoroalkyl substances in submerged and free-floating aquatic macrophytes: Results of mesocosm experiments with *Echinodorus horemanii* and *Eichhornia crassipes*. Water Research, 117, 167-174.
- Pickard, H. M., Criscitiello, A. S., Spencer, C., Sharp, M. J., Muir, D. C. G., Silva, A. O. D. & Young, C. J. (2018): Continuous non-marine inputs of per- and polyfluoroalkyl substances to the High Arctic: a multi-decadal temporal record. Atmospheric Chemistry and Physics, 18, 5045–5058.

- Pico, Y., Blasco, C., Farré, M. & Barceló, D. (2012): Occurrence of perfluorinated compounds in water and sediment of L'Albufera Natural Park (València, Spain). Environmental Science and Pollution Research, 19, 946-957.
- Poothong, S., Thomsen, C., Padilla-Sanchez, J. A., Papadopoulou, E. & Haug, L. S. (2017): Distribution of Novel and Well-Known Poly- and Perfluoroalkyl Substances (PFASs) in Human Serum, Plasma, and Whole Blood. Environmental Science & Technology, 51, 13388-13396.
- Post, G. B., Louis, J. B., Lippincott, R. L. & Procopio, N. A. (2013): Occurrence of Perfluorinated Compounds in Raw Water from New Jersey Public Drinking Water Systems. Environmental Science & Technology, 47, 13266–13275.
- Power, D. M., Llewellyn, L., Faustino, M., Nowell, M. A., Björnsson, B. T., Einarsdottir, I. E., Canario, A. V. M. & Sweeney, G. E. (2001): Thyroid hormones in growth and development of fish. Comparative Biochemistry and Physiology Part C, 130, 447-459.
- Qi, W., Clark, J. M., Timme-Laragy, A. R. & Park, Y. (2018): Perfluorobutanesulfonic acid (PFBS) potentiates adipogenesis of 3T3-L1 adipocytes. Food and Chemical Toxicology, 120, 340–345.
- Qian, J., Shen, M., Wang, P., Wang, C., Hou, J., Ao, Y., Liu, J. & Li, K. (2017): Adsorption of perfluorooctane sulfonate on soils: Effects of soil characteristics and phosphate competition. Chemosphere, 168, 1383-1388.
- Qu, B., Zhao, H. & Zhou, J. (2010): Toxic effects of perfluorooctane sulfonate (PFOS) on wheat (*Triticum aestivum* L.) plant. Chemosphere, 79, 555–560.
- Quinete, N., Orata, F., Maes, A., Gehron, M., Bauer, K.-H., Moreira, I. & Wilken, R.-D. (2010): Degradation Studies of New Substitutes for Perfluorinated Surfactants. Archives of Environmental Contamination and Toxicology, 59, 20-30.
- Rahman, M. F., Peldszus, S. & Anderson, W. B. (2014): Behaviour and fate of perfluoroalkyl and polyfluoroalkyl substances (PFASs) in drinking water treatment: A review. Water Research, 50, 318-340.
- Ramhøj, L., Axelstad, M., Boberg, J., Christiansen, S. & Hass, U. (2016): Developmental toxicity and endocrine disruption by PFHxS (perfluorohexane sulphonate) in rats. Reproductive Toxicology, 64, 15–21.
- Reemtsma, T., Berger, U., Arp, H. P. H., Gallard, H., Knepper, T. P., Neumann, M., Quintana, J. B. & Voogt, P. d. (2016): Mind the Gap: Persistent and Mobile Organic Compounds - Water Contaminants That Slip Through. Environmental Science & Technology, 50, 10308-10315.
- Ren, X.-M., Qin, W.-P., Cao, L.-Y., Zhang, J., Yang, Y., Wan, B. & Guo, L.-H. (2016): Binding interactions of perfluoroalkyl substances with thyroid hormone transport proteins and potential toxicological implications. Toxicology, 366-367, 32-42.
- Rodea-Palomares, I., González-García, C., Leganés, F. & Fernández-Piñas, F. (2009): Effect of pH, EDTA, and Anions on Heavy Metal Toxicity Toward a Bioluminescent Cyanobacterial Bioreporter. Archives of Environmental Contamination and Toxicology, 57, 477–487.
- Rosal, R., Rodea-Palomares, I., Boltes, K., Fernández-Piñas, F., Leganés, F. & Petre, A. (2010): Ecotoxicological assessment of surfactants in the aquatic environment:

- Combined toxicity of docusate sodium with chlorinated pollutants. Chemosphere, *81*, 288–293.
- Rosenmai, A. K., Ahrens, L., Godec, T. I., Lundqvist, J. & Oskarsson, A. (2018): Relationship between peroxisome proliferator-activated receptor alpha activity and cellular concentration of 14 perfluoroalkyl substances in HepG2 cells. Journal of Applied Toxicology, *38*, 219–226.
- Rostkowski, P., Yamashita, N., So, I. M. K., Taniyasu, S., Lam, P. K. S., Falandysz, J., Lee, K. T., Kim, S. K., Khim, J. S., Im, S. H., Newsted, J. L., Jones, P. D., Kannan, K. & Giesy, J. P. (2006): Perfluorinated compounds in streams of the Shihwa industrial zone and Lake Shihwa, South Korea. Environmental Toxicology and Chemistry, *25*, 2374-2380.
- Rotander, A., Toms, L.-M. L., Aylward, L., Kay, M. & Mueller, J. F. (2015): Elevated levels of PFOS and PFHxS in firefighters exposed to aqueous film forming foam (AFFF). Environment International, *82*, 28-34.
- Routti, H., Aars, J., Fuglei, E., Hanssen, L., Lone, K., Polder, A., Pedersen, Å. Ø., Tartu, S., Welker, J. M. & Yoccoz, N. G. (2017): Emission Changes Dwarf the Influence of Feeding Habits on Temporal Trends of Per- and Polyfluoroalkyl Substances in Two Arctic Top Predators. Environmental Science & Technology, *51*, 11996-12006.
- Routti, H., Gabrielsen, G. W., Herzke, D., Kovacs, K. M. & Lydersen, C. (2016): Spatial and temporal trends in perfluoroalkyl substances (PFASs) in ringed seals (*Pusa hispida*) from Svalbard. Environmental Pollution, *214*, 230-238.
- Rumpler, J., Das, K., Wood, C., Lau, C., Strynar, M. & Wambaugh, J., US EPA (2016) "Pharmacokinetic profiles of perfluorobutane sulfonate and activation of hepatic genes in mice." Accessed on 2019-01-31. Available at https://cfpub.epa.gov/si/si_public_record_Report.cfm?Lab=NHEERL&dirEntryId=311561
- Saez, M., Moreno, D. V., Jimenez, B. & Leeuwen, S. v. (2008): Uncommon PFC-profile in Arctic ice samples from Russia. Organohalogen Compounds, *70*, 1870-1873.
- Salihovic, S., Kärrman, A., Lind, L., Lind, P. M., Lindström, G. & Bavel, B. v. (2015): Perfluoroalkyl substances (PFAS) including structural PFOS isomers in plasma from elderly men and women from Sweden: Results from the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS). Environment International, *82*, 21-27.
- Sant, K. E., Jacobs, H. M., Borofski, K. A., Moss, J. B. & Timme-Laragy, A. R. (2017): Embryonic exposures to perfluorooctanesulfonic acid (PFOS) disrupt pancreatic organogenesis in the zebrafish, *Danio rerio*. Environmental Pollution, *220*, 807-817.
- Sant, K. E., Venezia, O. L., Sinno, P. P. & Timme-Laragy, A. R. (2019): Perfluorobutanesulfonic Acid Disrupts Pancreatic Organogenesis and Regulation of Lipid Metabolism in the Zebrafish, *Danio rerio*. Toxicological Sciences, *167*, 258–268.
- Scher, D. P., Kelly, J. E., Huset, C. A., Barry, K. M., Hoffbeck, R. W., Yingling, V. L. & Messing, R. B. (2018): Occurrence of perfluoroalkyl substances (PFAS) in garden produce at homes with a history of PFAS-contaminated drinking water. Chemosphere, *196*, 548-555.

- Schlabach, M., Bavel, B. v., Lomba, J. A. B., Borgen, A., Gabrielsen, G. W., Götsch, A., Halse, A.-K., Hanssen, L., Krogseth, I. S., Nikiforov, V., Nygård, T., Nizzetto, P. B., Reid, M., Rostkowski, P. & Samanipour, S., Norwegian Environment Agency (2018). Screening Programme 2017: AMAP Assessment compounds (M-1080). <http://www.miljodirektoratet.no/Documents/publikasjoner/M1080/M1080.pdf>
- Schröter-Kermani, C., Müller, J., Jürling, H., Conrad, A. & Schulte, C. (2013): Retrospective monitoring of perfluorocarboxylates and perfluorosulfonates in human plasma archived by the German Environmental Specimen Bank. *International Journal of Hygiene and Environmental Health*, 216, 633-640.
- Schwanz, T. G., Llorca, M., Farré, M. & Barceló, D. (2016): Perfluoroalkyl substances assessment in drinking waters from Brazil, France and Spain. *Science of the Total Environment*, 539, 143-152.
- Shan, G., Wei, M., Zhu, L., Liu, Z. & Zhang, Y. (2014): Concentration profiles and spatial distribution of perfluoroalkyl substances in an industrial center with condensed fluorochemical facilities. *Science of the Total Environment*, 490, 351–359.
- Shao, M., Ding, G., Zhang, J., Wei, L., Xue, H., Zhang, N., Li, Y., Chen, G. & Sun, Y. (2016): Occurrence and distribution of perfluoroalkyl substances (PFASs) in surface water and bottom water of the Shuangtaizi Estuary, China. *Environmental Pollution*, 216, 675-681.
- Shi, X., Liu, C., Wu, G. & Zhou, B. (2009): Waterborne exposure to PFOS causes disruption of the hypothalamus–pituitary–thyroid axis in zebrafish larvae. *Chemosphere*, 77, 1010–1018.
- Shi, Y., Vestergren, R., Nost, T. H., Zhou, Z. & Cai, Y. (2018): Probing the Differential Tissue Distribution and Bioaccumulation Behavior of Per- and Polyfluoroalkyl Substances of Varying Chain-Lengths, Isomeric Structures and Functional Groups in Crucian Carp. *Environmental Science & Technology*, 52, 4592–4600.
- Skaar, J. S., Ræder, E. M., Lyche, J. L., Ahrens, L. & Kallenborn, R. (2019): Elucidation of contamination sources for poly- and perfluoroalkyl substances (PFASs) on Svalbard (Norwegian Arctic). *Environmental Science and Pollution Research*, 26, 7356–7363.
- Skarbøvik, E., Allan, I., Sample, J. E., Greipsland, I., Selvik, J. R., Schanke, L. B., Beldring, S., Stålnacke, P. & Kaste, Ø., Norwegian Environment Agency (2017). Riverine inputs and direct discharges to Norwegian coastal waters – 2016 (M-862).
- Skutlarek, D., Exner, M. & Färber, H. (2006): Perfluorinated Surfactants in Surface and Drinking Waters. *Environmental Science and Pollution Research*, 13, 299-307.
- Slotkin, T. A., MacKillop, E. A., Melnick, R. L., Thayer, K. A. & Seidler, F. J. (2008): Developmental Neurotoxicity of Perfluorinated Chemicals Modeled in Vitro. *Environmental Health Perspectives*, 116, 716-722.
- Smit, L. A. M., Lenters, V., Høyer, B. B., Lindh, C. H., Pedersen, H. S., Liermontova, I., Jönsson, B. A. G., Piersma, A. H., Bonde, J. P., Toft, G., Vermeulen, R. & Heederik, D. (2015): Prenatal exposure to environmental chemical contaminants and asthma and eczema in school-age children. *Allergy*, 70, 653–660.
- Song, X., Tang, S., Zhu, H., Chen, Z., Zang, Z., Zhang, Y., Niu, X., Wang, X., Yin, H., Zeng, F. & He, C. (2018): Biomonitoring PFAAs in blood and semen samples:

- Investigation of a potential link between PFAAs exposure and semen mobility in China. Environment International, 113, 50–54.
- Spaan, K., Haigis, A.-C., Weiss, J. & Legradi, J. (2019): Effects of 25 thyroid hormone disruptors on zebrafish embryos: A literature review of potential biomarkers. Science of the Total Environment, 656, 1238–1249.
- Stefani, F., Rusconi, M., Valsecchi, S. & Marziali, L. (2014): Evolutionary ecotoxicology of perfluoroalkyl substances (PFASs) inferred from multigenerational exposure: A case study with *Chironomus riparius* (Diptera, Chironomidae). Aquatic Toxicology, 156, 41-51.
- Stock, N. L., Furdui, V. I., Muir, D. C. G. & Mabury, S. A. (2007): Perfluoroalkyl Contaminants in the Canadian Arctic: Evidence of Atmospheric Transport and Local Contamination. Environmental Science & Technology, 41, 3529-3536.
- Swedish Chemicals Agency (2015). Förekomst och användning av högfluorerade ämnen och alternativ (6/15). <http://www.kemi.se/global/rapporter/2015/rapport-6-15-forekomst-och-anvandning-av-hogfluorerade-amnen-och-alternativ.pdf>
- Taniyasu, S., Yamashita, N., Yamazaki, E., Petrick, G. & Kannan, K. (2013): The environmental photolysis of perfluorooctanesulfonate, perfluorooctanoate, and related fluorochemicals. Chemosphere, 90, 1686-1692.
- Torre, A. d. I., Navarro, I., Sanz, P. & Martínez, M. d. I. Á. (2019): Occurrence and human exposure assessment of perfluorinated substances in house dust from three European countries. Science of the Total Environment, 685, 308-314.
- Tröger, R., Klöckner, P., Ahrens, L. & Wiberg, K. (2018): Micropollutants in drinking water from source to tap - Method development and application of a multiresidue screening method. Science of the Total Environment, 627, 1404–1432.
- Ulhaq, M., Carlsson, G., Örn, S. & Norrgren, L. (2013a): Comparison of developmental toxicity of seven perfluoroalkyl acids to zebrafish embryos. Environmental Toxicology and Pharmacology, 36, 423-426.
- Ulhaq, M., Örn, S., Carlsson, G., Morrison, D. A. & Norrgren, L. (2013b): Locomotor behavior in zebrafish (*Danio rerio*) larvae exposed to perfluoroalkyl acids. Aquatic Toxicology, 144-145, 332-340.
- Ullah, S., Alsberg, T. & Berger, U. (2011): Simultaneous determination of perfluoroalkyl phosphonates, carboxylates, and sulfonates in drinking water. Journal of Chromatography A, 1218, 6388– 6395.
- UNEP (2006). Stockholm Convention on Persistent Organic Pollutants - Persistent Organic Pollutants Review Committee: Risk profile on perfluorooctane sulfonate. http://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=2ahUKewj1zsHBpdffAhWctYsKHYTVDAAQFjAAegQIARAC&url=http%3A%2F%2Fchm.pops.int%2FPortals%2F0%2Fdownload.aspx%3Fd%3DUNEP-POPS-POPRC.2-17-Add.5.English.PDF&usq=AOvVaw2-I7TQ27_jryZP_nopwY01
- Valsecchi, S., Rusconi, M., Mazzoni, M., Viviano, G., Pagnotta, R., Zaghi, C., Serrini, G. & Polesello, S. (2015): Occurrence and sources of perfluoroalkyl acids in Italian river basins. Chemosphere, 129, 126-134.
- Van de Vijver, K. I., Hoff, P., Das, K., Brasseur, S., Dongen, W. V., Esmans, E., Reijnders, P., Blust, R. & Coen, W. D. (2005): Tissue Distribution of Perfluorinated Chemicals

- in Harbor Seals (*Phoca vitulina*) from the Dutch Wadden Sea. Environmental Science & Technology, 39, 6978-6984.
- Van de Vijver, K. I., Holsbeek, L., Das, K., Blust, R., Joiris, C. & Coen, W. D. (2007): Occurrence of Perfluorooctane Sulfonate and Other Perfluorinated Alkylated Substances in Harbor Porpoises from the Black Sea. Environmental Science & Technology, 41, 315-320.
- Vecitis, C. D., Park, H., Cheng, J., Mader, B. T. & Hoffmann, M. R. (2009): Treatment technologies for aqueous perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA). Frontiers of Environmental Science & Engineering in China, 3, 129-151.
- Veillette, J., Muir, D. C. G., Antoniades, D., Spencer, C., Loewen, T. N., Babaluk, J. A., Reist, J. D. & Vincent, W. F. (2012): Perfluorinated Chemicals in Meromictic Lakes on the Northern Coast of Ellesmere Island, High Arctic Canada. Arctic, 65, 245-256.
- Verreault, J., Berger, U. & Gabrielsen, G. W. (2007): Trends of Perfluorinated Alkyl Substances in Herring Gull Eggs from Two Coastal Colonies in Northern Norway: 1983-2003. Environmental Science & Technology, 41, 6671-6677.
- Vierke, L., Möller, A. & Klitzke, S. (2014): Transport of perfluoroalkyl acids in a water-saturated sediment column investigated under near-natural conditions. Environmental Pollution, 186, 7-13.
- Vongphachan, V., Cassone, C. G., Wu, D., Chiu, S., Crump, D. & Kennedy, S. W. (2011): Effects of Perfluoroalkyl Compounds on mRNA Expression Levels of Thyroid Hormone-Responsive Genes in Primary Cultures of Avian Neuronal Cells. Toxicological Sciences, 120, 392-402.
- Wang, Z., MacLeod, M., Cousins, I. T., Scheringer, M. & Hungerbühler, K. (2011): Using COSMOtherm to predict physicochemical properties of poly- and perfluorinated alkyl substances (PFASs). Environmental Chemistry, 8, 389-398.
- Wang, F. & Shih, K. (2011): Adsorption of perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) on alumina: Influence of solution pH and cations. Water Research, 45, 2925-2930.
- Wang, F., Liu, C. & Shih, K. (2012): Adsorption behavior of perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) on boehmite. Chemosphere, 89, 1009-1014.
- Wang, Z., Cousins, I. T., Scheringer, M. & Hungerbühler, K. (2015): Hazard assessment of fluorinated alternatives to long-chain perfluoroalkyl acids (PFAAs) and their precursors: Status quo, ongoing challenges and possible solutions. Environment International, 75, 172-179.
- Wang, Z., Cousins, I. T. & Scheringer, M. (2015a): Comment on "The environmental photolysis of perfluorooctanesulfonate, perfluorooctanoate, and related fluorochemicals". Chemosphere, 122, 301-303.
- Wang, F., Shih, K. & Leckie, J. O. (2015b): Effect of humic acid on the sorption of perfluorooctane sulfonate (PFOS) and perfluorobutane sulfonate (PFBS) on boehmite. Chemosphere, 118, 213-218.

- Wang, B., Chen, Q., Shen, L., Zhao, S., Pang, W. & Zhang, J. (2016a): Perfluoroalkyl and polyfluoroalkyl substances in cord blood of newborns in Shanghai, China: Implications for risk assessment. Environment International, 97, 7-14.
- Wang, P., Lu, Y., Wang, T., Zhu, Z., Li, Q., Meng, J., Su, H., Johnson, A. C. & Sweetman, A. J. (2016b): Coupled production and emission of short chain perfluoroalkyl acids from a fast developing fluorochemical industry: Evidence from yearly and seasonal monitoring in Daling River Basin, China. Environmental Pollution, 218, 1234-1244.
- Wang, B., Zhang, R., Jin, F., Lou, H., Mao, Y., Zhu, W., Zhou, W., Zhang, P. & Zhang, J. (2017): Perfluoroalkyl substances and endometriosis-related infertility in Chinese women. Environment International, 102, 207-212.
- Wania, F. (2003): Assessing the Potential of Persistent Organic Chemicals for Long-Range Transport and Accumulation in Polar Regions. Environmental Science & Technology, 37, 1344-1351.
- Wei, S., Chen, L. Q., Taniyasu, S., So, M. K., Murphy, M. B., Yamashita, N., Yeung, L. W. Y. & Lam, P. K. S. (2007): Distribution of perfluorinated compounds in surface seawaters between Asia and Antarctica. Marine Pollution Bulletin, 54, 1813-1838.
- Wei, C., Song, X., Wang, Q. & Hu, Z. (2017): Sorption kinetics, isotherms and mechanisms of PFOS on soils with different physicochemical properties. Ecotoxicology and Environmental Safety, 142, 40-50.
- Weiss, J. M., Andersson, P. L., Lamoree, M. H., Leonards, P. E. G., Leeuwen, S. P. J. v. & Hamers, T. (2009): Competitive Binding of Poly- and Perfluorinated Compounds to the Thyroid Hormone Transport Protein Transthyretin. Toxicological Sciences, 109, 206-216.
- Wen, B., Li, L., Liu, Y., Zhang, H., Hu, X., Shan, X.-q. & Zhang, S. (2013): Mechanistic studies of perfluorooctane sulfonate, perfluorooctanoic acid uptake by maize (*Zea mays* L. cv. TY2). Plant Soil, 370, 345-354.
- Wen, B., Li, L., Zhang, H., Ma, Y., Shan, X.-Q. & Zhang, S. (2014): Field study on the uptake and translocation of perfluoroalkyl acids (PFAAs) by wheat (*Triticum aestivum* L.) grown in biosolids-amended soils. Environmental Pollution, 184, 547-554.
- Wilhelm, M., Bergmann, S. & Dieter, H. H. (2010): Occurrence of perfluorinated compounds (PFCs) in drinking water of North Rhine-Westphalia, Germany and new approach to assess drinking water contamination by shorter-chained C4-C7 PFCs. International Journal of Hygiene and Environmental Health, 213, 224-232.
- Wójcik, A., Perczyk, P., Wydro, P. & Broniatowski, M. (2018): Effects of water soluble perfluorinated pollutants on phospholipids in model soil decomposer membranes. BBA - Biomembranes, 1860, 2576-2587.
- Wolf, C. J., Takacs, M. L., Schmid, J. E., Lau, C. & Abbott, B. D. (2008): Activation of Mouse and Human Peroxisome Proliferator-Activated Receptor Alpha by Perfluoroalkyl Acids of Different Functional Groups and Chain Lengths. Toxicological Sciences, 106, 162-171.
- Wong, F., Shoeib, M., Katsoyiannis, A., Eckhardt, S., Stohl, A., Bohlin-Nizzetto, P., Lid, H., Fellind, P., Su, Y. & Hung, H. (2018): Assessing temporal trends and source regions of per- and polyfluoroalkyl substances (PFASs) in air under the Arctic Monitoring and Assessment Programme (AMAP). Atmospheric Environment, 172, 65-73.

- Yang, X., Ye, C., Liu, Y. & Zhao, F.-J. (2015): Accumulation and phytotoxicity of perfluorooctanoic acid in the model plant species *Arabidopsis thaliana*. Environmental Pollution, 206, 560-566.
- Yeung, L. W. Y., Dassuncao, C., Mabury, S., Sunderland, E. M., Zhang, X. & Lohmann, R. (2017): Vertical Profiles, Sources, and Transport of PFASs in the Arctic Ocean. Environmental Science & Technology, 51, 6735–6744.
- Yeung, L. W.-Y. & Mabury, S. A. (2016): Are humans exposed to increasing amounts of unidentified organofluorine? Environmental Chemistry, 13, 102-110.
- Yi, L. B., Chai, L. Y., Xie, Y., Peng, Q. J. & Peng, Q. Z. (2016): Isolation, identification, and degradation performance of a PFOA-degrading strain. Genetics and Molecular Research, 15, 1-12.
- Yin, T., Chen, H., Reinhard, M., Yi, X., He, Y. & Gin, K. Y.-H. (2017): Perfluoroalkyl and polyfluoroalkyl substances removal in a full-scale tropical constructed wetland system treating landfill leachate. Water Research, 125, 418-426.
- Yu, W.-G., Liu, W., Jin, Y.-H., Liu, X.-H., Wang, F.-Q., Liu, L. & Nakayama, S. F. (2009): Prenatal and Postnatal Impact of Perfluorooctane Sulfonate (PFOS) on Rat Development: A Cross-Foster Study on Chemical Burden and Thyroid Hormone System. Environmental Science & Technology, 43, 8416–8422.
- Zareitalabad, P., Siemens, J., Hamer, M. & Amelung, W. (2013): Perfluorooctanoic acid (PFOA) and perfluorooctanesulfonic acid (PFOS) in surface waters, sediments, soils and wastewater – A review on concentrations and distribution coefficients. Chemosphere, 91, 725–732.
- Zeilmaker, M. J., Fragki, S., Bokkers, B., Verbruggen, E. & Lijzen, J., National Institute for Public Health and the Environment (RIVM) (2018). Mixture exposure to PFAS: A Relative Potency Factor approach. <https://www.rivm.nl/bibliotheek/rapporten/2018-0070.pdf>
- Zeng, X.-W., Qian, Z., Emo, B., Vaughn, M., Bao, J., Qin, X.-D., Zhu, Y., Li, J., Lee, Y. L. & Dong, G.-H. (2015): Association of polyfluoroalkyl chemical exposure with serum lipids in children. Science of the Total Environment, 512-513.
- Zhang, L., Ren, X.-M. & Guo, L.-H. (2013a): Structure-Based Investigation on the Interaction of Perfluorinated Compounds with Human Liver Fatty Acid Binding Protein. Environmental Science & Technology, 47, 11293–11301.
- Zhang, Y., Beesoon, S., Zhu, L. & Martin, J. W. (2013b): Biomonitoring of Perfluoroalkyl Acids in Human Urine and Estimates of Biological Half-Life. Environmental Science & Technology, 47, 10619–10627.
- Zhang, L., Ren, X.-M., Wan, B. & Guo, L.-H. (2014): Structure-dependent binding and activation of perfluorinated compounds on human peroxisome proliferator-activated receptor γ . Toxicology and Applied Pharmacology, 279, 275-283.
- Zhang, H., Liu, W., He, X., Wang, Y. & Zhang, Q. (2015): Uptake of perfluoroalkyl acids in the leaves of coniferous and deciduous broad-leaved trees. Environmental Toxicology and Chemistry, 34, 1499–1504.
- Zhang, Q., Liu, W., Niu, Q., Wang, Y., Zhao, H., Zhang, H., Song, J., Tsuda, S. & Saito, N. (2016): Effects of perfluorooctane sulfonate and its alternatives on long-term

- potentiation in the hippocampus CA1 region of adult rats in vivo. Toxicological Research, 5, 539-546.
- Zhao, L., Bian, J., Zhang, Y., Zhu, L. & Liu, Z. (2014): Comparison of the sorption behaviors and mechanisms of perfluorosulfonates and perfluorocarboxylic acids on three kinds of clay minerals. Chemosphere, 114, 51-58.
- Zhao, S., Zhu, L., Liu, L., Liu, Z. & Zhang, Y. (2013): Bioaccumulation of perfluoroalkyl carboxylates (PFCAs) and perfluoroalkane sulfonates (PFSA) by earthworms (*Eisenia fetida*) in soil. Environmental Pollution, 179, 45-52.
- Zhao, W., Zitzow, J. D., Weaver, Y., Ehresman, D. J., Chang, S.-C., Butenhoff, J. L. & Hagenbuch, B. (2017): Organic Anion Transporting Polypeptides Contribute to the Disposition of Perfluoroalkyl Acids in Humans and Rats. Toxicological Sciences, 156, 84-95.
- Zhao, Z., Xie, Z., Möller, A., Sturm, R., Tang, J., Zhang, G. & Ebinghaus, R. (2012): Distribution and long-range transport of polyfluoroalkyl substances in the Arctic, Atlantic Ocean and Antarctic coast. Environmental Pollution, 170, 71-77.
- Zheng, X.-M., Liu, H.-L., Shi, W., Wei, S., Giesy, J. P. & Yu, H.-X. (2012): Effects of perfluorinated compounds on development of zebrafish embryos. Environmental Science and Pollution Research, 19, 2498–2505.
- Zhou, Q., Deng, S., Yu, Q., Zhang, Q., Yu, G., Huang, J. & He, H. (2010): Sorption of perfluorooctane sulfonate on organo-montmorillonites. Chemosphere, 78, 688-694.
- Zhou, Y., Bao, W.-W., Qian, Z. M., Geiger, S. D., Parrish, K. L., Yang, B.-Y., Lee, Y. L. & Dong, G.-H. (2017): Perfluoroalkyl substance exposure and urine CC16 levels among asthmatics: A case-control study of children. Environmental Research, 159, 158–163.
- Zhou, Y., Hu, L.-W., Qian, Z. M., Chang, J.-J., King, C., Paul, G., Lin, S., Chen, P.-C., Lee, Y. L. & Dong, G.-H. (2016): Association of perfluoroalkyl substances exposure with reproductive hormone levels in adolescents: By sex status. Environment International, 94, 189-195.
- Zhou, Z., Liang, Y., Shi, Y., Xu, L. & Cai, Y. (2013): Occurrence and Transport of Perfluoroalkyl Acids (PFAAs), Including Short-Chain PFAAs in Tangxun Lake, China. Environmental Science & Technology, 47, 9249-9257.
- Zhou, Z., Shi, Y., Vestergren, R., Wang, T., Liang, Y. & Cai, Y. (2014): Highly Elevated Serum Concentrations of Perfluoroalkyl Substances in Fishery Employees from Tangxun Lake, China. Environmental Science & Technology, 48, 3864–3874.
- Zhu, Y., Qin, X.-D., Zeng, X.-W., Paul, G., Morawska, L., Su, M.-W., Tsai, C.-H., Wang, S.-Q., Lee, Y. L. & Dong, G.-H. (2016): Associations of serum perfluoroalkyl acid levels with T-helper cell-specific cytokines in children: By gender and asthma status. Science of the Total Environment, 559, 166–173.
- Zoeller, R. T., Tan, S. W. & Tyl, R. W. (2007): General Background on the Hypothalamic-Pituitary-Thyroid (HPT) Axis. Critical Reviews in Toxicology, 37, 11-53.





Part II

7. Registration and C&L notification status

7.1. Registration status





Per 20 August 2019 PFBS and seven different salt forms of PFBS had been registered in the ECHA database. PFBS is registered by one registrant, whereas KPFBs, the most used salt form, is registered by four registrants. The registration status for PFBS and its salts is provided in Table 41.

Table 41: Registration status for PFBS and some salt forms¹¹

	ID No.	Chemical structure	Volume
Perfluorobutane sulfonic acid, PFBS	CAS No. 375-73-5 EC No. 206-793-1		Full registration < 10 tons
Potassium perfluorobutane sulfonate, KPFBs	CAS No. 29420-49-3 EC No. 249-616-3		Full registration 100-1000 tons
Tetraethylammonium perfluorobutane sulfonate	CAS No. 25628-08-4 EC No. NA		Full registration 0-10 tons
Tetrabutylphosphonium perfluorobutane sulfonate (also registered as NONS)	CAS No. 220689-12-3 EC No.		Full registration 1+ tons

¹¹ Information collected from ECHA's dissemination page: <https://echa.europa.eu/information-on-chemicals/registered-substances> (accessed 20 August 2019)

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC

	444-440-5		
Triphenylsulfanium perfluorobutane sulfonate (also registered as triphenylsulfonium nonafluorobutanesulfonate, EC 477-300-7)	CAS No. 144317-44-2 EC No. 478-340-8		Registered as NONS ¹² Conf.
Dimethyl(phenyl)sulfanium perfluorobutane sulfonate	CAS No. 220133-51-7 EC No. 452-310-4		Registered as NONS Conf.
Bis(4-t-butylphenyl) iodonium perfluorobutane sulfonate	CAS No. 194999-85-4 EC No. 432-660-4		Registered as NONS Conf.
1-(4-Butoxy-1-naphthalenyl)tetrahydrothiophenium 1,1,2,2,3,3,4,4,4-nonafluoro-1-butanesulfonate	CAS No. NA EC No. 468-770-4		Registered as NONS Conf.

¹² NONS = Notification of new substances

7.2. CLP notification status

In addition to the seven salt forms of PFBS registered under REACH, four salt forms have been notified under CLP with self-classifications. Notified self-classifications for PFBS and the most widely used salt, KPFBS, are presented in Table 42 and Table 43, respectively. Information on the classification of the remaining registered salt forms are listed in Table 44, while the classification status for the notified PFBS salts may be found in Table 45.

Table 42: Notified self-classifications for perfluorobutane sulfonic acid, PFBS, according to CLP criteria¹³

Hazard Class and Category Code(s)	Hazard Statement Code(s)	Hazard Statement Code(s)	Number of notifiers
Acute Tox. 4	H302	H302	23
Skin Corr. 1B	H314	H314	
Acute Tox. 4	H302	H302	2
Skin Corr. 1B	H314	H314	
Met. Corr. 1	H290	H290	1
Acute Tox. 4	H302	H302	
Skin Corr. 1C	H314	H314	
Eye Dam. 1	H318	H318	





¹³ Information collected from the C&L Inventory database: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database> (accessed 20 August 2019)

Table 43: Notified self-classifications for potassium perfluorobutane sulfonate, KPFBS, according to CLP criteria¹⁴

Hazard Class and Category Code(s)	Hazard Statement Code(s)	Hazard Statement Code(s)	Number of notifiers
Eye Dam. 1	H318	H318	35
Skin Irrit. 2	H315	H315	23
Eye Irrit. 2	H319	H319	
STOT SE 3	H335	H335	
Skin Irrit. 2	H315	H315	18
Eye Irrit. 2	H319	H319	
STOT SE 3	H335 (Lung)	H335	
Not Classified			3
Skin Irrit. 2	H315	H315	3
Eye Irrit. 2	H319	H319	
STOT SE 3	H335 (respiratory sys...)	H335	
Skin Irrit. 2	H315	H315	2
Eye Irrit. 2	H319	H319	
Skin Irrit. 2	H315	H315	1
Eye Dam. 1	H318	H318	
STOT SE 3	H335 (respiratory tra...)		

¹⁴ Information collected from the C&L Inventory database: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database> (accessed 20 August 2019)

Table 44: Notified self-classifications and harmonised classifications of additional salts of PFBS that are registered under REACH¹⁵

	Notifi- cations	Self classifi cation	Harmo nised classifi cation
Tetraethylammonium perfluorobutane sulfonate, CAS No. 25628-08-4 	32	Eye Irrit.2; H319	-
Tetrabutylphosphonium perfluorobutane sulfonate, CAS no. 220689-12-3 	42	Acute Tox.4; H302 Aquatic chronic 3; H412	Acute Tox. 4; H302 Aquatic chronic 3; H412 ATP01
Triphenylsulfanium perfluorobutane sulfonate, CAS No. 144317-44-2 	71	Skin Irrit. 2; H315 Eye Irrit. 2; H319 STOT SE 3 H335	-
Dimethyl(phenyl)sulfanium perfluorobutane sulfonate, CAS No. 220133-51-7 	No results found	No results found	-

¹⁵ Information collected from the C&L Inventory database: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database> (accessed 20 August 2019)

ANNEX XV – IDENTIFICATION OF PFBS AND ITS SALTS AS SVHC







<p>Bis(4-t-butylphenyl) iodonium perfluorobutane sulfonate, EC No. 432-660-4</p> 	<p>1</p>	<p>Eye Dam.1; H318</p> <p>Aquatic chronic. 2; H411</p>	
<p>1-(4-Butoxy-1-naphthalenyl)tetrahydrothiophenium 1,1,2,2,3,3,4,4,4-nonfluoro-1-butanesulfonate, EC No. 468-770-4</p> 	<p>1</p>	<p>Eye Irrit. 2; H319</p>	

Table 45: Notified self-classifications according to CLP criteria of salts of PFBS that are not registered under REACH¹⁶

	ID No.	Notifi- cations	Self classification
Ammonium perfluorobutane sulfonate 	CAS No. 68259-10-9 EC No. 269-513-7	17	Not classified
Lithium perfluorobutane sulfonate 	CAS No. 131651-65-5 EC No. NA	1	Skin Irrit. 2; H315 Eye Irrit. 2; H319 STOT SE 3; H335 (not specified)
Magnesium perfluorobutane sulfonate 	CAS No. 507453-86-3 EC No. NA	1	Skin Irrit. 2; H315 Eye Irrit. 2; H319 STOT SE 3; H335 (not specified)
Morpholinium perfluorobutane sulfonate 	CAS No. 503155-89-3 EC No. NA	1	Skin Irrit. 2; H315 Eye Irrit. 2; H319 STOT SE 3; H335 (not specified)

The worldwide manufacture and use of PFBS and related substances were estimated and discussed by Lassen et al. (2017). In summary, the report concluded that the total global production of PFBS increased from 23 to 27 tonnes/year in the period 2011 – 2015. Of the 27 tonnes produced in 2015, 17 tonnes were consumed outside of China. The production of PBSF, the starting material for the manufacture of PFBS-related substances, increased from 287 to 317 tonnes/year in the same period. Hence, the global manufacture of PBSF is about ten times higher compared to PFBS. Of the 317 tonnes of PBSF manufactured globally in 2015, 299 tonnes (94%) were manufactured in China and 216 tonnes consumed

¹⁶ Information collected from the C&L Inventory database: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database> (accessed 20 August 2019)

in China (Lassen et al., 2017). The 83 tonnes of PBSF exported from China may be used as intermediates for further manufacture of fluorinated chemicals in the EEA or in other countries (Lassen et al., 2017).

Total tonnage of the substance

As of 20 August 2019, based on the information presented in Table 41, PFBS is manufactured and imported to the EU in a volume of <10 tonnes/year. The added manufactured and imported volume of the registered PFBS salt forms is 101-1011 tonnes/year in the EU, of which KPFBS contributes with 100-1000 tonnes/year. In addition to this, the volumes of the four salt forms with confidential tonnages contribute an unknown amount.

8. Information on uses of the substances

Specific applications for PFBS, as well as its potassium, tetraethylammonium and terabutylphosphonium salts, are discussed in more detail below. The applications of the other salts, are not reported or known. Information about the main applications of the PFBS-related substances is also provided.

PFBS

PFBS is registered in the tonnage band <10 tonnes/year in the EU, with uses registered as catalyst in polymer manufacture and as catalyst in chemical synthesis (one registrant per 2019-08-20)¹⁷.

KPFBS

KPFBS is the salt form of PFBS that is manufactured and used in the largest volumes. It is registered at 100-1000 tonnes/year manufacture and import in the EU. The uses indicated are as an additive/reactant in polymerisation processes in e.g. polycarbonate production, use as an intermediate in the synthesis of chemicals, use in articles and use in laboratories (three registrants per 20 August 2019)¹⁸. From the general literature it is known that the main application of KPFBS is as a flame retardant in polycarbonate, used mainly in electrical and electronic equipment. One European manufacturer states that their production capacity is several tonnes per year. The worldwide manufacture of KPFBS in 2015 has been estimated at 46 tonnes (Lassen et al., 2017). As the majority of electrical and electronic equipment used in Europe is imported from Asia, a considerable amount of KPFBS enters the EU in imported products (Lassen et al., 2017).

Tetraethylammonium perfluorobutane sulfonate (CAS No. 25628-08-4)

The substance is registered in the 1-10 tonnes/year tonnage band in the EU for use in metal (chromium) plating¹⁹. It is not indicated whether the substance is used for hard or

¹⁷ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/25546> (accessed 20 August 2019).

¹⁸ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/22432> (accessed 20 August 2019).

¹⁹ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/6361> (accessed 20 August 2019).

decorative plating. The product Bayowet FT 248 liquid (50% solution) is used as a spray mist inhibitor for the chromium galvanic industry and contains this substance (Lassen et al., 2017).

Tetrabutylphosphonium perfluorobutane sulfonate (CAS No. 220689-12-3)

The substance is registered at 1+ tonnes in the EU. Two registrations exist: one full registration by Miteni SpA (inactive) and SABIC Innovative Plastics BV (active)²⁰, and one NONS (Notification of New Substances) registration, containing only confidential information regarding registrants, uses and volumes²¹. In the full registration, the use indicated is as an additive, while its use is reported to lead to the inclusion into/onto articles, like plastic articles. From the literature it is known that the substance is used as an anti-static additive for plastics. The substance has been found in the mixture Anti-Stat FC-1 Fluor surfactant from DuPont, in the literature also referred to as Zonyl FASP-1 (Lassen et al., 2017).

The three remaining substances among the registered salt forms, the triphenylsulfanium, the dimethyl(phenyl)sulfanium, and the bis(4-t-butylphenyl) iodonium salts, are registered as notified new substances (NONS) and the tonnage band is confidential.

PFBS-related substances

The main application of PFBS-related substances in the EU is as water and stain repellent protection for leather, textiles and carpets, and in many cases the specific substances used are side-chain fluorinated polymers. According to Lassen et al. (2017), 3M is the only supplier of polymers based on PFBS-chemistry. These side-chain fluorinated polymer products were originally based on PFOS-related substances, but in 2003 PFOS-related substances were replaced by PFBS-related substances (Chu and Letcher, 2014). PFBS-related substances may be used in the manufacture of articles and are also available in consumer products such as aerosols for protective treatment of textiles, shoes and carpets. The total tonnage of PFBS-moieties in PFBS-related substances in articles and in mixtures for consumer applications on the EU market is estimated at 20-60 tonnes/year (Lassen et al., 2017).

A second major application of PFBS-related substances, including polymers, is as surface treatment for porous hard surfaces, like concrete, grout, unglazed tile, granite, clay, slate, limestone, marble and terracotta. A volume of 5-10 tonnes/year of PFBS-moieties for this application on the EU market was estimated by Lassen et al. (2017).

PFBS-related substances are registered under REACH with volumes adding up to 130-1300 tonnes/year:

- N-Methyl perfluorobutane sulfonamidoethanol (MeFBSE, EC 252-043-1, CAS no. 34454-97-2) is registered at 10-100 tonnes/year for industrial use as an intermediate for manufacture of other substances²²

²⁰ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/5653> (accessed 20 August 2019)

²¹ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/5004> (accessed 20 August 2019)

²² ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/5299> (accessed 20 August 2019)

- N-Methyl perfluorobutane sulfonamidoethyl acrylate (EC 266-733-5, CAS No. 67584-55-8) is registered at 100-1000 tonnes/year for use as a laboratory agent and for industrial polymerization²³
- N-Methyl perfluorobutane sulfonamidoethyl methacrylate (EC 266-737-7, CAS No. 67584-59-2) is registered at 10-100 tonnes/year for use in industrial polymerisation²⁴
- Perfluorobutane sulfonyl fluoride (PBSF, EC 206-792-6, CAS No. 375-72-4) is registered without tonnage information as an intermediate for the manufacture of MeFBSE, which is further used as an intermediate for manufacture of other substances²⁵
- N,N-Bis(2-hydroxyethyl) perfluorobutane-1-sulfonamide (EC 252-044-7, CAS No. 34455-00-0) is registered at 10-100 tonnes/year for use in polymerisation processes at industrial sites and for use by professional workers for coating²⁶

9. Additional information

9.1. Substances with similar hazard and use profiles on the Candidate List

PFASs that are listed on the Candidate List or that are subject to restriction or substance evaluation are listed in Table 46.

²³ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/11481> (accessed 20 August 2019)

²⁴ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/25633> (accessed 20 August 2019)

²⁵ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/10698> (accessed 20 August 2019)

²⁶ ECHA dissemination page: <https://echa.europa.eu/registration-dossier/-/registered-dossier/27023> (accessed 20 August 2019)

Table 46: PFASs subjected to regulatory activities under REACH

Substance	Regulatory activity under REACH	Status of regulatory activity
-		
C11-C14 PFCAs	SVHC-identification	Included in the Candidate List, reason for inclusion vPvB
C8 PFCA (PFOA) and its NH ₄ ⁺ salt		Included in the Candidate List, reason for inclusion vPvB and CMR
C9 PFCA (PFNA) and its Na/NH ₄ ⁺ salts		Included in the Candidate List, reason for inclusion vPvB and CMR
C10 PFCA (PFDA) and its Na/NH ₄ ⁺ salts		Included in the Candidate List, reason for inclusion vPvB and CMR
C6 PFSA (PFHxS)		Included in the Candidate List, reason for inclusion vPvB
HFPO-DA (GenX)		Included in the Candidate List, reason for inclusion ELoC
C8 PFCA and related substances		Restriction
C6 PFSA (PFHxS and related substances)	Ongoing	
C9-C14 PFCA and related substances	Ongoing	

Alternatives to the use of PFBS salts as flame retardants in polycarbonate

Levchik and Weil (2005) reviewed the recent development in the flame retardancy of polycarbonate and concluded that polycarbonate can pass stringent flame-retardant tests with very low additions of e.g. potassium diphenylsulfone sulfonate or some silicone-based flame retardants.

Alternatives to the use of PFBS salts as anti-static additive for plastics

Markarian (2008) sums up the new developments in antistatic and conductive additives for plastics. She explains that migrating antistatics are used widely in packaging such as film, thermoformed containers, and PET-bottles. The group of migrating antistatics includes long-chain alkyl phenols, ethoxylated amines, and glycerol esters like glycerol monostearate. For more long-term antistatic protection, permanent or conductive antistatic additives are used. The inherently dissipative polymers form a conductive polymer matrix or interpenetrating network within the base polymer, offering non-leaching, permanent static dissipation at a faster static decay rate than migrating antistatics. There are different solutions at the molecular level within this category, but the exact chemical nature is often considered confidential business information. Carbon black and conductive fibres, like graphite and metals, have applications as antistatic additives. Carbon nanotubes may be used at very low loadings and do not interfere with polymer properties. Graphene sheets are one of the most conductive materials available and may find increased use as an antistatic additive for plastics.

Alternatives to the use of PFBS salts in metal (chromium) plating

The Swedish Chemicals Agency (2015) discussed alternatives to the use of PFASs in metal plating. They pointed out that PFOS was once used in decorative chrome plating, but a new technology that uses chromium(III) instead of chromium(VI) does not have the same need for highly fluorinated substances as mist suppressants.

For hard metal plating, the chromium(III) technology is not equally applicable. Aminoethanol (EC 205-483-3, CAS 141-43-5) and alkyl sulfonates may be used as mist suppressants in metal plating, but they appear to be less effective and may degrade in the chrome bath. Hence, fresh material will need to be added during the process. However, alternative technologies, such as foam blankets and other barriers that can be used to inhibit misting, are also available. In combination with increased ventilation, these technologies reduce mist formation so much that the need for mist suppressants is greatly reduced.

Alternatives to the use of PFBS-related substances as water and stain repellent protection for leather, textiles and carpets and porous hard surfaces

There are a number of alternative chemical agents that may be used to induce water repellency to a product, while stain or oil repellency is more difficult to achieve. However, in many cases the stain/oil repellency is less important. The Swedish Chemicals Agency (2015) discussed the alternatives to PFASs in detail and listed the following alternatives as water repellent agents:

- Silicone-based agents, for example, high molecular weight polydimethylsiloxanes (PDMS).

- Mixtures of silicones and stearamidomethyl pyridine chloride, sometimes together with carbamide (urea) and melamine resins.
- Waxes and paraffins, which usually consist of modified melamine-based resins.
- Dendrimers, that were developed to imitate the ability of the lotus blossom to repel water. These structures often occur in nanosize (1 to 100 nm) and generally comprise hydrophobic, modified, highly branched polymers, including polyurethane.

In addition, non-chemical alternatives are available for some applications, like the use of tightly woven fabrics for water repellent textiles, and the use of high-density paper which prevents the passage of grease through it.

PFASs and non-fluorinated finishes for outdoor textiles were compared in a study by Hill et al. (2017). Among the tested finishes were C8- and C6- PFAS technology, as well as fat-modified resin, dendrimers and dendrimers with polyurethane backbone. It was found that water repellency ratings were similar across the range of all finished fabrics tested, all demonstrating a high level of resistance to wetting, and several non-fluorinated repellent fabrics provided similar water repellency as PFAS-based textile finishes. Non-fluorinated finishes provided no oil repellency and were clearly inferior in this property to PFAS-finished fabrics that demonstrated good oil-resistance. However, the primary repellency function required in outdoor apparel is water repellency, and it was suggested that the use of PFAS chemistry for such garments is over-engineering, providing oil repellency that exceeds user requirements.

9.2. Existing EU legislation

In the Proposal for a Directive of the European Parliament and of the Council on the quality of water intended for human consumption (recast)²⁷, the European Commission proposes, among a number of changes, the addition of limit values for certain chemicals. New limit values for PFAS are suggested at 0.1 µg/l for an individual PFAS ($C_nF_{2n+1}-R$, which includes PFBS) and 0.5 µg/l for PFASs in total (sum of individual PFASs, $C_nF_{2n+1}-R$). In the proposal it is argued that as PFOS and PFOA have been substituted by similar PFASs, often with shorter chains, PFOS and PFOA are likely no longer representative of this group of anthropogenic persistent chemicals. This indicates that the concern for PFAS has shifted from PFOS and PFOA as individual representatives of the group, to the group as a whole, with special mention of the short-chain PFASs.

9.3. Previous assessments by other authorities

The dossier submitter has previously performed an RMO analysis for PFBS, its salts and related substances (Norwegian Environment Agency, 2018).

The Swedish National Food Agency has recommended limits for drinking water based on the presence of 11 PFASs (PFBS, PFHxS, PFOS, 6:2 FTSA, PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFNA and PFDA)²⁸. If the sum of these 11 PFASs occurs at concentrations greater

²⁷ <https://eurlex.europa.eu/legalcontent/EN/TXT/?qid=1519210589057&uri=CELEX:52017PC0753>

²⁸ <https://www.livsmedelsverket.se/en/food-and-content/oonskade-amnen/miljogifter/pfas-in-drinking-water-fish-risk-management?AspxAutoDetectCookieSupport=1> (accessed 20 August 2019)

than 0.09 µg/l, the Agency recommends that measures are taken as soon as possible to reduce the pollution. That PFBS is specified as one of the 11 PFASs covered by the limit value, demonstrates that there is specific concern associated with this particular compound.

In 2018 the US EPA released a draft toxicity assessment for PFBS and its potassium salt KPFBS²⁹.

²⁹ <https://www.epa.gov/pfas/genx-and-pfbs-draft-toxicity-assessments> (accessed 20 August 2019)

References for Part II

- Chu, S. & Letcher, R. J. (2014): In vitro metabolic formation of perfluoroalkyl sulfonamides from copolymer surfactants of pre- and post-2002 Scotchgard fabric protector products. *Environmental Science & Technology*, 48, 6184-6191.
- Hill, P. J., Taylor, M., Goswami, P. & Blackburn, R. S. (2017): Substitution of PFAS chemistry in outdoor apparel and the impact on repellency performance. *Chemosphere*, 181, 500-507.
- Lassen, C., Brinch, A. & Jensen, A. A., COWI A/S (2017). Investigation of sources to PFBS in the environment (M-759). <http://www.miljodirektoratet.no/Documents/publikasjoner/M759/M759.pdf>
- Levchik, S. V. & Weil, E. D. (2005): Overview of recent developments in the flame retardancy of polycarbonates. *Polymer International*, 54, 981-998.
- Markarian, J. (2008): New developments in antistatic and conductive additives. *Plastics Additives & Compounding*, 10, 22-25.
- Norwegian Environment Agency (2018). Risk Management Option Analysis Conclusion Document. Substance Name: Perfluorobutane sulfonic acid (PFBS), its salts and related substances. <https://echa.europa.eu/documents/10162/1cb252bc-3952-3d5e-48b3-e78d5f5f0255>
- Swedish Chemicals Agency (2015). Occurrence and use of highly fluorinated substances and alternatives (7/15). <http://www.kemi.se/global/rapporter/2015/rapport-6-15-forekomst-och-anvandning-av-hogfluorerade-amnen-och-alternativ.pdf>

Annex I: Physical-chemical properties of substances used in read-across

Table 47 presents physical-chemical properties of the sulfonic acids trifluoromethane sulfonic acid, perfluorobutane sulfonic acid (PFBS) and perfluorooctane sulfonic acid (PFOS) collected from the OECD QSAR toolbox v4.2. These substances are compared in a read-across approach in the assessment of the persistence of PFBS.

Table 47: Physical-chemical properties of substances used in read-across (from OECD QSAR toolbox v4.2)

	Trifluoromethane sulfonic acid	PFBS	PFOS
C-chain length	C1	C4	C8
CAS No	1493-13-6	375-73-5	1763-23-1
SMILES	<chem>OS(=O)(=O)C(F)(F)F</chem>	<chem>OS(=O)(=O)C(F)(F)C(F)(F)C(F)(F)F</chem>	<chem>OS(=O)(=O)C(F)(F)C(F)(F)C(F)(F)C(F)(F)C(F)(F)C(F)(F)F</chem>
pK _a	-3.43	-3.31	-3.32
Log K _{ow}	-0.49	1.82	4.49
Log K _{oA}	4.90	5.05	4.84
Log K _{oc} (log K _{ow} method)	0.65	1.93	3.41
Water solubility (mg/L)	1000000	8860	0.050
Vapour pressure (mm Hg)	0.545	0.0518	0.0064
Boiling point (°C)	166	211	NA

Annex II: Modelling nominal biota concentration development of PFBS according to the Crookes and Fisk model

Crookes and Fisk (2018) investigated how the concentrations of mobile and persistent chemicals in the environment develop over time and compared their findings with the bioaccumulation criteria of the Stockholm Convention. They found that substances that are both persistent and mobile in the environment have the potential to be transported long distances from the point of emission. If such substances accumulate over time in remote regions, they can reach levels that may have effects on both ecosystems and human health.

Crookes and Fisk (2018) modelled the expected time-trend for concentrations in biota of substances with a certain combination of half-life in water and and bioconcentration factors, see Chapter 4 and Figure 3 in their report and reproduction in Figure 5 below. The authors found that for a given mass emission rate, as the degradation half-life increases the steady state concentration predicted in biota for substances with relatively low BCF values can, over extended periods of time, approach that of a substance that is considered to have a high bioaccumulation potential, and the steady state concentration ultimately reached is a function of both the persistence and BCF.

Figure 5 is a reproduction of Figure 3 i Crookes and Fisk (2018) showing time trends to accumulation in biota. The model shows that the nominal biota concentration for a substance with half-life 365 days and BCF 800 will approach the biota concentration for a substance with half-life 60 days and BCF 5000, after 2000 days.




Figure 5: Reproduction of model for biota concentration development from Crookes and Fisk (2018)

The nominal biota concentration calculations were repeated for PFBS and compared with some relevant model substances. A half-life in water of 10 years for PFBS was assumed, and the calculations performed for the bioaccumulation values reported in this dossier: BCF Fish: 23.5 (Chen et al., 2016); BAF crab 110 (Naile et al., 2013) and BAF fish 1736 (Campo et al., 2015). The outcome of the modelling of development of biota concentrations for PFBS over time is shown in Figure 6. The model substances (A, B, C and D) have combinations of half-life and BCF as shown in the figure.




Figure 6: Modelling of development of nominal biota concentrations for PFBS over time

It is evident from Figure 6 that when considering an appropriate time scale, a long half-life for a substance may lead to high nominal biota concentrations, even when the BCF is only moderate. The red line represents a BCF for PFBS of 1736 reported in fish (Campo et al., 2015) and demonstrates the effect of a long half-life and in combination with a relatively high BCF. However, as is outlined in Section 3.5.3 we consider this BCF to be an overestimate, and the red line is disregarded in this evaluation. The green line represents a BAF of 110 measured in crab (Naile et al., 2013). The graph in Figure 6 shows that this moderate BCF in combination with a half-life of 10 years, may lead to very high concentrations in biota over time. The green line even crosses the dark blue line, representing a substance with half-life in water of 41 days and a BCF of 2100, i.e. a substance defined as P and B according to the REACH Annex XIII criteria.

Hence, when the model from the Crookes and Fisk (2018) report is used for PFBS with an assumed half-life in water of 10 years and a reported BAF in crab of 110, the concentrations of PFBS in biota may be expected to exceed the biota concentrations for a persistent and bioaccumulative substance over time.