

Summary of product characteristics for a biocidal product

Product name: BIOPREN Ant Killer Gel

Product type(s): PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Authorisation number: (10-14 17.5 Mr)BPR-495(A-18VAP564428-23-260)

R4BP 3 asset reference number: LT-0031878-0000

Table Of Contents

| | |
|---|----|
| Administrative information | 1 |
| 1.1. Trade names of the product | 1 |
| 1.2. Authorisation holder | 1 |
| 1.3. Manufacturer(s) of the biocidal products | 1 |
| 1.4. Manufacturer(s) of the active substance(s) | 2 |
| 2. Product composition and formulation | 2 |
| 2.1. Qualitative and quantitative information on the composition of the biocidal product | 2 |
| 2.2. Type of formulation | 3 |
| 3. Hazard and precautionary statements | 3 |
| 4. Authorised use(s) | 3 |
| 5. General directions for use | 10 |
| 5.1. Instructions for use | 10 |
| 5.2. Risk mitigation measures | 11 |
| 5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment | 11 |
| 5.4. Instructions for safe disposal of the product and its packaging | 11 |
| 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage | 11 |
| 6. Other information | 11 |

Administrative information

1.1. Trade names of the product

| |
|-----------------------------|
| BIOPREN hangyairtó gél |
| PROTECT hangyairtó csalétek |
| Biopren ant gel |
| Biopren ant gel |
| Biopren ant gel |
| Biopren ant gel |
| Protect ant gel |
| Protect ant gel |
| Protect ant gel |
| Protect ant gel |
| Protect ant gel |
| Protect ant gel |
| Protect ant gel |
| Protect ant gel |

1.2. Authorisation holder

| | | |
|---|--|---|
| Name and address of the authorisation holder | Name | Bábolna Bioenvironmental Centre Private Limited Company |
| | Address | Szállás u. 6. H-1107 Budapest Hungary |
| Authorisation number | (10-14 17.5 Mr)BPR-495(A-18VAP564428-23-260) | |
| R4BP 3 asset reference number | LT-0031878-0000 | |
| Date of the authorisation | 20/11/2023 | |
| Expiry date of the authorisation | 05/08/2026 | |

1.3. Manufacturer(s) of the biocidal products

| | |
|--|--|
| Name of the manufacturer | Mylva S.A. |
| Address of the manufacturer | Via Augusta, 48 08006 Barcelona Spain |
| Location of manufacturing sites | Sant Galderic 23 08395 Sant Pol de Mar Spain |

1.4. Manufacturer(s) of the active substance(s)

| | |
|--|---|
| Active substance | 37 - imidacloprid |
| Name of the manufacturer | Bayer S.A.S. |
| Address of the manufacturer | 16 rue Jean-Marie Leclair 90106 Lyon Cedex France |
| Location of manufacturing sites | 16 rue Jean-Marie Leclair 90106 Lyon Cedex France |

| | |
|--|---|
| Active substance | 1386 - S-Methoprene |
| Name of the manufacturer | Bábolna Bioenvironmental Centre Ltd. |
| Address of the manufacturer | H-1107 Budapest, Szállás u. 6. 1107 Szállás, Budapest Hungary |
| Location of manufacturing sites | H-1107 Budapest, Szállás u. 6. 1107 Szállás, Budapest Hungary |

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

| Common name | IUPAC name | Function | CAS number | EC number | Content (%) |
|--------------|--|------------------|-------------|-----------|-------------|
| imidacloprid | (2E)-1-[(6-chloropyridin-3-yl) methyl]-N-nitroimidazolidin-2-imine | Active Substance | 138261-41-3 | 428-040-8 | 0,01 |
| S-Methoprene | | Active Substance | 65733-16-6 | | 0,08 |

2.2. Type of formulation

RB - Bait (ready for use)

3. Hazard and precautionary statements

Hazard statements

Harmful to aquatic life with long lasting effects.

Precautionary statements

Keep out of reach of children.

Avoid release to the environment.

Collect spillage.

Dispose of contents to according to hazard residues regulation.

4. Authorised use(s)

4.1 Use description

Use 1 - Trained Professional Users

Product type

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Where relevant, an exact description of the authorised use

Imidacloprid 0.01 % and 0.08% S-Methoprene Gel is an indoor an outdoor gel bait insecticide against Monoronomium Pharaonis, Linepithema Humile and Lasius Niger. Imidacloprid 0.01% and 0.08% S-methoprene Gel is designed for the controlled placement of bait in the following situations:
-User: Trained professional, Non-trained professional and general public (Non-professional).

Target organism(s) (including development stage)

Scientific name: Monomorium Pharaonis
Common name: Pharaoh Ant
Development stage: Adults, Nymphs

Scientific name: Linepithema Humile
Common name: Argentine Ant
Development stage: Adults, Nymphs

Scientific name: Lasius Niger
Common name: Black Ant
Development stage: Adults, Nymphs

Field(s) of use

Indoor

Outdoor

Other

-Outdoor use:
Apply MAGNUM GEL HORMIGAS IGR PLUS within the ant nests, next to ant trails and where their presence is detected by avoiding the sun which can dry the product. The places where it is more likely to find ants are: eaves, cornices, drain holes, where wires and pipes enter the structure of the building, trees, stumps, roots, wood decks, fences, planters, etc...Apply in the bottom flanges of the window and in the access doors to prevent entry of ants in houses or buildings.
After several days, inspect the points of application and if is necessary repeat treatments to get full control of the colony. Remove alternative food source that may be in competition from near areas.
Although it contains a bitter substance that makes it unappetizing to people or pets, should be applied in areas inaccessible to pets.

-Indoor use:
Apply the gel as drops or lines near ant trails or the bait station, in areas where ants are present such a within their nests, or in cracks and crevices of areas where their presence is detected. The places where ants are likely to be present and where the gel should be applied are: hollow walls, cracks and crevices in the kitchen, bathroom, under the sink, lower edge of windows, vents, garage, attics, etc...

-Outdoor use:
Apply MAGNUM GEL HORMIGAS IGR PLUS in form of drops or the bait station within the ant nests, next to ant trails and where their presence is detected by avoiding the sun which can dry the product. The places where it is more likely to find ants are: eaves, cornices, drain holes, where wires and pipes enter the structure of the building, trees, stumps, roots, wood decks, fences, planters, etc...Apply in the bottom flanges of the window and in the access doors to prevent entry of ants in houses or buildings.
After several days, inspect the points of application and if is necessary repeat treatments to get full control of the colony. Remove alternative food source that may be in competition from near areas.
Although it contains a bitter substance that makes it unappetizing to people or pets, should be applied in areas inaccessible to pets.

Application method(s)

Method: Bait application
Detailed description:
Indoor use: Place spots or lines of gel or baitstation near ant trails, or where ants have been seen, in their nests, or cracksand crevices in the areas where ants are detected. Places where it is more probable to find ants are: hollowwalls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc.
Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight sothat the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes,where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc.Place on windowsills and doorways to avoid ant entering houses or buildings.After some days, inspect application points and repeat treatment if necessary until total colony control isachieved. Get rid of other near sources of food that could compete with the bait.

Application rate(s) and frequencies

Application Rate: Cartridge and syringe : Apply spots or lines of Gel on infested areas. Use 1-3 lines per bait placement. / Bait station: 5g/25m2
Dilution (%): 0
Number and timing of application:
Check the baits and replace the consumed ones daily during treatment.

Category(ies) of users

Trained professional

Professional

General public (non-professional)

Pack sizes and packaging material

Cartridge - Plastic: LDPE - 1,2,3,4,5,8,10,15,20,25,30,35,40,50,60,75,80,100,125,150,175,200,250,300,350,400,450 and 500 g

Syringe - Plastic: LDPE - 1,2,3,4,5,8,10,15,20,25,30,35,40,50,g

Bait Station - Plastic: PET - 1, 2,2.5, 3, 4, 5, 8, 10, 15, 20, 25, 30 g

4.1.1 Use-specific instructions for use

Indoor use: Place spots or lines of gel or bait station near ant trails, or where ants have been seen, in their nests, or cracks and crevices in the areas where ants are detected. Places where it is more probable to find ants are: hollow walls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc.

Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight so that the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes, where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc. Place on windowsills and doorways to avoid ant entering houses or buildings. After some days, inspect application points and repeat treatment if necessary until total colony control is achieved. Get rid of other near sources of food that could compete with the bait.

4.1.2 Use-specific risk mitigation measures

Avoid contact with eyes and skin. If direct manipulation and possible contact with the product exist, wear protective gloves when handling chemicals. Use appropriate protective equipment.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures: Remove the person from the contaminated area. Remove stained or splashed clothing. Wash eyes with plenty of water for at least 15 minutes. Remove contact lenses. Wash skin with plenty of water and soap without rubbing. Check breathing. If necessary administer artificial respiration. Transfer the person to a medical center and if possible take the label or container with you. NEVER LEAVE THE PATIENT UNATTENDED.

Recommendations for the physician: As a general rule, symptomatic treatment for compensating the observed effects.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Product disposal: - Respect the rules on clothing and protective measures. - To clean the floor and contaminated objects by this material, use carbonated or soapy water. Do not discharge into drains or the environment. Cover the product with absorbent material such as sand, ground, diatomaceous ground to prevent spreading. - The product should NOT be eliminated in municipal sewers, drains or rivers. - The treatment and/or disposal shall be made in accordance with local legislation. Spilled product cannot be reused and must be discarded. Dispose the product residue on an authorized point waste collection. Container disposal: Empty containers should be removed by an authorized company according to the national rules for residues management.

Product waste: Wear appropriate clothes and adopt relevant safety measures. Wash floor and contaminated objects with soap water. Do not empty into drains or the environment. Cover the product with absorbent material such as sand or any other absorbent material. Dispose in an authorized waste management plants.

Container disposal: Empty packages must be treated by an authorized agent.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Store in the original container tightly closed. Store in a dry, cool and well-ventilated place. It is recommended to store the product at a temperature preferably between 5°C and 45°C.

4.2 Use description

Use 2 - Non-Trained Professional User

| | |
|---|--|
| Product type | PT18 - Insecticides, acaricides and products to control other arthropods (Pest control) |
| Where relevant, an exact description of the authorised use | Imidacloprid 0.01 % and 0.08% S-Methoprene Gel is an indoor an outdoor gel bait insecticide against Monomorium Pharaonis, Linepithema Humile and Lasius Niger. Imidacloprid 0.01% and 0.08% S-methoprene Gel is designed for the controlled placement of bait in the following situations: -User: Trained professional, Non-trained professional and general public (Non-professional). |
| Target organism(s) (including development stage) | <p>Scientific name: Monomorium Pharaonis Common name: Pharaoh ant Development stage: Adults, Nymphs</p> <p>Scientific name: Linepithema Humile Common name: Argentine Ant Development stage: Adults, Nymphs</p> <p>Scientific name: Lasius Niger Common name: Black Ant Development stage: Adults, Nymphs</p> |
| Field(s) of use | <p>Indoor</p> <p>Outdoor</p> <p>Other</p> <p>-Outdoor use: Apply MAGNUM GEL HORMIGAS IGR PLUS within the ant nests, next to ant trails and where their presence is detected by avoiding the sun which can dry the product. The places where it is more likely to find ants are: eaves, cornices, drain holes, where wires and pipes enter the structure of the building, trees, stumps, roots, wood decks, fences, planters, etc...Apply in the bottom flanges of the window and in the access doors to prevent entry of ants in houses or buildings. After several days, inspect the points of application and if is necessary repeat treatments to get full control of the colony. Remove alternative food source that may be in competition from near areas. Although it contains a bitter substance that makes it unappetizing to people or pets, should be applied in areas inaccessible to pets.</p> <p>-Indoor use: Apply the gel as drops or lines near ant trails or the bait station, in areas where ants are present such a within their nests, or in cracks and crevices of areas where their presence is detected. The places where ants are likely to be present and where the gel should be applied are: hollow walls, cracks and crevices in the kitchen, bathroom, under the sink, lower edge of windows, vents, garage, attics, etc...</p> <p>-Outdoor use: Apply MAGNUM GEL HORMIGAS IGR PLUS in form of drops or the bait station within the ant nests, next to ant trails and where their presence is detected by avoiding the sun which can dry the product. The places where it is more likely to find ants are: eaves, cornices, drain holes, where wires and pipes enter the structure of the building, trees, stumps, roots, wood decks, fences, planters, etc...Apply in the bottom flanges of the window and in the access doors to prevent entry of ants in houses or buildings. After several days, inspect the points of application and if is necessary repeat treatments to get full control of the colony. Remove alternative food source that may be in competition from near areas. Although it contains a bitter substance that makes it unappetizing to people or pets, should be applied in areas inaccessible to pets.</p> |
| Application method(s) | <p>Method: Bait application</p> <p>Detailed description: Indoor use: Place spots or lines of gel or baitstation near ant trails, or where ants have been seen, in their nests, or cracksand crevices in the areas where ants are detected.</p> |

| | |
|---|--|
| | <p>Places where it is more probable to find ants are: hollowwalls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc. Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight so that the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes, where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc. Place on windowsills and doorways to avoid ant entering houses or buildings. After some days, inspect application points and repeat treatment if necessary until total colony control is achieved. Get rid of other near sources of food that could compete with the bait.</p> |
| <p>Application rate(s) and frequencies</p> | <p>Application Rate: Cartridge and syringe : Apply spots or lines of Gel on infested areas. Use 1-3 lines per bait placement. / Bait station: 5g/25m² Dilution (%): 0 Number and timing of application: Check the baits and replace the consumed ones daily during treatment.</p> |
| <p>Category(ies) of users</p> | <p>Trained professional Professional General public (non-professional)</p> |
| <p>Pack sizes and packaging material</p> | <p>Cartridge - Plastic: LDPE - 1,2,3,4,5,8,10,15,20,25,30,35,40,50,60,75,80,100,125,150,175,200,250,300,350,400,450 and 500 g Syringe - Plastic: LDPE - 1,2,3,4,5,8,10,15,20,25,30,35,40,50,g Bait Station - Plastic: PET - 1, 2,2.5, 3, 4, 5, 8, 10, 15, 20, 25, 30 g</p> |

4.2.1 Use-specific instructions for use

Indoor use: Place spots or lines of gel or bait station near ant trails, or where ants have been seen, in their nests, or cracks and crevices in the areas where ants are detected. Places where it is more probable to find ants are: hollowwalls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc.
Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight so that the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes, where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc. Place on windowsills and doorways to avoid ant entering houses or buildings. After some days, inspect application points and repeat treatment if necessary until total colony control is achieved. Get rid of other near sources of food that could compete with the bait.

4.2.2 Use-specific risk mitigation measures

Avoid contact with eyes and skin. If direct manipulation and possible contact with the product exist, wear protective gloves when handling chemicals. Use appropriate protective equipment.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures: Remove the person from the contaminated area. Remove stained or splashed clothing. Wash eyes with plenty of water for at least 15 minutes. Remove contact lenses. Wash skin with plenty of water and soap without rubbing. Check breathing. If necessary administer artificial respiration. Transfer the person to a medical center and if possible take the label or container with you. NEVER LEAVE THE PATIENT UNATTENDED.

Recommendations for the physician:As a general rule, symptomatic treatment for compensating the observed effects.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Product disposal:- Respect the rules on clothing and protective measures. - To clean the floor and contaminated objects by this material, use carbonated or soapy water. Do not discharge into drains or the environment. Cover the product with absorbent material such as sand, ground, diatomaceous ground to prevent spreading. - The product should NOT be eliminated in municipal sewers, drains or rivers. - The treatment and/or disposal shall be made in accordance with local legislation. Spilled product cannot be reused and must be discarded. Dispose the product residue on an authorized point waste collection. Container disposal: Empty containers should be removed by an authorized company according to the national rules for residues management.
Product waste: Wear appropriate clothes and adopt relevant safety measures. Wash floor and contaminated objects with soap water. Do not empty into drains or the environment. Cover the product with absorbent material such as sand or any other absorbent material. Dispose in an authorized waste management plants.
Container disposal: Empty packages must be treated by an authorized agent.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Store in the original container tightly closed. Store in a dry, cool and well-ventilated place. It is recommended to store the product at a temperature preferably between 5°C and 45°C.

4.3 Use description

Use 3 - Non-Professional User (General Public)

Product type

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Where relevant, an exact description of the authorised use

Imidacloprid 0.01 % and 0.08% S-Methoprene Gel is an indoor and outdoor gel bait insecticide against *Monomorium Pharaonis*, *Linepithema Humile* and *Lasius Niger*. Imidacloprid 0.01% and 0.08% S-methoprene Gel is designed for the controlled placement of bait in the following situations:
-User: Trained professional, Non-trained professional and general public (Non-professional).

Target organism(s) (including development stage)

Scientific name: *Monomorium Pharaonis*
Common name: Pharaoh ant
Development stage: Adults, Nymphs

Scientific name: *Lasius Niger*
Common name: Black Ant
Development stage: Adults, Nymphs

Scientific name: *Linepithema Humile*
Common name: Argentine Ant
Development stage: Adults, Nymphs

Field(s) of use

Indoor
Outdoor
Other

-Outdoor use:
Apply MAGNUM GEL HORMIGAS IGR PLUS within the ant nests, next to ant trails and where their presence is detected by avoiding the sun which can dry the product. The places where it is more likely to find ants are: eaves, cornices, drain holes, where wires and pipes enter the structure of the building, trees, stumps, roots, wood decks, fences, planters, etc...Apply in the bottom flanges of the window and in the access doors to prevent entry of ants in houses or buildings.
After several days, inspect the points of application and if necessary repeat

treatments to get full control of the colony. Remove alternative food source that may be in competition from near areas.
 Although it contains a bitter substance that makes it unappetizing to people or pets, should be applied in areas inaccessible to pets.

-Indoor use:
 Apply the gel as drops or lines near ant trails or the bait station, in areas where ants are present such as within their nests, or in cracks and crevices of areas where their presence is detected. The places where ants are likely to be present and where the gel should be applied are: hollow walls, cracks and crevices in the kitchen, bathroom, under the sink, lower edge of windows, vents, garage, attics, etc...

-Outdoor use:
 Apply MAGNUM GEL HORMIGAS IGR PLUS in form of drops or the bait station within the ant nests, next to ant trails and where their presence is detected by avoiding the sun which can dry the product. The places where it is more likely to find ants are: eaves, cornices, drain holes, where wires and pipes enter the structure of the building, trees, stumps, roots, wood decks, fences, planters, etc...Apply in the bottom flanges of the window and in the access doors to prevent entry of ants in houses or buildings.
 After several days, inspect the points of application and if is necessary repeat treatments to get full control of the colony. Remove alternative food source that may be in competition from near areas.
 Although it contains a bitter substance that makes it unappetizing to people or pets, should be applied in areas inaccessible to pets.

Application method(s)

Method: Bait application
 Detailed description:
 Indoor use: Place spots or lines of gel or baitstation near ant trails, or where ants have been seen, in their nests, or cracksand crevices in the areas where ants are detected. Places where it is more probable to find ants are: hollowwalls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc.
 Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight sothat the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes,where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc.Place on windowsills and doorways to avoid ant entering houses or buildings.After some days, inspect application points and repeat treatment if necessary until total colony control isachieved. Get rid of other near sources of food that could compete with the bait.

Application rate(s) and frequencies

Application Rate: Cartridge and syringe : Apply spots or lines of Gel on infested areas. Use 1-3 lines per bait placement. / Bait station: 5g/25m2
 Dilution (%): 0
 Number and timing of application:
 Check the baits and replace the consumed ones daily during treatment.

Category(ies) of users

Trained professional
 Professional
 General public (non-professional)

Pack sizes and packaging material

Cartridge - Plastic: LDPE - 1,2,3,4,5,8,10,15,20,25,30,35,40,50,60,75,80,100,125,150,175,200,250,300,350,400,450 and 500 g
 Syringe - Plastic: LDPE - 1,2,3,4,5,8,10,15,20,25,30,35,40,50,g
 Bait Station - Plastic: PET - 1, 2,2.5, 3, 4, 5, 8, 10, 15, 20, 25, 30 g

4.3.1 Use-specific instructions for use

Indoor use: Place spots or lines of gel or bait station near ant trails, or where ants have been seen, in their nests, or cracks and crevices in the areas where ants are detected. Places where it is more probable to find ants are: hollow walls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc.

Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight so that the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes, where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc. Place on windowsills and doorways to avoid ant entering houses or buildings. After some days, inspect application points and repeat treatment if necessary until total colony control is achieved. Get rid of other near sources of food that could compete with the bait.

4.3.2 Use-specific risk mitigation measures

Avoid contact with eyes and skin. If direct manipulation and possible contact with the product exist, wear protective gloves when handling chemicals. Use appropriate protective equipment.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures: Remove the person from the contaminated area. Remove stained or splashed clothing. Wash eyes with plenty of water for at least 15 minutes. Remove contact lenses. Wash skin with plenty of water and soap without rubbing. Check breathing. If necessary administer artificial respiration. Transfer the person to a medical center and if possible take the label or container with you. NEVER LEAVE THE PATIENT UNATTENDED.

Recommendations for the physician: As a general rule, symptomatic treatment for compensating the observed effects.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Product disposal: - Respect the rules on clothing and protective measures. - To clean the floor and contaminated objects by this material, use carbonated or soapy water. Do not discharge into drains or the environment. Cover the product with absorbent material such as sand, ground, diatomaceous ground to prevent spreading. - The product should NOT be eliminated in municipal sewers, drains or rivers. - The treatment and/or disposal shall be made in accordance with local legislation. Spilled product cannot be reused and must be discarded. Dispose the product residue on an authorized point waste collection. Container disposal: Empty containers should be removed by an authorized company according to the national rules for residues management.

Product waste: Wear appropriate clothes and adopt relevant safety measures. Wash floor and contaminated objects with soap water. Do not empty into drains or the environment. Cover the product with absorbent material such as sand or any other absorbent material. Dispose in an authorized waste management plants.

Container disposal: Empty packages must be treated by an authorized agent.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Store in the original container tightly closed. Store in a dry, cool and well-ventilated place. It is recommended to store the product at a temperature preferably between 5°C and 45°C.

5. General directions for use

5.1. Instructions for use

Indoor use: Place spots or lines of gel or baitstation near ant trails, or where ants have been seen, in their nests, or cracks and crevices in the areas where ants are detected. Places where it is more probable to find ants are: hollowwalls, kitchen cracks and crevices, bathrooms, sinks, windowsills, ventilation grilles, garages, attics, etc.
Outdoor use: Place gel inside their nests, or place gel or bait station next to ant trails and where ants are detected, avoiding sunlight so that the gel does not dry. Places where it is more probable to find ants are: eaves, cornices, draining holes, where cables and pipes enter the building structure, trees, stumps, roots, wood roofs, fences, plant pots, etc. Place on windowsills and doorways to avoid ant entering houses or buildings. After some days, inspect application points and repeat treatment if necessary until total colony control is achieved. Get rid of other near sources of food that could compete with the bait.

5.2. Risk mitigation measures

Avoid contact with eyes and skin. If direct manipulation and possible contact with the product exist, wear protective gloves when handling chemicals. Use appropriate protective equipment.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures: Remove the person from the contaminated area. Remove stained or splashed clothing. Wash eyes with plenty of water for at least 15 minutes. Remove contact lenses. Wash skin with plenty of water and soap without rubbing. Check breathing. If necessary administer artificial respiration. Transfer the person to a medical center and if possible take the label or container with you. NEVER LEAVE THE PATIENT UNATTENDED.

Recommendations for the physician: As a general rule, symptomatic treatment for compensating the observed effects.

5.4. Instructions for safe disposal of the product and its packaging

Product elimination: Respect clothing norms and protection measures. To clean the floor and object contaminated by this product, use carbonated or soapy water. Do not dispose of the waste water in drains or the environment. Cover the product with an absorbent material such as sand, earth or diatomic earth to prevent spreading. The product MUST NOT be disposed of in municipal sewer system, drains or rivers. The treatment and/or elimination must be carried out according to the law. It must be eliminated at an authorized residual waste collecting point. The spilled product cannot be reused and must be disposed of.

Elimination of recipients: The empty recipients must be disposed of by an authorized body. "The end owner of residues from industrial and commercial containers must keep them in the conditions established in article 12 of the containers and Residues from Containers Law 11/1997".

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Store in the original container tightly closed. Store in a dry, cool and well-ventilated place. It is recommended to store the product at a temperature preferably between 5°C and 45°C.

6. Other information

x