

**Appendix: List of 'relevant data' as identified by the evaluating Competent Authority (eCA) in the Renewal Assessment Report of Flocoumafen PT14, see Article 95(7) of Regulation (EU) No 528/2012 (BPR)**

| Author(s) | Year | Section No / Reference No | Title.<br>Source (where different from company)<br>Company, Report No.<br>GLP (where relevant) / (Un)Published  | Data Protection Claimed (Yes/No) | Owner | Data Identified as 'relevant' by the eCA <sup>1</sup> (Yes/No) |
|-----------|------|---------------------------|---|----------------------------------|-------|--|
| ████████  | 2011 | II-9.1.4/01               | Fish, bioconcentration of Flocoumafen according to the OECD-Guideline 305 and EU method C.13<br>Fraunhofer Institute for Molecular Biology and Applied Ecology, Schmallingenberg, Germany,<br>Report no. EBR-003/4-10<br>BASF ID: 2010/1178749<br>GLP / unpublished | Yes                              | BASF  | Yes  |
| ████████  | 2007 | II-9.2.1/01               | Soil microorganisms: Effects of Flocoumafen on nitrogen and carbon transformation<br>Fraunhofer Institute for Molecular Biology and Applied Ecology, Schmallingenberg, Germany,<br>Report no. EBR-003/3-35<br>BASF ID: 2007/1022271<br>GLP / unpublished            | Yes                              | BASF  | Yes  |
| ████████  | 2007 | II-9.2.2/01               | Earthworm acute toxicity test: acute toxicity of Flocoumafen on <i>Eisenia fetida</i><br>Fraunhofer Institute for Molecular Biology and Applied Ecology, Schmallingenberg, Germany,<br>Report no. EBR-003/3-08<br>BASF ID: 2007/1028165<br>GLP / unpublished        | Yes                              | BASF  | Yes  |
| ████████  | 2007 | II-9.2.3/01               | Terrestrial plants, growth test: Effect of Flocoumafen on the seedling emergence and growth of <i>Avena sativa</i> , <i>Lactuca sativa</i> ,  | Yes                              | BASF  | Yes  |

<sup>1</sup> For further information, and criteria of "relevant", please see, please see [CA-Sept20-Doc.7.1.b - Relevant Renewal Data under Article 95\\_FINAL](#)

| Author(s)  | Year | Section No / Reference No | Title.<br>Source (where different from company)<br>Company, Report No.<br>GLP (where relevant) / (Un)Published  | Data Protection Claimed (Yes/No) | Owner | Data Identified as 'relevant' by the eCA <sup>1</sup> (Yes/No) |
|------------|------|---------------------------|---|----------------------------------|-------|--|
|            |      |                           | <p><i>Phaseolus aureus, Raphanus sativus, Sinapis alba, and Triticum aestivum</i><br/>Fraunhofer Institute for Molecular Biology and Applied Ecology, Schmallenberg, Germany,<br/>Report no. EBR-003/4-40<br/>BASF ID: 2007/1033910<br/>GLP / unpublished</p>       |                                  |       |  |
| ██████████ | 2012 | II-9.7/01                 | <p>Bioaccumulation in terrestrial oligochaetes - Uptake and elimination of flocoumafen (BAS 322 I) in <i>Eisenia fetida</i><br/>Fraunhofer Institute (IME), Schmallenberg, Germany<br/>Report no. EBR-003/3-25,<br/>BASF ID: 2011/1284061<br/>GLP / Unpublished</p> | Yes                              | BASF  | Yes  |