

Annex XV report

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57

Substance Name: Triphenyl phosphate (TPhP)

EC Number: 204-112-2

CAS Number: 115-86-6

Submitted by: France

Date: 23 Feb 2024

This document has been prepared according to template: TEM-0049.04

CONTENTS

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57	8
PART I	10
JUSTIFICATION	10
1. IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES	10
Name and other identifiers of the substance	10
Composition of the substance	10
Identity and composition of degradation products/metabolites relevant for the SVHC assessment	10
Identity and composition of structurally related substances (used in a grouping or read-across approach).....	10
Physicochemical properties	11
2. HARMONISED CLASSIFICATION AND LABELLING	12
3. ENVIRONMENTAL FATE PROPERTIES	12
Degradation.....	12
<i>Abiotic degradation</i>	12
<i>Biodegradation</i>	12
Environmental distribution	13
<i>Adsorption/desorption</i>	13
<i>Volatilisation</i>	13
<i>Distribution modelling</i>	13
<i>Environmental occurrence data</i>	13
<i>Human occurrence</i>	21
<i>Summary on occurrence and environmental distribution</i>	21
Bioaccumulation	21
<i>Bioaccumulation in aquatic organisms (pelagic & sediment organisms)</i>	21
<i>Bioaccumulation in terrestrial organisms (soil dwelling organisms, vertebrates)</i>	21
Summary.....	21
4. HUMAN HEALTH HAZARD ASSESSMENT	21
5. ENVIRONMENTAL HAZARD ASSESSMENT.....	22
Aquatic compartment (including sediment).....	22
<i>Fish</i>	22
<i>Aquatic invertebrates</i>	22
<i>Algae and aquatic plants</i>	23
<i>Sediment organisms</i>	23
<i>Other aquatic organisms</i>	23
Terrestrial compartment.....	23
<i>Toxicity to soil macro-organisms</i>	23
<i>Toxicity to terrestrial plants</i>	23
<i>Toxicity to soil micro-organisms</i>	23
<i>Toxicity to other terrestrial organisms</i>	24

Atmospheric compartment.....	24
Microbiological activity in sewage treatment systems.....	24
Toxicity to birds.....	24
Mammalian wildlife.....	24
6. ENDOCRINE DISRUPTION (ENVIRONMENT)	25
General approach for the assessment of endocrine properties.....	25
<i>Framework of the evaluation</i>	25
<i>Information sources and strategy for endocrine disruptor identification</i>	25
<i>In vitro</i> information indicative of endocrine activity.....	26
<i>In vivo</i> mechanistic data with regard to an endocrine activity.....	34
<i>In vivo</i> adverse effect data with regard to an endocrine mode of action.....	46
Conclusion regarding ED properties relevant for environment.....	62
<i>Adverse effects relevant for ED identification</i>	62
<i>Endocrine activity</i>	65
<i>Plausible link between adverse effects and endocrine activity</i>	73
6. CONCLUSIONS ON THE SVHC PROPERTIES.....	77
CMR assessment.....	77
PBT and vPvB assessment.....	77
Assessment under Article 57(f).....	78
<i>Summary of the data on the intrinsic/hazardous properties (providing scientific evidence of probable serious effects to HH and/or ENV)</i>	78
<i>Equivalent level of concern assessment</i>	78
<i>Conclusion on the Article 57(f) assessment</i>	84
PART II	86
7. REGISTRATION AND C&L NOTIFICATION STATUS	86
Registration status.....	86
CLP notification status.....	86
8. TOTAL TONNAGE OF THE SUBSTANCE.....	87
9. INFORMATION ON USES OF THE SUBSTANCE.....	87
10. INFORMATION ON STRUCTURE OF THE SUPPLY CHAIN	88
11. ADDITIONAL INFORMATION	88
REFERENCES FOR PART I.....	89

TABLES

Table 1: Substance identity	10
Table 2: Overview of physicochemical properties¹	11
Table 3: Results for Surface water compartment	14
Table 4: Results for sediment compartment	15
Table 5: Results for waste water	15
Table 6: Results for sewage sludge	16
Table 7: Results for biota	16
Table 8: Descriptive data analysis for Naiades (suite)	17
Table 9: Descriptive data analysis from Literature review	18
Table 10: Summary table of <i>in vitro</i> mechanistic data (chronologic order)	26
Table 11: Summary table of <i>in vivo</i> mechanistic data (chronologic order)	34
Table 12: Summary table of <i>in vivo</i> adverse effect data (chronologic order)	46
Table 13: Summary table of survival data - Japanese medaka (<i>Oryzias latipes</i>)	60
Table 14: Summary table of survival data - Zebrafish (<i>Danio rerio</i>)	61
Table 15: Line of evidence in relation to TPhP sexual dysfunction in <i>in-vivo</i> studies	64
Table 16: Line of evidence in relation to TPhP EAS activity	67
Table 17: Dose-response and temporal concordance between the key events for male fish	74
Table 18: ELoC summary reporting	82
Table 19: Registration status	86
Table 20: CLP notifications	86
Table 21: Tonnage status	87
Table 22: Uses	87

ABBREVIATIONS:

adcy3 : adenylyl Cyclase 3
ALDH: aldehyde dehydrogenase
AO: adverse outcome
AR: androgen receptor
3 β -HSD: 3 β -hydroxysteroid dehydrogenase
17 β -HSD: 17 β -hydroxysteroid dehydrogenase
BBzP: benzyl butyl phthalate
BCF: bioconcentration factor
BOD: biochemical oxygen demand
cAMP: cyclic adenosine monophosphate
CF: condition factor
CHO: chinese hamster ovary cells
C&L: classification and labelling
COS-7: simian kidney cells
CYP11A1: cholesterol side-chain cleavage
CYP11B2: aldosterone synthase
CYP17: 17 α -hydroxylase
CYP19: cytochrome P450 aromatase
DEGs: differentially expressed genes
DEP: diethyl phthalate
DEPs: differentially expressed proteins
DHEA: dehydroepiandrosterone
DiBP: diisobutyl phthalate
Dio1: thyroxine deiodinase, type I
Dio2: thyroxine deiodinase, type II
dpf/h: days post fertilisation/hatch
DT50: degradation half-life time
dw: dry weight
E2: 17 β -Estradiol
E2-F: fluorescent derivative of 17 β -Estradiol
EAS: Estrogen/Androgen/ Steroidogenesis (modalities)
EATS: Estrogen/Androgen/Thyroidal/Steroidogenesis (modalities)
EC ED EAG: Expert Advisory Group of the European Commission on Endocrine Disruptor
EC10: 10% effective concentration
EC20: 20% effective concentration
EC50: half maximal effective concentration
ECHA: European Chemicals Agency
ED: endocrine disruptor
EDC: endocrine disrupting chemical
EDC-WG: ANSES' Thematic Working group on Endocrine Disruptors
EFSA: European Food Safety Authority
EGFR: epidermal growth factor receptor
ELoC: equivalent level of concern
ER: estrogen receptor
ERE: estrogen response element
FRTL: rat thyroid follicular cell strain
FSDT: fish sexual development test
G15: GPR30 inhibitor
GC/MS: gas chromatography/ mass spectrometry
GH3: rat pituitary-derived cell line that expresses the Pit-1 transcription factor.
GPER: G protein-coupled estrogen receptor
GR: glucocorticoid receptor
GS: Gosner stage
GSI: gonadosomatic index (gonadal weight/body weight x 100)
H295R: Human adrenocarcinoma cell line
HEK293: immortalised human embryonic kidney cells

HepG2: Human liver cancer cell line
HG5LN: HeLa cells with the (GAL4RE) 5-betaGlobin-Luciferase-SV40-Neomycin plasmid
HMGR: 3-Hydroxy-3-methylglutaryl-coenzyme A reductase
hpf: hours post-fertilisation
HPLC: high performance liquid chromatography
HPSE: Heparanase
HSD3 β 2: 3 β -hydroxysteroid dehydrogenase type 2
HSI: hepatosomatic index
IC50: concentration required to inhibit the cell viability by 50%
JRC: Joint Research Centre
KEs: key events
Koc: organic carbon-water partition coefficient
Kow: octanol/water partition coefficient
kPa: kilopascal
11-KT: 11-ketotestosterone
LC: liquid chromatography
LC10: concentration inducing 10% lethality
LC50: concentration inducing 50% lethality
Lhcgr: luteinising hormone/choriogonadotropin receptor
LOD: limit of detection
LOEC: lowest observed effect concentration
LOQ: limit of quantification
MA-10: mouse Leydig cell line tumor
MAPK: mitogen-activated protein kinase
MBzP: monobenzyl phthalate
MCF-7: breast cancer cell line (Michigan Cancer Foundation-7)
MEP: monoethyl phthalate
MIE: molecular initiating event
MoA: mode of action
MR: mineralocorticoid receptor
MS: mass spectrometry
MSCA: member state competent authority
MTC: maximum tolerated concentration
MVLN ER α -positive and hormone-responsive human breast carcinoma cell line derived from the MCF-7 cell line, stably transfected with an ER-controlled luciferase reporter gene construct
nd: not detected
nER: nuclear estrogen receptor
NIS: sodium/iodide symporter
Nkx2.1: NK2 homeobox 1 also known as thyroid transcription factor 1 (TTF-1).
NO(A)EL: no observed (adverse) effect level
NOEC: no observed effect concentration
NR: not reported.
OEL: occupational exposure limit
Pax8: paired box gene 8
PI3K-Akt: phosphoinositide 3-kinase/protein kinase B
POMC: proopiomelanocortin receptor
PPAR γ : peroxisome proliferator-activated receptor- γ
PXR: pregnane X receptor
QSAR: Quantitative Structure-Activity Relationship
RT-PCR: real-time polymerase chain reaction
TWA: time weighted average
RA: retinoic acid
RAR: retinoic acid receptor
RBA: relative binding affinity
RDH: retinal dehydrogenase
REC/RIC20: response equal to 20 % agonist/inhibitory activity
RGA: reporter gene assay

RPE: relative proliferative effect
SKBR3: human breast cancer cell line isolated by the Memorial Sloan-Kettering Cancer Center in 1970
SOD: superoxide dismutase
SPM: suspended particulate matter
SRC: steroid receptor co-activators
SSC: secondary sex characteristics
StAR: steroidogenic acute regulatory protein
SULT: sulfotransferase
SVHC: substance of very high concern
T3 : 3,5,3-triiodo-L-thyronine
T4: L-thyroxine
T: testosterone
Tg: thyroglobulin
TPhP: triphenyl phosphate
TPO: thyroperoxidase
TR: thyroid hormone receptor
TSH: thyroid-stimulating hormone (thyrotropin)
TSHR: thyroid-stimulating hormone receptor (thyrotropin receptor)
TSPO: translocator protein
TTR: transthyretin receptor
UHPLC-MS/MS: ultra-high performance liquid chromatography tandem mass spectrometry
US: United States of America
US EPA: Environmental Protection Agency of the United States of America
VTG: vitellogenin (protein)
vtg: vitellogenin (gene)
WHO/IPCS: International Program on Chemical Safety of the World Health Organisation
ww: wet weight

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57

Substance name: Triphenyl phosphate (TPhP)

EC number: 204-112-2

CAS number: 115-86-6

- It is proposed to identify the substance as a substance of equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of Regulation (EC) No 1907/2006 (REACH) according to Article 57(f) of REACH Regulation.

Summary of how the substance meets the criteria set out in Article 57 of the REACH Regulation

TPhP is proposed to be identified as a substance of very high concern in accordance with Article 57(f) of Regulation (EC) 1907/2006 (REACH) because it is a substance with endocrine disrupting properties for which there is scientific evidence of probable serious effects to the environment, which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 REACH.

The available *in vitro* information demonstrates the capacity of TPhP to competitively bind and activate nuclear estrogen receptors ER α and ER β of several vertebrate species including rat, mouse, fish, chicken, frog and turtle as evidenced by ER transactivation in reporter cell lines. In addition, TPhP can induce ER-regulated gene expression, and related physiological cell responses (e.g., increased cell proliferation). Two recent studies show that TPhP can also bind and activate GPER. The available H295R assays on human adrenal carcinoma cells show that TPhP affects steroidogenesis by increasing estrogen levels (17 β -estradiol) and by increasing expression of genes involved in this pathway like CYP19 and 3 β -HSD2. *In vivo* fish studies indicate that CYP19A is significantly upregulated by exposure to TPhP. Significant alteration of plasmatic concentrations of E2 and E2/T ratio (more specific observation in females) and E2/11-KT ratio (more specific observation in males) can result from this modification in the steroidogenesis pathway. The degree of perturbation of circulating steroid concentrations depends on the fish developmental stage, species and tested concentrations. The observations of VTG concentrations, that are consistent with perturbation of E2 concentrations, suggest an EAS activity of TPhP in female and male zebrafish, with altered plasmatic concentration of VTG. Therefore, TPhP exerts an effect on the endocrine balance in fish. It has EAS activity as clearly shown both *in vitro* and *in vivo*.

In vivo studies on fish reproduction show an alteration of gametogenesis in both sexes with a disruption of testis and ovary maturation followed by a decrease in reproductive success in terms of fecundity (decrease in egg production, spawning events fertility (impaired hatchability) and fertilisation rate). Therefore, TPhP shows adverse effects on fertility and reproduction in fish, observed in several studies at levels without concurrent systemic effects.

The consistency between the observed adverse effects and EAS activity provides evidence that EAS modalities are plausibly biologically linked to the adverse effects. Depending on the developmental stage, species and concentration, antagonist and agonist effects are observed in organisms, leading *in vivo* to perturbations of circulating steroid concentrations in most of the analysed studies.

The effect observed on reproduction in fish (fecundity and fertility) can affect population stability and is considered as an adverse effect relevant at population level.

Based on all available scientific evidence, it can be concluded that TPhP fulfils the WHO/IPCS definition of an endocrine disruptor.

The very high concern raised by this property is substantiated by the severity and irreversibility of the effects on organisms and populations that may have long term consequences, the large variety of species that may be adversely affected and the difficulties to quantify a safe level of exposure with regard to the endocrine mediated effects.

In conclusion, there is scientific evidence that TPhP causes probable serious effects to the environment due to its endocrine disrupting properties, which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of the REACH Regulation.

Registration dossiers submitted for the substance: Yes

PART I

Justification


1. Identity of the substance and physical and chemical properties

Name and other identifiers of the substance

Table 1: Substance identity

EC number:	204-112-2
EC name:	triphenyl phosphate
CAS number (in the EC inventory):	115-86-6
CAS number:	115-86-6
IUPAC name:	triphenyl phosphate
Index number in Annex VI of the CLP Regulation	None
Molecular formula:	C ₁₈ H ₁₅ O ₄ P
Molecular weight range:	326.28
Synonyms:	TPhP

Structural formula:



Composition of the substance

Name: Triphenyl phosphate

Description: Colourless, odourless solid.

Substance type: mono-constituent

Identity and composition of degradation products/metabolites relevant for the SVHC assessment

Not relevant for the SVHC assessment of the substance.

Identity and composition of structurally related substances (used in a grouping or read-across approach)

Not applicable.

Physicochemical properties

Table 2: Overview of physicochemical properties

Property	Description of key information	Value [Unit]	Reference/source of information
Physical state at 20°C and 101.3 kPa	Visual inspection	Colourless, odourless solid.	Data quoted from the FR conclusion document ¹ published in: https://echa.europa.eu/documents/10162/916779d9-ec10-07fa-f178-9562bdd7dedc
Melting/freezing point	Non-guideline publication	49.5 - 50 °C. Melting point values within the range 49 - 50.5 °C are reported in a variety of secondary sources.	
Boiling point	Extrapolated according to the Clausius-Clapeyron equation using experimentally derived parameters - non-guideline publication.	414 °C at 101.3 kPa. One publication reported decomposition at or near the boiling point.	
Vapour pressure	Non-guideline publication using an isoteniscope under a nitrogen atmosphere. Results extrapolated using the Clausius-Clapeyron equation.	0.000853 Pa at 25°C	
Density	Non-guideline publication.	The relative density at 50°C is given as 1.21g/cm ³ .	
Water solubility	Non-guideline publication.	1.9 mg.l ⁻¹ at 20°C. Other supporting publications/reference sources give similar values. No modern guideline study.	
Partition coefficient n-octanol/water (log value)	Non-guideline but similar to shake-flask.	Log Pow 4.63. Values between 4.5 and 4.7 are reported in various publications	
Granulometry	OECD Guideline 110 (Particle Size Distribution / Fibre Length and Diameter Distributions).	All particles with a mean diameter < 100 µm have a mass fraction of 0.41 %. 0.023 % / 0.019 % (spherical/ cubical) particles of this mass fraction have a mean diameter < 4 µm.	

¹ Physicochemical properties quoted in the Conclusion document were based on the analysis of physicochemical properties extracted from the registration dossier of the lead registrant as disseminated in ECHACHEM (https://chem.echa.europa.eu/100.003.739/dossier-view/9b2f054a-c82b-4cc1-9393-415a4f7781c1/4c37f1df-d04d-4911-8afc-4a9bf3b52bf0_4c37f1df-d04d-4911-8afc-4a9bf3b52bf0?searchText=204-112-2).

2. Harmonised classification and labelling

No current harmonised classification for TPhP.

3. Environmental fate properties

The environmental fate data presented in this section are extracted from the registration dossier of the lead registrant as disseminated in ECHACHEM¹ and are provided as contextual information for the SVHC assessment of the substance that focus on endocrine disruptor (ED) properties for the environment. These data have not been further evaluated.

Degradation

Abiotic degradation

Hydrolysis

The main study gave half-lives of 3 days at pH 9, 19 days at pH 7 and >28 days at pH 5 at 25°C (Mayer et al., 1981). One study indicates that 100 % of the TPhP is hydrolysed after 10 minutes at pH 13 (Ishikawa et al., 1985). Another study indicates that at pH 4 the rate of disappearance is too slow to measure (Howard et al., 1979). All studies show that under alkaline conditions TPhP is rapidly degraded, while under acidic conditions TPhP is relatively stable.

Phototransformation in air

From AOPWIN v1.92, a calculation based on a 12 hour-day and a concentration of 1.5×10^6 OH/cm³ gave a half-life of TPhP of 11.85 hours.

Phototransformation in water

Two studies (Ishikawa et al., 1992; Wan and Wong, 1994) are cited for phototransformation in water, and the registrant concluded that TPhP was rapidly degraded when irradiated at 254 nm. The key test was performed at pH 3 and pH 10. However, since the stratospheric ozone layer prevents ultraviolet light of less than 290 nm from reaching the earth's surface, only light at wavelengths between 290 and 750 nm can result in photochemical transformations in the environment.

Phototransformation in soil

No relevant information available.

Biodegradation

Biodegradation: screening tests

Eight biodegradation screening studies are summarised in the registration dossier. These include tests of both ready and inherent biodegradation, as well as several river and pond sediment assays. The key study is CITI (1992), which is an OECD 301C test where biodegradation between 83-94% (based on BOD after 28 days was observed).

¹ https://chem.echa.europa.eu/100.003.739/dossier-view/9b2f054a-c82b-4cc1-9393-415a4f7781c1/4c37f1df-d04d-4911-8afc-4a9bf3b52bf0_4c37f1df-d04d-4911-8afc-4a9bf3b52bf0?searchText=204-112-2

Biodegradation in water and sediment

An OECD 303A (sewage treatment plant simulation study) is included assessing aerobic degradation (Unpublished study report, 1982). The registrant concludes that TPhP is readily degraded in water/sediment systems.

Biodegradation in sediment

No data available

Biodegradation in soil

Two simulation tests for biodegradation in soil (aerobic and anaerobic conditions) are summarised in the registration dossier (Anderson et al., 1993). They calculated a half-life (DT50) of 37 days in aerobic conditions and 21 days in anaerobic conditions.

Environmental distribution

Adsorption/desorption

Four studies in the registration dossier provide information on adsorption/desorption (Anderson et al., 1993; Mayer et al., 1981; Boethling & Cooper, 1985; Huckins et al., 1991). The registrant notes that the range of values (Koc between 2514 and 5500) indicates that the substance will be relatively immobile in soil. The key data is Anderson et al. (1993) using three soils, which gave measured Koc values between 2414 and 3561.

Volatilisation

The Henry's law constant has been calculated by QSAR using HENRYWIN v3.2. The estimated Henry's law constant is 0.004 Pa.m³/mol.

Distribution modelling

The registrant includes Mackay Level 1 modelling (Unpublished study report, 2002), which provides the following distribution:

Air:	0.7%
Water:	14.3%
Soil:	43.9%
Sediment:	41%
Suspended sediment:	0.07%
Biota:	0.03%
Aerosol:	0%

The following inputs were used: temperature = 25 °C, vapour pressure = 8.35 x 10E-4 Pa, water solubility = 1.90 g/m³, log Kow = 4.59.

Environmental occurrence data

This section integrates data from i) the Norman EMPODAT database, ii) the French Naïades database and, iii) a review of the scientific literature.

Norman EMPODAT database


Norman EMPODAT is a database of geo-referenced monitoring and bio-monitoring data in Europe on emerging substances in the following matrices: water, sediment, biota, SPM, soil, sewage sludge and air. This database cannot be used as representative of contaminated areas associated to industrial activities, but rather as an initial overview of the global state of various environmental media.

Surface water - results

24 580 samples, mainly from three countries (France, Germany and Netherlands) are provided in this database. It should be noted that the variability of the detection and of the quantification limits restrains the data exploitation. 8% of data was reported with no information about the limit of quantification, 85% of values with $LOQ \geq 0.01 \mu\text{g.l}^{-1}$ and 24% of values with $LOQ \geq 0.1 \mu\text{g.l}^{-1}$.

Table 3: Results for Surface water compartment

Descriptive data analysis for surface water – (2002 – 2020) Total individual values > LOQ: 1835 (7.47% of total analysis (24580))	
Parameter	Concentration Value ($\mu\text{g.l}^{-1}$)
90 th Percentile	5.42E-02
Mean	2.79E-02
Median	1.50E-02
Max	1.74

**Figure 1: Percentage of surface water data (Total number of analysis) distribution per country**

Sediment - results

Forty samples from eight countries (Moldova, Ukraine, Bulgaria, Georgia, Germania, Hungary, Romania, and Montenegro) are provided in this database for TPhP with 4 values above the LOD (1.5 µg/kg_{dw}). The other 36 samples had a LOD of 5 µg/kg_{dw}.

Table 4: Results for sediment compartment

Descriptive data analysis for sediment – values above the LOD								
COUNTRY	Ecosystem / Matrix: sediment	Concentration value	Unit	Month	Year	River basin name	Limit of Detection (LOD): [µg/kg _{dw}]	Analytical method
Germany	Sediments - River water	6	µg/kg dry weight	7	2019	Danube	1.5	HPLC-MS or MS/MS
Hungary		9						
Bulgaria		9						
Ukraine		5						

Waste water – results

The data for waste water covers the year 2007 and 3 consecutive years (2017-2018-2019). The range of the detection limits and the quantification limits in µg.l⁻¹ are [8E-05; 5E-02] and [2.5E-04; 6E-02] respectively.

Table 5: Results for waste water

Descriptive data analysis for waste water Total individual value (> LOQ): 29 (28.2% of total analysis) Total number of analysis: 103	
Parameter	Concentration Value (µg.l ⁻¹)
90 th Percentile	7.86E-02
Mean	3.52E-02
Median	1.77E-02
Max	1.96E-01




Figure 2: Percentage of waste water data (total number of analysis) distribution per country

Sewage sludge - results

Eight samples from Sweden only are provided in this database for TPhP. No information about the limit of quantification or detection is available.

Table 6: Results for sewage sludge

Descriptive data analysis for sewage sludge						
Country	Type	Value	Unit	Month	Year	Analytical method
Sweden	Sewage sludge - Mixed	54	µg/kg dry weight	11	2007	GC-MS or MS/MS
	Sewage sludge - Mixed	265	µg/kg dry weight	10	2007	GC-MS or MS/MS
	Sewage sludge - Mixed	54	µg/kg dry weight	10	2007	GC-MS or MS/MS
	Sewage sludge - Mixed	136	µg/kg dry weight	10	2007	GC-MS or MS/MS
	Sewage sludge - Municipal	107	µg/kg dry weight	10	2007	GC-MS or MS/MS
	Sewage sludge - Industrial	32	µg/kg dry weight	2	2008	GC-MS or MS/MS
	Sewage sludge - Mixed	104	µg/kg dry weight	10	2007	GC-MS or MS/MS
	Sewage sludge - Municipal	1306	µg/kg dry weight	10	2007	GC-MS or MS/MS

Biota - results

Among the 205 samples available in this database for TPhP, 2 samples on marine mammals and raptors are above the LOD (range value [0.00125; 5] µg.kg_{ww}⁻¹).

Table 7: Results for biota

Descriptive data analysis for biota (205 samples with 2 samples > LOD)						
Country	Sample matrix	Value	Unit	Year	Species group	Species name
Belgium	Biota - Territorial (marine) water	0.99	µg/kg wet weight	2021	Marine mammals	<i>Phocoena phocoena</i>
Belgium	Biota - Terrestrial	0.81	µg/kg body weight	2021	Raptors	<i>Buteo</i>

Naiades database

Naiades is a database collecting French data only. It is an observatory on the quality of river and water bodies. 4705 samples are included in the database for TPhP between 2020 and 2022, including 4647 values below the limit of quantification (LOQ) and 58 values above the LOQ (range: 0.02-0.1 $\mu\text{g.l}^{-1}$). These values ($>$ LOQ) were measured in a single French region and by a single analysis laboratory.

Table 8: Descriptive data analysis for Naiades (suite)

Descriptive data analysis for value > LOQ (58 values)	
Parameter	Raw water concentration ($\mu\text{g.l}^{-1}$)
90 th Percentile	0.12
Mean	0.06
Median	0.046
Max	0.22

Scientific literature related to occurrence data

A scientific literature review was conducted up to May 2023. The systematic literature search was performed in Scopus database. A single concept strategy search was applied to retrieve all relevant information on TPhP by using its Chemical Abstracts Service Registry Number (CAS No 115-86-6), scientific chemical names, and common names, as recommended in the EDC guidance (ECHA/EFSA, 2018). 812 entries were recorded from 23 studies that included monitoring data in an environmental media (water, sediment, biota) in Europe.

Table 9: Descriptive data analysis from Literature review

Descriptive data analysis from Literature review								
Country	Type of environmental media	Type of value	Value	Unit	LOD	Analytical method	Year	REF
WATER								
Spain	Nalon river	Mean (min-max)	1.7 (1.6-2.4)	ng.l ⁻¹	1.6 ng.l ⁻¹	GC-MS	2012 (11 samples)	Cristale et al., 2013
Spain	Arga river	Mean (min-max)	2.8 (1.6-7.2)	ng.l ⁻¹	1.6 ng.l ⁻¹	GC-MS	2012 (8 samples)	Cristale et al., 2013
Spain	Besos river	Mean (min-max)	12.5 (01.6-35)	ng.l ⁻¹	1.6 ng.l ⁻¹	GC-MS	2012 (13 samples)	Cristale et al., 2013
Sweden	21 rivers	Mean	< LOD	ng.l ⁻¹	37 ng.l ⁻¹	GC-MS-MS	2013 (25 samples)	Gustavsson et al., 2018
Greece	River water Dissolved fraction	Mean (min-max) Detection frequencies	137 (40-258) 100%	ng.l ⁻¹	NR	GC-MS-MS	2019-2020 (12 samples)	Pantelaki et al., 2021
Greece	River water Particulate fraction	Mean (min-max) Detection frequencies	100 (46-219) 100%	ng.l ⁻¹	NR	GC-MS-MS	2019-2020 (12 samples)	Pantelaki et al., 2021
Greece	Coastal water Dissolved fraction	Mean (min-max) Detection frequencies	135 (41-260) 100%	ng.l ⁻¹	NR	GC-MS-MS	2019-2020 (18 samples)	Pantelaki et al., 2021
Greece	Coastal water Particulate fraction	Mean (min-max) Detection frequencies	47 (34-81) 100%	ng.l ⁻¹	NR	GC-MS-MS	2019-2020 (18 samples)	Pantelaki et al., 2021
Greece	Streams Dissolved fraction	Mean (min-max) Detection frequencies	279 (45-1142) 100%	ng.l ⁻¹	NR	GC-MS-MS	2019-2020 (12 samples)	Pantelaki et al., 2021
Greece	Streams Particulate fraction	Mean (min-max) Detection frequencies	281 (38-980) 100%	ng.l ⁻¹	NR	GC-MS-MS	2019-2020 (12 samples)	Pantelaki et al., 2021
SEDIMENT								
United Kingdom	Worcester-Birmingham Canal	Mean (min-max)	4 (0.1-26)	ng/g dry weight	NR	GC-MSD	2019-2021 (12 samples)	Onoja et al., 2023
United Kingdom	River Severn	Mean (min-max)	1 (0.3-8)	ng/g dry weight	NR	GC-MSD	2019-2021 (12 samples)	Onoja et al., 2023
United Kingdom	River Sowe	Mean (min-max)	1 (0.4-2)	ng/g dry weight	NR	GC-MSD	2019-2021 (12 samples)	Onoja et al., 2023

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

United Kingdom	River Tame	Mean (min-max)	2 (0.3-9)	ng/g dry weight	NR	GC-MSD	2019-2021 (12 samples)	Onoja et al., 2023
Spain	Nalon	Mean (min-max)	15 (15-15)	µg/kg dry weight	15 µg/kg dw	GC-MS	2012 (8 samples)	Cristale et al., 2013
Spain	Arga	Mean (min-max)	22.17 (15-44)	µg/kg dry weight	15 µg/kg dw	GC-MS	2012 (6 samples)	Cristale et al., 2013
Spain	Besos	Mean (min-max)	37.71 (19- 63)	µg/kg dry weight	15 µg/kg dw	GC-MS	2012 (7 samples)	Cristale et al., 2013
Italy	Adige River	Max	9.69	ng/g dry weight	0.08 ng/g dw	Pressurised liquid extraction-Turbulent Flow Chromatography-LC-MS/MS	2015 (20 samples)	Giulivo et al., 2017
Greece	Evrotas River	Max	0.67	ng/g dry weight	0.08 ng/g dw	Pressurised liquid extraction-Turbulent Flow Chromatography-LC-MS/MS	2014-2015 (12 samples)	Giulivo et al., 2017
Slovenia	Sava River	Max	< LOD	ng/g dry weight	0.08 ng/g dw	Pressurised liquid extraction-Turbulent Flow Chromatography-LC-MS/MS	2014-2015 (20 samples)	Giulivo et al., 2017
France	Gulf of Lion	Mean (Min-Max)	1.57 (0-5.17)	ng/g dry weight	NR	GC-MS	2018 (12 samples)	Alkan et al., 2021
BIOTA								
France	Mussels (<i>Mytilus spp</i>) and oysters (<i>Crassostrea gigas</i>)	Mean or Median (Min-Max)	1.12 (0.17-8.01)	ng/g dry weight	53 pg/g dw	LC-ESI-MS/MS	2014-2021	Aminot et al., 2023
Italy	Adige River (Fish)	Max	30.3	ng/g lipid weight	1.30 ng/g lw	Pressurised liquid extraction-Turbulent Flow Chromatography-LC-MS/MS	2015 (13 samples)	Giulivo et al., 2017
Greece	Evrotas River (Fish)	Max	< LOD	ng/g lipid weight	1.30 ng/g lw	Pressurised liquid extraction-Turbulent Flow Chromatography-LC-MS/MS	2015 (4 samples)	Giulivo et al., 2017
Slovenia	Sava river (Fish)	Max	< LOD	ng/g lipid weight	1.30 ng/g lw	Pressurised liquid extraction-Turbulent Flow Chromatography-LC-MS/MS	2015 (10 samples)	Giulivo et al., 2017

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

Denmark	Western Iceland (Denmark Strait) Fin Whale	Mean	59.2	ng/g lipid weight	NR	Ultra-sonication extraction-LC-MS/MS	2015 (20 whales)	Garcia-Garin et al. 2020
Western Mediterranean area	Western Mediterranean Sea <i>Sardina Pilchardus</i>	Mean (Min-Max)	2.42 (ND-9.39)	ng/g wet weight	0.12 ng/g ww (LOQ)	Ultra-sonication extraction-TFC-HPLC-MS/MS	2019	Sala et al., 2022
Western Mediterranean area	Western Mediterranean Sea <i>Engraulis Encrasicolus</i>	Mean (Min-Max)	0.66 (ND-2.28)	ng/g wet weight	0.12 ng/g ww (LOQ)	Ultra-sonication extraction-TFC-HPLC-MS/MS	2019	Sala et al., 2022
Western Mediterranean area	Western Mediterranean Sea <i>Merluccius Merluccius</i>	Mean (Min-Max)	ND	ng/g wet weight	0.12 ng/g ww (LOQ)	Ultra-sonication extraction-TFC-HPLC-MS/MS	2019	Sala et al., 2022
Mediterranean sea (Catalan coast)	Turtle	Mean (Min-Max)	0.15 (ND-1.08)	ng/g wet weight	0.12 ng/g ww (LOQ)	Ultra-sonication extraction-TFC-HPLC-MS/MS	2014-2017	Sala et al., 2021
Mediterranean sea (Balearic Island)	Turtle	Mean (Min-Max)	0.31 (ND-1.34)	ng/g wet weight	0.12 ng/g ww (LOQ)	Ultra-sonication extraction-TFC-HPLC-MS/MS	2014-2017	Sala et al., 2021
Mediterranean sea (Balearic Island)	Prey of turtle	Mean (Min-Max)	1.25	ng/g wet weight	0.12 ng/g ww (LOQ)	Ultra-sonication extraction-TFC-HPLC-MS/MS	2014-2017	Sala et al., 2021

NR: not reported.

Human occurrence

In addition to indirect exposure *via* the environment, humans are exposed *via* several consumer articles or in the workplace (widespread uses), in formulation or repacking, on industrial sites and in manufacturing. TPhP is also present as an impurity in many other organophosphate flame retardants or as a constituent of this family of compounds.

Human biomonitoring data demonstrate also the presence of TPhP or its metabolites in human milk (Sundkvist et al., 2010; Kim et al., 2014; Zheng et al., 2021), human placenta (Ding et al., 2016; Zhao et al., 2017), blood serum (Ya et al., 2019; Zhao et al., 2016) and urine (Carignan et al., 2016; Li N. et al., 2019a). TPhP has also frequently been detected in indoor house dust samples in Europe (Belgium and Sweden) and outside Europe (Japan, Philippines and USA) with detection frequencies up to 98% (Kanazawa et al., 2010; Kim et al., 2013; Marklund et al., 2003; Stapleton et al., 2009; Van den Eede et al., 2011). It can be concluded that TPhP reaches diverse environmental compartments and biota of remote areas.

Summary on occurrence and environmental distribution

Several literature studies demonstrate that TPhP can be found in Europe and that it is ubiquitous in several environmental compartments (water, sediment, sewage sludge, indoor dust/air). Therefore, many environmental species and humans are exposed more or less continuously to TPhP and exposure cannot be avoided.

Bioaccumulation

Bioaccumulation in aquatic organisms (pelagic & sediment organisms)

The registration dossier of the Lead Registrant cites 9 fish bioaccumulation studies (Muir et al., 1980; Mayer et al., 1981; Sasaki et al., 1981; Muir et al., 1982; Sasaki et al., 1982; Muir et al., 1983; Boethling & Cooper, 1985; Kuehl and Haebler, 1995; Lo et al. 2000), although only three were assessed to be valid of which two are conventional tests (the third was blubber samples from dead Common bottlenose dolphins). These two conventional tests provide bioconcentration factor (BCF values of 110 and 144.

Bioaccumulation in terrestrial organisms (soil dwelling organisms, vertebrates)

No relevant information available.

Summary

Considering data coming from the registration dossier of the lead registrant, the substance is rapidly degraded in water under alkaline conditions, TPhP photodegrades rapidly in the atmosphere (DT50= 11.85 h) and TPhP would be readily biodegraded in water/sediment systems. The data provided in the registration dossier of the Lead registrant would indicate that TPhP is relatively immobile in soil.

4. Human health hazard assessment

Human health data for TPhP were not reviewed and thus not included in this document.

5. Environmental hazard assessment

Please note that the environmental toxicity data presented in this section are extracted from the registration dossier of the Lead Registrant as disseminated in ECHACHEM² and are provided as contextual information for the SVHC assessment of the substance that focuses on ED properties for the environment. These data have not been further evaluated. However, the data used as a basis for the environmental assessment of the potential endocrine disrupting properties of TPhP have been evaluated and are described in section 5.

Aquatic compartment (including sediment)

Fish

Short-term toxicity to fish

Twelve studies are reported in the registration dossier. All of these studies are old (generally performed more than thirty years ago) and mostly used static exposure conditions. Three include chemical analysis, although the results are only reported for one.

The registrant judges the majority of the tests to be Klimisch reliability 2 with two considered unreliable (Klimisch 3 and 4). The reliable studies report a range of LC₅₀ values between >0.32 and 1.26 mg.l⁻¹, with the majority below 1 mg.l⁻¹ (Mayer et al., 1981; Palawski et al., 1983; Huckins et al., 1991; Sasaki et al., 1981). The lowest, reliable value is a 96-h LC₅₀ of 0.36 mg.l⁻¹ for Rainbow Trout *Oncorhynchus mykiss* (Palawski et al., 1983). However, the key study selected by the registrant is Mayer et al. (1981), which, reported a 96-h LC₅₀ of 0.40 mg.l⁻¹ for *O. mykiss* based on nominal concentrations.

Long-term toxicity to fish

There are five long-term studies in the registration dossier, all of which are judged to be Klimisch reliability 2 by the registrant. However, two of these are 96 h tests which should not be considered as part of the chronic dataset. Of the remaining three tests, the key study for long-term toxicity to fish (Sitthichaikasem 1978) started with 10-days old sac fry and lasted 30 days, and is considered part of an Early Life Stage Test. No NOEC was obtained with the TPhP concentrations tested.

Aquatic invertebrates

Short-term toxicity to aquatic invertebrates

A test on the acute toxicity of TPhP to *Daphnia magna* was conducted according to the US guideline EPA-660/3-75-009. Static exposure for 96 h resulted in a LC₅₀ value of 1.0 mg.l⁻¹ (nominal concentration). Lower effective concentrations were found for another crustacean, *Mysidopsis bahia*, in the same test on acute toxicity. Static exposure for 96 h resulted in a LC₅₀ value of 0.18 mg.l⁻¹ (nominal concentration) (Mayer et al., 1981).

Further invertebrate toxicity studies investigated the acute effects of TPhP on scud (*Gammarus pseudolimnaeus*) and midge larvae (*Chironomus riparius*) (Huckins et al., 1991). Static exposure for 96 h resulted in a LC₅₀ value of 0.25 mg.l⁻¹ and 0.36 mg.l⁻¹ respectively (nominal concentrations). Further, Lo et al., 2000 investigated the toxicity of TPhP to golden apple snail

² https://chem.echa.europa.eu/100.003.739/dossier-view/9b2f054a-c82b-4cc1-9393-415a4f7781c1/4c37f1df-d04d-4911-8afc-4a9bf3b52bf0_4c37f1df-d04d-4911-8afc-4a9bf3b52bf0?searchText=204-112-2

(*Pomacea canaliculata*) and reported a relatively lower toxicity than identified for crustaceans in other studies (72 hr, LC₅₀ 38.2 mg.l⁻¹).

Long-term toxicity to aquatic invertebrates

The chronic toxicity of TPhP to aquatic invertebrates (*Daphnia magna*) was tested according to OECD Guideline 211 (*Daphnia magna* Reproduction test). After 21 days a NOEC of 0.254 mg.l⁻¹ (mean measured concentration) was obtained based on reproduction (Unpublished study report, 2000).

Algae and aquatic plants

The registrant reports nine tests. Eight of these tests appear to be the same as those reported in Environment Agency (2009). The registrant judges all of the tests to be Klimisch reliability 2, which is consistent with Environment Agency (2009).

The registrant considers the test (three species and three different media) conducted by Millington et al. (1988) according to an OECD guideline 201 to be the key study. The authors of the test only reported LOEC values. The most critical values were obtained for *Desmodesmus subspicatus* and *Pseudokirchneriella subcapitata*. The data reports LOECs between 0.5 and 5.0 mg.l⁻¹ depending on the growth medium. Mayer et al. (1981) determined the toxicity of TPhP to *Desmodesmus subspicatus* and a 96h-EC₅₀ was determined to be 2 mg.l⁻¹.

Sediment organisms

No relevant information available.

Other aquatic organisms

No relevant information available.

Terrestrial compartment

Toxicity to soil macro-organisms

No relevant information available.

Toxicity to terrestrial plants

No relevant information available.

Toxicity to soil micro-organisms

No relevant information available.

Toxicity to other terrestrial organisms

No relevant information available.

Atmospheric compartment

No relevant information available.

Microbiological activity in sewage treatment systems

The registrant used an activated sludge assay as key study, according to OECD guideline 301C (CITI, 1992). Activated sludge (30 mg.l⁻¹ as concentration of suspended solid) in a volume of 300 mL was exposed to TPhP (100 mg.l⁻¹) for 28 days at 25°C. Degradation was determined by BOD. Under the used conditions, TPhP did not adversely affect the microorganisms present, since TPhP biodegraded to the extent of 83 - 94 %. The key values should be understood as > 100 mg.l⁻¹ as no effect was observed up to the test concentration of 100 mg.l⁻¹.

Toxicity to birds

No relevant information available.

Mammalian wildlife

No relevant information available.

6. Endocrine disruption (Environment)

General approach for the assessment of endocrine properties

Framework of the evaluation

To evaluate whether or not TPhP fulfils the WHO/IPCS definition (WHO/IPCS, 2002) of an endocrine disruptor as interpreted by the EC ED EAG (JRC, 2013), both *in vitro* and *in vivo* data were taken into account, in order to demonstrate:

- Adverse effects
- Endocrine mode of action
- Biological plausible link between adverse effects and endocrine mode of action
-

As highlighted in EDC guidance developed by ECHA and the European Food Safety Authority (EFSA) to identify EDC under the plant protection products and the biocidal product regulations published in 2018 (ECHA/EFSA, 2018), the 'endocrine mode of action' in the second bullet point should be interpreted as 'endocrine activity', i.e., the substance has the potential to alter the function(s) of the endocrine system. Consequently, the third bullet point should be interpreted as biological plausible link between adverse effects and endocrine activity.

Specificity, i.e., that adverse effects were not non-specific secondary consequences of general toxic effects, and population relevance were also considered in the assessment.

The structure and the assessment of data are mainly based on the OECD Revised Guidance Document 150 on standardised test guidelines for evaluating chemicals for endocrine disruption (OECD, 2018).

Two different aspects are assessed separately:

- Evidence for endocrine activity
- Effects on apical endpoints that provide evidence that a substance exerts adverse effects owing to its endocrine activity.

Information sources and strategy for endocrine disruptor identification

Literature review

A literature review was performed following the principles displayed in the EDC guidance (ECHA/EFSA, 2018). The EDC guidance provides a tiered approach to assess the adversity of chemicals on vertebrates, and to link it with an estrogenic (E), androgenic (A), thyroid hormone (T), or steroidogenesis-related (S) mode of action (the so-called EATS modalities). The evidence is first assembled by using a literature review and weight of evidence approach. Then, the EATS-mediated adversity and the endocrine activity are assessed. The detailed methodology is presented in the following sections.

A literature review was conducted up to May 2023. The literature review on TPhP endocrine disruptive properties was focused on fish, amphibian and *in vitro* mechanistic studies. The literature search was performed in Scopus database.

A single concept strategy search was applied to retrieve all relevant information on TPhP by using its Chemical Abstracts Service Registry Number (CAS No 115-86-6), scientific chemical names, and common names, as recommended in the EDC guidance (ECHA/EFSA, 2018).

Studies were included in the literature review based on their relevance when they met all of the following criteria: a) peer-reviewed research articles or primary reports of research findings that presented original data; b) endocrine activity or adversity assessed in *in vitro*, *ex vivo*, or *in vivo* studies in vertebrate species; and c) English-language articles. The relevance filtering was first

based on title and abstract screening, and second, on full-text screening. When checking title and abstract was insufficient to decide if the paper was relevant and should be included in the review, full-text screening was applied (e.g., TPhP not explicitly mentioned in the abstract). Two distinct reviewers shared the two screening phases during the literature search and resolved any conflicts or discrepancies by complementary full-text screening and by discussion.

Assessment of the evidence

The present analysis was performed in collaboration with the ANSES Thematic Working group on Endocrine Disruptors (EDC-WG)³. The studies were retrieved from the literature review as well as from the registration dossier and were considered on the basis of their relevance (see criteria of selection based on relevance above), reliability and adequacy for the analysis. Studies were qualitatively weighted based on expert judgement to produce a conclusion on the selected adverse effects and their ED MoA.

Analysis of the results

The data were grouped into three categories in accordance with the OECD conceptual framework (OECD, 2018) and EU EDC guidance (ECHA/EFSA, 2018):

- a) *in vitro* mechanistic parameters;
- b) *in vivo* mechanistic parameters;
- c) parameters providing information on adversity.

Based on the adverse effects identified, results were further integrated into lines of evidence, defined as a “set of relevant information grouped to assess a hypothesis,” using a weight-of-evidence approach (ECHA/EFSA, 2018).

In vitro information indicative of endocrine activity

Eighteen studies gathered in the literature and included in the review provide *in vitro* mechanistic information on the capacity of TPhP to interact with the endocrine system. Most of the studies focus on TPhP estrogenic activity.

Table 10: Summary table of *in vitro* mechanistic data (chronologic order)

STUDIES INVESTIGATING ENDOCRINE DISRUPTION PROPERTIES OF TPhP		
Methodology	Results (statistically significant)	Reference
AR competitive binding assay (18 to 20h) (Rat AR protein) Dose: 4.28 nM to 0.428 mM.	IC ₅₀ : 1.50E-05 M AR Relative binding affinity (RBA): 0.0205% (moderate binding)	Fang et al., 2003
H295R cell bioassay (48h) Dose: 0 – 0.001 – 0.01 – 0.1 – 1 mg.l ⁻¹	<u>Hormone synthesis</u> 17β-estradiol (E2): ↑ at 1 mg.l ⁻¹ Testosterone (T): ↑ at 1 mg.l ⁻¹ E2/T Ratio: ↑ at 0.1 and 1 mg.l ⁻¹	Liu et al., 2012
Dose: 0 – 0.01 – 0.1 – 1 mg.l ⁻¹	<u>Steroidogenic genes</u> CYP11A1: ↑ at 1 mg.l ⁻¹ CYP11B2: ↑ at 1 mg.l ⁻¹ CYP19A1: ↑ at 1 mg.l ⁻¹	

³ <https://www.anses.fr/en/content/endocrine-disruptors>.

<p>MVLN cell binding bioassay (72h) Dose: 0 – 0.001 – 0.01 – 0.1 – 1 – 10 mg.l⁻¹</p>	<p>HSD3β2: no effect</p> <p><u>SULT genes</u> SULT1E1 ↓ at 1 mg.l⁻¹ SULT2A1 ↓ at 1 mg.l⁻¹</p> <p>↓ binding affinity of E2 for hER (0.001 mg.l⁻¹)</p>	
<p>Cell-based transactivation Assay</p> <p>CHO-K1 cells (24h) Dose: 0.10 to 30 μM.</p> <p>COS-7 cells (24h) Dose: 0.10 to 30 μM.</p>	<p>Era: REC₂₀ = 4.9 μM (weak agonist) Erβ: REC₂₀ = 6.5 μM (weak agonist) AR: RIC₂₀ = 17 μM (weak antagonist) GR: RIC₂₀ = 15 μM (weak antagonist)</p> <p>PPARγ: weak agonist (15% at concentration of 30 μM) PXR: REC₂₀ = 2.8 μM (agonist)</p>	<p>Kojima et al., 2013</p>
<p>Dual Luciferase Reporter Gene Assay / CHO-K1 cells (24h) Dose: 0.1 nM to 1 μM.</p> <p>Yeast two-hybrid assay (4h) Dose: 0.1 nM to 1 μM.</p> <p>E-Screen Assay / MCF-7 cells (5 days) Dose: 0.1 nM to 1 μM.</p> <p>Molecular Docking and Molecular Dynamic Simulation using Discovery Studio 2.5/Ligand fit module.</p>	<p>Era: REC₂₀ of 0.27 μM (↑ activation of Era in a dose-dependent manner)</p> <p>Era: REC₂₀ of 0.65 μM (↑ activation of Era in a dose-dependent manner)</p> <p>Maximum cell proliferation: 1 μM Relative proliferative effect: 37.5%</p> <p>Tight binding affinity for hERα in docking approach (agonist effect)</p>	<p>Zhang et al., 2014</p>
<p>GH3 (rat pituitary) cell assay (48h)</p> <p>Dose (μg.l⁻¹) : 0 – 1 (0.0003 μM) – 10 (0.031 μM) – 100 (0.31 μM)</p> <p>FRTL-5 (rat thyroid follicular) cell assay (24h)</p> <p>Dose (mg.l⁻¹) : 0 – 1 (3.1 μM) – 3 (9.2 μM) – 10 (31 μM)</p>	<p><i>Tshβ</i> ↑ at 0.31 μM <i>Tra</i> ↑ at 0.031 & 0.31 μM <i>trβ</i> ↑ at 0.31 μM <i>dio1</i> ↑ at 0.31 μM <i>dio2</i> no effect</p> <p><i>nis</i> ↑ at 9.2 & 31 μM <i>tg</i> ↓ at 3.1 μM <i>tpo</i> ↑ at 31 μM</p>	<p>Kim et al., 2015</p>

	<p><i>tshr</i> ↓ at 3.1 µM <i>nkx2.1</i> ↓ at 3.1 & 31 µM <i>pax8</i> no effect</p>	
<p>Cell-based transcriptional Assay</p> <p>CHO-K1 cells (24h) Dose: 0.1 to 30 µM</p> <p>COS-7 cells (24h) Dose: 1 to 30 µM</p>	<p>Era: REC₂₀ = 4.6 µM (weak agonist) Erβ: REC₂₀ = 7.3 µM (weak agonist) AR: RIC₂₀ = 11 µM (weak antagonist) GR: RIC₂₀ = 12 µM (weak antagonist)</p> <p>PPARγ: weak agonist (15% at concentration of 30 µM) PXR: REC₂₀ = 2.8 µM (agonist)</p>	Kojima et al., 2016
MCF-7 flow-cytometric proliferation assay (24h) Dose: 10 µM to 1 mM	Relative proliferative effect (RPE) = 87% (E2 100%) EC ₂₀ = 88 µM (E2 8E-07 µM)	Krivoshiev et al., 2016
MA-10 mouse Leydig tumour cells (48h) Dose: 1 to 100 µM	<p>qRT-PCR <i>(Expression of Key Genes Involved in Steroidogenesis)</i> Lhcgr: no effect Adcy3: no effect Star: no effect TSPO: no effect Cyp11A1: no effect Hsd3β: no effect</p> <p>Progesterone production: no effect</p>	Schang et al., 2016
Transthyretin (TTR) binding experiment (2h) Dose: approx. 6 nM to 32 µM	IC ₅₀ : 1083 nM (significant binding ability to TTR compared to T4 = IC ₅₀ of 291 nM)	Zhao et al., 2017
Yeast two-hybrid assay with Japanese medaka Era (mERα) Dose: approx. 0.1 to 100 µM (4h)	IC ₅₀ = 29 µM (antagonist activity)	Li et al., 2018
<p>Yeast two-hybrid assay with human Era (4h) 0.5 nM to 0.1µM.l⁻¹</p> <p>MVLN cell assay with human ERE (72h) 0.1 nM to 20µM.l⁻¹</p>	<p>hERα: Antagonist activity (RIC₂₀ = 2E-05 mol/L) compete with E2 for binding to Era</p> <p>hERE: Agonist response (EC₅₀ = 1.45E-05 mol/L) = Era agonist to activate the ERE pathway</p> <p>Promotes cell proliferation = agonist activity (REC₂₀ = 5.01E-07 M/L, max 77.08%). This proliferative effect was significantly inhibited when cells were pretreated with G15 (a GP30 inhibitor).</p>	Ji et al., 2020

E-SCREEN assay (proliferation MCF7)		
Multiplexed reporter assay (the Ecotox FACTORIAL) 40 µM (24h)	<p>Activation of hPPARγ, mPPARγ, zPPARγ</p> <p>Activation of Zebrafish ER2β, frogER1, turtleER1, chickenER1, hERα</p> <p>Activation of chickenAR</p> <p>No effect on ZebrafishER1 and ER2α, frogER2, hERβ, AR (fish, frog, turtle, human), fishTRα and TRβ, frogTRα, turtleTRα, hTRα and hTRβ</p>	Medvedev et al., 2020
HG5LN cell assay (24h) Dose (µM): 0.1 to 10 µM	<p><u>Transcriptional activation bioassay:</u></p> <p>Activation of hPPARγ (EC50: nd; % max. activity: 28%)</p> <p>Activation of mPPARγ (EC50: nd; % max. activity: 24%)</p> <p>Activation of xPPARγ: (EC50: nd; % max. activity: 21%)</p> <p>Activation of zfPPARγ (EC50 = 3.5 µM; % max. activity: 104%)</p>	Garoche et al., 2021
Multiplexed reporter assay (the Ecotox FACTORIAL) 22.2 µM	Agonist activity on mPPAR γ , hPPAR γ , zPPAR γ	Houck et al., 2021
Reporter gene assay: HEK293 cell assay (39h) with medaka estrogen receptor 1(mEsr1) HepG2 cell assay (39h) with medaka androgen receptor β (mAr β)	<p>mEsr1:</p> <ul style="list-style-type: none"> - Agonist activity (EC₅₀ = 9.7E-06 M). - No antagonist activity up to 1.0 E-05 M. <p>mArβ:</p> <ul style="list-style-type: none"> - No agonist or antagonist activity up to 1.0 E-05 M. 	Kawashima et al., 2021
Molecular dynamics analysis Transcriptomic analysis on L02 cells (HL-7702, human normal liver cells) for 24h. 20 and 200µM	<p>TPhP disturbed conformational changes to affect the structural stability of TRβ ligand binding domain leading to potential disruption of the transcriptional activity of TR.</p> <p><u>Gene expression</u></p> <p>↑ THRβ, RXRA, DIO2 and PIK3R3 (from 20 µM)</p> <p>↓ TP53, NOTCH11 (from 20 µM)</p> <p>↓ AKT3, DIO2, THRA, SLCO1C1 and CASP9 (200 µM)</p> <p>↑ MDM2, BAD (200 µM)</p>	Wang et al., 2021
Molecular docking (CDOCKER)	<p>With ERα, TPhP formed a pi-sigma bond with Arg394 as E2 which in addition bond to His524 and Glu353.</p> <p>With GPER, TPhP formed a pi-sigma bond with Tyr142 while E2 formed hydrogen bonds with Tyr123 and Glu275, and Van der Waals forces with Cys205. G1, a GPER</p>	Ji et al., 2022

<p>H295R cell assay (48h)</p> <p>Dose (μM) : 0.1 – 1 – 2 and 10</p> <p>MVLN (hERα +) cell assay (72h)</p> <p>Dose (μM) : 0.1 – 0.2 – 1 – 2 – 10 – 20</p> <p>SKBR3 (ERα-, GPER+) cell assay (24h)</p> <p>Dose (μM) : 0.1 – 1 and 10</p> <p>ERα- SRC (steroid receptor co-activators) recruitment assay (1 h)</p> <p>Dose 0.01 μM to 1000 μM.</p>	<p>agonist, had a hydrogen bond and a pi-pi bond with Gln138 and Phe208. The CDOCKER interaction energy with TPhP was - 29.60 kcal/mol.</p> <p><u>Hormone levels:</u> <i>E2</i> ↑ at 1, 2 and 10 μM <i>E2/T ratio</i> ↑ at 1, 2 and 10 μM <i>Progesterone</i> ↑ at 2 and 10 μM <i>Estrone</i> ↓ at 2 and 10 μM <i>Testosterone</i> ↓ at 1, 2 and 10 μM</p> <p><u>Transcriptional genes⁴</u> <i>CYP17</i> ↑ at 2 and 10 μM <i>CYP19</i> ↑ at 1, 2 and 10 μM <i>3β-HSD2</i> ↑ at 2 and 10 μM <i>17β-HSD1</i> ↑ at 1, 2 and 10 μM <i>HMGR⁵</i> ↑ at 1, 2 and 10 μM <i>StAR⁶</i> ↑ at 10 μM <i>SULT2A1</i> ↓ at 1, 2 and 10 μM <i>CYP11A1</i> No sign. effect <i>17β-HSD4</i> No sign. effect</p> <p><u>Receptor activation</u> hERα Agonist activity EC50 = 1.2E-05 M</p> <p>GPER-cAMP activation - Interaction with GPER at 1.0E-05 μM <i>via</i> cAMP formation - The agonist effects of TPhP on cAMP production were significantly inhibited when TPhP was co-exposed with G15. - The E2-control group responded as expected.</p> <p>ERα agonist interaction with SRC1–1: EC50 of 2.93E-04 mol/L ERα agonist interaction with SRC3–3: EC50 of 1.30E-05 mol/L, No effect on the other 8 SRCs peptides</p>	
<p>Molecular dynamics simulation</p> <p>Competitive binding assay:</p>	<p>Signs of TPhP binding to GPER leading to the change of its structural conformation for a possible activation</p> <p>TPhP inhibits the binding of E2-F to GPER (probe) (IC50 4.2 μM)</p>	<p>Guan et al., 2022</p>

⁴ Fold above 1.75 are reported in this table.

⁵ HMGR: 3-Hydroxy-3-methylglutaryl-coenzyme A reductase.

⁶ StAR: Steroidogenic acute regulatory protein.

<p>SKBR3 cells (10 min) Dose: 1 nM to 100 µM</p> <p>Transcriptomic and proteomic analysis (SKBR3 cells)</p>	<p>64 genes were enriched for both mRNA and protein after TPhP exposure. Among these overlaps, 52 DEGs⁷/DEPs⁸ were found to be up-regulated or down-regulated together. Combined transcriptome and proteome analysis showed that TPhP could exert its estrogen-like proliferation-promoting effect through the interaction of the PI3K-Akt⁹ signalling pathway, MAPK¹⁰ signalling pathway, and EGFR¹¹ signalling pathway, which are downstream pathways mediated by GPER.</p>	
---	--	--

Estrogen pathway

Overall, 10 studies investigating the estrogenic activity of TPhP *in vitro* were identified and analysed. The main results are summarised below and further discussed in the conclusion section.

TPhP has been tested in different types of *in vitro* assay including binding assays, transactivation in reporter gene assays, regulated gene expression and G protein-coupled estrogen receptor (GPER) signalling pathways.

It appears that TPhP has **agonist activity on human ER α and / or ER β receptor and decreases the binding affinity of E2 to ER** (Liu et al., 2012; Kojima et al., 2013; Zhang et al., 2014; Kojima et al., 2016; Medvedev et al., 2020, Ji et al., 2022). The agonist activity is also reported for the medaka ER1 (Kawashima et al., 2021) and for the zebrafish ER2 β , frog ER1, turtle ER1 and chicken ER1 (Medvedev et al., 2020). In some yeast two-hybrid assays with human and medaka ER α , an antagonist effect is shown indicating **a competition between TPhP and E2 for binding to ER α** (Li et al., 2018; Ji et al., 2020). Zhang et al. (2014) also demonstrated a binding affinity of TPhP to hER α with a docking approach. Moreover, TPhP can induce the ERE pathway as demonstrated by a dose-response agonist effect in MVLN cells (transfected with the human ERE-luciferase reporter gene) (Ji et al., 2020). This estrogenic activity is supported by proliferation assays on MCF-7 cells that show TPhP promotes cell proliferation (Zhang et al., 2014; Krivoshev et al., 2016; Ji et al., 2020).

In Guan et al. (2022), the authors explored the effects of TPhP on the key downstream signalling pathways mediated by the G-coupled protein ER (GPER) through combined transcriptome and proteome analysis. These relationships were also tested on a SKBR3 cell-based fluorescent competitive binding assay. The authors report that **TPhP binds GPER** with an IC₅₀ of 4.2 µM. This is confirmed by the E2-F displacement experiments. Combined transcriptome and proteome analyses show that **TPhP could exert its estrogen-like proliferation-promoting effect** through the interaction of the PI3K-Akt signalling pathway, MAPK signalling pathway, and EGFR signalling pathway, which are downstream pathways mediated by GPER.

In Ji et al. (2022), the interaction of TPhP with GPER was confirmed by molecular docking and their activation on GPER-mediated signalling was investigated in SKBR3 cells. In fact, the assay on SKBR3 cells shows that **TPhP can interact with GPER and activate cAMP formation**, which is one of the rapid signalling pathways induced by GPER activation. However, this activation is 10-times less efficient than the reaction induced by E2. This activation *via* GPER is

⁷ DEGs: differentially expressed genes.

⁸ DEPs: differentially expressed proteins.

⁹ PI3K-Akt: phosphoinositide 3-kinase/protein kinase B.

¹⁰ MAPK: mitogen-activated protein kinase.

¹¹ EGFR: epidermal growth factor receptor.

also confirmed by the results obtained with a TPhP co-exposure with the specific GPER30 inhibitor, G15, which inhibits the effects of TPhP in the assay.

Conclusion: The available *in vitro* information demonstrate the capacity of TPhP to competitively bind and activate nuclear ER α and ER β estrogen receptors of several vertebrate species including human, rat, mouse, fish, chicken, frog and turtle as evidenced by ER transactivation in reporter cell lines, increased promoter occupancy, induction of ER-regulated gene expression and related physiological cell responses (e.g., increased cell proliferation). Two recent studies showed that TPhP could also bind and activate GPER.

Androgen pathway

TPhP binding capacity to the androgen receptor (AR) has been investigated in the study by Fang et al. (2003). The results indicate a moderate binding of TPhP to the AR.

Weak AR antagonist effects were observed in hamster CHO-K1 cells (Kojima et al., 2013; 2016).

Conclusions: The data available indicate that TPhP can moderately bind to the AR and could induce a weak anti androgenic effect in vertebrate cells.

Thyroid pathway

The expression of thyroid hormone-related gene expression was investigated in rat pituitary GH3 cells and in rat thyroid follicular FRTL-5 cells (Kim et al., 2015). In GH3 cells, TPhP exposure led to an up-regulation of *tra* from the medium tested dose (0.031 μ M) and of *tsh β* , *tr β* and *dio1* gene expression at the highest concentration tested (0.31 μ M). In FRTL-5 cells, TPhP exposure led to a down-regulation of *tg* and *tshr* expression at 3.1 μ M, an up-regulation of *nis* (9.2 μ M and 31 μ M), and *tpo* gene expression (31 μ M).

Conclusions: The results on GH3 cells and FRTL-5 cells show that TPhP can interact with thyroid hormone receptors in the rat pituitary or thyroid gland cells and could thus lead to stimulation of the synthesis of thyroid hormones.

Steroidogenesis

The steroidogenesis activity of TPhP was assessed in human adrenal cortico-carcinoma (H295R) cells (Liu et al., 2012; Ji et al., 2022). These two assays show that TPhP affects steroidogenesis by increasing estrogen levels (17 β -estradiol: E2) (see




Figure 3).




Figure 3: Proposed mechanism framework of estrogen disrupting effects for TPhP from the publication of Ji et al. (2022). Hormones in red indicates a significant increase of levels and blue indicates a significant decrease of levels (from Ji et al., 2022).

TPhP leads to an increase of E2 and E2/T ratio (Liu et al., 2012; Ji et al., 2022) and of progesterone (Ji et al., 2022) suggesting that **TPhP has also an estrogenic activity through an action on steroidogenesis**. Ji et al. (2022) shows that TPhP can promote cholesterol synthesis and its transfer through cellular membranes by a significant increase of HMGCR and StAR gene expression levels respectively. TPhP also induces an increase of the expression of the CYP genes (CYP11A1, CYP11B2, CYP19A1, CYP19) and 17β-HSD1 gene, which are linked to steroidogenesis. TPhP up-regulated CYP11A1 gene expression, which then can result in an increased conversion of cholesterol to pregnenolone. In addition, the up-regulated CYP17 gene expression may cause more androstenedione synthesis, and up-regulated 3β-HSD2 will impact the next step of the steroid synthesis. The authors hypothesised that the increased progesterone levels noted in the study may result from the up-regulation of CYP11A1 and 3β-HSD2 gene expression. An increased androstenedione synthesis due to CYP17 up-regulation gene expression will then be followed by an increased conversion of testosterone and estrone to E2 *via* the up-regulated CYP19 and 17β-HSD1 gene expression, which ultimately resulted in an increase of E2 concentrations (mechanisms confirmed in two other studies on 5 phthalates (DEP, MEP, BBzP, MBzP and DiBP) (Sohn et al., 2016) and on the trichloroethylene (Tachachartvanich et al., 2018). In Ji et al. (2022) a decrease of estrone is observed while CYP19 is upregulated and E2 level increases. The reason given by the authors is that the conversion of estrone to E2 by 17β-HSD1 is stronger than the activity of CYP19 to estrone synthesis. Lastly, inhibition of SULT2A1 expression observed in the two H295R assay may lead to more dehydroepiandrosterone (DHEA), participating in the production of E2 (Liu et al., 2012; Ji et al., 2022).

Conclusion: The two H295R assays performed with TPhP show that TPhP affects steroidogenesis by increasing estrogen levels (17β-estradiol) and increasing expression of CYP19 and 3β-HSD2 genes involved in this pathway.

Other endocrine pathways

TPhP is shown to activate the transcription of human, mouse, zebrafish or *Xenopus* PPARγ (peroxisome proliferator-activated receptor) (Garoche et al., 2021). Transcriptional activation of human, mouse and fish PPARγ is also demonstrated in a multiplexed reporter assay (Medvedev et al. 2020; Houck et al., 2021). Kojima et al. (2013; 2016) also report agonist activity of TPhP on human PPARγ (COS-7 cells).

In Kojima et al. (2013; 2016) the authors investigated interactions with PXR receptor from COS-7 cell-based reporter gene assay (to detect the agonist activity of chemicals against PXR). TPhP showed PXR agonist activity ($REC_{20} = 2.8 \mu\text{M}$).

In vivo mechanistic data with regard to an endocrine activity

Table 11: Summary table of *in vivo* mechanistic data (chronologic order)

STUDIES INVESTIGATING ENDOCRINE DISRUPTION PROPERTIES OF TPhP				
Methodology	Results (statistically significant change)		Remarks	Reference
Adult Zebrafish (<i>Danio rerio</i>) – 4 months old	<u>Female Plasma sex hormone</u> 17 β -estradiol (E2) \uparrow at 1 mg.l ⁻¹ Testosterone (T) No change 11-ketotestosterone (11-KT) No change		2 (reliable with restriction)	Liu et al., 2012
Nominal concentration: 0.04 – 0.2 – 1 mg.l ⁻¹	E2/T Ratio No change E2/11-KT Ratio \uparrow at 1 mg.l ⁻¹		Test material: TPhP	
No analytical measurement	<u>Male Plasma sex hormone</u> 17 β -estradiol (E2) \uparrow at 1 mg.l ⁻¹ Testosterone (T) \downarrow at 1 mg.l ⁻¹ 11-ketotestosterone (11-KT) \downarrow at 1 mg.l ⁻¹		Purity unknown	
Solvent < 0.01% DMSO (v/v)	E2/T Ratio \uparrow at 1 mg.l ⁻¹ E2/11-KT Ratio \uparrow at 1 mg.l ⁻¹		Non-guideline/exploratory study	
Exposure of male and female fish separately	<u>Female Transcriptional genes</u> <i>CYP17</i> ¹² \uparrow at 1 mg.l ⁻¹ <i>CYP19</i> \uparrow at 1 mg.l ⁻¹ <i>VTG1</i> ¹³ \downarrow at 1 mg.l ⁻¹		Not GLP	
Replicates: 3 with 3 fish for each replicate	<u>Male Transcriptional genes</u> <i>CYP17</i> \uparrow at 1 mg.l ⁻¹ <i>CYP19</i> \uparrow at 1 mg.l ⁻¹ <i>VTG1</i> \uparrow at 0.04; 0.2 and 1 mg.l ⁻¹			
Exposure duration: 14 days				
Photoperiod : 16h light – 8h dark				
Water temperature: 24°C \pm 2°C	<u>General toxicity</u> No significant mortality was observed (no details provided).			
Water pH: 7.2 to 7.8				
Dissolved oxygen: > 7 mg.l ⁻¹				
Exposure medium: renewal every other day				

¹² CYP17 and CYP19A were measured in gonad.

¹³ VTG 1 was measured in liver.

<p>Zebrafish embryos/larvae (<i>Danio rerio</i>) – 4hpf</p> <p>Nominal concentration: 0.02 - 0.2 - 2 mg.l⁻¹</p> <p>Measured concentration: 48 hpf: 0.02 – 0.19 – 1.80 mg.l⁻¹ 120 hpf: 0 – 0 – 1.39 mg.l⁻¹</p> <p>Solvent : 0.01% DMSO</p> <p>Replicates: Four separate wells with 20 eggs</p> <p>Exposure duration: up to 120 hpf</p> <p>Exposure medium: renewed at 48 h only.</p>	<p>Expression of mRNA for genes in six receptor (AhR, PPARα, ER1, TRα, G and MR) centered gene networks¹⁴</p> <p><i>AHRRB</i> ↑ at 0.02 mg.l⁻¹only <i>CYP1A1</i> ↑ at 2 mg.l⁻¹ <i>NCOR2</i> ↑ at 2 mg.l⁻¹ <i>CYP1B1</i> ↑ at 2 mg.l⁻¹ <i>PPARα</i> ↑ at 2 mg.l⁻¹ <i>PPARGC1A</i> ↑ at 2 mg.l⁻¹ <i>LPL</i> ↑ at 2 mg.l⁻¹ <i>IL6</i> ↑ at 2 mg.l⁻¹ <i>PPARγ</i> ↑ from 0.2 mg.l⁻¹ <i>ER2b</i> ↑ at 2 mg.l⁻¹ <i>TRα</i> ↑ at 2 mg.l⁻¹ <i>FUS</i> ↑ at 2 mg.l⁻¹ <i>GR</i> ↑ at 0.02 mg.l⁻¹ <i>RELA</i> ↑ at 2 mg.l⁻¹ <i>TGFβ1</i> ↑ at 2 mg.l⁻¹ <i>HSP90AA1</i> ↑ at 2 mg.l⁻¹ <i>11βHSD</i> ↑ from 0.2 mg.l⁻¹ <i>EGFR</i> ↑ at 2 mg.l⁻¹ <i>MR</i> ↓ at 2 mg.l⁻¹ <i>HPSE</i> ↓ at 2 mg.l⁻¹</p> <p><u>General toxicity</u> Rates of hatching and survival were >90% and rate of malformation (spinal curvature, pericardial edema and yolk sac edema) was <5% in all exposure groups including controls (no details provided).</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity unknown</p> <p>Non-guideline/exploratory study</p> <p>Not GLP</p>	<p>Liu et al., 2013a</p>
<p>Adult Zebrafish (<i>Danio rerio</i>) – 4/5 months old</p> <p>Nominal concentration: 0 – 0.04 – 0.2 – 1 mg.l⁻¹</p> <p>Measured concentration: 0h: 0.03 – 0.14 – 0.89 mg.l⁻¹ 48h: <LOD – <LOD – 0.38 mg.l⁻¹ (LOD = 0.12 µg.l⁻¹)</p> <p>Solvent: DMSO (0.005%)</p> <p>Replicates: 6 replicates with six pair of fish</p> <p>Exposure duration: 21 days</p>	<p><u>Female plasma sex hormone and VTG levels</u></p> <p>E2 level ↑ at 1 mg.l⁻¹ E2/11-KT ratio level ↑ at 1 mg.l⁻¹ E2/T ratio level ↑ at 0.2 and 1 mg.l⁻¹ VTG ↑ at 0.2 and 1 mg.l⁻¹ Testosterone level ↓ at 1 mg.l⁻¹ 11-KT ↓ at 1 mg.l⁻¹</p> <p><u>Male plasma sex hormone and VTG levels</u></p> <p>E2 level ↑ at 0.2 mg.l⁻¹ E2/11-KT level ↑ at 0.2 mg.l⁻¹ VTG level ↑ at 1 mg.l⁻¹ E2/T level ↓ at 0.04 and 0.2 mg.l⁻¹</p> <p><u>Transcriptional genes in ovaries</u></p> <p><i>LHR</i> ↑ at 1 mg.l⁻¹ <i>FSHR</i> ↑ at 1 mg.l⁻¹ <i>HMGRA</i> ↑ at 1 mg.l⁻¹ <i>STAR</i> ↑ at 0.2 mg.l⁻¹ <i>17βHSD</i> ↑ at 0.04 mg.l⁻¹ <i>CYP17A</i> ↑ at 1 mg.l⁻¹ <i>CYP19A</i> ↑ at 1 mg.l⁻¹ <i>HMGRB</i> ↓ at 1 mg.l⁻¹</p> <p><u>Transcriptional genes in testis</u></p> <p><i>CYP11A</i> ↑ at 0.2 mg.l⁻¹ <i>CYP17</i> ↑ at 1 mg.l⁻¹ <i>CYP19A</i> ↑ at 1 mg.l⁻¹ <i>LHR</i> ↓ at 0.04 mg.l⁻¹</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity: 99%</p> <p>Equivalent or similar to OECD TG 229</p> <p>Not GLP</p>	<p>Liu et al., 2013b</p>

¹⁴ Only upregulated or downregulated gene are reported in this table.

<p>Photoperiod : 14h light – 10h dark</p> <p>Water temperature: 27 °C ± 1 °C</p> <p>Exposure medium: renewed in every 48h.</p>	<p><i>HMGRA</i> ↓ at 1 mg.l⁻¹</p> <p><i>STAR</i> ↓ at 1 mg.l⁻¹</p> <p><i>17βHSD</i> ↓ at 0.2 mg.l⁻¹</p> <p><u>Female transcriptional genes in brain</u></p> <p><i>GnRH2</i> ↓ at 1 mg.l⁻¹</p> <p><i>GnRH3</i> ↓ at 0.2 mg.l⁻¹</p> <p><i>GnRHR3</i> ↓ at 0.2 mg.l⁻¹</p> <p><i>GnRHR4</i> ↑ at 1 mg.l⁻¹</p> <p><i>LHβ</i> ↑ at 0.2 mg.l⁻¹</p> <p><i>FSHβ</i> ↑ at 0.2 mg.l⁻¹</p> <p><i>CYP19B</i> ↑ at 0.2 mg.l⁻¹</p> <p><i>AR</i> ↑ at 0.2 mg.l⁻¹</p> <p><i>Era</i> ↑ at 1 mg.l⁻¹</p> <p><i>ER2β1</i> ↑ at 1 mg.l⁻¹</p> <p><u>Male transcriptional genes in brain</u></p> <p><i>GnRH2</i> ↑ at 0.04 mg.l⁻¹</p> <p><i>GnRH3</i> ↓ at 0.04 mg.l⁻¹</p> <p><i>GnRHR2</i> ↑ at 0.04 mg.l⁻¹</p> <p><i>GnRHR3</i> ↑ at 1 mg.l⁻¹</p> <p><i>FSHβ</i> ↑ at 1 mg.l⁻¹</p> <p><i>LHβ</i> ↓ at 0.04 mg.l⁻¹</p> <p><i>CYP19B</i> ↑ at 1 mg.l⁻¹</p> <p><i>AR</i> ↓ at 1 mg.l⁻¹</p> <p><i>Era</i> ↑ at 1 mg.l⁻¹</p> <p><i>ER2β1</i> ↑ at 0.2 mg.l⁻¹</p> <p><u>General toxicity</u></p> <p>No mortality was observed in any of the experimental concentrations during the exposure period (no details provided).</p>		
<p>Zebrafish embryos/larvae (<i>Danio rerio</i>) – 3hpf</p> <p>Nominal concentration: 0 – 40 – 200 – 500 µg.l⁻¹</p> <p>No analytical measurement</p> <p>Solvent: DMSO (0.005%)</p> <p>Replicates: 6 per treatment with 100 larvae per replicate</p> <p>Exposure duration: 7 days</p> <p>Photoperiod : 14h light – 10h dark</p> <p>Water temperature:</p>	<p><u>Hormone measurements in whole-body homogenate of 7 dpf Zebrafish larvae (100 larvae per replicate used)</u></p> <p>T3 level ↑ at 40, 200 and 500 µg.l⁻¹</p> <p>T4 level ↑ at 40, 200 and 500 µg.l⁻¹</p> <p><u>Gene expression in whole-body homogenate of 7 dpf Zebrafish larvae (20 larvae per replicate used)</u></p> <p><i>ttr</i> ↑ at 40, 200 and 500 µg.l⁻¹</p> <p><i>tra</i> ↑ at 200 µg.l⁻¹</p> <p><i>dio1</i> ↑ at 500 µg.l⁻¹</p> <p><i>nis</i> ↑ at 200 and 500 µg.l⁻¹</p> <p><i>tg</i> ↑ at 200 and 500 µg.l⁻¹</p> <p><i>ugt1ab</i> ↑ at 200 and 500 µg.l⁻¹</p> <p><i>crh</i> ↓ at 500 µg.l⁻¹</p> <p><i>trβ</i> ↓ at 500 µg.l⁻¹</p> <p><i>tshβ</i> No change</p> <p><i>tshr</i> No change</p> <p><i>nkx2.1</i> No change</p> <p><i>pax8</i> No change</p> <p><i>Dio2</i> No change</p> <p><u>General toxicity</u></p> <p>Larval survival at 7d: 91.8% (control) – 92.5% (40 µg.l⁻¹) – 92% (200 µg.l⁻¹) – 88.7% (500 µg.l⁻¹).</p> <p>Body weight were not affected by the experimental concentrations.</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity: 99%</p> <p>Non-guideline/exploratory study</p> <p>Not GLP</p>	<p>Kim et al., 2015</p>

<p>25°C ± 1°C</p> <p>Exposure medium: 50% renewed daily.</p>			
<p>Larvae, juvenile and adult Zebrafish (<i>Danio rerio</i>) – 4hpf</p> <p>Nominal concentration: 0 – 0.005 – 0.05 – 0.5 mg.l⁻¹</p> <p>Measured concentration: 24h: 0.0008 – 0.0055 – 0.131 mg.l⁻¹ 48h: < LOD - < LOD – 0.011 mg.l⁻¹ (LOD = 0.12 µg.l⁻¹)</p> <p>Solvent: DMSO (0.005%)</p> <p>Replicates: 3 per treatment with 100 embryos per replicate</p> <p>Exposure duration: 120 days</p> <p>Photoperiod : 14h light – 10h dark</p> <p>Water temperature: 27°C ± 1 °C</p> <p>Water pH: 7.2 to 7.8</p> <p>Dissolved oxygen: > 7 mg.l⁻¹</p> <p>Exposure medium: 50% renewed every 48h.</p>	<p><u>Female plasma sex hormone</u></p> <p>E2 level ↑ at 0.005 and 0.5 mg.l⁻¹</p> <p>Cortisol level ↑ at 0.5 mg.l⁻¹</p> <p>11-KT level ↓ at 0.5 mg.l⁻¹</p> <p>E2/11-KT ratio level ↑ at 0.5 mg.l⁻¹</p> <p>T3 level ↑ at 0.5 mg.l⁻¹</p> <p>T4 level ↑ at 0.5 mg.l⁻¹</p> <p><u>Male plasma sex hormone</u></p> <p>E2 level ↑ at 0.005 mg.l⁻¹</p> <p>Cortisol level ↓ from 0.05 mg.l⁻¹</p> <p>11-KT level ↓ from 0.005 mg.l⁻¹</p> <p>E2/11-KT level ↑ from 0.005 mg.l⁻¹</p> <p>T3 level No change</p> <p>T4 level No change</p> <p><u>Female transcriptional genes in brain</u></p> <p><i>FSHβ</i> ↑ at 0.05 and 0.5 mg.l⁻¹</p> <p><i>LHβ</i> ↑ at 0.005 and 0.5 mg.l⁻¹</p> <p><i>GnRH3</i> ↑ at 0.005 and 0.5 mg.l⁻¹</p> <p><i>Era</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>pomc</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>mr</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>trhr2</i> ↑ at 0.5 mg.l⁻¹</p> <p><u>Male transcriptional genes in brain</u></p> <p><i>LHβ</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>GnRH3</i> ↓ at 0.5 mg.l⁻¹</p> <p><i>pomc</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>trh</i> ↓ at 0.5 mg.l⁻¹</p> <p><i>FSHβ</i> No change</p> <p><i>Era</i> No change</p> <p><u>Transcriptional genes in ovaries</u></p> <p><i>LHR</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>STAR</i> ↑ at 0.05 and 0.5 mg.l⁻¹</p> <p><i>CYP19A</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>CYP17</i> No change</p> <p><i>FSHR</i> No change</p> <p><i>3β-HSD</i> No change</p> <p><i>17β-HSD</i> No change</p> <p><u>Transcriptional genes in testis</u></p> <p><i>STAR</i> ↓ at 0.5 mg.l⁻¹</p> <p><i>CYP17</i> ↓ at 0.5 mg.l⁻¹</p> <p><i>FSHR</i> ↑ at 0.005 mg.l⁻¹</p> <p><i>LHR</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>3β-HSD</i> ↑ at 0.5 mg.l⁻¹</p> <p><i>17β-HSD</i> ↑ at 0.050 mg.l⁻¹</p> <p><i>CYP19A</i> No change</p> <p><u>General toxicity</u></p> <p>None of the test concentrations caused any distinct malformation, and there were no significant differences</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity: 99%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	<p>Liu et al., 2016</p>

	in mortality among treatment groups during the exposure duration (no details provided).		
<p>Japanese medaka transgenic (<i>Oryzias latipes</i>) – o dph – Only male</p> <p>Nominal concentration: 1.6 – 8 – 40 µg.l⁻¹</p> <p>Measured concentration: 0.134 – 0.299 – 1.43 µg.l⁻¹</p> <p>Solvent: DMSO (0.001%)</p> <p>Replicates: 2 per treatment with 50 larvae per replicate</p> <p>Exposure duration: 100 days</p> <p>Photoperiod : 16h light – 8h dark</p> <p>Water temperature: 25°C ± 1 °C</p> <p>Water pH: 7.7 ± 0.2</p> <p>Dissolved oxygen: > 7.8 ± 0.3 mg.l⁻¹</p> <p>Number of tested fish: 16 males</p> <p>Exposure medium: flow-through system</p>	<p>17β-E2 level ↑ at 0.29 and 1.43 µg.l⁻¹</p> <p>T level ↓ at 1.43 µg.l⁻¹</p> <p>11-KT level ↓ at 1.43 µg.l⁻¹</p> <p><u>General toxicity</u> No data on systemic toxicity provided.</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity ≥ 95%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	<p>Li et al., 2018</p>
<p>Japanese medaka transgenic (<i>Oryzias latipes</i>) – only females</p> <p>Solvent: DMSO (0.001%)</p> <p>Photoperiod : 16h light – 8h dark</p> <p>Water temperature: 25°C ± 1 °C</p>	<p>Short-term exposure: 21 days</p> <p><u>Female plasma sex hormone</u></p> <p>17β-E2 level ↑ at 1725 ng.l⁻¹</p> <p>Testosterone level ↑ at 1725 ng.l⁻¹</p> <p><u>Female transcriptional genes in liver</u></p> <p>VTG2 ↓ at 1725 ng.l⁻¹</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity ≥ 95%</p>	<p>Li et al., 2019b</p>

<p>Water pH: 7.7 ±0.2</p> <p>Dissolved oxygen: > 7.8 ± 0.3 mg.l⁻¹</p> <p>Exposure medium: flow- through system</p> <p>Short-term exposure: adult exposed during 21 days</p> <p>Nominal concentration: 1.6 – 8 – 40 µg.l⁻¹</p> <p>Measured concentration: 243 – 589 – 1725ng.l⁻¹</p> <p>Replicates: 2 per treatment with 8 adult female per replicate</p> <p>Long-term exposure: larvae exposed during 100 days</p> <p>Nominal concentration: 1.6 – 8 – 40 µg.l⁻¹</p> <p>Measured concentration: 131 – 363 – 1773 ng.l⁻¹</p> <p>Replicates: 2 per treatment with 25 adult female per replicate</p>	<p>Long-term exposure: 100 days</p> <p><u>Female plasma sex hormone</u></p> <p>17β-E2 level ↓ at 1773 ng.l⁻¹ Testosterone level ↓ from 131 ng.l⁻¹ 11-KT level ↓ from 131 ng.l⁻¹</p> <p><u>Female transcriptional genes in liver</u></p> <p>VTG1 ↓ from 131 ng.l⁻¹ VTG2 ↓ from 363 ng.l⁻¹</p> <p><u>General toxicity</u> No data on systemic toxicity provided.</p>	<p>Non- guideline/e xploratory</p> <p>Not GLP</p>	
<p>Zebrafish adult (<i>Danio rerio</i>)</p> <p>Nominal Concentration: 0 – 0.04 – 0.2 – 1 mg.l⁻¹</p> <p>Measured concentration:</p>	<p><u>Female plasma sex hormone</u></p> <p>T4 level ↑ at 1 mg.l⁻¹ T3 level ↑ at 1 mg.l⁻¹</p> <p><u>Male plasma sex hormone</u></p> <p>T4 level ↓ at 1 mg.l⁻¹ T3 level ↓ at 1 mg.l⁻¹</p> <p><u>Female transcriptional genes in brain</u></p> <p><i>crh</i> ↓ at 1 mg.l⁻¹</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity 99%</p>	<p>Liu et al., 2019</p>

<p>0h: 0.03 – 0.15 – 0.87 mg.l⁻¹ 48h: < LOD - < LOD – 0.42 mg.l⁻¹ (LOD = 0.12 µg.l⁻¹) Solvent: DMSO (0.005%) Replicates: For males: 3 replicates with 5 males per replicate. For females: 3 replicates with 5 females per replicate. exposure duration: 14 days Photoperiod : 14h light – 10h dark Water pH: 7.2-7.8 Dissolved oxygen: > 7 mg.l⁻¹ Exposure medium: renewed at least three times per week</p>	<p><i>tshβ</i> ↓ at 1 mg.l⁻¹ <i>tra</i> ↑ at 0.2 and 1 mg.l⁻¹ <i>trβ</i> ↑ at 0.2 and 1 mg.l⁻¹</p> <p><u>Male transcriptional genes in brain</u> <i>crh</i> ↑ at 0.2 and 1 mg.l⁻¹ <i>tshβ</i> ↑ at 1 mg.l⁻¹ <i>trβ</i> ↑ at 0.2 and 1 mg.l⁻¹ <i>tra</i> No change</p> <p><u>Female transcriptional genes in thyroid</u> <i>trβ</i> ↓ at 1 mg.l⁻¹ <i>tra</i> ↑ at 1 mg.l⁻¹ <i>dio2</i> ↑ at 1 mg.l⁻¹ <i>tg</i> No change <i>tpo</i> No change <i>Dio1</i> No change</p> <p><u>Male transcriptional genes in thyroid</u> <i>trβ</i> ↓ at 0.2 and 1 mg.l⁻¹ <i>dio1</i> ↓ at 0.2 and 1 mg.l⁻¹ <i>dio2</i> ↓ at 0.2 and 1 mg.l⁻¹ <i>tra</i> No change <i>tg</i> No change <i>tpo</i> No change</p> <p><u>General toxicity</u> No mortality was observed over the exposure period (no details provided).</p>	<p>Non-guideline/exploratory study Not GLP</p>	
<p>Zebrafish adult (<i>Danio rerio</i>): 5-month adult <u>Sub-chronic toxicity (10% of LC₅₀)</u> Nominal concentration: 80 µg.l⁻¹ No analytical measurement TPhP Solvent: DMSO (0.01%) Replicates: 10 males and 10</p>	<p><u>Sex hormone and VTG levels in female whole-body homogenates</u> E2/T ratio level ↓ at 80 µg.l⁻¹ VTG level ↓ at 80 µg.l⁻¹ E2 No change T No change</p> <p><u>Sex hormone and VTG levels in male whole-body homogenates</u> E2/T ratio level No change VTG No change E2 No change T No change</p> <p><u>General toxicity</u> No data on systemic toxicity provided.</p>	<p>2 (reliable with restriction) Test material: TPhP Purity 99% Non-guideline/exploratory study Not GLP</p>	<p>He et al., 2021</p>

<p>females exposed separately</p> <p>Exposure duration: 21 days</p> <p>Water temperature: 25°C ± 1 °C</p> <p>Photoperiod : 14h light – 10h dark</p> <p>Water pH: 7.2 ± 0.2</p> <p>Dissolved oxygen: > 6.8 ± 0.2 mg.l⁻¹</p> <p>Exposure medium: renewed every day</p>			
<p>Japanese medaka (<i>Oryzias latipes</i>) – 16± 2 weeks old</p> <p>OECD 229</p> <p>Nominal concentration (µg.l⁻¹): 20 – 64 – 200 – 640</p> <p>Measured concentration (µg.l⁻¹): 2.13 – 7.19 – 17.1 – 44.9</p> <p>(LOQ = 0.03 µg.l⁻¹)</p> <p>Replicates: 4 replicates per treatment with 3 males and 3 females per replicate</p> <p>Exposure duration: 21 days</p> <p>Water temperature: 25°C ± 2 °C</p> <p>Photoperiod : 16h light – 8h dark</p>	<p><u>Female plasma VTG level</u></p> <p>VTG level ↓ at 7.19, 17.1 and 44.9 µg.l⁻¹</p> <p><u>General toxicity</u></p> <p>4.2% of mortality at 44.9 µg.l⁻¹ (1 female). No remarkable abnormal response (feeding activity, equilibrium, swimming...). No statistical analysis for length and weight parameters.</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity >97%</p> <p>According to OECD TG 229</p> <p>Not GLP</p>	<p>Kawashima et al., 2021</p>

<p>Water pH: 6.5-8.5</p> <p>Dissolved oxygen: > than 60% of air saturation</p> <p>Exposure medium: Flow-through (5 times/day)</p>													
<p>Zebrafish embryos (<i>Danio rerio</i>) – 72hpf</p> <p>Nominal concentrations: 0.32 – 0.97 – 3.2 – 9.8 mg.l⁻¹</p> <p>Solvent: DMSO (0.1% v/v)</p> <p>Replicates : 20 embryos per treatment</p> <p>Exposure duration : 24h</p>	<p><u>Transcriptional genes</u></p> <table border="0"> <tr> <td><i>tra</i></td> <td>↓ at 3.2 mg.l⁻¹</td> </tr> <tr> <td><i>ttr</i></td> <td>↓ at 3.2 mg.l⁻¹</td> </tr> <tr> <td><i>tshβ</i></td> <td>↓ at 9.8 mg.l⁻¹</td> </tr> <tr> <td><i>dio1</i></td> <td>↓ at 3.2 mg.l⁻¹</td> </tr> <tr> <td><i>dio2</i></td> <td>↑ at 3.2 mg.l⁻¹</td> </tr> </table> <p><u>General toxicity</u></p> <p>Significant concentration-dependent decrease in body length at 0.97, 3.2 and 9.8 mg.l⁻¹ (n=10 embryos per treatment). Mortality > 50% at 9.8 mg.l⁻¹. No significant mortality observed for other concentrations (no details provided).</p>	<i>tra</i>	↓ at 3.2 mg.l ⁻¹	<i>ttr</i>	↓ at 3.2 mg.l ⁻¹	<i>tshβ</i>	↓ at 9.8 mg.l ⁻¹	<i>dio1</i>	↓ at 3.2 mg.l ⁻¹	<i>dio2</i>	↑ at 3.2 mg.l ⁻¹	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 98%</p> <p>Non-guideline/exploratory study</p> <p>Not GLP</p>	<p>Lee et al., 2022</p>
<i>tra</i>	↓ at 3.2 mg.l ⁻¹												
<i>ttr</i>	↓ at 3.2 mg.l ⁻¹												
<i>tshβ</i>	↓ at 9.8 mg.l ⁻¹												
<i>dio1</i>	↓ at 3.2 mg.l ⁻¹												
<i>dio2</i>	↑ at 3.2 mg.l ⁻¹												
<p>OECD 234 (Fish sexual development test – FSDT)</p> <p>Zebrafish (<i>Danio rerio</i>)</p> <p>Nominal concentration: 1, 3.2, 10, 32, 100 µg.l⁻¹</p> <p>Measured concentration: 1.11 – 3.01 – 7.76 – 33.3 – 76.8 µg.l⁻¹ (TPhP in 1L acetone)</p> <p>Replicates: 4 replicates per concentration with 30 fertilised eggs per replicate.</p> <p>Water temperature: 27°C ± 2 °C</p>	<p><u>Female plasma sex hormone and VTG level*</u></p> <table border="0"> <tr> <td>VTG level</td> <td>↑ at 33.3 µg.l⁻¹</td> </tr> <tr> <td>17β-E2 level</td> <td>↑ at 3.01, 7.76, 33.3 and 76.8 µg.l⁻¹</td> </tr> </table> <p><u>Male plasma sex hormone and VTG level*</u></p> <table border="0"> <tr> <td>VTG level</td> <td>↑ at 33.3 µg.l⁻¹</td> </tr> <tr> <td>11-KT level</td> <td>↑ at 3.01 µg.l⁻¹</td> </tr> </table> <p><u>General toxicity</u></p> <p>Survival rates were 88% (control group) - 84% (1.11 µg.l⁻¹); 80% (3.01 µg.l⁻¹); 77% (7.76 µg.l⁻¹); 63% (33.3 µg.l⁻¹) - 51% (76.8 µg.l⁻¹)</p> <p>*Statistical unit: all individual measurements per concentration (not the mean of replicates per concentration in contrast to analysis provided in the study report).</p>	VTG level	↑ at 33.3 µg.l ⁻¹	17β-E2 level	↑ at 3.01, 7.76, 33.3 and 76.8 µg.l ⁻¹	VTG level	↑ at 33.3 µg.l ⁻¹	11-KT level	↑ at 3.01 µg.l ⁻¹	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 99%</p> <p>OECD TG 234</p> <p>GLP</p>	<p>Unpublished study report, 2022</p>		
VTG level	↑ at 33.3 µg.l ⁻¹												
17β-E2 level	↑ at 3.01, 7.76, 33.3 and 76.8 µg.l ⁻¹												
VTG level	↑ at 33.3 µg.l ⁻¹												
11-KT level	↑ at 3.01 µg.l ⁻¹												

<p>Photoperiod : 12h light – 12h dark</p> <p>Water pH: 7.6-7.8</p> <p>Dissolved oxygen: > than 60% of air saturation</p> <p>Exposure duration: 73 days</p> <p>Exposure medium: flow-through system</p>			
<p>Zebrafish embryo (<i>Danio rerio</i>) – (0-2 hpf)</p> <p>Nominal concentration: 0 – 20 – 50 – 100 – 500 and 1000 µg.l⁻¹</p> <p>No analytical measurement</p> <p>Solvent: DMSO (0.005% v/v)</p> <p>Water temperature: 28°C ± 1 °C</p> <p>Photoperiod : 14h light - 10h dark</p> <p>Replicates: 3 replicates per treatment with 40 embryos per replicate</p> <p>Exposure duration: 7 days</p> <p>Exposure medium: 90 % fresh exposure solution changed daily</p> <p>Measurement at 96 hpf and 7 dpf</p>	<p><u>Transcriptional genes in whole body (96 hpf)</u></p> <p><i>BCL2A</i> ↓ at 20 µg.l⁻¹</p> <p><i>CASP9</i> ↓ at 500 µg.l⁻¹</p> <p><i>MBPA</i> ↓ at 1000 µg.l⁻¹</p> <p><i>SYN2A</i> ↓ at 1000 µg.l⁻¹</p> <p><i>FOXO3A</i> ↓ at 1000 µg.l⁻¹</p> <p><i>PAX6A</i> ↓ at 500 and 1000 µg.l⁻¹</p> <p><i>ALDH1A2</i> ↑ at 50 µg.l⁻¹</p> <p><i>RDH1</i> ↓ at 1000 µg.l⁻¹</p> <p><i>CRABP1A</i> ↓ at 1000 µg.l⁻¹</p> <p><i>RBP2A</i> ↓ at 1000 µg.l⁻¹</p> <p><i>RARAA</i> ↓ at 1000 µg.l⁻¹</p> <p><i>CYP26A1</i> ↓ at 500 and 1000 µg.l⁻¹</p> <p><i>RBP5</i> ↓ at 100, 500 and 1000 µg.l⁻¹</p> <p><u>Transcriptional genes in whole body (7 dpf)</u></p> <p><i>ALDH1A2</i> ↑ at 100 and 500 µg.l⁻¹</p> <p><i>MBPA</i> ↑ at 20 and 50 µg.l⁻¹</p> <p><i>RDH1</i> ↑ at 500 µg.l⁻¹</p> <p><i>RBP2A</i> ↑ at 20 µg.l⁻¹</p> <p><u>General toxicity</u></p> <p>From 120 hpf, the survival rate of zebrafish larvae was significantly decreased at 500 µg.l⁻¹ and 1000 µg.l⁻¹. Survival rate: 30% at 500 µg.l⁻¹ and 70% at 1000 µg.l⁻¹ 7 dpf.</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 98%</p> <p>Non-guideline/exploratory study</p> <p>Not GLP</p>	<p>Zhang et al., 2023</p>

Thyroid pathway

Clear sex-dependent alterations in plasma TH levels were observed in zebrafish with an increase of the plasmatic concentration of T3 and T4 in females (Liu et al., 2016; Liu et al., 2019) and a decrease or no change of the plasmatic concentration of T3 and T4 in males (Liu et al., 2016; 2019). Kim et al. (2015) also reported an increase of the whole body homogenate concentration of T3 and T4 in zebrafish embryos/larvae.

EAS modalities

The *in vivo* mechanistic effects described in the literature indicate that TPhP has EAS endocrine activity.

In Liu et al. (2012), the transcription of steroidogenic genes (CYP17 and CYP19A) was measured in adult Zebrafish gonads. Exposure of male and female to 1 mg.l⁻¹ of TPhP significantly increased CYP17 and CYP19A gene transcription. In Liu et al. (2013b), exposure to TPhP led to significant transcriptional changes in both fish testes and ovaries. In testes, transcription levels of HMGRA, StAR, and 17 β -HSD were significantly down-regulated, while transcriptions of CYP11A, CYP17, and CYP19A were significantly up-regulated. In ovaries, the major steroidogenic genes, including HMGRA, StAR, 17 β -HSD, CYP17A, and CYP19A, were significantly up-regulated by exposure to TPhP.

Several studies demonstrated a significant increase of plasmatic concentrations of E2 and E2/11-KT ratio in both sexes of zebrafish at the top dose (Liu et al., 2012; 2013b; 2016). An increase of E2 level was also noted in female zebrafish (Unpublished study report, 2022), male medaka (Li et al., 2018) and female medaka (Li et al., 2019b). Nevertheless, in one study (Li et al., 2019b) carried out on Japanese medaka, a decrease of E2 was noted after 100-day exposure of female larvae (age of fish at exposure initiation), while an increase of E2 was observed after 21-days exposure of adult females (age of fish at exposure initiation). Finally in one study (He et al., 2021), no change in E2 level in male or female whole body homogenates was noted.

Concerning testosterone, an increase in the plasma level could be observed after a short term exposure in female medaka plasma, while a decrease was noted after a long-term exposure (Li et al., 2019b). In He et al. (2021), no change was recorded in female and male fish homogenates. A decrease of testosterone was observed in male medaka and zebrafish (Li et al., 2018; Liu et al., 2012) and in female zebrafish (Liu et al., 2013b). Decreased levels of the hormone 11-ketotestosterone (11-KT) were recorded in female fish after exposure to TPhP (Liu et al., 2013b; Liu et al., 2016; Li et al., 2018). The variation in 11-KT for males also followed a decreasing trend (Liu et al., 2012; Liu et al., 2016; Liu et al., 2018), except in the unpublished study report (2022), where the 11-KT increased in male fish. Moreover, E2/11-KT ratio increased in females and males (Liu et al., 2012; Liu et al., 2016; Li et al., 2018; Liu et al., 2013b). This trend could not be verified in the unpublished study report (2022), as E2 levels were not measured in males and consequently the E2/11KT ratio could not be calculated.

In the same way, an activity on the E2/T ratio was noted, but the direction of variation was not established, with an increase of E2/T reported in some studies for males (Liu et al., 2012) or females (Liu et al., 2013b), and a decrease or no change reported in others for females (Liu et al., 2012) or males (Liu et al., 2013b) or both (He et al., 2021).

The observations of VTG concentrations also suggest an estrogenic activity of TPhP in female and male zebrafish, with increased plasmatic concentrations of VTG reported in Liu et al. (2013b) and the Unpublished study report (2022). Contradictory results are also reported with decreased or no changed concentrations in plasma VTG of female medaka (Kawashima et al., 2021) or in whole-body homogenate of male and female zebrafish (He et al., 2021). VTG1 gene expression in liver is increased in male zebrafish in one assay (Liu et al., 2012), while the expression in liver of VTG1 and 2 are decreased in other assays (Liu et al., 2012, for females; Li et al., 2019b).

Conclusion: Transcription levels of key genes involved in the steroidogenesis was significantly upregulated by exposure to TPhP in several studies. Significant alteration of plasmatic concentrations of E2, and E2/T ratio (more specific observation in females) and E2/11-KT ratio (more specific observation in males) could result from this modification in the steroidogenesis pathway. These different perturbations of circulating steroid concentrations could depend on the fish developmental stage, species and tested concentrations. The observations of VTG concentrations, that were consistent with perturbation of E2 concentrations, suggested an EAS activity of TPhP in female and male zebrafish, with altered plasmatic concentration of VTG.

Other endocrine pathways

In Zhang et al. (2023), the authors investigated the effects of TPhP exposure on transcriptional levels in zebrafish at different developmental stages namely 96 hpf and 7 dpf. The expression level of different genes were affected in zebrafish and 96 hpf larvae were shown to be more sensitive than 7 dpf larvae. The expression of apoptotic-related genes (*bcl2a* and *Casp9*) was down-regulated in 96 hpf zebrafish at the concentration of 20 $\mu\text{g.l}^{-1}$ and at 1000 $\mu\text{g.l}^{-1}$ respectively. The expression of neurodevelopmental-related genes (*pax6a* and *mbpa*, *syn2a*, *foxo3a*) was also down-regulated from 500 $\mu\text{g.l}^{-1}$ and at 1000 $\mu\text{g.l}^{-1}$ respectively.

Regarding the retinoic acid (RA) metabolic-related gene expression, at 96 hpf, the expression of *aldh1a2* was significantly up-regulated in 50 $\mu\text{g.l}^{-1}$ TPhP. Exposure to 1000 $\mu\text{g/L}$ TPhP significantly downregulated the expressions of *rdh1*, *crabp1a*, *rbp2a* and *raraa*. Furthermore, the expression of *cyp26a1* was significantly decreased in the 500 and 1000 $\mu\text{g.l}^{-1}$ TPhP treated groups. The expression of *rbp5* was also decreased in 100, 500 and 1000 $\mu\text{g.l}^{-1}$ TPhP treated groups. Several genes in the RA metabolic pathway were also examined in zebrafish larvae at 7 dpf. The expression of *aldh1a2* was significantly up-regulated in 100 and 500 $\mu\text{g.l}^{-1}$ TPhP treated groups. And exposure to 500 $\mu\text{g.l}^{-1}$ TPhP significantly increased the expressions of *rdh1*. Exposure to 20, 50, 100 and 500 $\mu\text{g.l}^{-1}$ TPhP significantly up-regulated the transcriptions of *rbp2a*.

Although not extensively detailed here, it is also important to note that TPhP was previously shown to interact with the retinoic acid receptor (RAR). TPhP exhibits weak RAR α antagonist activity (Jia et al., 2022; Isales et al., 2015), which could lead to ocular and cardiovascular malformations. The RAR signaling pathway is essential for reproduction and embryonic development. Indeed, genetic studies in zebrafish embryos that are deficient in RA-generating enzymes or RARs revealed that retinoic acid (RA) signalling regulates development of many organs and tissues, including the body axis, spinal cord, forelimb buds, skeleton, heart, eye, pancreas, lung and spermatogenesis (Ghyselinck et al. 2019; Clagett-Dame and DeLuca, 2002). There are many additional functions for RA that are supported by *in vivo* genetic loss-of-function studies in zebrafish. Further studies are needed to identify the key genes regulated by RA signalling. The retinoid signaling pathway is an endocrine pathway considered to be susceptible to environmental endocrine disruption (OECD, 2021) and possibly related to endocrine effects on development.

In vivo adverse effect data with regard to an endocrine mode of action**Table 12: Summary table of *in vivo* adverse effect data (chronologic order)**

STUDIES INVESTIGATING ENDOCRINE DISRUPTION PROPERTIES OF TPhP				
Methodology	Results (statistically significant change)		Remarks	Reference
<p>Adult Zebrafish (<i>Danio rerio</i>) – 4/5 months old</p> <p>Nominal concentration: 0 – 0.04 – 0.2 – 1 mg.l⁻¹ No analytical measurement</p> <p>Measured concentration: 0h: 0.03 – 0.14 – 0.89 mg.l⁻¹ 48h: <LOD – <LOD – 0.38 mg.l⁻¹ (LOD = 0.12 µg.l⁻¹)</p> <p>Solvent: DMSO (0.005%)</p> <p>Replicates: 6 replicates with six pair of fish</p> <p>Exposure duration: 21 days</p> <p>Photoperiod : 14h light - 10h dark</p> <p>Water temperature: 27 °C ± 1 °C</p> <p>Exposure medium: renewed in every 48h.</p>	<p><u>Female Zebrafish</u></p> <p>Cumulative Egg number ↓ at 0.2 and 1 mg.l⁻¹ Spawning event ↓ at 0.2 and 1 mg.l⁻¹ Hatchability ↓ at 0.2 and 1 mg.l⁻¹ Fertilisation success No change</p> <p>Hepato-somatic index (HSI) No change Gonado-somatic index (GSI) No change</p> <p><u>General toxicity</u> No mortality was observed in any of the experimental concentrations during the exposure period (no details provided).</p>		<p>2 (reliable with restriction)</p> <p>test material: TPhP</p> <p>Purity: 99%</p> <p>Equivalent or similar to OECD TG 229</p> <p>Not GLP</p>	<p>Liu et al., 2013b</p>
<p>Zebrafish embryos/larvae (<i>Danio rerio</i>) – 3hpf</p> <p>Nominal concentration:</p>	<p>Malformation rate ↑ at 500 µg.l⁻¹</p> <p><u>General toxicity</u></p>		<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity: 99%</p>	<p>Kim et al., 2015</p>

<p>0 – 40 – 200 – 500 µg.l⁻¹ No analytical measurement</p> <p>Solvent: DMSO (0.005%)</p> <p>Replicates: 6 per treatment with 100 larvae per replicate</p> <p>Exposure duration: 7 days</p> <p>Photoperiod : 14h light - 10h dark</p> <p>Water temperature: 25°C ± 1°C</p> <p>Exposure medium: 50% renewed daily.</p>	<p>Larval survival at 7d: 91.8% (control) – 92.5% (40 µg.l⁻¹) – 92% (200 µg.l⁻¹) – 88.7% (500 µg.l⁻¹). Body weight were not affected by the experimental concentrations.</p>	<p>Non-guideline/exploratory</p> <p>Not GLP</p>													
<p>Larvae, juvenile and adult Zebrafish (<i>Danio rerio</i>)</p> <p>Nominal concentration: 0 – 0.005 – 0.05 – 0.5 mg.l⁻¹</p> <p>Measured concentration: First 24h: 0.0008 - 0.0055 - 0.131 mg.l⁻¹ First 48h: < LOD - < LOD - 0.011 mg.l⁻¹</p> <p>Solvent: DMSO (0.005%)</p> <p>Replicates: 3 per treatment with 100 embryos per replicate</p> <p>Exposure duration: 120 days</p>	<p><u>Female Zebrafish</u></p> <table border="0"> <tr> <td>Gonadosomatic index (GSI)</td> <td>↓ at 0.005 and 0.5 mg.l⁻¹</td> </tr> <tr> <td>Hepato-somatic index (HSI)</td> <td>No change</td> </tr> <tr> <td>Condition factor (CF)</td> <td>No change</td> </tr> </table> <p><u>Male Zebrafish</u></p> <table border="0"> <tr> <td>Gonadosomatic index (GSI)</td> <td>No change</td> </tr> <tr> <td>Hepato-somatic index (HSI)</td> <td>No change</td> </tr> <tr> <td>Condition factor (CF)</td> <td>↓ at 0.5 mg.l⁻¹</td> </tr> </table> <p><u>General toxicity</u></p> <p>None of the test concentrations caused any distinct malformation, and there were no significant differences in mortality among treatment groups during the exposure duration (no details provided).</p>	Gonadosomatic index (GSI)	↓ at 0.005 and 0.5 mg.l ⁻¹	Hepato-somatic index (HSI)	No change	Condition factor (CF)	No change	Gonadosomatic index (GSI)	No change	Hepato-somatic index (HSI)	No change	Condition factor (CF)	↓ at 0.5 mg.l ⁻¹	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity: 99%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	<p>Liu et al., 2016</p>
Gonadosomatic index (GSI)	↓ at 0.005 and 0.5 mg.l ⁻¹														
Hepato-somatic index (HSI)	No change														
Condition factor (CF)	No change														
Gonadosomatic index (GSI)	No change														
Hepato-somatic index (HSI)	No change														
Condition factor (CF)	↓ at 0.5 mg.l ⁻¹														

<p>Photoperiod : 14h light - 10h dark</p> <p>Water temperature: 27°C ± 1 °C</p> <p>Water pH: 7.2 to 7.8</p> <p>Dissolved oxygen: > 7 mg.l⁻¹</p> <p>Exposure medium: 50% renewed every 48h.</p>			
<p>Japanese medaka transgenic (<i>Oryzias latipes</i>) - < 4hpf</p> <p>Nominal concentration: 5, 25, 125, 625 µg.l⁻¹</p> <p>No analytical measurement</p> <p>Solvent: DMSO (0.01%)</p> <p>Replicates: 3 replicates per concentration</p> <p>Water temperature: 25°C ± 1 °C</p> <p>Photoperiod : 14h light - 10h dark</p> <p>Exposure duration: 14 d</p> <p>Exposure medium: renewed daily (at least 90%)</p>	<p>Hatchability (%) at 96h ↓ at 625 µg.l⁻¹ Time of hatching at 96h ↑ at 625 µg.l⁻¹</p> <p><u>General toxicity (14 d)</u> Heart rates ↓ at 125 and 625 µg.l⁻¹ Body length ↓ at 125 and 625 µg.l⁻¹ Gross abnormality ↑ at 625 µg.l⁻¹ Mortality No information given</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 99%</p> <p>Non- guideline/exploratory</p> <p>Not GLP</p>	<p>Sun et al., 2016</p>
<p>Japanese medaka transgenic (<i>Oryzias latipes</i>) - 0 dph - Only male</p>	<p><u>Male Medaka:</u></p> <p>Gonadal intersex incidence ↑ at 1.43 µg.l⁻¹</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p>	<p>Li et al., 2018</p>

<p>Nominal concentration: 1.6 – 8 – 40 µg.l⁻¹</p> <p>Measured concentration: 0.134 – 0.299 – 1.43 µg.l⁻¹</p> <p>Solvent: DMSO (0.001%)</p> <p>Replicates: 2 per treatment with 50 larvae per replicate</p> <p>Exposure duration: 100 days</p> <p>Photoperiod : 16h light - 8h dark</p> <p>Water temperature: 25°C ± 1 °C</p> <p>Water pH: 7.7 ± 0.2</p> <p>Dissolved oxygen: > 7.8 ± 0.3 mg.l⁻¹</p> <p>Number of tested fish: 16 males</p> <p>Exposure medium: flow-through system</p>	<p>Abnormal chasing behaviours ↑ at 0.29 and 1.43 µg.l⁻¹</p> <p>Courtship behaviour ↑ at 0.29 µg.l⁻¹</p> <p>Chasing time ↑ at 0.29 µg.l⁻¹</p> <p>Successful mating rate ↓ at 1.43 µg.l⁻¹</p> <p><u>General toxicity</u> No data on systemic toxicity provided.</p>	<p>Purity ≥ 95%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	
<p>Japanese medaka transgenic (<i>Oryzias latipes</i>) – only females</p> <p>Solvent: DMSO (0.001%)</p> <p>Photoperiod : 16h light - 8h dark</p> <p>Water temperature: 25°C ± 1 °C</p>	<p><u>Female medaka:</u></p> <p>Long-term exposure: 100 days</p> <p>Number of mature oocyte ↓ at 0.13, 0.36 and 1.77 µg.l⁻¹</p> <p>Egg production ↓ at 0.36 and 1.77 µg.l⁻¹</p> <p><u>General toxicity</u> No data on systemic toxicity provided</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity ≥ 95%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	<p>Li et al., 2019b</p>

<p>Water pH: 7.7 ±0.2</p> <p>Dissolved oxygen: > 7.8 ± 0.3 mg.l⁻¹</p> <p>Exposure medium: flow-through system</p> <p><u>Long-term exposure:</u> larvae exposed during 100 days</p> <p>Nominal concentration: 1.6 – 8 – 40 µg.l⁻¹</p> <p>Measured concentration: 131 – 363 – 1773 ng.l⁻¹</p> <p>Replicates: 2 per treatment with 25 adult female per replicate</p>															
<p>Japanese medaka (<i>Oryzias latipes</i>) – 16± 2 weeks old</p> <p>OECD 229</p> <p>Nominal concentration (µg.l⁻¹): 20 – 64 – 200 – 640</p> <p>Measured concentration (µg.l⁻¹): 2.13 – 7.19 – 17.1 – 44.9</p> <p>Replicates: 4 replicates per treatment with 3 males and 3 females per replicate</p>	<table border="0"> <tr> <td>Secondary sex characteristics</td> <td>No change</td> </tr> <tr> <td>Egg number (female)</td> <td>↓ at 44.9 µg.l⁻¹</td> </tr> <tr> <td>Fertilised egg number</td> <td></td> </tr> <tr> <td>Fertility rate</td> <td>↓ at 44.9 µg.l⁻¹</td> </tr> <tr> <td>HSI (male)</td> <td>No change</td> </tr> <tr> <td></td> <td>↑ at 7.19, 17.1, 44.9 µg.l⁻¹</td> </tr> </table> <p><u>General toxicity</u> 4.2% of mortality at 44.9 µg.l⁻¹ (1 female). No remarkable abnormal response (feeding activity, equilibrium, swimming...). No statistical analysis for length and weight parameters</p>	Secondary sex characteristics	No change	Egg number (female)	↓ at 44.9 µg.l ⁻¹	Fertilised egg number		Fertility rate	↓ at 44.9 µg.l ⁻¹	HSI (male)	No change		↑ at 7.19, 17.1, 44.9 µg.l ⁻¹	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity >97%</p> <p>According to OECD TG 229</p> <p>Not GLP</p>	<p>Kawashima et al., 2021</p>
Secondary sex characteristics	No change														
Egg number (female)	↓ at 44.9 µg.l ⁻¹														
Fertilised egg number															
Fertility rate	↓ at 44.9 µg.l ⁻¹														
HSI (male)	No change														
	↑ at 7.19, 17.1, 44.9 µg.l ⁻¹														

<p>Exposure duration: 21 days</p> <p>Water temperature: 25°C ± 2 °C</p> <p>Photoperiod : 16h light - 8h dark</p> <p>Water pH: 6.5-8.5</p> <p>Dissolved oxygen: > than 60% of air saturation</p> <p>Exposure medium: Flow-through (5 times/day)</p>			
<p>Zebrafish adult (<i>Danio rerio</i>): 5-month adult</p> <p><u>Sub-chronic toxicity (10% of LC₅₀)</u></p> <p>Nominal concentration: 80 µg.l⁻¹</p> <p>No analytical measurement</p> <p>Solvent: DMSO (0.01%)</p> <p>Replicates: 10 males and 10 females exposed separately</p> <p>Exposure duration: 21 days</p> <p>Water temperature: 25°C ± 1 °C</p>	<p><u>Female Zebrafish:</u></p> <p>Hepato-somatic index (HSI) ↑ at 80 µg.l⁻¹</p> <p>Gonado-somatic index (GSI) ↑ at 80 µg.l⁻¹</p> <p>Ovary maturation stages ↓ at 80 µg.l⁻¹ (oocytes less mature with more oocytes at the first two stages (perinuclear and corticolar alveolar oocytes). Structure of oocytes more irregular. Inhibition of the ovarian development.</p> <p><u>Male Zebrafish:</u></p> <p>Hepato-somatic index (HSI) ↑ at 80 µg.l⁻¹</p> <p>Gonado-somatic index (GSI) ↓ at 80 µg.l⁻¹</p> <p>Spermatocyte maturation stages ↓ at 80 µg.l⁻¹ (more immature spermatocytes (spermatogonia and spermatocyte) and less mature spermatocytes (spermatids and spermatozoa).</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity 99%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	<p>He et al., 2021</p>

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

<p>Photoperiod : 14h light - 10h dark</p> <p>Water pH: 7.2 ± 0.2</p> <p>Dissolved oxygen: > 6.8 ± 0.2 mg.l⁻¹</p> <p>Exposure medium: renewed every day TPhP</p>	<p>Fecundity/fertility</p> <p>↓ at 80 µg.l⁻¹ of fertilisation rate, cumulative eggs, spawning, hatching rate.</p> <p><u>General toxicity</u> No data on systemic toxicity provided.</p>		
<p>Amphibian tadpoles (<i>Rana zhenhaiensis</i>)</p> <p>Solvent: DMSO (0.01%)</p> <p>Nominal concentration: 0.02 and 0.1 mg.l⁻¹</p> <p>No analytical measurement</p> <p>Exposure duration: 30 days</p> <p>Replicates: 6 replicate aquaria, 60 tadpoles in each treatment group</p> <p>Stage GS26 (Gosner stage)</p> <p>Water temperature: 25°C ± 1°C</p> <p>Photoperiod : 12h light - 12h dark</p>	<p>Developmental stages</p> <p>↑ at 0.1 mg.l⁻¹ (more advanced developmental stage)</p> <p><u>General toxicity</u> No death occurred in any treatment groups throughout the exposure duration.</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 99%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p> <p>Non-standard species</p>	<p>Lu et al., 2021</p>
<p>Amphibians tadpoles (<i>Hoplobatrachus rugulosus</i>)</p>	<p>Survival (30d)</p> <p>Metamorphosis rates (30d)</p> <p>↓ at 50 µg.l⁻¹</p> <p>↓ at 50 µg.l⁻¹</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p>	<p>Chen et al., 2022</p>

<p>Stage GS35 (Gosner stage)</p> <p>Solvent: DMSO (0.5 ‰ v/v)</p> <p>Nominal concentration: 0- solvent control - 10 - 50 - 100 - 200 µg.l⁻¹</p> <p>Exposure duration: 30 days</p> <p>Replicates: 3 replicates per treatment with 30 tadpoles per replicate</p> <p>Water temperature: 29°C ± 2 °C</p> <p>Photoperiod : 14h light - 10h dark</p> <p>Water pH: 7.2 ± 0.2</p> <p>Dissolved oxygen: > 5.78 ± 0.3 ppm</p> <p>Exposure medium: renewed daily</p>	<p>Metamorphosis time ↓ at 100 µg.l⁻¹ (30d)</p> <p><u>General toxicity</u> Survival rate: 87.8% (control group) – 87.8% (solvent control group) – 76.7% (10 µg.l⁻¹) – 66.7% (50 µg.l⁻¹) – 63.3% (100 µg.l⁻¹) – 61.1% (200 µg.l⁻¹)</p>	<p>Purity > 99%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	
<p>OECD 234 (Fish sexual development test – FSdT)</p> <p>Zebrafish (<i>Danio rerio</i>)</p> <p>Nominal concentration: 1, 3.2, 10, 32, 100 µg.l⁻¹</p> <p>Measured concentration: 1.11 – 3.01 – 7.76 – 33.3 – 76.8 µg.l⁻¹ (TPhP in 1L acetone)</p>	<p><u>Sex ratio</u> No statistically significant change % of females: 64.8% (control) – 52.6% (1.11 µg.l⁻¹) – 55.8% (3.01 µg.l⁻¹); 59.1% (7.76 µg.l⁻¹) – 67.5% (33.3 µg.l⁻¹) – 65.4% (76.8 µg.l⁻¹) Note: high proportion of females in the control group (64.8%)</p> <p><u>Male Zebrafish:</u></p> <p>Testis maturation stages ↑ at 76.8 µg.l⁻¹</p> <p><u>General toxicity</u> Survival rates were 88% (control group) - 84% (1.11 µg.l⁻¹); 80% (3.01 µg.l⁻¹); 77% (7.76 µg.l⁻¹); 63% (33.3 µg.l⁻¹) - 51% (76.8 µg.l⁻¹)</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 99%</p> <p>OECD TG 234</p> <p>GLP</p>	<p>Unpublished study report, 2022</p>

<p>Replicates: 4 replicates per concentration with 30 fertilised eggs per replicate.</p> <p>Water temperature: 27°C ± 2 °C</p> <p>Photoperiod : 12h light - 12h dark</p> <p>Water pH: 7.6-7.8</p> <p>Dissolved oxygen: > than 60% of air saturation</p> <p>Exposure duration: 73 days</p> <p>Exposure medium: flow-through system</p>			
<p>Zebrafish embryos/larvae (<i>Danio rerio</i>) – 2hpf</p> <p>Nominal concentration: 8, 24, 72, 144 µg.l⁻¹</p> <p>No analytical measurement</p> <p>Replicates: 3 replicates per concentration</p> <p>Water temperature: 28°C ± 0.5 °C</p> <p>Photoperiod : 14h light - 10h dark</p> <p>Exposure duration: 120 h</p>	<p>Hatching rates ↓ at 72 and 144 µg.l⁻¹</p> <p><u>General toxicity</u></p> <p>Heart rates ↓ at 72 and 144 µg.l⁻¹</p> <p>Malformation rates ↑ at 72 and 144 µg.l⁻¹</p> <p>Survival rates ↓ at 144 µg.l⁻¹</p> <p>Body length ↓ at 144 µg.l⁻¹</p> <p><u>Survival rates</u></p> <p>86.5% (control group) – 86% (8 µg.l⁻¹) - 84% (24 µg.l⁻¹) – 82% (72 µg.l⁻¹) – 80.5% (144 µg.l⁻¹)</p>	<p>2 (reliable with restriction)</p> <p>Test material: TPhP</p> <p>Purity > 99%</p> <p>Non-guideline/exploratory</p> <p>Not GLP</p>	<p>Fan et al., 2022</p>

Exposure medium: renewed daily			
Zebrafish embryo (<i>Danio rerio</i>) – (0-2 hpf) Nominal concentration: 0 – 20 – 50 – 100 – 500 and 1000 µg.l ⁻¹ No analytical measurement Solvent: DMSO (0.005% v/v) Water temperature: 28°C ± 1 °C Photoperiod : 14h light - 10h dark Replicates: 3 replicates per treatment with 40 embryos per replicate Exposure duration: 7 days	Survival rate ↓ at 1000 µg.l ⁻¹ (6 dpf) ↓ at 500 µg.l ⁻¹ (7 dpf) Hatching rate ↓ at 50 µg.l ⁻¹ at 48 hpf ↓ at 500 µg.l ⁻¹ at 72 hpf Heart rate ↑ at 100 and 1000 µg.l ⁻¹ at 48 hpf ↓ at 500 µg.l ⁻¹ at 96 hpf Behaviour changes Swimming distance ↑ at 20; 100; 500 and 1000 µg.l ⁻¹ Movement trajectory ↑ at 20; 100; 500 and 1000 µg.l ⁻¹ Number of apoptotic cells in the head, heart and intestine ↑ with the increase of TPhP concentration <u>General toxicity</u> From 120 hpf, the survival rate of zebrafish larvae was significantly decreased at 500 µg.l ⁻¹ and 1000 µg.l ⁻¹ .	2 (reliable with restriction) Test material: TPhP Purity > 98% Non-guideline/exploratory Not GLP	Zhang et al., 2023

Thyroid pathway

Lu et al. (2021) studied exposure of amphibian tadpoles (Zhenhai brown frog, *Rana zhenhaiensis*) throughout the developmental period to assess physiological and metabolic impacts of TPhP on amphibian larvae. The exposure started with tadpoles at the Gosner stage (GS) 26. A total of 180 tadpoles were randomly selected and allocated to different aquaria, each containing the solvent control (DMSO), 0.02 and 0.1 mg TPhP.l⁻¹ (6 replicate aquaria, 60 tadpoles in each treated group). After 30-days exposure the tadpoles were euthanised. Snout-vent length (SVL), tail length (TL), body weight (BW) and developmental stage were recorded. No death occurred in any treated groups throughout the exposure duration. Tadpoles from 0.1 mg.l⁻¹-treated group had more advanced developmental stages than those from the other two groups. However, there were no significant differences in other traits measured in this study.

Chen et al. (2022) studied the sub-chronic toxicity of TPhP to amphibian tadpoles (*Hoplobatrachus rugulosus*). GS35 tadpoles were divided in four TPhP-treated groups (three replicates/group, 30 tadpoles/replicate/30 L): 10, 50, 100, and 200 µg.l⁻¹. After two weeks of continuous TPhP exposure, the dechlorinated tap water culture was switched until the tadpoles completed metamorphosis (GS46). The endpoints were survival, metamorphosis, and

development. All tadpoles were observed every 24 h until they underwent complete metamorphosis. The survival and metamorphosis rates were determined within 30 days of sub-chronic toxicity test, and the metamorphosis time was defined as the interval between the beginning of TPhP exposure and complete metamorphosis. When the tadpoles completed metamorphosis, their snout–vent lengths (SVLs) and body masses (BMs) were measured. The threshold concentration that affected survival and metamorphosis rate was 50 µg/L, while that affecting the metamorphosis time was 100 µg.l⁻¹. No significant differences were observed in the body size and condition of the tadpole after metamorphosis; however, BM and TPhP concentration were negatively correlated.

Conclusion: although indications of effects on thyroid exist, available information are scarce and possible effects of TPhP on the thyroid pathway were not further considered in the context of the SVHC identification.

EAS modalities

Based on the available data (Liu et al., 2013b; Li et al., 2019b; Kawashima et al., 2021; He et al., 2021; Zhang et al., 2023), it was considered that TPhP impairs fertility and fecundity in fish.

In Liu et al. (2013b) the effects of TPhP on reproductive endpoints were studied in adult zebrafish (*Danio rerio*) over 21 days. The study was performed with 12 fish per concentration, exposed to 0, 0.04, 0.2 and 1 mg.l⁻¹ nominal TPhP concentrations based on the results of a range-finding test. No mortality occurred at any concentration. There was a statistically significant decrease at 0.2 and 1 mg.l⁻¹ of the cumulative egg numbers, the number of spawning events and the hatchability.

Liu et al. (2016) studied the effects of TPhP on zebrafish (*Danio rerio*) during long-term exposure covering larval, juvenile and adult life stages. Embryos were collected with 4-h post fertilisation and exposed to 0, 0.005, 0.05 and 0.5 mg.l⁻¹ nominal concentrations of TPhP for 120 days post-fertilisation (dpf). There were 3 replicates per treatment, and each replicate contained 100 embryos and 300 ml of test solution. Half of the exposure solution was renewed every 48 h throughout exposure. Mortality, condition factor (CF), gonadosomatic index (GSI) and hepatosomatic index (HSI) were assessed at 120 dpf. No distinct malformations were observed at any concentration, and the authors concluded that there were no significant differences in mortality between treated groups (although the level of mortality observed, including controls, was not specified). The CF and GSI were unaffected up to 0.5 mg.l⁻¹ for the larval (14 dpf) and juvenile (40 dpf) fish. In adult fish (n= 5 per sex), CF (i.e., growth) in male fish was significantly affected at 0.5 mg.l⁻¹. The gonadosomatic index (GSI) in female zebrafish exposed for 120 days to TPhP was significantly decreased at 0.005 and 0.5 mg/L with no change in the hepatosomatic index, suggesting a negative impact on female gonad development.

In Sun et al. (2016), larvae (Japanese medaka) collected in the first hours post-hatching (< 4 hpf) were exposed to TPhP at nominal concentrations of 5, 25, 125, 625 µg.l⁻¹ (no analytical measurement during the test was reported) until 14 d. The media was renewed every 24 h (at least 90%). Embryos were observed 3 times daily, and the dead larvae were removed. At 60 h, their heart rates were monitored. At the end of the exposure duration, the body lengths of the larval medaka were measured. Exposure to TPhP significantly decreased hatchability, delayed time to hatch, and increased the occurrence of gross abnormalities at the highest exposure concentration (625 µg.l⁻¹) compared with controls. Moreover, embryo heart rate and body length were also decreased relatively to the controls in the TPhP exposure groups with concentrations over 125 µg.l⁻¹.

In Li et al. (2018), larvae (Japanese medaka, male only) collected on the first day post-hatching (0 dph) were exposed to TPhP at nominal concentrations of 1.6, 8, and 40 µg.l⁻¹ (0.134, 0.299 and 1.43 µg.l⁻¹ measured respectively) until 100 dph. A flow-through system with a 2-fold volume of water flowing through every 24 h was used. After exposure, 16 males were randomly selected and mated to non-exposed females in clean water for reproductive behaviours and

success tests prior to observation of gonadal intersex. A significant gonadal intersex incidence of 26% in male medaka was observed at $1.43 \mu\text{g.l}^{-1}$ (indicating a potentially feminisation risk for male fish). In this study chasing trajectory and time, courtship frequency and mating number were recorded to evaluate the effects of TPhP on reproductive behaviours. After exposure, the incidences of male fish with abnormal chasing behaviour indicated a dose-dependent increase. Additionally, significant decreases in the fertilisation rate after $1.43 \mu\text{g.l}^{-1}$ TPhP exposure and in the hatching rate after 0.299 and $1.43 \mu\text{g.l}^{-1}$ TPhP exposure were observed in the study.

Li et al. (2019b) studied the effects of TPhP on the reproduction of Japanese medaka by an exposure of 100 days following hatching (FSDT-like test). Medaka (50 individuals, duplicate experiments) were exposed to TPhP in a continuous flow system at 1.6, 8 and $40 \mu\text{g.l}^{-1}$ (measured concentrations 1.3 , 3.6 and $17.7 \mu\text{g.l}^{-1}$). The reproduction test was performed with 6 exposed females paired with control males. Oocyte maturation was quantified by a double measurement of oocyte size and *osp1* gene expression by fluorescence. After 100 days of exposure, oocytes underwent a delay in maturation characterised by a significant dose-dependent decrease in the number of mature and pre-vitellogenic oocytes with more than one third of females affected. The highest doses induced more substantial effects involving the absence of oocyte II (stage 3). Cumulative egg production over three days decreased in a dose-dependent manner and was significantly affected at the two highest doses (with 39 and 51% decrease respectively). It should be noted that the concentration of TPhP was higher in the liver and the ovary than in muscle and brain.

In Kawashima et al. (2021), Japanese medaka (*Oryzias latipes*) were exposed to TPhP in a flow-through system for 21 days with 4 replicates per treatment. The age of fish at exposure initiation was 14 weeks old. Six fish (3 males and 3 females) were used per treatment. The endpoints measured were mortality, growth (length and weight), fecundity (number of eggs, number of fertile eggs, fertility rate), GSI, HSI and secondary sexual characteristics. To prevent excessive lethal effects, the highest test concentration was determined based on acute toxicity on Japanese medaka. The nominal concentrations were $0 - 20 - 64 - 200 - 640 \mu\text{g.l}^{-1}$ (measured concentration were respectively $0 - 2.13 - 7.19 - 17.1 - 44.9 \mu\text{g.l}^{-1}$). Regarding the mortality, only one female fish died at the highest concentration ($44.9 \mu\text{g.l}^{-1}$), therefore it is considered that the validity criteria was met. The secondary sex characteristics (SSC) were not changed after exposure to TPhP. A statistically significant decrease in number of total and fertilised eggs was reported at $44.9 \mu\text{g.l}^{-1}$. However, no significant effect on fertility rate (the ratio of the number of fertilised eggs to the number of total eggs) was observed.

He et al. (2021) studied acute toxicity with some endocrine indices and effects on reproduction of TPhP on adult zebrafish. The study was conducted with a single dose corresponding to less than 10% of the LC50 (based on a range-finding study), *i.e.*, $80 \mu\text{g.l}^{-1}$. The test included groups of 10 separate male or female fish for 21-day exposure. Exposure to TPhP induced a statistically significant increase in the hepatosomatic index (HSI) by a factor of 1.8 and 2.2 for males and females, respectively, as well as a decrease in the gonadosomatic index (GSI) in males and an increase in females (not quantified). These factors were accompanied by histological changes. The testes and ovaries showed an increase in the early stages of sex cells, and conversely, a decrease in the more developed stages in both sexes indicating an inhibition of gametogenesis (based on qualitative histological observations). Moreover, TPhP significantly reduced the egg production, the number of spawning, the fertilisation rate and the hatching rate. In conclusion, the study indicated that TPhP ($80 \mu\text{g.l}^{-1}$) altered gametogenesis in both sexes of adult zebrafish resulting in decreased fecundity.

An unpublished study report (2022) aimed to assess the potential endocrine activity and adverse effects of continuous exposure to TPhP for 73 days *i.e.*, on the early life stages and sexual differentiation of zebrafish (OECD 234). Thirty fertilised eggs were used for each test with 4 replicates per concentration. Endpoints were determined including hatching success and rates, mortalities during early life stage and juvenile growth, and maturation phase. Sex ratio was determined macroscopically and by histological examination of the gonads. Measurements were taken at the end of the study. Additionally, a histopathological examination was performed on

the gonads and liver tissues. A maximum tolerated concentration (MTC) of $88.3 \mu\text{g.l}^{-1}$ (where no effect on hatching success, but reduced fish numbers were observed) has been defined by a range-finding study. In the main study, the exposure doses showed a significant deviation (*i.e.*, 80% or 120%) from the nominal concentrations, and the authors reported the results in mean measured concentrations of 0; 1.11; 3.01; 7.76; 33.3 and $76.8 \mu\text{g.l}^{-1}$. According to the OECD 234 guidelines, all the validity criteria were met.

The survival rates observed were 88%; 84%; 80.0%; 77%; 63% and 51% (control group; 1.11; 3.01; 7.76; 33.3 and $76.8 \mu\text{g.l}^{-1}$ respectively). The low survival rate observed at the two highest doses in the main study was not expected at this level of dose in light of the range finding results. Mortality was mainly observed during the early life stage of zebrafish until 35 days post-fertilisation (dpf), with only few cases of mortality between 35 dpf and test termination (73 dpf). It should be noted that according to the OECD 234 guideline, if exposure-related mortality occurs, the number of replicates should be reduced appropriately so that fish density between treatment levels is kept as equal as possible, as fish density is extremely important for growth and development. Unfortunately, this reduction of replicate was not performed at the two highest concentrations where mortality occurred. During the early life stage, reduced growth in terms of total length was observed at the highest tested concentrations. This reduced size was likely due to the general systemic toxicity of TPhP. However, at 73 dpf, a subsequent exposure to TPhP led to the increase in mass of the exposed fish in a concentration-dependent manner with a statistically significant difference for the highest dose in males and the two highest concentrations in females. This mass increase was likely due to reduction of fish density in treatment conditions, resulting from the reduced post-hatch survival rates. The increase in mass should have been avoided by the reduction of the number of replicates, which was not performed. The variability of the environmental conditions among the groups could impair the ability of the test to show an adverse effect in terms of population development or sex-ratio.

Regarding the apical effect of the OECD 234, namely sex ratio, no significant difference was observed, but it should be noted that there was a high proportion of females in the control group (64.8%) compared to exposure groups; even if the acceptance criteria related to proportion of sex in control group at termination of the test (30% males – 70% females) was fulfilled. The percentage of females for all the exposed groups increased from 52.6 to 65.4% and the percentage of males decreased from 46.3 to 33.3% in relation to increasing doses. The number of undifferentiated intersex was not significantly different (there was only one undifferentiated fish at the 1.11 and $76.8 \mu\text{g.l}^{-1}$ concentrations). It is worth noting that the survival rate might have altered the statistical power of the study to detect a significant effect on sex ratio in these groups. Given the high mortality rate at high doses and the low percentage of males in those samples, a deficit of males was observed for these high doses, which might have altered the representativeness and therefore the interpretation of the measurements performed, in particular in males at these doses. It was also not possible to determine whether the mortality affected one sex more specifically (no sex indication was reported for the dead fish), which may have interfered with the assessment of sex ratio.

The histopathological analysis of fish gonad revealed that with increasing concentrations, the gonads reached a mature stage more rapidly (for males and females but statistically significant at the highest concentration for males only). The proportion of stage 2 (the most advanced stage found) increased from 55 to 90% for testes. Stage 4 in females increased from 15 (control) to 30% (highest dose). The acceleration in gonadal maturation is consistent with the elevated circulating steroid levels.

Concerning the histology of female gonads analysed at 3.33 and $76.8 \mu\text{g.l}^{-1}$ and in the control group, a dose-dependent increase in all ovarian pathologies (oocyte atresia, egg debris, granulomatous inflammation) was observed but without statistical significance in any group. These pathologies are indicative of the alteration (acceleration) of the oocyte maturation process and oocyte quality, although not statistically significant.

Regarding males, only the testis-ova was measured and was slightly decreased (not statistically significant) with combined acceleration of maturation of the gonad. However, the percentage of testis-ova males in the control was high (11%), which makes difficult the interpretation of the other doses. Moreover, although testis-ova will likely lead to a decrease in fertility or hatching success, the design of the test (FSDT, *i.e.*, OECD 234) precludes this type of examination since FSDT *per se* does not include the production of a next generation.

In addition to gonad histopathological analysis, liver and heart were examined. Liver from all test groups was analysed while for the heart, only two groups were examined (control and 76.8 µg.l⁻¹). No statistically significant effect on the severity of the hepatic lesions was observed. Nevertheless, liver histopathological analysis revealed a dose-dependent decrease in hepatocyte lipid inclusions in females. In males, a dose-dependent increase in bile duct proliferation and inflammatory foci was observed. The analysis of the liver may reveal a toxic effect or a more specific effect of lipid metabolism seen in females. Cardiac lesions were also observed in treated TPhP groups but no statistics could be performed due to the limited number of replicates and test concentrations.

In Fan et al. (2022) the authors explored the effects of TPhP alone and/or combined to nano-titanium dioxide (nano-TiO₂) on the neurodevelopment of zebrafish larvae. Zebrafish embryos (within 2 h after fertilisation) were exposed to concentrations of 0, 8, 24, 72 and 144 µg.l⁻¹ (no analytical measurement) of TPhP alone for 120h. Zebrafish embryos/larvae were kept in an incubator at 28 ± 0.5 °C, with a light-dark ratio of 14 h/10 h. During the period of exposure, the solutions were renewed daily to keep the concentrations constant. No developmental toxicity was detected by treating with TPhP at 8 and 24 µg.l⁻¹, while higher concentrations of TPhP (72 µg.l⁻¹ and 144 µg.l⁻¹) significantly altered the hatching rate, the heart rate, the malformation rate, and the body length in zebrafish larvae (only at 144 µg.l⁻¹). Note that at the highest dose the inhibitions on the hatching rate, heartbeat and body length in zebrafish larvae were significantly promoted in the co-exposure groups where TiO₂ was added, compared to TPhP alone.

In Zhang et al. (2023) the authors investigated the effects of TPhP exposure on developmental parameters, locomotor behaviour, oxidative stress, apoptosis and transcriptional levels in zebrafish at different developmental stages. Zebrafish embryos (0 – 2 hpf) were exposed to concentrations of 0, 20, 50, 100, 500 and 1000 µg.l⁻¹ of TPhP for 7 days (based on a range finding study with a 96h LC₅₀ value of TPhP on zebrafish embryos of 5.13 mg.l⁻¹). The results showed that the survival rate decreased significantly compared to the control at the concentration of 500 µg/L from 120 hpf. In the same way, the hatching rate decreased at 50µg/L at 48 hpf and 500 µg/L at 72 hpf compared to the control. TPhP also significantly increased the heart rate of zebrafish larvae at 48 hpf (100 and 1000 µg.l⁻¹) and at 60 hpf (500 and 100 µg.l⁻¹) but significantly decreased at 96 hpf (500 µg.l⁻¹). Microscopic observations also showed that the exposure of TPhP led to several malformation symptoms such as yolk sac edema, spinal curvature, abnormal body pigmentation, pericardial edema, small eyes, etc. Concerning the behaviour changes, TPhP could affect the embryonic spontaneous movement. Regarding the locomotor behaviour and the neurotoxicity, all TPhP treated groups caused statistically significant increase in swimming distance and movement trajectory of zebrafish larvae except for the 50 µg.l⁻¹ groups. The authors also found that TPhP increased the number of apoptotic cells in the head, heart and intestine of zebrafish. In the head region, the number of apoptotic cells were localised in the eyeball of zebrafish larvae.

Conclusion: *in vivo* studies on fish reproduction show an alteration of gametogenesis in both sexes with a disruption of testis and ovary maturation followed by a decrease in reproductive success in terms of fecundity (decrease in egg production, spawning events) and fertility (impaired hatchability and fertilisation rate). *In vivo* studies on amphibian development show that TPhP could affect tadpole survival and metamorphosis.

Analysis of systemic toxicity in relation to the observation of EAS-related adverse effects

As recommended in the EDC guidance (ECHA/EFSA, 2018), adverse effects that are non-specific secondary consequences of other toxic effects shall not be considered for the identification of the substance as endocrine disruptor. The top concentration selected for the conduction of the ecotoxicological studies should provide information on substance toxicity at an exposure of the tested agent that should be tolerated without inducing significant chronic physiological dysfunctions, be compatible with animal survival and permits data interpretation in the context of the use of the study. It is recommended that a Maximum Tolerated Concentration (MTC) should be considered for the evaluation of changes which could be due to excessive systemic toxicity. In ecotoxicology, the MTC is defined as the highest test concentration of the chemical which results in less than 10% mortality.

For each study, information on mortality when available has been reported in the Table above. Data do not systematically allow to determine a MTC value.

Japanese medaka (*Oryzias latipes*)

Four studies (Sun et al., 2016 - Li et al., 2018 - Li et al., 2019b - Kawashima et al., 2021) of interest are available. The only study to provide information on the survival rate of fish after exposure of TPHP is Kawashima et al., 2021.

Table 13: Summary table of survival data - Japanese medaka (*Oryzias latipes*)

Ref.	Age at study start	Duration of exposure	Media renewal	Nominal conc. ($\mu\text{g.l}^{-1}$)	Measured conc. ($\mu\text{g.l}^{-1}$)	Survival rate
Kawashima et al. 2021	16±2 weeks old	21d	Flow-through (5 times/day)	Control group	Control group	0%
				20	2.13	0%
				64	7.19	0%
				200	17.1	0%
				640	44.9	95.8% (1 female)

It is therefore not possible to determine a lowest observed effect concentration (LOEC) on mortality from this study. However, MTC was not exceeded. It is therefore not possible to determine a systemic toxicity threshold from survival rate for Japanese medaka (*Oryzias latipes*) from these studies.

Zebrafish (*Danio rerio*)

Eleven studies (Liu et al., 2012 - Liu et al., 2013a - Liu et al., 2013b - Kim et al., 2015 - Liu et al., 2016 - Liu et al., 2019- He et al., 2021 - Lee et al., 2022 - Unpublished study report, 2022 - Fan et al., 2022 - Zhang et al., 2023) of interest are available and information on mortality is summarised below.

Table 14: Summary table of survival data - Zebrafish (*Danio rerio*)

	Age at study start	Duration of exposure	Media renewal	Nominal conc. (mg.l ⁻¹)	Measured conc. (mg.l ⁻¹)	Survival rate
Liu et al., 2012	4 months	14d	Every other day	Control group		No significant mortality (no value provided)
				0.04	NAM	
				0.2		
				1		
No value for survival rate. MTC was not exceeded						
Liu et al., 2013b	4-5 months	21d	Every 48h	Control group		No mortality (no value provided)
				0.04	<LOD (48h)	
				0.2	<LOD (48h)	
				1	0.38 (48h)	
No mortality at 0.38 mg.l ⁻¹ . MTC was not exceeded						
Liu et al., 2019	4-5 months	14d	at least three times per week	Control group		No mortality (no value provided)
				0.04	0.03 (0h) <LOD (48h)	
				0.2	0.15 (0h) <LOD (48h)	
				1	0.87 (0h) 0.42 (48h)	
No mortality at 0.42 mg.l ⁻¹ . MTC was not exceeded						
He et al., 2021	Adult	21d	Every day	0.08	NAM	No information
No information. Impossible to determine a systemic toxicity threshold from survival rate						
Liu et al., 2013a	4hpf	120hpf	at 48h only	Control group		
				0.02	0 (120hpf)	> 90%
				0.2	0 (120hpf)	> 90%
				2	1.39 (120hpf)	> 90%
Mortality < 10% at 1.39 mg.l ⁻¹ . MTC was not exceeded.						
Kim et al., 2015	3hpf	7d	50% daily	Control group		91.8%
				0.04	NAM	92.5%
				0.2		92%
				0.5		88.7%
Mortality > 10% at 0.5 mg.l ⁻¹ but not > 10% compared to mortality in the control group. MTC is not considered to have been exceeded.						
Liu et al., 2016	4hpf	120d	50% every 48h	Control group		No significant mortality (no value provided)
				0.005	0.0008 (24h) < LOD (48h)	
				0.05	0.0055 (24h) < LOD (48h)	
				0.5	0.131 (24h)	

					0.011 (48h)	
Impossible to determine a correct range of exposure concentrations. No value for survival rate. Impossible to determine a systemic toxicity threshold from survival rate. MTC was not exceeded.						
Lee et al., 2022	72hpf	24h	No information	Control group		No significant mortality (no value provided) < 50%
				0.32	NAM	
				0.97		
				3.2		
9.8						
Low survival rate at 9.8 mg.l ⁻¹ . No value for survival rate. MTC may have been exceeded.						
Unpublished study report, 2022	<12hpf	60d	Flow-through system	Control group		88%
				0.001	0.001	84%
				0.0032	0.003	80%
				0.01	0.008	77%
				0.032	0.033	63%
0.1	0.077	51%				
Mortality in the control group > 10%. Mortality > 10% compared to mortality in the control group above 0.008 mg/l. Observation of mortality in controls demonstrate that TPhP exposure is not the only driver for mortality.						
Fan et al., 2022	2hpf	120h	daily	Control group		87%
				0.008	NAM	86%
				0.024		84%
				0.072		82%
				0.144		81%
Mortality in the control group > 10% but not > 10% compared to mortality in the control group. Observation of mortality in controls demonstrate that TPhP exposure is not the only driver for mortality. MTC is not considered to have been exceeded.						
Zhang et al. 2023	2hpf	7d	90% daily	Control group		>80%
				0.02	NAM	90%
				0.05	NAM	90%
				0.1	NAM	90%
				0.5	NAM	30%
1	NAM	20%				
No exact value for the survival rate (observations from a graph). Mortality in the control group > 10%. Mortality > 10% compared to mortality in the control group above 0.5 mg/l. Observation of mortality in controls demonstrate that TPhP exposure is not the only driver for mortality.						

NAM: no analytical measurement

Data do not systematically allow to determine a MTC value. It is noted that due to limitations in the studies (absence of measured concentrations) and differences in experimental design (stage and duration of exposure), it is impossible to determine a systemic toxicity threshold from survival rate for Zebrafish (*Danio rerio*).

Overall, mortality and its possible interference with EAS-adverse effects shall be interpreted in each study independently.

Conclusion regarding ED properties relevant for environment

Adverse effects relevant for ED identification

The available data indicate that exposure to TPhP damages fish reproduction by impacting processes and events during gametogenesis (oocyte growth, ovary and testis maturation), disrupting fish fecundity (eggs production and spawning) and fertility (fertilisation and hatching). Changes in fecundity and fertility are apical effects, which inform about potential adverse effects at fish population level.

The decrease in mature spermatocytes (He et al., 2021) suggests a potential delay in spermatogenesis. A significant gonadal intersex incidence of 26% in male medaka was observed at $1.43 \mu\text{g}\cdot\text{l}^{-1}$ (Li et al., 2018), indicating a potential feminisation risk for male fish. A disruption (dose-dependent increase) of male-typical reproductive behaviours before spawning (chasing trajectory and time, courtship frequency and mating number) observed by Li et al. (2018) also supports the reproduction impairment of male fish after exposure to TPhP. This effect can explain the observed decrease in fertilisation rate.

Data show an inhibition of the ovarian development in female fish (He et al., 2021) and alterations of the ovaries (Li et al., 2019b; He et al., 2021). These alterations and development retardation can affect the fecundity of female fish with the observed decrease in egg production and spawning.

Despite the observation of some effects in the FSDT study, it should be noted that the quality of this study did not allow to evaluate properly EAS mediated parameters.

The decrease in egg production (Liu et al., 2013b; Li et al., 2019b, He et al., 2021; Kawashima et al., 2021) is a clear effect indicator of the impaired reproductive ability, reinforced by a decrease of spawning events (Liu et al., 2013b; He et al., 2021), an insufficient fertilisation rate (Li et al., 2018; He et al., 2021) and a decrease in hatchability (Liu et al., 2013b; Li et al., 2018; He et al., 2021), disrupting the generation of new individuals. This impact on fish reproduction could affect fish population stability and is considered as an adverse effect relevant at population level.

Moreover, it is important to note that several adverse effects on altered reproductive capacity of fish are observed in several studies at environmentally relevant levels without concurrent systemic effects. Indeed, no specific mention of mortality was reported in several studies (He et al., 2021; Li et al., 2018; Li et al., 2019b) although the absence of specific information on this parameter leaves some uncertainties. Most importantly, effects on hatching rate, fecundity and fertility were clearly observed without any concurrent mortality in Liu et al. (2013b) and Kawashima et al. (2021).

TPhP shows adverse effects on fertility and fecundity of fish at environmentally relevant levels. The impact on fish reproduction could affect fish population stability and is considered as an adverse effect relevant at population level. These effects cannot be considered as a secondary non-specific consequence of general toxicity.

Data (*in vivo*) providing scientific evidence of an adverse effect of TPhP on the reproductive system are summarised in Table 13 below.

Table 15: Line of evidence in relation to TPhP sexual dysfunction in *in-vivo* studies

Endpoint	Biological model	Species	Exposure duration	Adult or juvenile (when exposed)	Lowest significant concentration	Observed effects	Reference
Hatching rate	-	Zebrafish	21 days	Adult	0.2 mg.l ⁻¹	↓	Liu et al., 2013b
Hatching rate	-	Zebrafish	21 days	Adult	80 µg.l ⁻¹	↓ [#]	He et al., 2021
Hatching rate	-	Japanese Medaka	100 days	Juvenile	0.299 µg.l ⁻¹	↓ [#]	Li et al., 2018
Sex ratio	-	Japanese Medaka	100 days	Juvenile	1.43 µg.l ⁻¹	↑ [#] intersex incidence in male fish	Li et al., 2018
Sex ratio	-	Zebrafish	73 days	Juvenile	-	##No change	Unpublished study report, 2022
Reproductive behaviour	-	Japanese Medaka	100 days	Juvenile	1.43 µg.l ⁻¹	[#] Abnormal chasing behaviour in male fish	Li et al., 2018
Fecundity	-	Zebrafish	21 days	Adult	0.2 mg.l ⁻¹	↓ egg production, spawning event	Liu et al., 2013b
Fecundity	-	Japanese Medaka	100 days	Female juvenile	0.36 µg.l ⁻¹	[#] ↓ egg production	Li et al., 2019b
Fecundity/fertility	-	Zebrafish	21 days	Adult	-	Fertility rate: no change	Liu et al., 2013b
Fecundity/fertility	-	Japanese Medaka	100 days	Juvenile	1.43 µg.l ⁻¹	↓ [#] fertilisation rate	Li et al., 2018
Fecundity/fertility	-	Zebrafish	21 days	Adult	80 µg.l ⁻¹	↓ [#] fertilisation rate, cumulative eggs, spawning event	He et al., 2021
Fecundity/fertility	-	Japanese Medaka	21 days	Adult	44.9 µg.l ⁻¹	↓ egg production and fertilised eggs*	Kawashima et al., 2021
Histology	Testes	Zebrafish	73 days	Male juvenile	76.8 µg.l ⁻¹	↑ ^{##} testis maturation stages	Unpublished study report, 2022
Histology	Ovaries	Japanese Medaka	100 days	Female juvenile	0.13 µg.l ⁻¹	↓ [#] number of mature oocytes	Li et al., 2019b
Histology	Ovaries	Zebrafish	21 days	Female adult	80 µg.l ⁻¹	↓ [#] ovary maturation stages	He et al., 2021
Histology	Testes	Zebrafish	21 days	Male adult	80 µg.l ⁻¹	↓ [#] testis maturation stages	He et al., 2021

*: however no significant effect on fertility rate (ratio)

No information on mortality

Mortality > 10% observed, with mortality > 10% also in the control group

MTC not exceeded for the other studies

Endocrine activity

In vitro and *in vivo* studies are available supporting an **estrogenic activity** of TPhP, as summarised in Table 14 below.

Molecular Initiating events (MIEs)

- Nuclear estrogen receptors

The estrogenic activity is mainly mediated *via* nuclear estrogen receptors (nERs), which function as transcription factors upon ligand binding for genomic responses. In most vertebrates, including mammals, birds, and some lower vertebrates, two estrogen receptors exist: the nuclear intracellular receptors ER α and ER β , considered to mediate the effect of E2 and VTG in liver of most fish species. Data show that TPhP has an agonist activity on human ER α and ER β receptors and decrease the binding affinity of ER (Liu et al., 2012; Kojima et al., 2013; Zhang et al., 2014; Kojima et al., 2016; Medvedev et al., 2020; Ji et al., 2022). The agonist activity is also reported for the medaka ER1 (Kawashima et al., 2021) and for the fish ER2 β , frog ER1, turtle ER1 and chicken ER1 (Medvedev et al., 2020).

In some yeast two-hybrid assays with human and medaka ER α , an antagonist effect is shown indicating a competition between TPhP and E2 for binding to ER α (Ji et al., 2020; Li et al., 2018). Zhang et al. (2014) demonstrated a binding affinity of TPhP to hER α with a docking approach. Moreover, TPhP can induce the ERE pathway demonstrated by a dose-response of the agonist effect in MVLN cells (transfected with the human ERE-luciferase reporter gene) (Ji et al., 2020). This estrogenic activity is supported by proliferation assays on MCF-7 cells that show TPhP promotes cell proliferation (Zhang et al., 2014; Krivoshev et al., 2016; Ji et al., 2020).

- GPER (membrane estrogen receptor)

The estrogenic activity can also be mediated *via* membrane estrogen receptors as G protein-coupled membrane receptors (GPER). The role of GPER in oocyte maturation is well defined in zebrafish (Thomas, 2017; Chen et al., 2019).

In Guan et al. (2022), the authors report that TPhP binds GPER with an IC₅₀ of 4.2 μ M. This is confirmed by E2-F displacement experiments. Combined transcriptome and proteome analysis show that TPhP can exert its estrogen-like proliferation-promoting effect through the interaction of the PI3K-Akt signalling pathway, MAPK signalling pathway, and EGFR signalling pathway, which are downstream pathways mediated by GPER. In Ji et al. (2022), the assay on SBRK3 cells shows that TPhP can interact with GPER and activate cAMP formation, which is one of the rapid signalling pathways induced by GPER activation. However, this activation is 10-times less efficient than the reaction induced by E2. Nevertheless, this activation *via* GPER is also confirmed by the results obtained with a TPhP co-exposure with the specific GPER30 inhibitor, G15, which inhibits the effects of TPhP in the assay.

- Aromatase activity

Aromatase (CYP19) catalyses the conversion of androgenic steroids to estrogenic steroids (Conley and Hinshelwood, 2001). The CYP19 plays a role in signaling pathways critical for sexual development and fertility. TPhP induced an increase of the expression of the CYP19 gene (Liu et al., 2012; Liu et al., 2013b; Liu et al. 2016; Ji et al., 2022) and 17 β -HSD1 gene (Liu et al., 2013b; Liu et al., 2016; Ji et al., 2022), which is linked to the steroidogenic pathway. In Ji et al. (2022), a decrease of estrone was observed while CYP19 is upregulated and E2 level increased. The reason given by the authors was that the conversion activity of estrone to E2 by 17 β -HSD1 was stronger than the activity of CYP19 to estrone synthesis. TPhP affected steroidogenesis by increasing the estrogen levels (17 β -estradiol) *via* the up-regulation of CYP19.

Steroid hormone level and Vitellogenin

During medium-term exposure, TPhP induces an increase in circulating E2 concentrations in zebrafish and medaka (Liu et al., 2012; Liu et al., 2013b; Liu et al., 2016, Li et al., 2018; Li et al., 2019b, Unpublished study report, 2022) and a decrease in 11-KT (Liu et al., 2012; Liu et al., 2013b; Liu et al., 2016; Li et al., 2018; Li et al., 2019b). TPhP also induces an increase of VTG concentrations in Liu et al. (2013b) and Unpublished study report (2022), which support the estrogenic activity of TPhP.

Nevertheless, in one study (Li et al., 2019b) carried out on Japanese medaka, a decrease of E2 level was noted after 100-day exposure of female larvae (age of fish at exposure initiation), while an increase of E2 was observed after 21-day exposure of adult females (age of fish at exposure initiation).

The observations of VTG concentrations suggested an estrogenic activity of TPhP in female and male zebrafish, with increased plasmatic concentration of VTG (Liu et al., 2013b) and Unpublished study report (2022). Some results were also reported with decreased or unchanged concentration in plasma VTG of female medaka (Kawashima et al., 2021) or in whole-body homogenates of male and female zebrafish (He et al., 2021). Vtg1 gene expression in liver was increased in male zebrafish in one assay (Liu et al., 2012), while the expression in liver of vtg1 and 2 were decreased in other assays (Liu et al., 2012 (for females); Li et al., 2019b). These different perturbations of circulating steroid concentrations relate to the fish developmental stage, species and tested concentrations.

Data therefore provide *in vitro* and *in vivo* evidence that TPhP has EAS activity. In absence of mortality reported in several studies, this activity cannot be considered as a secondary non-specific consequence of general toxicity.

Table 16: Line of evidence in relation to TPhP EAS activity

Assay category	Species/ Endpoint	Biological model	Exposure time	Adult or Juvenile (when exposed)	Parameter	Lowest significant concentration	Observed effects	Reference
In vitro endocrine activity								
Transactivation	hER α	CHO-K1 cells	24h	-	REC20	-	Agonist: 0.27 μ M	Zhang et al., 2014
Transactivation	hER α	Yeast cells	-	-	REC20	-	Agonist: 0.65 μ M	Zhang et al., 2014
Transactivation	hER α	MCF-7 cells	60h	-	REC20	-	Agonist: 0.1 μ M	Zhang et al., 2014
Transactivation	hER α	CHO-K1 cells	24h	-	REC20	-	Weak agonist: 4.9 μ M	Kojima et al., 2013
Transactivation	hER α	CHO-K1 cells	24h	-	REC20	-	Weak agonist: 4.6 μ M	Kojima et al., 2016
Transactivation	hER β	CHO-K1 cells	24h	-	REC20	-	Weak agonist: 6.5 μ M	Kojima et al., 2013
Transactivation	hER β	CHO-K1 cells	24h	-	REC20	-	Weak agonist: 7.3 μ M	Kojima et al., 2016
Hormone measurement	Estradiol level	H295R cells	48h	Adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2 level	H295R cells	48h	-	b	1 μ M	↑	Ji et al., 2022
Hormone measurement	Testosterone level	H295R cells	48h	Adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	Testosterone level	H295R cells	48h	-	b	1 μ M	↓	Ji et al., 2022
Hormone measurement	E2/T ratio level	H295R cells	48h	Adult	b	0.1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2/T ratio	H295R cells	48h	-	b	1 μ M	↑	Ji et al., 2022
Gene expression	CYP19A1	H295R cells	48h	Adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

Transactivation	ER	MCF-7 cells (MVLN cells)	72h	Adult	b	0.001 mg.l ⁻¹	Antagonism	Liu et al., 2012
Transactivation	mERα	MVLN cell	24h	-	b	IC50	Antagonist activity (IC 50 = 29 μM)	Li et al., 2018
Transactivation	ER	MCF-7 cells (MVLN cells)	72h	-	EC50	EC50	Agonist activity (EC50 = 1.2 x 10 ⁻⁵ M)	Ji et al., 2022
Binding	-	SBRK3 cells	24h	-	b	-	Interaction with GPER via cAMP formation	Ji et al., 2022
Gene expression	Cyp19	H295R cells	48h	-	b	1 μM	↑	Ji et al., 2022
Transactivation	mEsr1	-	-	-	b	-	Agonist activity	Kawashima et al., 2021
<i>In vivo</i> endocrine activity								
Hormone measurement	E2 level	Zebrafish plasma	120 days	Male embryo	b	5 μg.l ⁻¹	↑	Liu et al., 2016
Hormone measurement	E2 level	Medaka plasma	100 days	Male embryo	b	0.299 μg.l ⁻¹	↑ [#]	Li et al., 2018
Hormone measurement	E2 level	Zebrafish plasma	14 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2 level	Zebrafish plasma	21 days	Male adult	b	0.2 mg.l ⁻¹	↑	Liu et al., 2013b
Hormone measurement	E2 level	Zebrafish plasma	14 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2 level	Zebrafish plasma	21 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Hormone measurement	E2 level	Medaka plasma	21 days	Female adult	b	1.72 μg.l ⁻¹	↑ [#]	Li et al., 2019b
Hormone measurement	E2 level	Zebrafish plasma	73 days	Female embryo	b	3.01 μg.l ⁻¹	↑ ^{##}	Unpublished study report, 2022
Hormone measurement	E2 level	Zebrafish plasma	120 days	Female embryo	b	5 μg.l ⁻¹	↑	Liu et al., 2016
Hormone measurement	E2 level	Medaka plasma	100 days	Female embryo	b	1.77 μg.l ⁻¹	↓ [#]	Li et al., 2019b

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

Hormone measurement	E2 level	Zebrafish plasma	21 days	Male adult	b	-	No change	He et al., 2021
Hormone measurement	E2 level	Zebrafish plasma	21 days	Female adult	b	-	#No change	He et al., 2021
Hormone measurement	T level	Zebrafish plasma	14 days	Male adult	b	1 mg.l ⁻¹	↓	Liu et al., 2012
Hormone measurement	T level	Zebrafish plasma	14 days	Female adult	b	-	No change	Liu et al., 2012
Hormone measurement	T level	Zebrafish plasma	21 days	Female adult	b	1 mg.l ⁻¹	↓	Liu et al., 2013b
Hormone measurement	T level	Medaka plasma	100 days	Male embryo	b	1.43 µg.l ⁻¹	↓#	Li et al., 2018
Hormone measurement	T level	Medaka plasma	21 days	Female adult	b	1.72 µg.l ⁻¹	↑#	Li et al., 2019b
Hormone measurement	T level	Medaka plasma	100 days	Female embryo	b	0.13 µg.l ⁻¹	↓#	Li et al., 2019b
Hormone measurement	T level	Zebrafish plasma	21 days	Male adult	b	-	#No change	He et al., 2021
Hormone measurement	T level	Zebrafish plasma	21 days	Female adult	b	-	#No change	He et al., 2021
Hormone measurement	E2/T level	Zebrafish plasma	21 days	Female adult	b	0.2 mg.l ⁻¹	↑	Liu et al., 2013b
Hormone measurement	E2/T level	Zebrafish plasma	21 days	Female adult	b	80 µg.l ⁻¹	↓#	He et al., 2021
Hormone measurement	E2/T level	Zebrafish plasma	21 days	Male adult	b	-	#No change	He et al., 2021
Hormone measurement	E2/T level	Zebrafish plasma	21 days	Male adult	b	0.04 mg.l ⁻¹	↓	Liu et al., 2013b
Hormone measurement	E2/T level	Zebrafish plasma	14 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2/T level	Zebrafish plasma	14 days	Female adult	b	-	No change	Liu et al., 2012

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

Hormone measurement	11-KT level	Zebrafish plasma	14 days	Female adult	b	-	No change	Liu et al., 2012
Hormone measurement	11-KT level	Zebrafish plasma	120 days	Female embryo	b	500 µg.l ⁻¹	↓	Liu et al., 2016
Hormone measurement	11-KT level	Medaka plasma	100 days	Female embryo	b	0.13 µg.l ⁻¹	↓ [#]	Li et al., 2019b
Hormone measurement	11-KT level	Zebrafish plasma	120 days	Male embryo	b	5 µg.l ⁻¹	↓	Liu et al., 2016
Hormone measurement	11-KT level	Medaka plasma	100 days	Male embryo	b	1.43 µg.l ⁻¹	↓ [#]	Li et al., 2018
Hormone measurement	11-KT level	Zebrafish plasma	73 days	Male embryo	b	3.01 µg.l ⁻¹	↑ ^{##}	Unpublished study report, 2022
Hormone measurement	11-KT level	Zebrafish plasma	21 days	Female adult	b	1 mg.l ⁻¹	↓	Liu et al., 2013b
Hormone measurement	11-KT level	Zebrafish plasma	14 days	Male adult	b	1 mg.l ⁻¹	↓	Liu et al., 2012
Hormone measurement	E2/11-KT level	Zebrafish plasma	120 days	Female embryo	b	500 µg.l ⁻¹	↑	Liu et al., 2016
Hormone measurement	E2/11-KT level	Zebrafish plasma	120 days	Male embryo	b	5 µg.l ⁻¹	↑	Liu et al., 2016
Hormone measurement	E2/11-KT level	Zebrafish plasma	14 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2/11-KT level	Zebrafish plasma	21 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Hormone measurement	E2/11-KT level	Zebrafish plasma	14 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Hormone measurement	E2/11-KT level	Zebrafish plasma	21 days	Male adult	b	0.2 mg.l ⁻¹	↑	Liu et al., 2013b
Hormone measurement	VTG level	Zebrafish plasma	73 days	Female embryo	b	33.3 µg.l ⁻¹	↑ ^{##}	Unpublished study report, 2022
Hormone measurement	VTG level	Zebrafish plasma	73 days	Male embryo	b	33.3 µg.l ⁻¹	↑ ^{##}	Unpublished study report, 2022

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

Hormone measurement	VTG level	Zebrafish plasma	21 days	Female adult	b	0.2 mg.l ⁻¹	↑	Liu et al., 2013b
Hormone measurement	VTG level	Zebrafish plasma	21 days	Female adult	b	80 µg.l ⁻¹	↓ [#]	He et al., 2021
Hormone measurement	VTG level	Zebrafish plasma	21 days	Male adult	b	-	[#] No change	He et al., 2021
Hormone measurement	VTG level	Japanese Medaka plasma	21 days	Female adult	b	7.19 µg.l ⁻¹	↓	Kawashima et al., 2021
Hormone measurement	VTG level	Zebrafish plasma	21 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	vtg1	Medaka liver	100 days	Female embryo	b	0.13 µg.l ⁻¹	↓ [#]	Li et al., 2019b
Gene expression	vtg1	Zebrafish plasma	14 days	Female adult	b	1 mg.l ⁻¹	↓	Liu et al., 2012
Gene expression	vtg1	Zebrafish plasma	14 days	Male adult	b	0.04 mg.l ⁻¹	↑	Liu et al., 2012
Gene expression	vtg2	Medaka liver	100 days	Female embryo	b	0.36 µg.l ⁻¹	↓ [#]	Li et al., 2019b
Gene expression	vtg2	Medaka liver	21 days	Female adult	b	1.72 µg.l ⁻¹	↓ [#]	Li et al., 2019b
Gene expression	cyp19a mRNA	Zebrafish plasma	14 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Gene expression	cyp19a mRNA	Zebrafish plasma	14 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2012
Gene expression	ER2b	Zebrafish	5 days	Embryos	b	2 mg.l ⁻¹	↑	Liu et al., 2013a
Gene expression	CYP19A	Zebrafish – ovaries	21 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	CYP19b	Zebrafish – brain	21 days	Female adult	b	0.2 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	ERα	Zebrafish – brain	21 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	ER2β1	Zebrafish – brain	21 days	Female adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	CYP19A	Zebrafish – testis	21 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

Gene expression	CYP19B	Zebrafish – brain	21 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	ERα	Zebrafish – brain	21 days	Male adult	b	1 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	ER2β1	Zebrafish – brain	21 days	Male adult	b	0.2 mg.l ⁻¹	↑	Liu et al., 2013b
Gene expression	ERα	Zebrafish – brain	120 days	Female embryo	b	500 µg.l ⁻¹	↑	Liu et al., 2016
Gene expression	CYP19A	Zebrafish – ovaries	120 days	Female embryo	b	500 µg.l ⁻¹	↑	Liu et al., 2016

b: qualitative assessment only, no parameter calculated.

No information on mortality

Mortality > 10% observed, with mortality > 10% also in the control group

MTC not exceeded for the other studies

Plausible link between adverse effects and endocrine activity

The available data indicate that an exposure of TPhP damages fish reproduction by impacting some processes and events during gametogenesis, disrupting fish fecundity (egg production and spawning) and fertility (hatching and fertilisation). Changes in fecundity and fertility are apical effects, which inform about potential adverse effects at the population level.

Although these reproduction parameters are sensitive to substances interfering with the sex hormone system, they are not considered as "EATS-mediated" as they might be influenced by non-endocrine factors such as systemic toxicity. Nevertheless, they can be used in a weight of evidence approach to draw a conclusion on a specific endocrine pathway.

These adverse effects are related to the disturbances in steroid synthesis and gametogenesis. In fact, all available *in vitro* and *in vivo* studies demonstrate that TPhP exerts an action on the endocrine hormone balance in fish. Depending on the developmental stage, species and concentration, antagonist and agonist effects on nuclear and membrane estrogen receptors are observed in organisms as well as on transcription of genes involved in steroidogenesis, leading *in vivo* to perturbations of circulating steroid concentrations. Consistency between endocrine perturbation and adverse effect on reproduction is generally observed in the available studies. The data were analysed separately for males and females as the major steroid hormones in the regulation of fish gametogenesis is the estrogen E2 in females and the androgen 11KT in males.

Reproductive dysfunction in male fish

Based on the available information for males and lines of evidence for adversity and endocrine activity, the following potential sequence of causality linked events at different levels of biological organisation, that potentially lead to an adverse ecotoxicological effect (reproductive dysfunction in fish), is represented in the following figure.




Figure 4: Potential sequences of linked events at different levels of biological organisation that potentially lead to an adverse ecotoxicological effect (reproductive dysfunction in male fish)

The decreased amount of mature spermatocyte cells (He et al., 2021) suggests a potential delay of spermatogenesis following the exposure to TPhP. A disruption (dose-dependent increase) of male-typical reproductive behaviours before spawning (chasing trajectory and time, courtship frequency and mating number) observed in Li et al. (2018) also supports the reproduction impairment of male fish by exposure to TPhP.

CYP19A and CYP19B were significantly upregulated in the testis and the brain respectively (Liu et al., 2012; 2013 b) and could increase conversion of T to E2. Moreover, estrogen receptors (ER α , ER2 β 1) were up-regulated in the brain (Liu et al., 2013b).

Generally, after TPhP exposure of male fish, plasma E2 increased significantly, plasma 11-KT decreased and E2/11-KT ratio increased (Liu et al., 2012; Liu et al., 2016; Li et al., 2018; Liu et al., 2013b). However, this trend could not be verified in the unpublished study report (2022), as E2 levels were not measured in males and consequently the E2/11KT ratio could not be calculated. 11-KT plays an important role in the spermatogonial proliferation. A decreased 11-KT level could stop meiosis and germ cell maturation (Schulz et al., 2010). The decrease of 11-KT and the increase in E2 concentrations in plasma revealed by E2/11KT ratio (indicative of estrogenicity) can be associated with reduced semen production and density in male fish. Finally, the increase of the expression in liver of *VTG1* gene (Liu et al., 2012) and the increase of plasma VTG levels in male fish exposed to TPhP (Liu et al., 2013b; Unpublished study report, 2022) are consistent with the observed changes in sex hormones (increased E2 synthesis). These events support the reproduction impairment of male fish observed after exposure to TPhP. The following table highlights these relationships. It has to be noted that the reported concentrations are those for which the measured parameters are significant.

Table 17: Dose-response and temporal concordance between the key events for male fish

[Zebrafish – Japanese Medaka - male] potential dose–response and temporal concordance between the key events and AO							
Dose ($\mu\text{g}\cdot\text{l}^{-1}$)	Estradiol level	11-KT level	E2/11KT ratio	VTG level	Testis histopathology (abnormalities)	Observed Adverse effect	Reference
0.299 [#]	↑ (100 days)					↓ hatching rate	Li et al., 2018
1.43 [#]	↑ (100 days)	↓ (100 days)				↑ intersex incidence, abnormal chasing behaviour ↓ fertilisation rate.	Li et al., 2018
3.01 ^{##}		↑ (73 days)*					Unpublished study report, 2022
5	↑ (120 days)	↓ (120 days)	↑ (120 days)				Liu et al., 2016
7.76 ^{##}		↑ (73 days)*					Unpublished study report, 2022
33.3 ^{##}		↑ (73 days)*		↑ (73 days)			Unpublished study report, 2022
40				↑ (14 days)***			Liu et al., 2012
44.9						↓ Fertilised eggs**	Kawashima et al., 2021
50		↓ (120 days)	↑ (120 days)				Liu et al., 2016
76.8 ^{##}		↑ (73 days)*			↑ testis maturation stage (73 days)		Unpublished study report, 2022

80 [#]					↑ immature spermatocytes (spermatogonia and spermatocyte) ↓ mature spermatocytes (spermatids and spermatozoa)	↓ hatching rate ↓ fertilisation rate	He et al., 2021
200				↑ (14 days)***			Liu et al., 2012
200	↑ (21 days)		↑ (21 days)			↓ hatching rate	Liu et al., 2013b
500		↓ (120 days)	↑ (120 days)				Liu et al., 2016
1000	↑ (14 days)	↓ (14 days)	↑ (14 days)	↑ (14 days)***			Liu et al., 2012
1000				↑ (21 days)			Liu et al., 2013b

* Unpublished study report (2022), where E2 levels were not measured in males and consequently the E2/11KT ratio could not be calculated.

** However no significant effect on fertility rate (ratio).

*** Transcriptional gene *VTG1*.

No information on mortality

Mortality > 10% observed, with mortality > 10% also in the control group

MTC not exceeded for the other studies

Blank cells: parameter not investigated at this concentration.

Dose–response and temporal concordance between the key events and AO are often observed at concentration where systemic toxicity was not observed. It cannot be considered as a secondary non-specific consequence of general toxicity.

Reproductive dysfunction in female fish

Impairment of female fish fecundity is also a clear adverse effect observed after TPhP exposure with a decreased number of eggs and spawning events.

In fact, an inhibition of the ovarian development in female fish (He et al. 2021) and alterations of ovaries (Li et al., 2019b; He et al., 2021) were observed. Ovary maturation retardation affected the fecundity of female fish.

TPhP exposure subsequently leads to:

- a decrease in egg production (Liu et al., 2013b; Li et al., 2019b, He et al., 2021; Kawashima et al., 2021),
- a decrease in spawning events (Liu et al., 2013b; Li et al., 2018; He et al., 2021).

TPhP induced an increase in circulating E2 concentrations in female zebrafish and female medaka (Liu et al., 2012; Liu et al., 2013b; Liu et al., 2016; Li et al., 2019b; Unpublished study report, 2022). Nevertheless, no change was observed in one study (He et al., 2021). In another study (Li et al., 2019b) carried out on Japanese medaka, a decrease of E2 level was noted after 100-day exposure of female larvae (age of fish at exposure initiation), while an increase of E2 was observed after 21-day exposure of adult females (age of fish at exposure initiation). The authors hypothesised that anti-ER activity would retard ovary development, which would further decrease the E2 level in the long term. To support this hypothesis, the authors showed that the metabolite 4-OH-TPhP (but not TPhP) was able to inhibit E2 activity in the transgenic medaka model.

A perturbation in maturation-inducing steroids could cause impairment in oocyte maturation and quality thereby resulting in a decrease in cumulative fecundity.

VTG, which induces oocyte maturation in fish is synthesised in the liver in response to estradiol stimulation. TPhP induced a down regulation of the expression of *VTG1* and *VTG2* gene in liver (Liu et al., 2012; Li et al., 2019b) and perturbations of VTG concentrations (Liu et al., 2013b;

He et al., 2021; Kawashima et al., 2021; Unpublished study report, 2022), associated with a concurrent perturbation of E2. Thus, TPhP activates the estrogenic pathway and/or perturbs the steroidogenic pathway, which can explain the synthesis perturbation of E2 and VTG after exposure to TPhP.

The following table reinforces the hypothesis that TPhP disturbs steroid synthesis leading to impairment in oocyte maturation and subsequently in egg production and spawning by emphasising the consistency of the effects observed in each study. It has to be noted that all the measurements were performed at the adult stage and the reported concentrations are those for which the measured parameters are significant.

Table 16: Dose-response and temporal concordance between the key events and AO for female fish

[Zebrafish – Japanese Medaka - female] potential dose–response and temporal concordance between the key events and AO					
Dose (µg/L)	Estradiol level	VTG level	Gonad histopathology impairment	Observed Adverse effect	Reference
0.131 [#]		↓ (100 days)**	↓ number of mature oocyte (100 days)		Li et al., 2019b
0.299 [#]				↓ hatching rate	Li et al., 2018
0.363 [#]		↓ (100 days)***	↓ number of mature oocyte (100 days)	↓ egg production	Li et al., 2019b
1.43 [#]				↓ fertilisation rate	Li et al., 2018
1.77 [#]	↑ (21 days) ↓ (100 days)	↓ (21 days)***	↓ number of mature oocyte (100 days)	↓ egg production	Li et al., 2019b
3.01 ^{##}	↑ (73 days)				Unpublished study report, 2022
5	↑ (120 days)				Liu et al., 2016
7.19		↓ (21 days)			Kawashima et al., 2021
7.76 ^{##}	↑ (73 days)				Unpublished study report, 2022
17		↓ (21 days)			Kawashima et al., 2021
33.3 ^{##}	↑ (73 days)	↑ (73 days)			Unpublished study report, 2022
44.9		↓ (21 days)		↓ egg production ↓ fertilised eggs*	Kawashima et al., 2021
76.8 ^{##}	↑ (73 days)	↑ (73 days)			Unpublished study report, 2022
80 [#]		↓ (21 days)	↓ number of mature oocyte (21 days)	↓ hatching rate ↓ fertilisation rate ↓ cumulative eggs ↓ spawning event	He et al., 2021
200		↑ (21 days)		↓ hatching rate ↓ eggs production ↓ spawning event	Liu et al., 2013b
500	↑ (120 days)				Liu et al., 2016
1000	↑ (14 days)	↓ (14 days)**			Liu et al., 2012

1000	↑ (21 days)	↑ (21 days)		↓ hatching rate ↓ egg production ↓ spawning event	Liu et al., 2013b
------	-------------	-------------	--	---	-------------------

* However no significant effect on fertility rate (ratio).

** Transcriptional gene *VTG1*.

*** Transcriptional gene *VTG2*.

No information on mortality

Mortality > 10% observed, with mortality > 10% also in the control group

MTC not exceeded for the other studies

Blank cells: parameter not investigated at this concentration

Dose–response and temporal concordance between the key events and AO are often observed at concentration where systemic toxicity was not observed. It cannot be considered as a secondary non-specific consequence of general toxicity.

Conclusion

TPhP shows endocrine activity in fish with adverse effects on fecundity, fertility and subsequently to reproduction. These adverse effects can be related to the disturbances in steroid hormone concentrations and impairment of gametogenesis. The resulting perturbation in maturation-inducing steroids could cause impairment in oocyte and testis maturation. Changes in fecundity and fertility are apical outcomes, which inform about potential adverse effects at the population level. Moreover, it is important to note that several adverse effects on altered reproductive capacity of fish are observed at environmentally relevant levels without concurrent systemic effect.

Overall, based on all available scientific evidence, it can be concluded that TPhP fulfils the WHO/IPCS definition of an endocrine disruptor:

- **It shows population relevant adverse effects on fertility and reproduction in fish.**
- **It has EAS activity as clearly shown both *in vitro* and *in vivo*.**
- **Consistency in the effect and EAS activity observed provides evidence that EAS modalities are plausibly biologically linked to the adverse effects.**

6. Conclusions on the SVHC Properties

CMR assessment

Not relevant for the identification of the substance as SVHC in accordance with Article 57 (a) to (f) of the REACH Regulation.

PBT and vPvB assessment

Not relevant for the identification of the substance as SVHC in accordance with Article 57 (a) to (f) of the REACH Regulation.

Assessment under Article 57(f)

Summary of the data on the intrinsic/hazardous properties (providing scientific evidence of probable serious effects to HH and/or ENV)

The available *in vitro* information demonstrates the capacity of TPhP to competitively bind and activate the nuclear estrogen receptors ER α and ER β of several vertebrate species including rat, mouse, fish, chicken, frog and turtle as evidenced by ER transactivation in reporter cell lines (Liu et al., 2012; Kojima et al., 2013; Zhang et al., 2014; Kojima et al., 2016; Li et al., 2018; Ji et al., 2020; Medvedev et al., 2020; Kawashima et al., 2021; Ji et al., 2022). In addition, TPhP can induce ER-regulated gene expression, and related physiological cell responses (e.g., increased cell proliferation in Zhang et al. (2014); Krivoshiev et al. (2016); Ji et al. (2020)). Two recent studies show that TPhP can also bind and activate GPER (Guan et al., 2022 and Ji et al., 2022). The available H295R assays (Liu et al., 2012; Ji et al., 2022) on human adrenal carcinoma cells show that TPhP affects steroidogenesis by increasing estrogen levels (17 β -estradiol) and by increasing expression of genes involved in this pathway like CYP19 and 3 β -HSD2. *In vivo* fish studies indicate that CYP19A is significantly up-regulated by exposure to TPhP (Liu et al., 2016). Significant alteration of plasmatic concentrations of E2 and E2/T ratio (more specific observation in females) and E2/11-KT ratio (more specific observation in males) can result from this modification in the steroidogenesis pathway. The degree of perturbation of circulating steroid concentrations depends on the fish developmental stage, species and tested concentrations. The observations of VTG concentrations, that are consistent with perturbation of E2 concentrations, suggest an EAS activity of TPhP in female and male zebrafish, with altered plasmatic concentration of VTG (Liu et al., 2013b; Unpublished study report, 2022). Therefore, TPhP exerts an effect on the endocrine balance in fish. It has EAS activity as clearly shown both *in vitro* and *in vivo*.

In vivo studies on fish reproduction show an alteration of gametogenesis in both sexes with a disruption of testis (He et al., 2021; Unpublished study report, 2022) and ovary maturation (Li et al., 2019b; He et al., 2021) followed by a decrease in reproductive success in terms of fecundity: decrease in egg production (Li et al., 2019b; Kawashima et al., 2021), spawning events (He et al., 2021; Liu et al., 2013b), fertility (impaired hatchability in Liu et al. (2013b); Li et al. (2018); He et al., (2021)), and fertilisation rate (Li et al., 2018; He et al., 2021). Therefore, TPhP shows adverse effects on fertility and reproduction in fish, observed in several studies at levels without concurrent systemic effects.

The consistency between the observed adverse effects and EAS activity provides evidence that EAS modalities are plausibly biologically linked to the adverse effects. Depending on the developmental stage, species and concentration, antagonist and agonist effects are observed in organisms, leading *in vivo* to perturbations of circulating steroid concentrations in most of the analysed studies.

The effect observed on reproduction in fish (fecundity and fertility) can affect population stability and is considered as an adverse effect relevant at population level.

Based on all available scientific evidence, it can be concluded that TPhP fulfils the WHO/IPCS definition of an endocrine disruptor.

Equivalent level of concern assessment

Environment

Triphenyl phosphate shall be identified as a substance of very high concern in accordance with Article 57(f) of Regulation (EC) 1907/2006 (REACH) because of its endocrine disrupting properties for which there is scientific evidence of probable serious effects to environment, which

gives rise to an equivalent level of concern to those substances listed in points (a) to (e) of Article 57 REACH.

In order to assess equivalent level of concern (ELoC), some of the factors identified in ECHA (2012) to evaluate ELoC for sensitisers, which are considered relevant for the present case, are also included in the list below:

- Characteristics of the effects:
 - *Probable adverse environmental effects related to:*
 - *Severity, irreversibility and delayed effects.*
 - *Broad environmental relevance.*
 - *Other factors:*
 - *Wide occurrence and environmental distribution.*
 - *Concern related to co-exposure and combined effects.*
 - *Societal concern.*
 - *Is derivation of 'safe concentration' possible?*

The following description includes the elements relevant to the analysis in an environmental context.

Probable serious ecotoxicological effects

Severity, irreversibility and delayed effects

TPhP induces adverse effects on reproductive organs and fecundity, and affects the reproductive ability of fish. Changes in fecundity and fertility are apical outcomes, which inform about potential adverse effects at the population level, which can lead to serious effects on ecosystems. Moreover, it is important to note that several adverse effects on altered reproductive capacity of fish are observed at environmentally relevant levels without concurrent systemic effect. Exposure during the juvenile period of fish may cause gonad histopathological changes, leading to delayed effects on survival and growth of offspring. There is evidence that a short time exposure may be sufficient to provoke long-term effects.

Broad environmental relevance

Effects on sexual development and reproductive ability *via* endocrine EAS MoA has a broad environmental relevance. As data on only a small proportion of the existing species are available, mainly in fish, potential effects on other organisms remain unknown. Adverse effects are thus not expected to be restricted to certain taxonomic groups or species. As the main endocrine systems are known to be largely conserved among vertebrate species in various environments, it is likely that a wide range of wildlife species with different functions in ecosystems could be affected.

Wide occurrence and environmental distribution

Several literature studies demonstrate that TPhP can be found in the environment and TPhP has a widespread use, which leads to a ubiquitous occurrence. TPhP may enter the environment via emissions from various sources, which is supported by occurrence and monitoring studies (water, sediment, sewage sludge, indoor dust/air). Therefore, environmental species including predators and vulnerable species (Giulivo et al., 2017; Garcia-Garin et al., 2020; Sala et al., 2021; Sala et al., 2022) are exposed to TPhP. As long as emissions of TPhP do not cease, exposure of species from the environment cannot be avoided. In addition, human biomonitoring data demonstrate also the presence of TPhP or its metabolites in human milk (Sundkvist et al., 2010; Kim et al., 2014; Zheng et al., 2021), placenta (Ding et al., 2016; Zhao et al., 2017) and urine (Carignan et al., 2016; Li et al., 2019a) confirming human exposure to this substance from the very early stages of life. Moreover, TPhP has also frequently been detected in indoor house dust samples in Europe, including Belgium and Sweden, and outside Europe, including Japan, Philippines and USA (Kanazawa et al., 2010; Kim et al., 2013; Marklund et al., 2003; Stapleton et al., 2009; Van den Eede et al., 2011). Thus, it can be concluded that TPhP reaches diverse environmental compartments and biota of remote areas including humans.

Concern related to co-exposure and combined effects

TPhP is a constituent or an impurity of several registered substances¹⁵. Thus, mixture effects, where substances act additively or with synergistic effects, cannot be excluded and this might impact the threshold of toxicity. Besides, environmental occurrence and human biomonitoring data (see section 3.2) show that TPhP is detected in the environment, in environmental species as well as in human fluids together with other organophosphate flame retardants. Typical examples are sewage plant effluents and human fluids where TPhP occurs jointly with organophosphate flame retardants (Carignan et al., 2016; Ding et al., 2016; Kim et al., 2014; Li et al., 2019; Sundkvist et al., 2010; Ya et al., 2019; Zhao et al., 2016; Zhao et al., 2017; Zheng et al., 2021).

Societal concern

In relation to the environment, the impairment of fertility can be an issue regarding species survival. There is an increasing concern related to the preservation of biodiversity and increasing evidence that it is threatened due to various causes including global warming and excessive pressure due to human activities (Jenssen, 2006). EDCs may also contribute to the pressure on the survival of endangered species (Tubbs and McDonough, 2018). Preserving and restoring ecosystems and biodiversity is one of the key aims of the European Green Deal (European Commission, 2019) that is an integral part of the European Commission's strategy to implement the United Nations' 2030 Agenda and the sustainable development goals¹⁶.

¹⁵ <https://echa.europa.eu/documents/10162/c7a3c8b0-3fd3-3a4a-e791-2b4ae6a696d5>

¹⁶ <https://sustainabledevelopment.un.org/post2015/transformingourworld>

Is derivation of 'safe concentration' possible?

Endocrine regulation, which is set up during critical life stages in vertebrates, is a complex feedback process. Any disturbance of this regulation during transient but vulnerable life stages can lead to irreversible effects for the entire lifetime or even in the following generations which will also be dependent on the organism group. Moreover, based on the available ED specific test guidelines, it is difficult to assess the latency of the effects. Therefore, prediction of future effects and derivation of safe concentration for the environment is associated with large uncertainties. Thus, effects may be overlooked, not expressed or equivocal. For TPHP, for example, no data is available for most of the trophic levels, which makes it difficult to derive a safe exposure level in the environment. Another reason may be that low effect concentrations are difficult to determine definitively, as effects may only be observed in certain life stages or time windows. Additionally, seasonal effects may lead to difficulty in predicting the impact on the development of different groups of organisms.

The ELoC elements are summarised in Table 16 below.

Table 18: ELoC summary reporting

	Adverse effects on reproduction	Overall conclusion
Probable serious effects?*	<ul style="list-style-type: none"> ▪ Adverse effects on reproductive organs of fish exposed to TPhP. ▪ Adverse effect on fecundity and reproductive ability in fish exposed to TPhP. ▪ Several adverse effects on altered reproductive capacity of fish observed at environmentally relevant levels without concurrent systemic effect. 	<p>YES</p> <ul style="list-style-type: none"> ▪ Pattern of ED-related effects associated with serious dysfunction in fish (fecundity, reproductive ability, gonad histopathological effects).
▪ Delayed and irreversibility of effects?	<ul style="list-style-type: none"> ▪ Exposure during juvenile period of fish may cause gonad histopathological changes, leading to delayed effects on survival and growth of offspring. ▪ There is evidence that a short time TPhP exposure may be sufficient to provoke long-term effects on survival and growth of offspring. ▪ Irreversible effects during the entire lifetime or even in the following generations are expected following disruption of endocrine regulation during transient but vulnerable life stages. 	<p>YES</p> <ul style="list-style-type: none"> ▪ Exposure during the juvenile period of fish has been shown to have consequences in relation to ED-related effects later in life. ▪ Short time TPhP exposure may be sufficient to provoke long-term effects on survival and growth of offspring. ▪ Due to conservation of the reproductive endocrine system in a wide range of taxa in different ecosystems, other species or taxa can be impacted as well.
▪ Broad environmental relevance	<ul style="list-style-type: none"> ▪ Broad environmental relevance due to the effects observed on sexual development and reproductive ability <i>via</i> endocrine EAS MoA. ▪ As data on only a small proportion of the existing species are available, mainly fish, potential effects on other organisms remain unknown. ▪ It is very likely that a wide range of wildlife species with different functions in ecosystems could be affected. 	<p>YES</p> <ul style="list-style-type: none"> ▪ ED MoA with broad environmental relevance.

ANNEX XV – IDENTIFICATION OF TRIPHENYL PHOSPHATE AS SVHC

<p>Wide occurrence and environmental distribution</p>	<ul style="list-style-type: none"> ▪ Widespread use of TPhP, which leads to an ubiquitous occurrence. ▪ Environmental species including predators and vulnerable species (fin whale) can be exposed more or less continuously to TPhP and exposure can thus not be avoided. ▪ Concern supported by occurrence and environmental monitoring data (water, sediment, sewage sludge, indoor dust/air) and human studies demonstrating the presence of TPhP or its metabolites in human milk, placenta and urine. 	<p>YES</p> <ul style="list-style-type: none"> ▪ Wide occurrence and environmental distribution.
<p>Potential to cause combined effects (co-exposure)</p>	<ul style="list-style-type: none"> ▪ TPhP is a constituent or an impurity of several registered substances¹⁷. ▪ Mixture effects cannot be excluded and may impact the threshold of toxicity. <p>TPhP and organophosphate flame retardant co-exposure have been documented in sewage plant effluents and in human fluids.</p>	<p>YES</p> <ul style="list-style-type: none"> ▪ Combined exposure cannot be avoided and thus combined effects cannot be ruled out.
<p>Societal concern?</p>	<ul style="list-style-type: none"> ▪ Fertility impairment can be an issue regarding environmental species survival. ▪ Increasing concern about biodiversity preservation. ▪ EDC may also contribute to the challenge of survival of endangered species. 	<p>YES</p> <ul style="list-style-type: none"> ▪ Major societal environmental concerns.
<p>Uncertainties in deriving safe concentration limits</p>	<ul style="list-style-type: none"> ▪ Endocrine regulation, which is set up during critical life stages in vertebrates, is a very complex feedback process. ▪ This complex ED feedback process prevents to predict potential future effects and thus safe exposure levels for the environment. ▪ For TPhP, data is not available for all trophic levels, which makes it difficult to derive a safe exposure level in the environment. 	<p>YES</p> <ul style="list-style-type: none"> ▪ Derivation of safe concentration associated with large uncertainties.

**This factor is intended to discuss the severity of the effects and not their probability*

¹⁷ <https://echa.europa.eu/documents/10162/c7a3c8b0-3fd3-3a4a-e791-2b4ae6a696d5>

Summary of the ELoC assessment

Triphenylphosphate (TPhP) fulfils the WHO/IPCS definition of an endocrine disruptor relevant for the environment. The very high concern raised by this property is substantiated by the severity and irreversibility of the effects observed on organisms. These effects are relevant at the population level and may have long term consequences. A large variety of species may be adversely affected and there are difficulties to quantify a safe level of exposure with regard to the endocrine mediated effects.

In conclusion, there is scientific evidence that TPhP causes probable serious effects to the environment due to its endocrine disrupting properties, which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of the REACH Regulation.

TPhP is proposed to be identified as a substance of very high concern in accordance with Article 57(f) of Regulation (EC) 1907/2006 (REACH) because it is a substance with endocrine disrupting properties for which there is scientific evidence of probable serious effects to the environment, which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 REACH.

Conclusion on the Article 57(f) assessment

TPhP is proposed to be identified as a substance of very high concern in accordance with Article 57(f) of Regulation (EC) 1907/2006 (REACH) because it is a substance with endocrine disrupting properties for which there is scientific evidence of probable serious effects to the environment, which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 REACH.

The available *in vitro* information demonstrates the capacity of TPhP to competitively bind and activate nuclear estrogen receptors ER α and ER β of several vertebrate species including rat, mouse, fish, chicken, frog and turtle as evidenced by ER transactivation in reporter cell lines (Liu et al., 2012; Kojima et al., 2013; Zhang et al., 2014; Kojima et al., 2016; Li et al., 2018; Ji et al., 2020; Medvedev et al., 2020; Kawashima et al., 2021; Ji et al., 2022). In addition, TPhP can induce ER-regulated gene expression, and related physiological cell responses (e.g., increased cell proliferation in Zhang et al., 2014; Krivoshiev et al., 2016; Ji et al., 2020). Two recent studies show that TPhP can also bind and activate GPER (Guan et al., 2022; Ji et al., 2022). The available H295R assays (Liu et al., 2012; Ji et al., 2022) on human adrenal carcinoma cells show that TPhP affects steroidogenesis by increasing estrogen levels (17 β -estradiol) and by increasing expression of genes involved in this pathway like CYP19 and 3 β -HSD2. *In vivo* fish studies indicate that CYP19A is significantly up-regulated by exposure to TPhP (Liu et al., 2016). Significant alteration of plasmatic concentrations of E2 and E2/T ratio (more specific observation in females) and E2/11-KT ratio (more specific observation in males) can result from this modification in the steroidogenesis pathway. The degree of perturbation of circulating steroid concentrations depends on the fish developmental stage, species and tested concentrations. The observations of VTG concentrations, that are consistent with perturbation of E2 concentrations, suggest an EAS activity of TPhP in female and male Zebrafish, with altered plasmatic concentration of VTG (Liu et al., 2013b; Unpublished study report 2022). Therefore, TPhP exerts an effect on the endocrine balance in fish. It has EAS activity as clearly shown both *in vitro* and *in vivo*.

In vivo studies on fish reproduction show an alteration of gametogenesis in both sex with a disruption of testis (He et al., 2021; Unpublished study report 2022) and ovary maturation (Li et al., 2019b; He et al., 2021) followed by a decrease in reproductive success in terms of fecundity : decrease in egg production (Li et al., 2019b; Kawashima et al., 2021), spawning events (He et al., 2021; Liu et al., 2013b), fertility (impaired hatchability in Liu et al., 2013b;

Li et al., 2018; He et al., 2021), and fertilisation rate (Li et al., 2018; He et al., 2021). Therefore, TPhP shows adverse effects on fertility and reproduction in fish, observed in several studies at levels without concurrent systemic effect.

The consistency between the observed adverse effects and EAS activity provides evidence that EAS modalities are plausibly biologically linked to the adverse effects. Depending on the developmental stage, species and concentration, antagonist and agonist effects are observed in organisms, leading *in vivo* to perturbations of circulating steroid concentrations in most of the analysed studies.

The effect observed on reproduction in fish (fecundity and fertility) can affect population stability and is considered as an adverse effect relevant at population level.

Based on all available scientific evidence, it can be concluded that TPhP fulfils the WHO/IPCS definition of an endocrine disruptor. The very high concern raised by this property is substantiated by the severity and irreversibility of the effects on organisms and populations that may have long term consequences, the large variety of species that may be adversely affected and the difficulties to quantify a safe level of exposure with regard to the endocrine mediated effects.

In conclusion, there is scientific evidence that TPhP causes probable serious effects to the environment due to its endocrine disrupting properties, which give rise to an equivalent level of concern to those of other substances listed in points (a) to (e) of Article 57 of the REACH Regulation.

Part II

7. Registration and C&L notification status

Registration status

Table 19 Registration status

From the ECHA dissemination site ¹⁸	
Registrations	<input checked="" type="checkbox"/> Full registration(s) (Art. 10) <input type="checkbox"/> Intermediate registration(s) (Art. 17 and/or 18)

CLP notification status

Table 20: CLP notifications

	CLP Notifications ¹⁹
Number of aggregated notifications	25
Total number of notifiers	1440

¹⁸ Substance specific entry on the dissemination site: [Registration Dossier - ECHA \(europa.eu\)](https://echa.europa.eu/registration-dossier) (accessed 14-11-2023).

¹⁹ C&L Inventory database, <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database> (accessed 14-11-2023).

8. Total tonnage of the substance

Table 21: Tonnage status

Total tonnage band for the registered substance (excluding the volume registered under Art 17 or Art 18) ²⁰	≥ 100 to < 1 000 t/pa
Tonnage information from public sources other than registration dossiers (if available) ²¹	No information

9. Information on uses of the substance

TPhP is used as a flame retardant and plasticiser in polymer formulations, adhesives and sealants, cosmetics and personal care products. TPhP is used by consumers, in articles, by professional workers (widespread uses), in formulation or re-packing, at industrial sites and in manufacturing. TPhP is present as an impurity in many other organophosphate flame retardants or as a constituent of this family of compounds.

Other releases to the environment of this substance are likely to occur from: indoor use and outdoor use resulting in inclusion into or onto materials (e.g. binding agent in paints and coatings or adhesives).

The following information is extracted from the ECHA dissemination website.

Table 22: Uses

	Use(s)	Registered use	Use <u>likely</u> to be in the scope of Authorisation
Uses as intermediate	-	Yes	No
Formulation or repacking	Formulation of plastic (compounding) and rubber preparations (compounding), including coatings and adhesive products. Formulation of flame retardant/plasticiser preparations and cosmetics.	Yes	Yes
Uses at industrial sites	Production of plastic and rubber articles (conversion)	Yes	Yes

²⁰ ECHA's dissemination site <https://echa.europa.eu/fr/information-on-chemicals/registered-substances/-/disreg/substance/100.003.739> (accessed 14 11 2023, last updated 21-07-2022).

Uses by professional workers	Use of adhesives and sealants, outdoor. Use of adhesives and sealants, indoor. Use as laboratory chemicals.	Yes	Yes
Consumer uses	Use of adhesives and sealants, outdoor. Use of adhesives and sealants, indoor. Use of cosmetic products containing triphenyl phosphate. Other release to the environment of this substance is likely to occur from: indoor use and outdoor use resulting in inclusion into or onto materials (e.g. binding agent in paints and coatings or adhesives).	Yes	Yes
Article service life	Release to the environment of this substance can occur from: - Waste stage: hazardous waste incineration, municipal and industrial. - Waste stage: shredding. - Waste stage: municipal waste incineration. - Municipal landfill.	Yes	No

10. Information on structure of the supply chain

No information available.

11. Additional information

No additional information available.

References for Part I

- Alkan, N., Alkan, A., Castro-Jiménez, J., Royer, F., Papillon, L., Ourgaud, M., & Sempéré, R. (2021). Environmental occurrence of phthalate and organophosphate esters in sediments across the Gulf of Lion (NW Mediterranean Sea). *Science of the Total Environment*, 760, 143412. <https://doi.org/10.1016/j.scitotenv.2020.143412>
- Aminot, Y., Tao, L., Héas-Moisán, K., Pollono, C., O'Loghlin, M., & Munsch, C. (2023). Organophosphate esters (OPEs) in the marine environment : Spatial distribution and profiles in French coastal bivalves. *Chemosphere*, 330, 138702. <https://doi.org/10.1016/j.chemosphere.2023.138702>
- Anderson, C., Wischer, D., Schmieder, A., & Spiteller, M. (1993). Fate of triphenyl phosphate in soil. *Chemosphere*, 27(5), 869-879. [https://doi.org/10.1016/0045-6535\(93\)90017-y](https://doi.org/10.1016/0045-6535(93)90017-y)
- Boethling RS & Cooper JC (1985). Environmental fate and effects of triaryl and trialkyl/aryl phosphates. *Residue Reviews* 94: 49-99.
- Carignan, C. C., Fang, M., Stapleton, H. M., Heiger-Bernays, W., McClean, M. D., & Webster, T. F. (2016). Urinary biomarkers of flame retardant exposure among collegiate U.S. gymnasts. *Environment International*, 94, 362-368. <https://doi.org/10.1016/j.envint.2016.06.030>
- Chen, Y., Tang, H., He, J., Wu, X., Wang, L., Liu, X., & Lin, H. (2019). Interaction of nuclear ERs and GPER in vitellogenesis in zebrafish. *The Journal Of Steroid Biochemistry And Molecular Biology*, 189, 10-18.
- Chen, J., Hu, H., Feng, L., & Ding, G. (2022). Ecotoxicity assessment of triphenyl phosphate (TPHP) exposure in *Hoplobatrachus rugulosus* tadpoles. *Chemosphere*, 292, 133480. <https://doi.org/10.1016/j.chemosphere.2021.133480>
- CITI (1992). Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL Japan. Compiled under the Supervision of Chemical Products Safety Division, Basic Industries Bureau MITI, Ed. by Japan Chemical Industry Ecology-Toxicology & Information Centre CITI.
- Clagett - Dame, M., & DeLuca, H. F. (2002). THEROLE OF VITAMIN A IN MAMMALIAN REPRODUCTION AND EMBRYONIC DEVELOPMENT. *Annual Review Of Nutrition*, 22(1), 347-381. <https://doi.org/10.1146/annurev.nutr.22.010402.102745e>
- Conley, A., and Hinshelwood M. (2001). "Mammalian aromatases." *Reproduction* 121 (5): 685-95. <https://doi.org/10.1530/rep.0.1210685>.
- Cristale, J., García - Vázquez, A., Barata, C., & Lacorte, S. (2013). Priority and emerging flame retardants in rivers : Occurrence in water and sediment, *Daphnia magna* toxicity and risk assessment. *Environment International*, 59, 232-243.
- Ding, J., Zhang, X., Huang, W., Feng, L., & Yang, F. (2016). Organophosphate ester flame retardants and plasticizers in human placenta in Eastern China. *Science Of The Total Environment*, 554-555, 211-217. <https://doi.org/10.1016/j.scitotenv.2016.02.171>
- ECHA (2012). Identification of substances as SVHCs due to equivalent level of concern to CMRs (Article 57(f)) – sensitizers as an example. [ECHA proposed approach - \(europa.eu\)](https://echa.europa.eu/europa)
- ECHA/EFSA (2018) European Chemicals Agency and European Food Safety Authority with the technical support of the Joint Research Centre (JRC), "Guidance for the identification of endocrine disruptors in the context of Regulations (EU) No 528/2012 and (EC) No 1107/2009." *EFSA Journal* 16 (6):e05311. doi: doi:10.2903/j.efsa.2018.5311.

- Environment Agency (2009). Environmental risk evaluation report: Triphenyl phosphate (CAS no. 115-86-6).
- European Commission (2019). The European Green Deal. Communication from the Commission to the European Parliament, the European Council, the Council, the European Economic and Social Committee and the Committee of the Regions. Brussels, 11.12.2019. [resource.html \(europa.eu\)](#)
- Fan, B., Dai, L., Liu, C., Sun, Q., Yu, L. (2022) Nano-TiO₂ aggravates bioaccumulation and developmental neurotoxicity of triphenyl phosphate in zebrafish larvae. *Chemosphere*, 287, 132161. <https://doi.org/10.1016/j.chemosphere.2021.132161>
- Fang, H., Tong, W., Branham, W. S., Moland, C. L., Dial, S. L., Hong, H., Xie, Q., Perkins, R., Owens, W. A., & Sheehan, D. M. (2003). Study of 202 Natural, Synthetic, and Environmental Chemicals for Binding to the Androgen Receptor. *Chemical Research In Toxicology*, 16(10), 1338-1358. <https://doi.org/10.1021/tx030011g>
- Garcia-Garin, O., Sala, B., Aguilar, Á., Vighi, M., Víkingsson, G. A., Chosson, V., Eljarrat, E., & Borrell, A. (2020). Organophosphate contaminants in North Atlantic fin whales. *Science Of The Total Environment*, 721, 137768. <https://doi.org/10.1016/j.scitotenv.2020.137768>
- Garoche, C., Boulahtouf, A., Grimaldi, M., Chiavarina, B., Toporová, L., Broeder, M. J. D., Legler, J., Bourguet, W., & Balaguer, P. (2021). Interspecies Differences in Activation of Peroxisome Proliferator-Activated Receptor γ by Pharmaceutical and Environmental Chemicals. *Environmental Science & Technology*, 55(24), 16489-16501. <https://doi.org/10.1021/acs.est.1c04318>
- Ghyselinck, N. B., & Duester, G. (2019). Retinoic acid signaling pathways. *Development*, 146(13).
- Giulivo, M., Capri, E., Kalogianni, E., Milačić, R., Majone, B., Ferrari, F., Eljarrat, E., & Barceló, D. (2017). Occurrence of halogenated and organophosphate flame retardants in sediment and fish samples from three European river basins. *Science Of The Total Environment*, 586, 782-791. <https://doi.org/10.1016/j.scitotenv.2017.02.056>
- Guan, R., Li, N., Wang, W., Liu, W., Li, X., & Zhao, C. (2022). The adverse outcome pathway (AOP) of estrogen interference effect induced by triphenyl phosphate (TPP) : Integrated multi-omics and molecular dynamics approaches. *Ecotoxicology And Environmental Safety*, 234, 113387. <https://doi.org/10.1016/j.ecoenv.2022.113387>
- Gustavsson, J., Wiberg, K., Ribéli, E., Nguyen, M. A., Josefsson, S., & Ahrens, L. (2018). Screening of organic flame retardants in Swedish river water. *Science Of The Total Environment*, 625, 1046-1055. <https://doi.org/10.1016/j.scitotenv.2017.12.281>
- He, J., Yang, X., & Liu, H. (2021). Enhanced toxicity of triphenyl phosphate to zebrafish in the presence of micro- and nano-plastics. *Science Of The Total Environment*, 756, 143986. <https://doi.org/10.1016/j.scitotenv.2020.143986>
- Houck, K. A., Simha, A., Bone, A. J., Doering, J. A., Vliet, S. M., LaLone, C. A., Medvedev, A. V., & Makarov, S. S. (2021). Evaluation of a multiplexed, multispecies nuclear receptor assay for chemical hazard assessment. *Toxicology In Vitro*, 72, 105016. <https://doi.org/10.1016/j.tiv.2020.105016>
- Howard, P. H., & Deo, P. G. (1979). Degradation of aryl phosphates in aquatic environments. *Bulletin Of Environmental Contamination And Toxicology*, 22(1), 337-344. <https://doi.org/10.1007/bf02026952>

- Huckins, J. N., Fairchild, J. F., & Boyle, T. P. (1991). Role of exposure mode in the bioavailability of triphenyl phosphate to aquatic organisms. *Archives Of Environmental Contamination And Toxicology*, 21(4), 481-485. <https://doi.org/10.1007/bf01183868>
- Isales, G. M., Hipszer, R. A., Raftery, T. D., Chen, A., Stapleton, H. M., & Volz, D. C. (2015). Triphenyl phosphate-induced developmental toxicity in zebrafish : Potential role of the retinoic acid receptor. *Aquatic Toxicology*, 161, 221-230. <https://doi.org/10.1016/j.aquatox.2015.02.009>
- Ishikawa S, Shigezumi K, Yasuda K & Shigemori N (1985a). Behaviors of organic phosphate esters in several water treatment processes. *Japan J Water Pollut Res (Suishitsu Odaku Kenkyo, Tokyo)* 8 (12): 799-807.
- Ishikawa S, Uchimura Y, Baba K, Eguchi Y, Kido K (1992). Photochemical Behavior of Organic Phosphate Esters in Aqueous Solutions Irradiated with a Mercury Lamp. *Bull. Environ. Contam. Tox.* 49: 368-374.
- Jenssen, B. M. (2006). Endocrine-Disrupting Chemicals and Climate Change : A Worst-Case Combination for Arctic Marine Mammals and Seabirds ? *Environmental Health Perspectives*, 114(Suppl 1), 76-80. <https://doi.org/10.1289/ehp.8057>
- Jia, Y., Zhou, H., Hu, W., Wang, L., Kang, Q., Liu, J., Nakanishi, T., Hiromori, Y., Kimura, T., Tao, S., & Hu, J. (2022). Discovery of contaminants with antagonistic activity against retinoic acid receptor in house dust. *Journal Of Hazardous Materials*, 426, 127847. <https://doi.org/10.1016/j.jhazmat.2021.127847>
- Ji, X., Li, N., Ma, M., Rao, K., & Wang, Z. (2020). In vitro estrogen-disrupting effects of organophosphate flame retardants. *Science Of The Total Environment*, 727, 138484. <https://doi.org/10.1016/j.scitotenv.2020.138484>
- Ji, X., Li, N., Ma, M., Li, X., Zhu, K., Rao, K., Wang, Z., Wang, J., & Fang, Y. (2022). Comparison of the mechanisms of estrogen disrupting effects between triphenyl phosphate (TPhP) and tris(1,3-dichloro-2-propyl) phosphate (TDCIPP). *Ecotoxicology And Environmental Safety*, 229, 113069. <https://doi.org/10.1016/j.ecoenv.2021.113069>
- Joint Research Centre (2013). Key scientific issues relevant to the identification and characterisation of endocrine disrupting substances. Report of the Endocrine Disrupters Expert Advisory Group. Joint Research Center. <https://op.europa.eu/en/publication-detail/-/publication/4b84ccc2-422d-4bd1-97da-1f414ad52c27/language-en>
- Kanazawa, A., Saito, I., Araki, A., Takeda, M., Ma, M., Saijo, Y., Kishi, R., (2010). Association between indoor exposure to semi-volatile organic compounds and building-related symptoms among the occupants of residential dwellings. *Indoor Air* 20, 72–84.
- Kawashima, Y., Onishi, Y., Tatarazako, N., Yamamoto, H., Koshio, M., Oka, T., Horie, Y., Watanabe, H., Nakamoto, T., Yamamoto, J., Ishikawa, H., Satō, T., Yamazaki, K., & Iguchi, T. (2021). Summary of 17 chemicals evaluated by OECD TG229 using Japanese Medaka, *Oryzias latipes* EXTEND 2016. *Journal Of Applied Toxicology*, 42(5), 750-777. <https://doi.org/10.1002/jat.4255>
- Kim, J.-W., Isobe, T., Sudaryanto, A., Malarvannan, G., Chang, K.-H., Muto, M., Prudente, M., Tanabe, S., (2013). Organophosphorus flame-retardants in house dust from the Philippines: occurrence and assessment of human exposure. *Environ. Sci. Pollut. Res.* 20, 812–822.
- Kim, J.W., Isobe, T., Muto, M., Tue, N.M., Katsura, K., Malarvannan, G., et al., (2014). Organophosphorus flame-retardants (pfrs) in human breast milk from several asian countries. *Chemosphere* 116, 91–97.

- Kim, Sujin, Joeun Jung, Inae Lee, Dawoon Jung, Hyewon Youn, and Kyungho Choi. (2015). Thyroid Disruption by Triphenyl Phosphate, an Organophosphate Flame Retardant, in Zebrafish (*Danio Rerio*) Embryos/Larvae, and in GH3 and FRTL-5 Cell Lines. *Aquatic Toxicology* 160 (March): 188–96. doi:10.1016/j.aquatox.2015.01.016.
- Kojima H, Takeuchi S, Itoh T, Iida M, Kobayashi S, Yoshida T. (2013). In vitro endocrine disruption potential of organophosphate flame-retardants via human nuclear receptors. *Toxicology* 314 (2013) 76– 83.
- Kojima H, Takeuchi S, Van den Eede N, Covaci A. (2016). Effects of primary metabolites of organophosphate flame retardants on transcriptional activity via human nuclear receptors. *Toxicology Letters* 245 (2016) 31–39.
- Krivoshiev B.V., Dardennea, F., Covacib, A., Blusta, R., Hussona, S.J., (2016): Assessing in vitro estrogenic effects of currently used flame retardants, *Toxicology in vitro*, 33, 153-162
- Kuehl DW & Haebler R (1995). Organochlorine, organobromine, metal, and selenium residues in bottlenose dolphins (*Tursiops truncatus*) collected during an unusual mortality event in the Gulf of Mexico, 1990. *Arch environ Contamin Tox* 28, 494 - 499.
- Lee, J. S., Y. K. Kawai, Y. Morita, A. Covaci, et A. Kubota. (2022). "Estrogenic and growth inhibitory responses to organophosphorus flame retardant metabolites in zebrafish embryos." *Comp Biochem Physiol C Toxicol Pharmacol* 256:109321. doi: 10.1016/j.cbpc.2022.109321.
- Li, Y., C. Wang, F. Zhao, S. Zhang, R. Chen, et J. Hu. (2018). "Environmentally Relevant Concentrations of the Organophosphorus Flame Retardant Triphenyl Phosphate Impaired Testicular Development and Reproductive Behaviors in Japanese Medaka (*Oryzias latipes*)." *Environmental Science and Technology Letters* 5 (11):649--654. doi: 10.1021/acs.estlett.8b00546.
- Li, N., Ho, W., Sun Wu, R.S., Ying, G.G., Wang, Z., Jones, K., et al., 2019a. Organophosphate flame-retardants and bisphenol a in children's urine in hong kong: has the burden been underestimated? *Ecotoxicol. Environ. Saf.* 183, 109502.
- Li, Y., R. Chen, J. He, H. Ma, F. Zhao, S. Tao, J. Liu, et J. Hu. 2019b. "Triphenyl Phosphate at Environmental Levels Retarded Ovary Development and Reduced Egg Production in Japanese Medaka (*Oryzias latipes*)." *Environmental Science and Technology* 53 (24):14709--14715.
- Liu, Xiaoshan, Kyungho Ji, and Kyungho Choi. (2012). Endocrine Disruption Potentials of Organophosphate Flame Retardants and Related Mechanisms in H295R and MVLN Cell Lines and in Zebrafish. *Aquatic Toxicology* 114–115: 173–81. doi:10.1016/j.aquatox.2012.02.019.
- Liu, Chunsheng, Qiangwei Wang, Kang Liang, Jingfu Liu, Bingsheng Zhou, Xiaowei Zhang, Hongling Liu, John P. Giesy, and Hongxia Yu. 2013a. "Effects of Tris(1,3-Dichloro-2-Propyl) Phosphate and Triphenyl Phosphate on Receptor-Associated mRNA Expression in Zebrafish Embryos/Larvae." *Aquatic Toxicology* 128–129 (March): 147–57. doi:10.1016/j.aquatox.2012.12.010.
- Liu, Xiaoshan, Kyungho Ji, Areum Jo, Hyo-Bang Moon, and Kyungho Choi. 2013b. "Effects of TDCPP or TPP on Gene Transcriptions and Hormones of HPG Axis, and Their Consequences on Reproduction in Adult Zebrafish (*Danio Rerio*)." *Aquatic Toxicology* 134–135: 104–11. doi:10.1016/j.aquatox.2013.03.013.

- Liu, Xiaoshan, Dawoon Jung, Areum Jo, Kyunghee Ji, Hyo-Bang Moon, and Kyungho Choi. 2016. "Long-Term Exposure to Triphenylphosphate Alters Hormone Balance and HPG, HPI, and HPT Gene Expression in Zebrafish (*Danio Rerio*).*" Environmental Toxicology and Chemistry 35 (9): 2288–96. doi:10.1002/etc.3395.*
- Liu, X., Y. Cai, Y. Wang, S. Xu, K. Ji, et K. Choi. (2019). "Effects of tris(1,3-dichloro-2-propyl) phosphate (TDCPP) and triphenyl phosphate (TPP) on sex-dependent alterations of thyroid hormones in adult zebrafish." *Ecotoxicol Environ Saf 170:25-32. doi: 10.1016/j.ecoenv.2018.11.058.*
- Lo CC and Hsieh TT (2000). Acute Toxicity to the Golden Apple Snail and Estimated Bioconcentration Potential of Triphenylphosphine Oxide and Series of Related Compounds. *Bull. Environ. Contam. Toxicol. 65: 104-111.*
- Lu, H., Hu, YC., Xu CQ., Dang W., (2021). "Developmental and liver metabolite changes induced by TPhP exposure in Brown frog (*Rana zhenhaiensis*) tadpoles." *Asian herpetological research 12(1): 135-142.*
- Marklund, A., Andersson, B., Haglund, P., 2003. Screening of organophosphorus compounds and their distribution in various indoor environments. *Chemosphere 53, 1137–1146.*
- Mayer FL, Adams WJ, Finley MT, Michael PR, Mehrle PM & Saeger VW (1981). Phosphate Ester Hydraulic Fluids: An Aquatic Environmental Assessment of Pydrauls 50E and 115E. *Aquatic Toxicology and Hazard Assessment, Fourth Conference. ASTM STP 737, D. R. Branson and K. L. Dickson, Eds.: 103-123.*
- Medvedev, A. V., et al. (2020). "Harmonized Cross-Species Assessment of Endocrine and Metabolic Disruptors by Ecotox FACTORIAL Assay." *Environmental Science & Technology 54(19): 12142-12153.*
- Millington LA, Goulding KH & Adams N (1988). The Influence of Growth Medium Composition on the Toxicity of Chemicals to Algae. *Water Res. 22: 1593-1597.*
- Muir DCG, Grift NP, Blouw AP & Lockhart WL (1980). Environmental dynamics of phosphate esters. 1. Uptake and bioaccumulation of triphenyl phosphate by Rainbow Trout. *Chemosphere 9: 525-532.*
- Muir DCG, Grift NP & Lockhart WL (1982). Comparison of laboratory and field results for prediction of the environmental behaviour of phosphate esters. *Environ. Toxicol. Chem. 1: 113-119.*
- Muir DCG, Yarechewski AL & Grift NP (1983). Environmental Dynamics of Phosphate Esters. III. Comparison of the Bioconcentration of Four Triaryl Phosphates by fish. *Chemosphere 12: 155-166.*
- OECD, 2018. Revised Guidance Document 150 on Standardised Test Guidelines for Evaluating Chemicals for Endocrine Disruption, OECD Series on Testing and Assessment, n° 150, Éditions OECD, Paris, <https://doi.org/10.1787/9789264304741-en>. OECD, 2021. *Detailed Review Paper on Retinoid Pathway Signalling*. Éditions OECD, Paris, [one.oecd.org/document/ENV/CBC/MONO\(2021\)20/en/pdf](https://one.oecd.org/document/ENV/CBC/MONO(2021)20/en/pdf)
- Onoja, S., Abdallah, M. A., & Harrad, S. (2023). Concentrations, spatial and seasonal variations of Organophosphate esters in UK freshwater Sediment. *Emerging Contaminants, 9(3), 100243. https://doi.org/10.1016/j.emcon.2023.100243*
- Palawski, D. U., Buckler, D. R., & Mayer, F. L. (1983). Survival and condition of rainbow trout (*Salmo gairdneri*) after acute exposures to methyl parathion, triphenyl phosphate, and DEF. *Bulletin Of Environmental Contamination And Toxicology, 30(1), 614-620. https://doi.org/10.1007/bf01610183*

- Pantelaki, I., & Voutsas, D. (2021). Organophosphate esters in inland and coastal waters in northern Greece. *Science Of The Total Environment*, 800, 149544. <https://doi.org/10.1016/j.scitotenv.2021.149544>
- Sala, B., Balasch, A., Eljarrat, E., & Cardona, L. (2021). First study on the presence of plastic additives in loggerhead sea turtles (*Caretta caretta*) from the Mediterranean Sea. *Environmental Pollution*, 283, 117108. <https://doi.org/10.1016/j.envpol.2021.117108>
- Sala, B., Giménez, J., Fernández-Arribas, J., Bravo, C., Lloret-Lloret, E., Esteban, A., Bellido, J. M., Coll, M., & Eljarrat, E. (2022). Organophosphate ester plasticizers in edible fish from the Mediterranean Sea : Marine pollution and human exposure. *Environmental Pollution*, 292, 118377. <https://doi.org/10.1016/j.envpol.2021.118377>
- Sasaki K, Takeda M & Uchiyama M (1981). Toxicity, Absorption and Elimination of Phosphoric Acid Triesters by Killifish and Goldfish. *Bull Environ Contam Toxicol* 27: 775 - 782.
- Sasaki, K. Suzuki T, Takeda M & Uchiyama M (1982). Bioconcentration and Excretion of Phosphoric Acid Triesters by Killifish (*Oryzeas latipes*). *Bull. Environ. Contam. Toxicol.* 28: 752-759.
- Schang G, Robaire B, Hales BF (2016): Organophosphate Flame Retardants Act as Endocrine-Disrupting Chemicals in MA-10 Mouse Tumor Leydig Cells. *Toxicological Sciences* 150(2), 499-509.
- Schulz R. W., de França L. R., Lareyre J. J., Le Gac F., Chiarini-Garcia H., Nobrega R. H., and Miura T. (2010). "Spermatogenesis in fish." *Gen Comp Endocrinol* 165 (3): 390-411. <https://doi.org/10.1016/j.ygcen.2009.02.013>.
- Sitthichai Kasem S (1978). Some toxicological effects of phosphate esters on Rainbow Trout and Bluegill. Iowa State University Dissertation. Testing laboratory: Dept. of Animal Ecology, Iowa State University.
- Sohn, J., Kim, S., Koschorreck, J., Kho, Y., Choi, K.(2016). Alteration of sex hormone levels and steroidogenic pathway by several low molecular weight phthalates and their metabolites in male zebrafish (*Danio rerio*) and/or human adrenal cell (h295r) line. *J. Hazard. Mater.* 320, 45–54.
- Stapleton, H.M., Klosterhaus, S., Eagle, S., Fuh, J., Meeker, J.D., Blum, A., Webster, T.F., (2009). Detection of organophosphate flame-retardants in furniture foam and US house dust. *Environ. Sci. Technol.* 43, 7490–7495.
- Sundkvist, A. et al. Organophosphorus flame-retardants and plasticizers in marine and fresh water biota and in human milk. *Journal of Environmental Monitoring*, 2010, 12(4), 943-951.
- Tachachartvanich, P., Sangsuwan, R., Ruiz, H.S., Sanchez, S.S., Durkin, K.A., Zhang, L., et al., 2018. Assessment of the endocrine-disrupting effects of trichloroethylene and its metabolites using in vitro and in silico approaches. *Environ. Sci. Technol.* 52, 1542–1550.
- Thomas, P. (2017). Role of G-protein-coupled estrogen receptor (GPER/GPR30) in maintenance of meiotic arrest in fish oocytes. *Journal of Steroid Biochemistry and Molecular Biology*. Vol. 167, p 153-161. <https://doi.org/10.1016/j.jsbmb.2016.12.005>
- Tubbs, C. W., and McDonough C. E. (2018). "Reproductive Impacts of Endocrine-Disrupting Chemicals on Wildlife Species: Implications for Conservation of Endangered Species." *Annu Rev Anim Biosci* 6: 287-304. <https://doi.org/10.1146/annurev-animal-030117-014547>.

- Unpublished study report (1982): Bio elimination of TPP in the Simulation Test - Aerobic Sewage, OECD Coupled Units Test No. 303A.
- Unpublished study report, 2000: Reproduction Test of Triphenyl Phosphate with *Daphnia magna*.
- Unpublished study report, 2002: Calculation of the environmental distribution of triphenyl phosphate acc. to Mackay Fugacity Model Level I, v2.11 (2001).
- Unpublished study report, 2002: Biologische Wikttests - polare Spurenstoffe.
- Unpublished study report, 2022. Zebrafish (*Danio rerio*), Sexual development Test.
- Van den Eede, N., Dirtu, A.C., Neels, H., Covaci, A., 2011. Analytical developments and preliminary assessment of human exposure to organophosphate flame retardants from indoor dust. *Environ. Int.* 37, 454–461.
- Wan HB and Wong MK (1994). Comparative study on the quantum yields of direct photolysis of organophosphorus pesticides in aqueous solution. *J. Agric Food Chem.* 42: 2625-2630.
- Wang, X., et al. (2021). "Integration of Computational Toxicology, Toxicogenomics Data Mining, and Omics Techniques to Unveil Toxicity Pathways." *ACS Sustainable Chemistry & Engineering* 9(11): 4130-4138.
- WHO/IPCS. 2002. *IPCS Global assessment of the state-of-the-science of endocrine disruptors.* edited by T. Damstra, Barlow, S., Bergman, A., Kavlock, R, Van der Kraak, G.
- Ya, M., Yu, N., Zhang, Y., Su, H., Tang, S., Su, G., (2019). Biomonitoring of organophosphate triesters and diesters in human blood in jiangsu province, eastern china: occurrences, associations, and suspect screening of novel metabolites. *Environ. Int.* 131, 105056.
- Zhang, Quan, Meiya Lu, Xiaowu Dong, Cui Wang, Chunlong Zhang, Weiping Liu, and Meirong Zhao. 2014. Potential Estrogenic Effects of Phosphorus-Containing Flame Retardants. *Environmental Science & Technology* 48 (12): 6995–7001. doi:10.1021/es5007862.
- Zhang, Q., et al. (2023). "Neurodevelopmental toxicity of organophosphate flame retardant triphenyl phosphate (TPhP) on zebrafish (*Danio rerio*) at different life stages." *Environment International* 172.
- Zhao, F., Wan, Y., Zhao, H., Hu, W., Mu, D., Webster, T.F., et al., (2016). Levels of blood organophosphorus flame retardants and association with changes in human sphingolipid homeostasis. *Environ. Sci. Technol.* 50, 8896–8903.
- Zhao, F.R., Chen, M., Gao, F.M., Shen, H., Hu, J.Y., (2017). Organophosphorus flame-retardants in pregnant women and their transfer to chorionic villi. *Environ. Sci. Technol.* 51, 6489–6497.
- Zheng, G.M., Schreder, E., Dempsey, J.C., Uding, N., Chu, V., Andres, G., et al., (2021). Organophosphate esters and their metabolites in breast milk from the United States: breastfeeding is an important exposure pathway for infants. *Environ. Sci. Tech. Lett.* 8, 224–230.