

Summary of product characteristics for a biocidal product

Product name: AGRI IODE FLUIDE

Product type(s): PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

Authorisation number: EU-0018497-0000

R4BP 3 asset reference number: EU-0018497-0005

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	2
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	3
2.1. Qualitative and quantitative information on the composition of the biocidal product	3
2.2. Type of formulation	3
3. Hazard and precautionary statements	3
4. Authorised use(s)	4
5. General directions for use	8
5.1. Instructions for use	8
5.2. Risk mitigation measures	8
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	9
5.4. Instructions for safe disposal of the product and its packaging	9
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	9
6. Other information	9

Administrative information

1.1. Trade names of the product

AGRI IODE FLUIDE
HYGIA IODE FLUIDE
IODE FLUIDE
IOSPRAY FLUIDE
PULVE IODE
IODO FL55
IODOLIN
ADIEL IODE FLUIDE
BOVI IODE SPRAY
PRO ROBO I
PROXA IODE FLUIDE
HELIODE FLUIDE
TECNO FLUIDE
DR15
GANAIODE FLUIDE
PLUSAIOD
PROTECTION FLUIDE IODE
BREIZH-IODE FLUIDE
BRETAIODE LQ
IODE NET FLUIDE
LACTOIODE FLUIDE
MAG IODE FLUIDE
NETAIODE FLUIDE
NETAIODE

1.2. Authorisation holder

**Name and address of the
authorisation holder**

Name	PRODHYNET SA
Address	11 avenue des châtelets 22440 PLOUFRAGAN France

Authorisation number

EU-0018497-0000 1-5

R4BP 3 asset reference number

EU-0018497-0005

Date of the authorisation

11/10/2018

Expiry date of the authorisation

30/09/2028

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer

PRODHYNET SA

Address of the manufacturer

11, Avenue des Châtelets 22440 PLOUFRAGAN France

Location of manufacturing sites

PRODHYNET SA - 11, Avenue des Châtelets 22440 PLOUFRAGAN France

1.4. Manufacturer(s) of the active substance(s)

Active substance

1319 - Iodine

Name of the manufacturer

COSAYACH : SCM Cía. Cosayach Minera Negreiros, Rut. N°96.625.710-5

Address of the manufacturer

Terrenos de Elena S/N Terrenos de Elena S/N Huara, Región de Tarapacá Chile

Location of manufacturing sites

Mined at : S.C.M. Cía. Minera Negreiros, S.C.M. Cosayach Soledad. Refined at : S.C.M. Cía. Minera Negreiros. Pozo Almonte Chile

Active substance	1319 - Iodine
Name of the manufacturer	ACF MINERA SA
Address of the manufacturer	San Martín 499 Iquique Chile
Location of manufacturing sites	Faena Lagunas KM. 1.722 Ruta A-5, Pozo Almonte Chile
Active substance	1319 - Iodine
Name of the manufacturer	SOCIEDAD QUIMICA y MINERA SA
Address of the manufacturer	Los Militares 4290 SANTIAGO DE CHILE Chile
Location of manufacturing sites	Pedro de Valdivia (PV) Route B 180 Antofagasta Chile
	Nueva Victoria (NV) Route 5 North, Km 1925 Pozo Almonte Chile

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Iodine		Active Substance	7553-56-2	231-442-4	0,55
Alcohols, C12-14, ethoxylated (11 mol EO average molar ratio)	Poly(oxy-1,2-ethanediyl), -C12-14-(even numbered)-alkyl--hydroxy		68439-50-9		4,69

2.2. Type of formulation

AL - Any other liquid

3. Hazard and precautionary statements

Hazard statements	Causes serious eye irritation. Harmful to aquatic life with long lasting effects.
--------------------------	--

Precautionary statements

Keep out of reach of children.

Wash hands thoroughly after handling.

Wear protective gloves.

Wear protective clothing.

Wear eye protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice.

Get medical

4. Authorised use(s)

4.1 Use description

Use 1 - Use # 5.1 – Manual or automated dipping, foaming or spraying before milking

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

-

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage:

Scientific name: Yeasts
Common name: Yeasts
Development stage:

Field(s) of use

Indoor

Disinfection of teats of milk producing animals by dipping, foaming or spraying before milking

Application method(s)

Dipping, foaming, spraying -

Manual or automated disinfection of teats by dipping, foaming or spraying before milking

Dipping cup, foaming cup, teat sprayer, automated dipping machine, automated foaming machine or automated spraying machine

Application rate(s) and frequencies

- cows and buffaloes (3 to 10ml : 5 to 8 ml recommended) - sheep (1.5 to 5 ml : 1.5 to 3 ml recommended) - goats (2.5 to 6 ml : 2.5 to 4 ml recommended) - -
Frequency : 2 to 3 times per day

Category(ies) of users

Professional

Pack sizes and packaging material

HDPE JERRYCAN 5, 10, 22 L
HDPE DRUM 60, 120, 220 L

HDPE CONTAINER 1000L

4.1.1 Use-specific instructions for use

Fill the dipping / foaming cup / sprayer manually or automatically with the ready to use product.

Eliminate all visible dirt before applying the product.

Manual or automatic dipping / foaming / spraying on animal's teats on the full length of the teat before milking.

Let the product act at least one minute.

Use the teat cleaning and wiping method systematically before attaching the milking cluster.

See also general instructions for use of meta SPC5.

4.1.2 Use-specific risk mitigation measures

For manual dipping/foaming application : Wear protective chemical resistant gloves (glove material to be specified by the authorisation holder within the product information)/ eye protection.
For manual spraying applications wear protective chemical resistant gloves (glove material to be specified by the authorisation holder within the product information)/protective clothing /eye protection.

In case a combination of pre- and post-milking disinfection is necessary, using another biocidal product not containing iodine has to be considered for post-milking disinfection.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of meta SPC 5

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of meta SPC 5

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of meta SPC 5

4.2 Use description

Use 2 - Use # 5.2 – Manual or automated dipping, foaming or spraying after milking

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

-

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage:

Scientific name: Yeasts
Common name: Yeasts
Development stage:

Scientific name: Algae
Common name: Algae
Development stage:

Scientific name: Viruses
Common name: Viruses
Development stage:

Field(s) of use

Indoor

Disinfection of teats of milk producing animals by dipping, foaming or spraying after milking

Application method(s)

Dipping, foaming, spraying -

Manual or automated disinfection of teats by dipping, foaming or spraying after milking. Dipping cup, foaming cup, teat sprayer, automated dipping machine, automated foaming machine or automated spraying machine.

Application rate(s) and frequencies

- cows and buffaloes (3 to 10ml : 5 to 8 ml recommended) - sheep (1.5 to 5 ml : 1.5 to 3 ml recommended) - goats (2.5 to 6 ml : 2.5 to 4 ml recommended) - 0 -
Frequency : 2 to 3 times per day

Category(ies) of users

Professional

Pack sizes and packaging material

HDPE JERRYCAN 5, 10, 22 L
HDPE DRUM 60, 120, 220 L

HDPE CONTAINER 1000L

4.2.1 Use-specific instructions for use

Fill the dipping / foaming cup / sprayer manually or automatically with the ready to use product.

Manual or automatic dipping / foaming / spraying on animal's teats on the full length of the teat after milking.

Leave the product until next milking. Keep the cows standing until the product has dried. (at least 5 minutes)

At the next milking, use the teat cleaning and wiping method systematically before attaching the milking cluster.

See also general instructions for use of meta SPC 5.

4.2.2 Use-specific risk mitigation measures

For manual spraying applications wear protective chemical resistant gloves (glove material to be specified by the authorisation holder within the product information)/protective clothing/eye protection.

In case a combination of pre- and post-milking disinfection is necessary, using another biocidal product not containing iodine has to be considered for pre-milking disinfection.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of meta SPC 5

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of meta SPC 5

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of meta SPC 5

5. General directions for use

5.1. Instructions for use

See use-specific instructions for each use.

Always read the label or leaflet before use and follow all the instructions provided.
The product must be brought to a temperature above 20°C before use.
The use of a dosing pump for filling the product into the application equipment is recommended.
Where necessary repeat the application at each milking
Clean the application equipment regularly with warm water.

5.2. Risk mitigation measures

See use-specific risk mitigation measures for each use.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Take the contaminated clothes and shoes off immediately. Wash them before wearing them again.

FIRST AID INSTRUCTIONS

- In the event of inhalation : Bring to fresh air.

- In the event of contact with the skin : Wash with water.

- In the event of contact with the eyes :

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

- In the event of ingestion : Rinse mouth. Do NOT induce vomiting. Get medical advice.

Refer to the safety data sheet available for professional user : it contains an emergency phone number.

Large spillage : Mark out, dyke up with an inert absorbant and pump in an emergency tank. Keep in suitable, properly labelled and closed containers for disposal. Never return spills in original containers for re-use.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose unused product and the packaging in accordance with local requirements. Used product can be flushed to the municipal sewer or disposed to the manure deposit depending on local requirements. Avoid release to an individual waste water treatment plant.

The paper towels used for removing the product and drying the teats are disposed in the normal and domestic waste.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life : 2 years in HDPE

Do not store at a temperature higher than 30°C.

6. Other information