

Summary of product characteristics for a biocidal product

Product name: calgodip D 1200

Product type(s): PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

Authorisation number: EU-0018724-0000

R4BP 3 asset reference number: EU-0018724-0008

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	3
2.1. Qualitative and quantitative information on the composition of the biocidal product	3
2.2. Type of formulation	3
3. Hazard and precautionary statements	3
4. Authorised use(s)	4
5. General directions for use	15
5.1. Instructions for use	15
5.2. Risk mitigation measures	15
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	15
5.4. Instructions for safe disposal of the product and its packaging	15
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	16
6. Other information	16

Administrative information

1.1. Trade names of the product

calgodip D 1200
Jod-Dip S 12
Dip es SF 1200

1.2. Authorisation holder

Name and address of the authorisation holder	Name	CVAS Development GmbH
	Address	Dr. Albert Reimann Str. 16 a 68526 Ladenburg Germany
Authorisation number	EU-0018724-0000 1-7	

R4BP 3 asset reference number	EU-0018724-0008
Date of the authorisation	18/12/2018
Expiry date of the authorisation	30/11/2028

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	Calvatis GmbH
Address of the manufacturer	Dr. Albert Reimann Str. 16a 68526 Ladenburg Germany
Location of manufacturing sites	Dr. Albert Reimann Str. 16a 68526 Ladenburg Germany

1.4. Manufacturer(s) of the active substance(s)

Active substance	1319 - Iodine
Name of the manufacturer	Cosayach Nitratos S.A.
Address of the manufacturer	Amunategui 178 not applicable Santiago Chile
Location of manufacturing sites	S.C.M. Cosayach Cala Cala not applicable Pozo Almonte Chile

Active substance	1319 - Iodine
Name of the manufacturer	ACF Minera S.A.
Address of the manufacturer	San Martin No 499 not applicable Iquique Chile
Location of manufacturing sites	Lagunas mine not applicable Pozo Almonte Chile

Active substance	1319 - Iodine
Name of the manufacturer	SQM S.A.
Address of the manufacturer	Los Militares 4290, Piso 4 not applicable Las Condes Chile
Location of manufacturing sites	Nueva Victoria plant not applicable Pedro de Valdivia plant Chile

Active substance	1319 - Iodine
Name of the manufacturer	Nihon Tennen Gas Co., Ltd / Kanto Natural Gas Development Co., Ltd
Address of the manufacturer	661 Mobara 297-8550 Mobara City, Chiba Japan
Location of manufacturing sites	2508 Minami-Hinata 299-4205 Shirako-Machi, Chosei-Gun, Chiba Japan

Active substance	1349 - Polyvinylpyrrolidone iodine
Name of the manufacturer	Norkem Limited
Address of the manufacturer	Norkem House, Bexton Lane WA 16 9FB Knutsford, Cheshire United Kingdom
Location of manufacturing sites	Norkem House, Bexton Lane WA 16 9FB Knutsford, Cheshire United Kingdom

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Iodine		Active Substance	7553-56-2	231-442-4	0,14
Polyvinylpyrrolidone iodine		Active Substance	25655-41-8		0
Acetic acid	Acetic acid		64-19-7	200-580-7	0,15

2.2. Type of formulation

AL - Any other liquid

3. Hazard and precautionary statements

Hazard statements	Harmful to aquatic life with long lasting effects.
Precautionary statements	Avoid release to the environment. Dispose of contents to local/regional/national/international regulation. Dispose of container to local/regional/national/international regulation.

4. Authorised use(s)

4.1 Use description

Use 1 - Use #7.1 - Teat disinfection of milkable animals: Post-milking teat disinfection by manual dipping

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	not relevant
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: vegetative cells Scientific name: Yeasts Common name: Yeasts Development stage: vegetative cells
Field(s) of use	Indoor Teat disinfection for milkable animals (dairy cows) for use after milking
Application method(s)	Manual dipping using a dip cup - See instructions for use
Application rate(s) and frequencies	cows: 5 mL per treatment - 0% - Post-milking application: 2-3x/day (after each milking)
Category(ies) of users	Professional
Pack sizes and packaging material	Jerrycan (HDPE): 5 – 60 kg Drum (HDPE): 60 – 200 kg IBC (HDPE): 600 - 1000 kg

4.1.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
The use of a dosing pump for filling the product into the application equipment is recommended.
Fill the reservoir with the RTU product assuming 5 mL product per cow and screw the dip cup on top. Avoid discharge of surplus fluids.
Clean the teats carefully by wiping with a single service paper towel/cloth immediately before milking. After milking, squeeze the reservoir and put the dip cup over each teat from below making sure that about 3 cm of the teat are immersed into the disinfectant.
Refill the cup of the dipping unit with fresh disinfectant by squeezing the reservoir as needed. Refill the reservoir with fresh

disinfectant as needed.
Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.
After disinfection, empty the reservoir and clean reservoir and dip cup by rinsing with water.

4.1.2 Use-specific risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use.

4.2 Use description

Use 2 - Use #7.2 - Teat disinfection of milkable animals: Post-milking teat disinfection by manual foaming

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

not relevant

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage: vegetative cells

Scientific name: Yeasts
Common name: Yeasts
Development stage: vegetative cells

Field(s) of use

Indoor

Teat disinfection for milkable animals (dairy cows) for use after milking

Application method(s)	Manual foaming using a foam cup - See instructions for use
Application rate(s) and frequencies	Cows: 5 mL per treatment - 0% - Post-milking application: 2-3x/day (after each milking)
Category(ies) of users	Professional
Pack sizes and packaging material	Jerrycan (HDPE): 5 – 60 kg Drum (HDPE): 60 – 200 kg IBC (HDPE): 600 - 1000 kg

4.2.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
The use of a dosing pump for filling the product into the application equipment is recommended.
Fill the reservoir with the RTU product assuming 5 mL product per cow and screw the foam cup on top. Avoid discharge of surplus fluids.
Clean the teats carefully by wiping with a single service paper towel/cloth immediately before milking.
After milking, squeeze the reservoir and put the foam cup over each teat from below making sure that about 3 cm of the teat are immersed into the disinfectant.
Refill the cup of the foaming unit with fresh disinfectant by squeezing the reservoir as needed. Refill the reservoir with fresh disinfectant as needed.
Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.
After disinfection, empty the reservoir and clean reservoir and foam cup by rinsing with water.

4.2.2 Use-specific risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use.

4.3 Use description

Use 3 - Use #7.3 - Teat disinfection of milkable animals: Post-milking teat disinfection by manual spraying using a trigger sprayer

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	not relevant
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: vegetative cells Scientific name: Yeasts Common name: Yeasts Development stage: vegetative cells
Field(s) of use	Indoor Teat disinfection for milkable animals (dairy cows) for use after milking
Application method(s)	Manual spraying using a trigger sprayer - See instructions for use
Application rate(s) and frequencies	Cows: 5 mL per treatment - 0% - Post-milking application: 2-3x/day (after each milking)
Category(ies) of users	Professional
Pack sizes and packaging material	Jerrycan (HDPE): 5 – 60 kg Drum (HDPE): 60 – 200 kg IBC (HDPE): 600 - 1000 kg

4.3.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
The use of a dosing pump for filling the product into the application equipment is recommended.

Fill the reservoir with the RTU product assuming 5 mL product per cow and screw the top of the trigger sprayer on it. Avoid discharge of surplus fluids.

Clean the teats carefully by wiping with a single service paper towel/cloth immediately before milking.

After milking, spray the disinfectant on the teats using the trigger sprayer making sure that about 3 cm of the teat around the streak canal are covered with the disinfectant.

Refill the reservoir of the trigger sprayer with fresh disinfectant as needed.

Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.

After disinfection, empty the reservoir and clean reservoir and trigger sprayer by rinsing with water.

4.3.2 Use-specific risk mitigation measures

Use chemical resistant gloves (glove material to be specified by the authorisation holder within the product information) during post-milking teat disinfection by manual spraying using a trigger sprayer.

Avoid working in spray mist.

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use.

4.4 Use description

Use 4 - Use #7.4 - Teat disinfection of milkable animals: Post-milking teat disinfection by manual spraying using an electronic sprayer

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

not relevant

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage: vegetative cells

	<p>Scientific name: Yeasts Common name: Yeasts Development stage: vegetative cells</p>
Field(s) of use	<p>Indoor</p> <p>Teat disinfection for milkable animals (dairy cows) for use after milking</p>
Application method(s)	<p>Manual spraying using an electronic sprayer - See instructions for use</p>
Application rate(s) and frequencies	<p>cows: 5 mL per treatment - 0% - Post-milking application: 2-3x/day (after each milking)</p>
Category(ies) of users	<p>Professional</p>
Pack sizes and packaging material	<p>Jerrycan (HDPE): 5 – 60 kg Drum (HDPE): 60 – 200 kg IBC (HDPE): 600 - 1000 kg</p>

4.4.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
The use of a dosing pump for filling the product into the application equipment is recommended.
Open a can containing the RTU product assuming 5 mL product per cow and insert a sucking lance of the electronic sprayer. Avoid discharge of surplus fluids.
Clean carefully the teats by wiping with a single service paper towel/cloth before milking.
After milking, spray the disinfectant on the teats using the electronic sprayer making sure that about 3 cm of the teat around the streak canal are covered with the disinfectant.
Replace the empty can by a new can containing the RTU product as needed.
Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.
After disinfection, put the sucking lance system into a bucket of water and rinse the sprayer by pumping the water through the sprayer.

4.4.2 Use-specific risk mitigation measures

Use chemical resistant gloves (glove material to be specified by the authorisation holder within the product information) during post/milking teat disinfection by manual spraying using an electronic sprayer.
Avoid working in spray mist.
In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use.

4.5 Use description

Use 5 - Use #7.5 - Teat disinfection of milkable animals: Post-milking teat disinfection by automated dipping

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	not relevant
Target organism(s) (including development stage)	<p>Scientific name: Bacteria Common name: Bacteria Development stage: vegetative cells</p> <p>Scientific name: Yeasts Common name: Yeasts Development stage: vegetative cells</p>
Field(s) of use	<p>Indoor</p> <p>Teat disinfection for milkable animals (dairy cows) for use after milking</p>
Application method(s)	<p>Automated dipping -</p> <p>See instructions for use</p>
Application rate(s) and frequencies	<p>cows: 5 mL per treatment - 0% - Post-milking application: 2 - 3x/day (after each milking)</p>

Category(ies) of users

Professional

Pack sizes and packaging material

Jerrycan (HDPE): 5 – 60 kg
Drum (HDPE): 60 – 200 kg
IBC (HDPE): 600 - 1000 kg

4.5.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
Open a can containing the RTU product and insert a suction tube of the automated dipping-system. Avoid discharge of surplus fluids. After milking, the vacuum is shut off and the teat dip is injected into a manifold on the clawpiece. The teats are coated with ca. 5 mL of dip when the teat cup is withdrawn by the Automatic Cluster Removal (ACR). After the removal of the ACR, every liner of the automated dipping-system is thoroughly rinsed with water and blown out with compressed air.
In a final cleaning step after each milking session of the herd, the liners are disinfected (e.g. with a chlorine-based product) and blown out again with compressed air.
Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.
Afterwards, the milking system is ready for the next milking event.
The whole process is automated.

4.5.2 Use-specific risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use.

4.6 Use description

Use 6 - Use #7.6 - Teat disinfection of milkable animals: Post-milking teat disinfection by automated foaming

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	not relevant
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: vegetative cells Scientific name: Yeasts Common name: Yeasts Development stage: vegetative cells
Field(s) of use	Indoor Teat disinfection for milkable animals (dairy cows) for use after milking
Application method(s)	Automated foaming - see instructions for use
Application rate(s) and frequencies	cows: 5 mL per treatment - 0% - Post-milking application: 2 - 3x/day (after each milking)
Category(ies) of users	Professional
Pack sizes and packaging material	Jerrycan (HDPE): 5 – 60 kg Drum (HDPE): 60 – 200 kg IBC (HDPE): 600 - 1000 kg

4.6.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
Open a can containing the RTU product and insert a suction tube of the automated foaming-system. Avoid discharge of surplus fluids.

After milking, the vacuum is shut off and the teat disinfectant is injected into a manifold on the clawpiece. The teats are coated with ca. 5 mL of foam when the teat foam cup is withdrawn by the Automatic Cluster Removal (ACR). After the removal of the ACR, every liner of the automated foaming-system is thoroughly rinsed with water and blown out with compressed air.
 In a final cleaning step after each milking session of the herd, the liners are disinfected (e.g. with a chlorine-based product) and blown out again with compressed air.
 Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.
 Afterwards, the milking system is ready for the next milking event.
 The whole process is automated.

4.6.2 Use-specific risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use (section 5).

4.7 Use description

Use 7 - Use #7.7 - Teat disinfection of milkable animals: Post-milking teat disinfection by automated spraying by robot

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

not relevant

Target organism(s) (including development stage)

Scientific name: Bacteria
 Common name: Bacteria
 Development stage: vegetative cells

Scientific name: Yeasts
 Common name: Yeasts
 Development stage: vegetative cells

Field(s) of use	Indoor Teat disinfection for milkable animals (dairy cows) for use after milking
Application method(s)	Automated spraying by robot - See instructions for use
Application rate(s) and frequencies	cows: 5 mL per treatment - 0% - Post-milking application: 2 - 3x/day (after each milking)
Category(ies) of users	Professional
Pack sizes and packaging material	Jerrycan (HDPE): 5 – 60 kg Drum (HDPE): 60 – 200 kg IBC (HDPE): 600 - 1000 kg

4.7.1 Use-specific instructions for use

The product must be brought to a temperature above 20°C before use.
Open a can containing the RTU product and insert a suction tube of the robotic milking device. Avoid discharge of surplus fluids.
The teats are cleaned by robot with automatic brushes.
After robotic milking, the disinfectant is sprayed automatically onto teats from a cluster arm.
Leave the product on the teats until next milking. Keep the animals standing for at least 5 minutes after treatment.
Rinsing of the sprayer is automatic.

4.7.2 Use-specific risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

4.7.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use.

4.7.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use.

4.7.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use.

5. General directions for use

5.1. Instructions for use

See use specific instructions for use.

5.2. Risk mitigation measures

See use specific risk mitigation measures.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Mentioned in the MSDS

Description of first aid measures

After inhalation: Supply fresh air; consult doctor in case of symptoms.

After skin contact: Wash with water and soap.

After eye contact: Rinse opened eye for several minutes under running water. Then consult doctor.

After swallowing: Rinse out mouth and then drink plenty of water. Seek immediate medical advice.

Stability and reactivity

Possibility of hazardous reactions: Reaction with oxidant- and reducing agent.

Conditions to avoid: No further relevant information available.

Incompatible materials: No further relevant information available.

Hazardous decomposition products: iodine (when warming up).

Accidental release measures

Personal precautions, protective equipment and emergency procedures: No special measures required.

Environmental precautions: Do not allow product to reach sewage systems or water bodies in great quantities.

Methods and material for containment and cleaning up: Absorb with liquid-binding material (sand, diatomite, acid binders, universal binders, sawdust). Dispose of the material collected according to regulations.

5.4. Instructions for safe disposal of the product and its packaging

Mentioned in the MSDS
Waste treatment methods
Recommendation: Must be specially treated with regard to official regulations.
Waste disposal key number: Corresponding to the regulation of the European Waste catalogue the relation of the waste key numbers has to be made specific to industry and process.
European waste catalogue: Corresponding to the regulation of the EWC the relation of the waste key numbers has to be made specific to industry and process.

At the end of the treatment, dispose unused product and the packaging in accordance with local requirements. Used product can be flushed to the municipal sewer or disposed to the manure deposit depending on local requirements. Avoid release to an individual waste water treatment plant.
Recommended cleaning agent: Water, if necessary with cleaning agent.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Shelf-life: 24 months
Products need to be protected from frost, stored at temperatures not exceeding 30°C and away from direct sunlight.

6. Other information

Not provided