

Summary of product characteristics for a biocidal product family

Family name: FAMILY SALVESAFE E

Product type(s): PT03 - Veterinary hygiene (Disinfectants)

Authorisation number: EU-0020512-0000

R4BP 3 asset reference number: EU-0020512-0000

Table Of Contents

Part I.- First information level	1
1. Administrative information	1
2. Product family composition and formulation	3
Part II.- Second information level - meta SPC(s)	3
1. Meta SPC administrative information - Pre-milking teat disinfection	4
2. Meta SPC composition	4
3. Hazard and precautionary statements of the meta SPC	4
4. Authorised use(s) of the meta SPC	5
5. General directions for use of the meta SPC	9
6. Other information	10
7. Third information level: individual products in the meta SPC	10
1. Meta SPC administrative information - Pre-milking and post-milking teat disinfection	12
2. Meta SPC composition	12
3. Hazard and precautionary statements of the meta SPC	13
4. Authorised use(s) of the meta SPC	13
5. General directions for use of the meta SPC	18
6. Other information	19
7. Third information level: individual products in the meta SPC	19
1. Meta SPC administrative information - Post-milking teat disinfection	21
2. Meta SPC composition	22
3. Hazard and precautionary statements of the meta SPC	22
4. Authorised use(s) of the meta SPC	22
5. General directions for use of the meta SPC	27
6. Other information	28
7. Third information level: individual products in the meta SPC	28

Part I.- First information level

1. Administrative information

1.1. Family name

FAMILY SALVESAFE E

1.2. Product type(s)

PT03 - Veterinary hygiene (Disinfectants)

1.3. Authorisation holder

Name and address of the authorisation holder

Name	SALVECO S.A.S.
Address	Avenue Pierre Mendès-France Z.A. Hellieule 4 F 88100 SAINT DIE DES VOSGES France

Authorisation number

EU-0020512-0000

R4BP 3 asset reference number

EU-0020512-0000

Date of the authorisation

28/08/2019

Expiry date of the authorisation

28/08/2029

1.4. Manufacturer(s) of the biocidal products

Name of the manufacturer

SALVECO S.A.S

Address of the manufacturer

Avenue Pierre Mendès-France, Z.I. Hellieule 4 F-88100 SAINT DIE DES VOSGES France

Location of manufacturing sites

Avenue Pierre Mendès-France, Z.I. Hellieule 4 F-88100 SAINT DIE DES VOSGES France

Name of the manufacturer	MULTIFILL BV BULKONT VANGST
Address of the manufacturer	CONSTRUCTIEWEG 25A 3641 SB MIJDRECHT Netherlands
Location of manufacturing sites	CONSTRUCTIEWEG 25A 3641 SB MIJDRECHT Netherlands

Name of the manufacturer	Diversey Netherlands Production BV
Address of the manufacturer	Rembrandtlaan 414 7545 ZW Enschede Netherlands
Location of manufacturing sites	Rembrandtlaan 414 7545 ZW Enschede Netherlands

Name of the manufacturer	Diversey UK Production Ltd
Address of the manufacturer	Cotes Park Industrial Estate, Somercotes DE55 4PA Alfreton United Kingdom
Location of manufacturing sites	Cotes Park Industrial Estate, Somercotes DE55 4PA Alfreton United Kingdom

Name of the manufacturer	Diversey España Production S.L.U
Address of the manufacturer	Avenida Conde Duque 5, 7 y 9, Poligono Industrial La Postura 28343 Valdemoro (Madrid) Spain
Location of manufacturing sites	Avenida Conde Duque 5, 7 y 9, Poligono Industrial La Postura 28343 Valdemoro (Madrid) Spain

Name of the manufacturer	Diversey Italy Production Srl
Address of the manufacturer	Strada Statale 235 26010 Bagnolo Cremasco (CR) Italy
Location of manufacturing sites	Strada Statale 235 26010 Bagnolo Cremasco (CR) Italy

Name of the manufacturer	Diversey Germany Production oHG
Address of the manufacturer	Morschheimer Strasse 12 67292 Kirchheimbolanden Germany
Location of manufacturing sites	Morschheimer Strasse 12 67292 Kirchheimbolanden Germany

1.5. Manufacturer(s) of the active substance(s)

Active substance	1421 - Lactic acid
Name of the manufacturer	JUNGBUNZLAUER S.A
Address of the manufacturer	Z. I Portuaire BP 32 67390 Marckolsheim France
Location of manufacturing sites	Z. I Portuaire BP 32 67390 Marckolsheim France

2. Product family composition and formulation

2.1. Qualitative and quantitative information on the composition of the family

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38 - 2,38

2.2. Type(s) of formulation

AL - Any other liquid

Part II.- Second information level - meta SPC(s)

1. Meta SPC administrative information

1.1. Meta SPC identifier

Pre-milking teat disinfection

1.2. Suffix to the authorisation number

1-1

1.3 Product type(s)

PT03 - Veterinary hygiene (Disinfectants)

2. Meta SPC composition

2.1. Qualitative and quantitative information on the composition of the meta SPC

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38 - 2,38

2.2. Type(s) of formulation of the meta SPC

Formulation(s)

AL - Any other liquid

3. Hazard and precautionary statements of the meta SPC

Hazard statements

Safety data sheet available on request.

Precautionary statements

4. Authorised use(s) of the meta SPC

4.1 Use description

Use 1 - Ready-to-use non-medical pre-milking teat disinfection - spraying

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: Scientific name: Yeasts Common name: Yeasts Development stage:
Field(s) of use	Indoor Outdoor Pre-milking disinfection of teats of milk producing animals
Application method(s)	Method: Spraying Detailed description: Manual and automated spraying
Application rate(s) and frequencies	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking applications per day
Category(ies) of users	Professional
Pack sizes and packaging material	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

4.1.1 Use-specific instructions for use

4.1.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill a manual or automate spraying device with ready to use product from the original packaging and close with spraying head. Apply the product by spray to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk.

4.1.2 Use-specific risk mitigation measures

Not applicable

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

4.2 Use description

Use 2 - Ready-to-use non-medical pre-milking teat disinfection - foaming

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

-

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage:

Scientific name: Yeasts
Common name: Yeasts
Development stage:

Field(s) of use	Indoor Outdoor Pre-milking disinfection of teats of milk producing animals
Application method(s)	Method: Foam application Detailed description: Manual and automated foaming
Application rate(s) and frequencies	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking applications per day
Category(ies) of users	Professional
Pack sizes and packaging material	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

4.2.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill the foaming cup with ready to use product from the original packaging. Apply the product to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk.

4.2.2 Use-specific risk mitigation measures

Not applicable

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

4.3 Use description

Use 3 - Ready-to-use non-medical pre-milking teat disinfection - dipping

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: Scientific name: Yeasts Common name: Yeasts Development stage:
Field(s) of use	Indoor Outdoor Pre-milking disinfection of teats of milk producing animals
Application method(s)	Method: Dipping Detailed description: Manual and automated dipping
Application rate(s) and frequencies	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking applications per day
Category(ies) of users	Professional
Pack sizes and packaging material	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L

IBC, Plastic: HDPE 1000L

4.3.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill the manual or automated dipping cup with ready to use product from the original packaging. Apply the product to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk.

4.3.2 Use-specific risk mitigation measures

Not applicable

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

5. General directions for use of the meta SPC

5.1. Instructions for use

Please refer to use-specific instructions for use

5.2. Risk mitigation measures

Not applicable.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

In case of eye contact: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

In case of skin contact: Wash with plenty of water.

If swallowed: Rinse mouth. Do not induce vomiting. If symptoms occurs: Get medical advice/attention.

If inhaled: Get medical attention if symptoms occur.

5.4. Instructions for safe disposal of the product and its packaging

Dispose of contents/container to in accordance with national regulation.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Store in original container at room temperature. Protect from direct sun light. Protect from frost. Shelf-life: 2 years.

6. Other information

7. Third information level: individual products in the meta SPC

7.1 Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)

SALVESAFE E1

Market area: EU

Authorisation number

EU-0020512-0001 1-1

(R4BP 3 asset reference number - National Authorisation)

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E2

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0002 1-1

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E19

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0003 1-1

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

1. Meta SPC administrative information

1.1. Meta SPC identifier

Pre-milking and post-milking teat disinfection

1.2. Suffix to the authorisation number

1-2

1.3 Product type(s)

PT03 - Veterinary hygiene (Disinfectants)

2. Meta SPC composition

2.1. Qualitative and quantitative information on the composition of the meta SPC

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38 - 2,38

2.2. Type(s) of formulation of the meta SPC

Formulation(s)

AL - Any other liquid

3. Hazard and precautionary statements of the meta SPC

Hazard statements

Safety data sheet available on request.

Precautionary statements

4. Authorised use(s) of the meta SPC

4.1 Use description

Use 1 - Ready-to-use non-medical pre-milking and post-milking teat disinfection - spraying

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

-

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage:

Scientific name: Yeasts
Common name: Yeasts
Development stage:

Field(s) of use

Indoor
Outdoor

Application method(s)

Method: Spraying
Detailed description:
Manual and automated spraying

Application rate(s) and frequencies

Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface.
Dilution (%): Ready-to-use product
Number and timing of application:
Maximum 3 pre-milking and 3 post-milking applications per day

Category(ies) of users

Professional

Pack sizes and packaging material

Bottle, Plastic: HDPE 1L
Jerrycan, Plastic: HDPE 5 - 60L
Drum, Plastic: HDPE 220 L
IBC, Plastic: HDPE 1000L

4.1.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill a manual or automate spraying device with ready to use product from the original packaging and close with spraying head.
Pre-milking
Apply the product by spray to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk.
Post-milking
The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

4.1.2 Use-specific risk mitigation measures

Not applicable

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

4.2 Use description

Use 2 - Ready-to-use non-medical pre-milking and post-milking teat disinfection - foaming

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: Scientific name: Yeasts Common name: Yeasts Development stage:
Field(s) of use	Indoor Outdoor
Application method(s)	Method: Foam application Detailed description: Manual and automated foaming
Application rate(s) and frequencies	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking and 3 post-milking applications per day
Category(ies) of users	Professional
Pack sizes and packaging material	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

4.2.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill the foaming cup with ready to use product from the original packaging.
Pre-milking
Apply the product to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk. Leave the teats skin dry.
Post-milking
The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals

standing or walking at least 5 minutes.

4.2.2 Use-specific risk mitigation measures

Not applicable

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

4.3 Use description

Use 3 - Ready-to-use non-medical pre-milking and post-milking teat disinfection - dipping

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: Scientific name: Yeasts Common name: Yeasts Development stage:
Field(s) of use	Indoor Outdoor

Application method(s)	<p>Method: Dipping Detailed description: Manual and automated dipping</p>
Application rate(s) and frequencies	<p>Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking and 3 post-milking applications per day</p>
Category(ies) of users	<p>Professional</p>
Pack sizes and packaging material	<p>Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L</p>

4.3.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill the manual or automated dipping cup with ready to use product from the original packaging.

Pre-milking
Apply the product to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk. Leave the teats skin dry.

Post-milking
The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

4.3.2 Use-specific risk mitigation measures

Not applicable

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

5. General directions for use of the meta SPC

5.1. Instructions for use

Please refer to use-specific instructions for use

5.2. Risk mitigation measures

Not applicable.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

In case of eye contact: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

In case of skin contact: Wash with plenty of water.

If swallowed: Rinse mouth. Do not induce vomiting. If symptoms occurs: Get medical advice/attention.

If inhaled: Get medical attention if symptoms occur.

5.4. Instructions for safe disposal of the product and its packaging

Dispose of contents/container to in accordance with national regulation.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Store in original container at room temperature. Protect from direct sun light. Protect from frost. Shelf-life: 2 years.

6. Other information

N/A

7. Third information level: individual products in the meta SPC

7.1 Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)

SALVESAFE E7	Market area: EU
DEOSAN Triathlon	Market area: EU
Mole Valley LACTOCIDE	Market area: EU
Authorisation number <small>(R4BP 3 asset reference number - National Authorisation)</small>	EU-0020512-0004 1-2

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E8

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0005 1-2

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E20

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0006 1-2

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E22

Market area: EU

OSANIS PRO Ready-to-use disinfectant for teat

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0010 1-2

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

1. Meta SPC administrative information**1.1. Meta SPC identifier**

Post-milking teat disinfection

1.2. Suffix to the authorisation number

1-3

1.3 Product type(s)

PT03 - Veterinary hygiene (Disinfectants)

2. Meta SPC composition

2.1. Qualitative and quantitative information on the composition of the meta SPC

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38 - 2,38

2.2. Type(s) of formulation of the meta SPC

Formulation(s)

AL - Any other liquid

3. Hazard and precautionary statements of the meta SPC

Hazard statements

Safety data sheet available on request.

Precautionary statements

4. Authorised use(s) of the meta SPC

4.1 Use description

Use 1 - Ready-to-use non-medical post-milking teat disinfection - spraying

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

-

Target organism(s) (including development stage)

Scientific name: Bacteria
Common name: Bacteria
Development stage:

Scientific name: Yeasts
Common name: Yeasts
Development stage:

Field(s) of use

Indoor

Outdoor

Application method(s)

Method: Spraying
Detailed description:
Manual and automated spraying

Application rate(s) and frequencies

Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface.
Dilution (%): Ready-to-use product.
Number and timing of application:
Maximum 3 post-milking application per day

Category(ies) of users

Professional

Pack sizes and packaging material

Bottle, Plastic: HDPE 1L
Jerrycan, Plastic: HDPE 5 - 60L
Drum, Plastic: HDPE 220 L
IBC, Plastic: HDPE 1000L

4.1.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill a manual or automate spraying device with ready to use product from the original packaging and close with spraying head. The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

4.1.2 Use-specific risk mitigation measures

Not applicable

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

4.2 Use description

Use 2 - Ready-to-use non-medical post-milking teat disinfection - foaming

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: Scientific name: Yeasts Common name: Yeasts Development stage:
Field(s) of use	Indoor Outdoor
Application method(s)	Method: Foam application Detailed description: Manual and automated foaming
Application rate(s) and frequencies	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface.

	Dilution (%): Ready-to-use product. Number and timing of application: Maximum 3 post-milking applications per da
Category(ies) of users	Professional
Pack sizes and packaging material	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

4.2.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill the foaming cup with ready to use product from the original packaging. The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

4.2.2 Use-specific risk mitigation measures

Not applicable

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

4.3 Use description

Use 3 - Ready-to-use non-medical post-milking teat disinfection - dipping

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: Bacteria Common name: Bacteria Development stage: Scientific name: Yeasts Common name: Yeasts Development stage:
Field(s) of use	Indoor Outdoor
Application method(s)	Method: Dipping Detailed description: Manual and automated dipping
Application rate(s) and frequencies	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product. Number and timing of application: Maximum 3 post-milking applications per day
Category(ies) of users	Professional
Pack sizes and packaging material	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

4.3.1 Use-specific instructions for use

Before use, bring the product at least 20°C.
Fill the manual or automated dipping cup with ready to use product from the original packaging. The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

4.3.2 Use-specific risk mitigation measures

Not applicable

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

5. General directions for use of the meta SPC

5.1. Instructions for use

Please refer to use-specific instructions for use

5.2. Risk mitigation measures

Not applicable.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

In case of eye contact: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

In case of skin contact: Wash with plenty of water.

If swallowed: Rinse mouth. Do not induce vomiting. If symptoms occurs: Get medical advice/attention.

If inhaled: Get medical attention if symptoms occur.

--

5.4. Instructions for safe disposal of the product and its packaging

Dispose of contents/container to in accordance with national regulation.
--

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Store in original container at room temperature. Protect from direct sun light. Protect from frost. Shelf-life: 2 years.
--

6. Other information

N/A

7. Third information level: individual products in the meta SPC

7.1 Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	SALVESAFE E13	Market area: EU
Authorisation number <small>(R4BP 3 asset reference number - National Authorisation)</small>	EU-0020512-0007 1-3	

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E14

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0008 1-3

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

Trade name(s)

SALVESAFE E21

Market area: EU

Authorisation number

(R4BP 3 asset reference number - National Authorisation)

EU-0020512-0009 1-3

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38
