


Regulation (EU) No 528/2012 concerning the making available on the market and use of biocidal products

Evaluation of active substances

Assessment Report



Biphenyl-2-ol

Product-type PT 13
(Preventol O Extra & Preventol ON Extra
Preservative Solution)

March 2015

Spain

CONTENTS

1. STATEMENT OF SUBJECT MATTER AND PURPOSE	3
1.1. Procedure followed	3
1.2. Purpose of the assessment report	3
2. OVERALL SUMMARY AND CONCLUSIONS	4
2.1. Presentation of the Active Substance	4
2.1.1. Identity, Physico-Chemical Properties & Methods of Analysis.....	4
2.1.2. Intended Uses and Efficacy.....	4
2.1.3. Classification and Labelling.....	5
2.2. Summary of the Risk Assessment	6
2.2.1. Human Health Risk Assessment	6
2.2.1.1. Hazard identification.....	6
2.2.1.2. Effects assessment.....	15
2.2.1.3. Exposure assessment.....	16
2.2.1.4. Risk characterisation	17
2.2.2. Environmental Risk Assessment	19
2.2.2.1. Fate and distribution in the environment.....	19
2.2.2.2. Effects assessment.....	20
2.2.2.3. PBT and POP assessment.....	22
2.2.2.4. Exposure assessment.....	23
2.2.2.5. Risk characterisation	23
2.2.2.6. Assessment of endocrine disruptor properties	26
2.3. Overall conclusions	27
2.4. List of endpoints	27
APPENDIX I: LIST OF ENDPOINTS	28
Chapter 1: Identity, Physical and Chemical Properties, Classification and Labelling	28
Chapter 2: Methods of Analysis	30
Chapter 3: Impact on Human Health	31
Chapter 4: Fate and Behaviour in the Environment	36
Chapter 5: Effects on Non-target Species	38
Chapter 6: Other End Points	40
APPENDIX II: LIST OF INTENDED USES	41
APPENDIX III: LIST OF STUDIES	42

1. STATEMENT OF SUBJECT MATTER AND PURPOSE

1.1. Procedure followed

This assessment report has been established as a result of the evaluation of the active substance [Biphenyl-2-ol] as Product-type [13] (Working or cutting fluid preservatives), carried out in the context of the work programme for the review of existing active substances provided for in Article 89 of Regulation (EU) No 528/2012, with a view to the possible approval of this substance.

Biphenyl-2-ol (CAS no. 90-43-7) was notified as an existing active substance, by LANXESS Deutschland GmbH and DOW Benelux B. V., hereafter referred to as the applicant, in Product-type 13.

Commission Regulation (EC) No 1451/2007 of 4 December 2007¹ lays down the detailed rules for the evaluation of dossiers and for the decision-making process.

In accordance with the provisions of Article 7(1) of that Regulation, Spain was designated as Rapporteur Member State to carry out the assessment on the basis of the dossier submitted by the applicant. The deadline for submission of a complete dossier for Biphenyl-2-ol as an active substance in Product-type 13 was 31st July 2007, in accordance with Annex V of Regulation (EC) No 1451/2007.

On 12th July 2007, Spain competent authorities received a dossier from the applicant. The Rapporteur Member State accepted the dossier as complete for the purpose of the evaluation on 31st October 2008.

On 2nd June 2014, the Rapporteur Member State submitted to the Commission and the applicant a copy of the evaluation report, hereafter referred to as the competent authority report.

In order to review the competent authority report and the comments received on it, consultations of technical experts from all Member States (peer review) were organised by the Agency. Revisions agreed upon were presented at the Biocidal Products Committee and its Working Groups meetings and the competent authority report was amended accordingly.

1.2. Purpose of the assessment report

The aim of the assessment report is to support the opinion of the Biocidal Products Committee and a decision on the approval of [Biphenyl-2-ol] for Product-type 13, and, should it be approved, to facilitate the authorisation of individual biocidal products. In the evaluation of applications for product-authorisation, the provisions of Regulation (EU) No 528/2012 shall be applied, in particular the provisions of Chapter IV, as well as the common principles laid down in Annex VI.

For the implementation of the common principles of Annex VI, the content and conclusions of this assessment report, which is available from the Agency web-site shall be taken into account.

However, where conclusions of this assessment report are based on data protected under the provisions of Regulation (EU) No 528/2012, such conclusions may not be used to the benefit of another applicant, unless access to these data for that purpose has been granted to that applicant.

¹ Commission Regulation (EC) No 1451/2007 of 4 December 2007 on the second phase of the 10-year work programme referred to in Article 16(2) of Directive 98/8/EC of the European Parliament and of the Council concerning the placing of biocidal products on the market. OJ L 325, 11.12.2007, p. 3

2. OVERALL SUMMARY AND CONCLUSIONS

2.1. Presentation of the Active Substance

2.1.1. Identity, Physico-Chemical Properties & Methods of Analysis

This evaluation covers the use of Biphenyl-2-ol in Product-type 13, but it does not cover sodium 2-biphenylate and potassium 2-biphenylate. The most important mechanism is the interaction with bio-membranes. In the first step an adsorption of Biphenyl-2-ol to the cell membrane takes place. The greater the proportion of undissociated molecules of the biocide in the surrounding medium the stronger will be the adsorption. In further steps the function of membrane proteins is disturbed, substrate transport and ATP synthesis are inhibited. The cell membrane loses its semi-permeability and ions and organic molecules escape.

Specifications for the reference source are established.

The physico-chemical properties of the active substance and of the representative biocidal product have been evaluated and are deemed acceptable for the appropriate use, storage and transportation of the active substance and biocidal product.

Validated analytical methods are available for the determination of Biphenyl-2-ol as manufactured and for the analysis of impurities. Validated analytical methods are also available for the determination of Biphenyl-2-ol in soil, water, air and food/feeding stuffs matrices. Other analytical methods are not required because Biphenyl-2-ol is not classified as toxic or highly toxic.

2.1.2. Intended Uses and Efficacy

The assessment of the biocidal activity of the active substance demonstrates that it has a sufficient level of efficacy against the target organisms and the evaluation of the summary data provided in support of the efficacy of the accompanying product, establishes that the product may be expected to be efficacious.

Biphenyl-2-ol and Biphenyl-2-ol Solution (active substance Biphenyl-2-ol in alkaline solution) are a multi-site bactericides and fungicides with basic activity at the cell wall, disruption of membrane potentials and general membrane permeability of cytoplasmic membranes.

Biphenyl-2-ol and Biphenyl-2-ol in alkaline solution have a broad efficacy against potentially harmful germs as fungi and yeasts, e.g. *Acremonium strictum*, *Fusarium solani*, *Geotrichium candidum* and *Rhodotorula rubra*.

Efficacy against fungi and yeasts has been proven to protect cooling lubricants against microbial spoilage. The duration of the test was 5 weeks. The results show that the efficacy concentration of Biphenyl-2-ol is $\geq 0.15\%$.

The tests were considered relevant to the evaluation for PT 13. Efficacy against bacteria was not demonstrated and it should be shown at product authorisation stage.

Due to the unspecific mode of action (multi-site activity) a development of resistance against biocidal use of OPP is not expected.

Biphenyl-2-ol and Biphenyl-2-ol in alkaline solution are antimicrobial preservatives for aqueous products. Metal working fluids are usually emulsified or dissolved with water prior to being

used. The emulsions or solutions (milky fluid) are very sensitive to microbial contamination which can:

- Lower the pH by secretion of acidic metabolites (e.g.: lactic acid, H₂S, etc.)
- Destroy (split) the emulsion
- Shorten the emulsion working-life
- Produce malodorous smells
- Cause diseases (pathogenic bacteria such as *Mycobacteria* sp.)

Biocidal products containing Biphenyl-2-ol are used to control the detrimental microorganisms. The tests were performed according to internal methods of Lanxess:

- Trial to check the application-oriented effectiveness of preservatives and screening substances in a cooling lubricant.
- A method for the evaluation of biocidal compounds in aqueous based metal working fluids.

In addition, in order to facilitate the work of Member States in granting or reviewing authorisations, the intended uses of the substance, as identified during the evaluation process, are listed in [Appendix II](#).

2.1.3. Classification and Labelling

CURRENT CLASSIFICATION

Classification according to the CLP Regulation		
Hazard Class and Category Codes	Eye Irrit. 2	H319
	Skin Irrit. 2	H315
	STOT SE 3	H335
	Aquatic Acute 1	H400
Labelling		
Pictograms	GHS07 GHS09 Wng	
Signal Word	Warning	
Hazard Statement Codes	H319: Causes serious eye irritation H315: Causes skin irritation H335: May cause respiratory irritation H400: Very toxic to aquatic life	
Specific Concentration limits, M-Factors		

PROPOSED CLASSIFICATION

The proposed classification and labelling for Biphenyl-2-ol according to Regulation (EC) No 1272/2008 (CLP Regulation) is:

Classification according to the CLP Regulation		
Hazard Class and Category Codes	Eye Irrit. 2	H319
	Skin Irrit. 2	H315

	STOT SE 3 Carc 2 Aquatic Acute 1 Aquatic Chronic 1	H335 H351 H400 H410
Labelling		
Pictograms	GHS07 GHS09 Wng	
Signal Word	Warning	
Hazard Statement Codes	H319: Causes serious eye irritation H315: Causes skin irritation H335: May cause respiratory irritation H351: Suspected of causing cancer H400: Very toxic to aquatic life H410: Very toxic to aquatic life with long lasting effects	
Specific Concentration limits, M-Factors		

2.2. Summary of the Risk Assessment

2.2.1. Human Health Risk Assessment

2.2.1.1. Hazard identification

Toxicokinetics and metabolism

A study was conducted in six human volunteers (males) to determine the degree of dermal absorption (Selim 6.2-03). The mean total absorption was 43.19. For the purpose of risk assessment in this dossier 43.19% dermal absorption of OPP through the skin will be applied. The mean total absorption, defined as the compound-related radioactivity present in the urine, feces (excluding tape strips) was 43.15% (concentration 0.4% \cong 0.006 mg OPP/kg bw). This indicates that the ^{14}C -OPP derived radioactivity did not accumulate in the superficial layers of the skin.

A dermal study was conducted in six human volunteers (males) to obtain information on the metabolism of OPP (Bartels 6.2-01). Metabolites of OPP present in the urine samples from the study 6.2-03 were characterized. The major urinary metabolite was found to be the sulphate conjugate of OPP, accounting for 68.33% of the absorbed dose. Conjugation of OPP with glucuronic acid was less significant, accounting for only 3.46% of the absorbed dose. Hydroxylation of the phenol or phenyl ring, followed by conjugation was also shown to be significant, with phenylhydroquinoneglucuronide and 2,4'-dihydroxybiphenyl-sulfate representing 14.34% and 12.35% of the absorbed dose, respectively. Trace levels of unmetabolized parent compound (0.50% of absorbed dose) were found in early time interval samples only. No free phenylhydroquinone or phenylhydroquinone-sulphate were found in any of the urine samples (limit of detection = 0.25-0.59% absorbed dose). OPP, both free and conjugated, accounted for 73.0% of the total absorbed dose following dermal exposure to 0.4 mg test material for 8 h.

A study was conducted to determine the degree of oral absorption and to obtain information on the metabolism of ^{14}C -*ortho*-Phenylphenol (^{14}C -OPP) in the B6C3F1 mouse (██████████ 6.2-02). The mean total absorption for the mice treatment groups, defined as the compound-related radioactivity present in the urine, faeces, tissues and carcass was 95-104% (concentration 25mg/kg and 1000 mg/kg). This suggests a low potential for bioaccumulation. The excretion of ^{14}C -OPP was rapid and complete by 12 - 24 h post-dosing with 74 - 98% of the recovered radioactivity in the urine and 6 - 13% in the faeces

An ADME study was conducted to obtain information on the metabolism of ^{14}C -*ortho*-Phenylphenol (^{14}C -OPP) in the B6C3F1 mouse and Fischer rats (██████████ 6.2-02). In mice OPP was completely metabolized and rapidly eliminated via the urine predominantly as a sulphate and glucuronide conjugate of OPP. Qualitatively the extent of metabolism was comparable

between mice and rats, although quantitative differences in the extent of OPP sulphation and glucuronidation were seen between these species. Binding to macromolecules or conjugation with intracellular glutathione occurs very rapidly thereby preventing the substance from being detectable or appearing free in the plasma.

No specific study of inhalation absorption of OPP is available.

Products of degradation (photolysis) in laboratory simulated ground waters

In laboratory experimental tests, it was observed that bisphenol-2-ol is degraded by photolysis in water (See Doc IIA, point 4.1.1.1.2 and 4.4) Two products of degradation are formed, benzoic acid and a diketohydroxy-compound, being this the higher proportion (maximum observes 13.7% of the OPP at day 1. The presence of these products is expected to be transiently as they are also quickly photodegraded.

In a QSAR evaluation, the environmental formation was predicted and also predicted lower toxicity than for OPP to aquatic media. Therefore, exposure and adverse effects in the aquatic media have been considered to be negligible and that the risk covered by the risk evaluated for the OPP. The risk of exposure for OPP and metabolites is considered negligible to aquatic media. Therefore it is still less likely the exposure to human to the product of transformation via the drinking water. In any case, the risk may be covered by the assessment of the OPP parent compound.

Therefore, additional toxicological information of this "products of transformation" (photolysis) is in principle not required as exposure to human via drinking water is expected to be negligible and risk may be covered from the assessment of parent compound. Nevertheless, it may be reasonable requiring performing an assessment for predicting the relative toxicity by read across from other similar substances in mammals, if enough information from similar substance is available.

Oral, dermal and inhalation absorption

A study was conducted in six human volunteers (males) to determine the degree of dermal absorption (Selim 6.2-03). The mean total absorption was 43.19. For the purpose of risk assessment in this dossier 43% dermal absorption of OPP through the skin will be applied.

A study was conducted to determine the degree of oral absorption and to obtain information on the metabolism of ¹⁴C-*ortho*-Phenylphenol (¹⁴C-OPP) in the B6C3F1 mouse (████████ 6.2-02). The mean total absorption for the mice treatment groups, defined as the compound-related radioactivity present in the urine, faeces, tissues and carcass was 95-104% (concentration 25 mg/kg and 1000 mg/kg). For the purpose of risk assessment in this dossier 100% oral absorption of OPP will be applied.

No specific study to determine the inhalation absorption of OPP is available. For inhalation application of OPP 100% absorption is assumed for risk characterization.

Acute toxicity

The oral acute toxicity was evaluated in the available document Gilbert 6.1.1-01. Under the conditions of this study, the acute oral LD₅₀ of Dowicide 1 Antimicrobial (99.9% OPP) for male and female Fischer 344 rats was 2733 mg/kg (2730.3 mg OPP/kg), by nonlinear interpolation.

The dermal acute toxicity was evaluated in the available document Bomhard 6.1.2-01. The LD₅₀ values for male and female rats were greater than 2000 mg/kg body weight and were not exactly determined.

The acute inhalation toxicity was evaluated in the available document Landry 6.1.3-01a. The LD₅₀ values for male and female Fischer rats were greater than 36 mg/m³ (0.036 mg/L) and were not exactly determined because the highest test atmosphere that could be generated was 0.036 mg/L, which is too low to provide an accurate determination (Landry 6.1.3-01b).

Irritation and Corrosivity

OPP is currently classified as Skin Irrit. 2 (H315: Causes skin irritation). The skin irritation was evaluated in the available document Gilbert 6.1.4-01/1981a in New Zealand White rabbits.

OPP is currently classified as Eye Irrit. 2 (H319: Causes serious eye irritation). To investigate eye irritation properties of OPP a test in the eye of albino rabbit was performed (████████ 6.1.4-01/1981b).

Based on the weight of evidence from existing information, it can be reasonably concluded that the substance is moderately irritant to the eye and because of its proven irritant effects on mucosa, it can be reasonably assumed that OPP is irritating to the airways when inhaled in high concentrations (e.g. pure substance dust) then it is classified as STOT SE 3 (H335: May cause respiratory irritation).

Sensitisation

OPP was tested for its skin sensitisation potential in Buehler test on Guinea pigs (██████████ 6.1.5-01/1994b) with Dowicide 1 Antimicrobial (99.9% OPP). The animals were in apparent good health and gained weight over the study period. Therefore, under the conditions of this study, Dowicide 1 Antimicrobial (99.9% OPP) did not cause delayed contact hypersensitivity in guinea pigs.

A paper is submitted where OPP was tested for its skin sensitisation potential in Magnusson-Kligman test on Guinea pigs (Andersen 6.1.5-02) with Preventol O Extra (OPP concentration \geq 99.5 %). No animals were sensitized by Preventol O Extra.

In humans there are some case reports indicating positive patch test reactions in dermatological patients. Important data for humans is available from a volunteer study showing clearly negative results. See below section of "Human Data" and Table 2.2.1.1 1
The overall conclusion is that biphenyl-2-ol is not skin sensitizer in humans

Local effects

Based on the irritation effect of the assay dosing in the Screen Phase of the guinea pig sensitization study, a NOAEC of 7.5% is proposed.

No NOAEC/LOAEC may be deduced for medium or long term exposure. Therefore, only risk assessment may be performed for systemic effects for medium and long exposure.

Repeated dose toxicity

OPP was examined in a 21-day dermal study (██████████ 6.3.2-01a) in Fischer 344 rats, in a 28-day oral study with Dog Beagle (██████████ 6.3.1-01, 6.5-02), in a 91-day oral study (██████████ 6.4.1-01a) in male Fischer rats, in a 1-year oral study in dog (██████████ 6.3.1-01, 6.5-02) and a 2-years oral study in Fischer rats (██████████ 6.5-01a, 6.7-01a).

The NO(A)EL for dermal exposure in a 21-day dermal study in Fischer rat is 1000 mg/kg bw/day on the basis of the no systemic effects in any dose group.

The NO(A)EL for oral exposure in a 28-day oral study in dog Beagle is 300 mg/kg bw/day on the basis of the no adverse effects in any dose group.

The NO(A)EL for oral exposure in a 91-day oral study in male Fischer is 224 mg/kg/day (4000 ppm) on the basis of the urothelial hyperplasia and the necrotic foci in the bladders in the highest dose.

The NO(A)EL for oral exposure in a 1-year oral study in dog is 300 mg/kg/day on the basis of the no adverse effects in any dose group.

The NO(A)EL for oral exposure in a 2-year oral study in Fischer rats is 39 mg/kg/day on the basis of the increased incidence of simple urinary bladder hyperplasia in males and the increased incidence of urinary bladder transitional cell carcinoma in males.

No specific studies for subchronic and chronic dermal toxicity and for short, subchronic and chronic inhalation toxicity are available

Genotoxicity and carcinogenicity

Genotoxicity

In-vitro

The results of the Salmonella/Mammalian-Microsome Plate Incorporation Mutagenicity Assay (San 6.6.1-01) indicate that under the conditions of this study, a positive response was not observed with any of the tester strains either in the presence or absence of microsomal enzymes prepared from Aroclor induced rat and hamster liver.

The test substance Preventol O Extra (99.9 % OPP) is considered to be non mutagenic in the CHO-HGPRT Forward Mutation Assay, (Brendler 6.6.3-01) both with and without metabolic activation.

OPP was clastogenic in Chinese hamster ovary cells at cytotoxic concentrations. In the

presence of S9 mix, phenylhydroquinone (metabolite produced from OPP) is formed which has a higher cytotoxic and clastogenic potential than OPP (██████████ 6.6.2-01).

In-vivo

Preventol O Extra (99.9 % OPP) was evaluated as non-genotoxic in the in vivo comet assay in hepatocytes and kidney cells of male mice (██████████ 6.6.5-01).

Carcinogenicity

The carcinogenicity was examined in two combined chronic toxicity/oncogenicity testing studies:

- In the rat Fischer 344 (██████████ 6.5-01a, 6.7-01a), where the urinary bladder showed evidence of a compound-induced neoplasia in the highest doses (male animals only). It was considered border-line at 4000 ppm (200 mg/kg body wt/day) as there was only a marginal and non-statistical increase in both urinary bladder hyperplasia and transitional cell carcinoma when compared to controls or 800-ppm males (39 mg/kg body wt/day). Evidence of a compound-induced neoplasia was not observed in female animals at any dose tested.
- In B6C3F1 mice (██████████ 6.7-02a), where A statistically significant increased incidence of hepatocellular adenomas was observed in male mice of the 500 and 1000 mg/kgBW/day groups (in the middle and high dose groups) . There were no significant increases in tumours in female mice fed OPP.

For OPP there is convincing evidence that the carcinogenetic effects shown in rodents are threshold effects with an indirect and non-genotoxic mechanism and tumours observed in rodent species (liver tumours in mice and bladder tumours in rats) are not predictive of carcinogenicity for humans due to proven species differences. Based on the criteria for classification of Directive 2001/59/EC, liver tumours in sensitive strain of mice are not of relevance for classification.

In the WG and in the ad hoc follow up process for discussing the AF is was discussed the relevant of tumours for humans. Th no relevant of the liver tumours in mice was agreed. The bladder tumour observed in male rats has been discussed in deep in Doc IIA and considering the special studies related with the use of biphenol-2-ol in alkaline conditions There are evidences suggesting that these tumours in male rats are not relevant to human as the MOA is related with special sensitivity to alkalinization in male rat bladder. However three ad hoc follow-up participants considered that the mechanisms of bladder tumour formation is not completely known and the relevance of these tumours for humans cannot be completely excluded. Therefore, biphenyl-2-ol may be classified as carcinogen Cat 2.

Reproductive and developmental toxicity

The teratogenicity of the OPP is examined in two studies:

- (1) in Wistar rats (██████████ 6.8.1- 01)
- (2) in New Zealand White rabbits (██████████ 6.8.1-02).

The relevant NOAEL for **maternal toxicity** adopted was **100 mg/kg bw/day** on the basis of the increased mortality (13%) in New Zealand White rabbits, gross pathologic alterations (ulceration and haemorrhage of the gastric mucosa, haemolysed blood in the intestinal tract and decreased ingesta) and histopathologic alterations (renal tubular degeneration and inflammation). The relevant NOAEL for **teratogenic toxicity** adopted was **250 mg/kg bw/day** (the highest assayed dose).on the basis of no adverse embryonal/fetal effects were observed at any dose level tested in New Zealand White rabbits

Two two-generation studies examined the impact of OPP in fertility in Sprague-Dawley rats (██████████ 6.8.2-02a and ██████████ 6.8.2-01). The NOAEL for parental toxicity in rats is 35 mg/kg bw/d in males and females, based on the incidence of urothelial hyperplasia and calculi in the kidney and/or urinary bladder was increased in male rats. The NOAEL for development (F1) is 457 mg/kg bw/d in males and females, based on no adverse effects in any dose group

Neurotoxicity

OPP does not belong to a class of compounds for which a neurotoxic potential can be expected. In addition the available toxicity studies gave no indication of any relevant neurotoxic potential of the compound.

Human data

A short report entitled "Occupational medical experiences with *o*-Phenylphenol" is submitted (Heyne 6.12.1-01; no GLP). Occupational medical surveillance of workers exposed to *o*-Phenylphenol, performed every 3 years on a routine basis. The workers have been in the production of *o*-Phenylphenol in average for 13,9 years. During this period accidents with *o*-Phenylphenol or unwanted contamination with *o*-Phenylphenol haven't been recorded and consultations of the Medical Department due to work or contact with *o*-Phenylphenol haven't been required. The Phenol-levels in urine have always been far below German biological tolerance level of 200 mg/L (formerly 300 mg/L). *o*-Phenylphenol did not reveal any unwanted effects in the workers. Especially no sensitization of airways or skin to *o*-Phenylphenol has occurred. The examinations have included the above laboratory parameters as well as clinical and technical examinations.

A short communication is submitted (Adams 6.12.6-01) where it is described two cases of allergic contact dermatitis due to occupational contact with OPP containing products. In both patients the dermatitis was extensive and severe. In the case 1, a 34-year-old medical laboratory assistant applied a common over-the-counter "medicated" cream to various parts of his body for "dry skin". Patch testing with the cream and *o*-Phenylphenol in 0.5% and 1% concentrations showed strong positive reactions at 72 h. In the case 2, a 57-year-old male machinist had experienced a recurring dermatitis on the hands, arms, trunk, thighs and feet for 25 years. A patch testing revealed a positive reaction to 1% *o*-Phenylphenol in petrolatum, and a positive "provocative use test" from a suspected coolant which contained this preservative.

A short communication is submitted (Van Hecke 6.12.6-02) where it is described a case of allergic contact dermatitis due to occupational contact with OPP containing products. A 24-year-old machinist had had dermatitis of the hands for 10 months due to a coolant and a cleanser.

A paper is submitted (Schnuch 6.12.6-03) where it is examined the role of different preservatives in a large number of patients with suspected allergic contact dermatitis. Patch test data and data from the patients' history were collected from the 24 departments participating in the Information Network of Departments of Dermatology from 1 January 1990 to 31 December 1994. Patch test data from 28349 patients tested with preservatives of the standard series (SS), from 11485 patients tested additionally with a preservative series (PS), and from 1787 patients tested with an industrial biocide tray (IB) were evaluated. Nine of 24 centers applied patch tests for 24 h, the remainder (15 of 24) for 48 h. Readings were done at 72 h after application of the test chambers. The PS and IB contained OPP at a concentration of 1% in petrolatum. Of 11418 subjects tested, 59 showed an irritant or questionable result, 33 (0.3%) were positive in PS. Of 1785 subjects tested, 5 showed an irritant or questionable result, 5 (0.4%) were positive in IB.

A paper is submitted (Brasch 6.12.6-05) where the main purpose was to identify the most frequent contact allergens and reconsider the test concentrations. This study is a retrospective evaluation of patch test results with medical antimicrobials and preservatives, performed by eight centres of the IVDK (Informations verb und Dermatologischer Kliniken) from 1989 to 1991. It was evaluated the patch test results and questionnaires of 2059 patients tested with a preliminary series of medical antimicrobials and preservatives where OPP was included. This series was tested in patients clinically suspected to suffer from contact allergy to preservatives. Of 2043 subjects tested with OPP (at a concentration of 1% in petrolatum), 6 showed a medium positive reaction, 8 an equivocal reaction and one an irritant reaction.

A paper is submitted (Geier 6.12.6-04) where 1132 patients were patch tested with a variety of "antiseptics/industrial chemicals". OPP was one of the test compounds. OPP was applied as a 1% solution in petrolatum. Of 1131 patients tested with OPP, 5 individuals (0.4%) showed positive reactions. One individual showed ambiguous results.

Other no critic studies with complementary information which does not contradict the results of the key studies are included in the next table.

Table 2.2.1.1-1: Effects of OPP in Humans

Doc IIIA Section No.	Type	Description	Results	Reference
6.12.1 Key study	Surveillance of manufacturing plant personnel	Medical surveillance of personnel involved in OPP production No. of workers exposed: 73 (2 ♀, 71 ♂) in average 13.9 years of medical supervision	No adverse effects. No airway or skin sensitisation towards OPP has occurred.	Heyne 6.12.1 (01)
6.12.6 Key study	Clinical cases	Two cases of allergic contact dermatitis due to occupational contact with OPP containing products (1) germicidal agent (2) coolant	allergic contact dermatitis in both cases due to OPP	Adams 6.12.6 (01)
6.12.6 Key study	Clinical case	One case of sensitivity to OPP due to occupational contact to a coolant containing OPP	Contact sensitivity to OPP in a coolant	Van Hecke 6.12.6 (02)
6.12.6 Key study	Multi-centre study	Patch tests on patients with suspected contact dermatitis. 11485 patients were tested additionally with a preservative series (PS) and 1785 were tested with an industrial biocide tray (IB). Occupational exposure was suspected in 17% of the cases	59 of 11418: irritative or questionable result in PS 33 of 11418: positive reaction in PS 5 of 1785: irritative or questionable result in IB 7 of 1785: positive reaction in IB	Schnuch 6.12.6 (03)
6.12.6 Key study	Study	retrospective study patch tests 1 % OPP was applied	6 of 2043: medium positive reaction 8 of 2043: equivocal reaction 1 of 2043: irritant reaction	Brasch 6.12.6 (05)
6.12.6 Key study	epidemiological study	1132 patients were patch tested with a variety of "antiseptics/industrial chemicals". OPP was one of the test compounds.	Of 1131 patients tested with OPP, 5 individuals (0.4%) showed positive reactions. One individual showed ambiguous results	Geier 6.12.6 (04)
6.12.6	Epidemiological study	Epidemiological study on metal workers. Patch tests with 1% OPP. 40 workers were tested. 39 of them presented with dermatitis of hands and/or forearms. 5 had incidences of dermatitis in the past.	OPP was not a contact allergen in any of the cases.	De Boer 6.12.6 (08)

Table 2.2.1.1-1: Effects of OPP in Humans

Doc IIIA Section No.	Type	Description	Results	Reference
6.12.6	epidemiological study	Epidemiological study on 424 metalworkers who were exposed to metal working fluid. Patch tests with 1% OPP on 277 patients.	2 of 277: positive reaction	Uter 6.12.6 (06)
6.12.1	Surveillance of manufacturing plant personnel	Regular medical examination and urine biomonitoring.	Medicinal surveillance and biomonitoring did not reveal findings of concern.	6.12.1 (02)

Other/special studies

A paper is submitted (Fukushima 6.10-01/AIII 6.10-1) where the effects of sodium *o*-phenylphenate (OPP-Na) and OPP on two-stage urinary bladder carcinogenesis in male F344 rats initiated with *N*-butyl-*N*-(4-hydroxybutyl)nitrosamine (BBN) were investigated. OPP-Na acts as a tumour promoter in the urinary bladder following initiation by BBN. OPP-Na alone also induced tumour formation in the urinary bladder and can therefore be considered a weak initiator in the two-stage model of carcinogenesis and a complete carcinogen. OPP had no significant tumour-promoting or initiating effects. The increase in urinary pH caused by OPP-Na but not by OPP might cause the difference in the carcinogenic potential of the two compounds.

A paper is submitted (Fujii 6.10-03/ AIII 6.10-2) where the effects of an alkalizer or an acidifier on bladder carcinogenesis induced by OPP or OPP-Na were examined. The results indicate that the administration of an alkalizer enhanced the carcinogenicity of OPP and the administration of an acidifier inhibited the carcinogenicity of OPP-Na to the rat urinary bladder. This suggests that the earlier finding that OPP-Na was more carcinogenic than OPP resulted from the higher alkalinity of OPP-Na.

A study is submitted (██████████ 6.10-15/ AIII 6.10-3; no guideline; no GLP) where the possible role of prostaglandin-*H*-synthase (PGHS) in OPP-induced bladder tumour formation is investigated. OPP and phenylhydroquinone (PHQ) stimulate cyclooxygenase activity and are oxidised by PGHS. OPP, PHQ and 2-phenyl-1,4-benzo-quinone (PBQ) inhibit PGHS at higher concentrations.

Other no critic studies with complementary information which does not contradict the results of the key studies are included in the Table 2.2.1.1-2.

These effects of concern observed with Na/K salts (or OPP in alkaline condition) should be considered in the evaluation of the hazard and risk of products formulated or used in dilution in alkaline conditions.

Table 2.2.1.1-2: Other/special studies with OPP

Type of study	Dosage	Results	Reference
32-week, dietary, rats Key study	20000 ppm, with and without tumour initiator <i>ad libitum</i>	OPP had no significant tumour-promoting or -initiating effects in the urinary bladder.	██████████ 6.10 (01)/AIII 6.10 (1)
26-week, dietary, rats Key study	12500 ppm, with/without NaHCO ₃ <i>ad libitum</i>	Urinary bladder tumourigenesis of OPP is enhanced by NaHCO ₃ .	██████████ 6.10 (03)/ AIII 6.10 (2)

Table 2.2.1.1-2: Other/special studies with OPP

Type of study	Dosage	Results	Reference
<i>In-vitro</i> interaction with PGHS Key study	OPP, PHQ, PBQ: 100 µM	OPP and PHQ stimulate cyclooxygenase activity and are oxidised by PGHS. OPP, PHQ and PBQ inhibit PGHS at higher concentrations.	██████████ 6.10 (15)/ AIII 6.10 (3)
32-week, dietary, rats	12,500 ppm, with varying amounts of NaHCO ₃ <i>ad libitum</i>	Morphological changes of the bladder epithelium, correlating with increased urinary pH.	██████████ 6.10 (01)
32-week, dietary, rats	20,000 ppm, <i>ad libitum</i>	Reduced urinary osmolality. Increased pH and Na ⁺ correlate with tumourigenesis.	██████████ 6.10 (04)
12-week, dietary, rats	0, 2500, 5000, 10,000, 20,000 ppm, <i>ad libitum</i>	At 20,000 ppm: morphological changes of the bladder luminal surface evident by SEM	██████████ 6.10 (02)
90-day, dietary + acute DNA-binding study in rats	90-day study: OPP, SOPP: 2% in diet Acute assay: OPP, SOPP: 500 mg/kg	SOPP, but not OPP, caused regenerative hyperplasia of the urinary bladder. OPP-treated rats revealed renal damage. No interactions with DNA could be demonstrated for either compound.	██████████ 6.10 (06)
8-week, dietary, rats	OPP: 1.25% with or without NaHCO ₃ SOPP: 2% with or without NH ₄ Cl	Males are more sensitive to OPP than females under alkalinuric conditions with respect to bladder hyperplasia.	██████████ 6.10 (07)
1-week, dietary, rats	OPP, SOPP: 0.1-2.0%	OPP and SOPP caused a dose-dependent increase in agglutinability of bladder epithelial cells by Con A which is an indication for carcinogenic potential.	██████████ 6.10 (08)
Acute oral, rat	OPP, PHQ, PBQ: 700, 1400 mg/kg bw, single oral gavage, with or without inhibition of GSH synthesis	OPP treatment led to GSH depletion and eosinophilic degeneration of centrilobular hepatocytes. Inhibition of GSH synthesis aggravated hepatotoxicity of OPP.	██████████ 6.10 (09)
Cytotoxicity test in primary rat hepato-	OPP, PHQ: 0-1 mM	OPP cytotoxicity is enhanced by monooxygenase inhibition and GSH depletion. PHQ-induced cell death can be inhibited by	██████████ 6.10 (10)

Table 2.2.1.1-2: Other/special studies with OPP

Type of study	Dosage	Results	Reference
<i>cytes</i>		sulfhydryl compounds.	
<i>In-vitro and in-vivo macro-molecular binding assay</i>	¹⁴ C-OPP: 1 µCi In vivo: OPP, SOPP: 50-500 mg/kg, oral gavage, 16-18 h	A non-linear increase in macromolecular binding of OPP and SOPP was observed in vivo and in vitro. This may be caused by the saturation of detoxification pathways.	████████ 6.10 (11)
<i>In-vitro metabolism of OPP</i>	OPP: 1-100 µM	OPP is oxidised to PHQ and PHQ is oxidised to PBQ by cytochrome P-450. PBQ is reduced back to PHQ by cytochrome P-450 reductase (redox cycling).	Roy 6.10 (12)
<i>In-vivo assay of DNA synthesis in bladder</i>	OPP, SOPP: 2% in diet; 4-24 weeks	OPP and SOPP cause a proliferative response in renal pelvis and papilla when given at a dietary level of 2%.	████████ 6.10 (13)
<i>In-vitro and in-vivo GSH conjugation</i>	In-vitro study: 79 µg/mL In-vivo study: 1000 mg/kg, single oral dose	PHQ-GSH is excreted via the bile after OPP administration to rats. In vitro, PHQ-GSH can be formed non-enzymatically from PBQ and GSH or enzymatically from OPP and GSH.	████████ 6.10 (14)
<i>In-vivo assay of DNA and protein adducts in rats</i>	0, 15, 50, 125, 250, 500, 1000 mg/kg OPP, single oral gavage	OPP or its metabolites form protein, but not DNA, adducts in urinary bladder tissue.	████████ 6.10 (16)
<i>Ten-week feeding study in rats</i>	OPP: 1.25% in diet SOPP: 2.0% in diet 10 weeks	OPP and SOPP caused urothelial hyperplasia in rats as evident by histology and increased cell proliferation.	████████ 6.10 (17)
<i>7 and 14 days feeding study in male B6C3F1 mice</i>	0, 500, and 1000 mg/kg/day OPP in the diet for 7 and 14 days	The results indicate that OPP may be an agonist ligand for PPARα.	OPP_TOX_chronMaus_PPAR tumors_REPORT_2009-10

2.2.1.2. Effects assessment

The AELs were set as follows:

	Critical Study	Critical NOAEL	Assessment factor	AEL
Short exposure	teratogenicity oral study in New Zealand White rabbits	100 mg/kg bw/day	100	1 mg/kg bw/day
Mid exposure	2-years oral study	39 mg/kg/day for males	100	0.4 mg/kg bw/day
Long exposure	2-years oral study	39 mg/kg/day for males	100	0.4 mg/kg bw/day

Reasons for establishing critical endpoints

The acute AEL for risk characterization was deduced from a teratogenicity oral study in New Zealand White rabbits (██████████ 6.8.1-02). The relevant NOAEL for maternal toxicity adopted was 100 mg/kg bw/day on the basis of the increased mortality (13%), gross pathologic alterations and histopathologic alterations. Therefore, considering an assessment factor of 100, an AEL_{acute} of 1 mg/kg bw/day was calculated.

For mid and long term exposure, an Acceptable Exposure Level (AEL) value for repeated use is deduced from the NO(A)EL for chronic oral exposure in a 2-years oral study (██████████ 6.5-01a, 6.7-01a). The NOAEL is 39 mg/kg/day on the basis of the increased incidence of simple urinary bladder hyperplasia in males and the increased incidence of urinary bladder transitional cell carcinoma in males. . An AF=100 was established after a follow up discussion (See comment below). Therefore, considering an assessment factor of 100, an AEL_{medium} and AEL_{long} of 0.39 mg/kg bw/day was calculated.

Conclusion of the follow up discussion for establishing AF

In the combined chronic toxicity and carcinogenicity study of ██████████ (1996), the transitional cell carcinoma occurred in rats treated with biphenyl-2-ol at 200 mg/kg bw/d, while the same effect was reported in rats at 270 mg/kg bw/d after life span administration of sodium biphenylate (Fujii 1985). The NOAEL of 39 mg/kg bw/d from ██████████ study, to be used for the derivation of the reference values, would be 5-fold lower than the LOAEL of 200 mg/kg bw/d for transitional cell carcinoma. Overall, the rat seemed to be the most sensitive species, since the administration of biphenyl-2-ol to mice and dogs did not lead to adverse effects in the urinary bladder, and male rats appeared to be more susceptible to bladder tumours than the female rats. The male rat is in general considered much more susceptible to bladder changes including tumours related to local effects than other animal species and humans.

Three ad hoc follow-up participants considered that the mechanisms of bladder tumour formation is not completely known and the relevance of these tumours for humans cannot be excluded, therefore they proposed a margin of safety of 1000 from the LOAEL of 200 mg/kg bw/d, that would result in an additional assessment factor of 2.

However, given the bladder tumours species sensitivity, five participants agreed that an assessment factor of 100 applied to the conservative NOAEL of 39 mg/Kg bw/d would provide an adequate margin of safety for humans.

The eCA supported the majority view and an AF of 100 is applied.

The AELlong-term and AELmedium-term are rounded to 0.4 mg/kg bw/d

End points for Local effect assessment

For local effects, the NOAEC for short exposure is 7.5% on the basis of irritation effect of the assay dosing in the Screen Phase of the guinea pig sensitization study (██████████ 6.1.5-01/1994b).

No NOAEC/LOAEC may be deduced for medium or long term exposure.

Conclusion of classification for carcinogenicity

There are evidences suggesting that these tumours in male rats are not relevant to human as the MOA is related with special sensitivity to alkalinisation in male rat bladder. However, the mechanisms of bladder tumour formation is not completely known and the relevance of these tumours for humans cannot be completely excluded. Therefore, biphenyl-2-ol may be classified as carcinogen Cat 2.

2.2.1.3. Exposure assessment

The active substance, Biphenyl-2-ol as Preventol O Extra or Preventol ON Extra Preservative Solution can be used as a preservative in metal working industry. Therefore human exposure has been assessed both for the use of the biocidal product and use of preserved metal working fluids by professional users including an assessment of indirect exposure. Non professional use is not envisaged.

Preventol O Extra contains $\geq 99.5\%$ of the active substance (a.s.) Biphenyl-2-ol. The biocidal product is used by the manufacturer of the metal working fluid (MWF) concentrate and added at concentrations ranging within 4% to 6% w/w, or it can be directly used at metal working plants by the end-user in concentrations up to 3,000 ppm as tank side addition. In this case Biphenyl-2-ol is usually dissolved in a suitable solvent (i.e., propylene glycol), prior to be dosed in the functional fluid. The representative formulation Preventol ON Extra Preservative Solution (ca. 17.7% OPP, w/w) is an alkaline ready-for-use preservative solution. A ready-to-use MWF will contain 0.954 – 1.7% (w/w) of the b.p., equivalent to 0.159 – 0.30% (w/w) Biphenyl-2-ol.

The Human exposure to Biphenyl-2-ol was calculated based in the selected models and default values from TNsG on Human Exposure, 2002 & 2007, taking into account the User Guidance to report 2002 and HEEG's opinion on Human exposure assessment to biocidal products used in metalworking fluids (PT 13), (MOTA v5 2013, HEEG Opinion 5).

Human exposure assessment for industrial users

The exposure during the production of the active substance and the formulation of the biocidal product (b.p.) are not assessed by the Rapporteur under the requirements of the BPD. However the Rapporteur assumes that the production/formulation is performed in conformity with national and European occupational safety and health regulations.

Industrial exposure covers the dilution of the product Preventol O Extra to formulate the metal working fluid concentrate (MWF concentrate). The b.p. is automatically dosed in the mixing vessel. Exposure of industrial operators during formulation is addressed using Mix and Load model 7 (TNsG). For exposure assessment purposes 10 minutes each day is considered (Excel Data Base Human Exposure, TNsG 2007). Exposure via inhalation route should be reduced as much as possible due to the hazard properties of solid Biphenyl-2-ol. The use of RPE is considered to reduce inhalation exposure by a factor of 10 (equivalent to the efficacy of technical measures reducing 10-fold airborne levels of OPP). RPE would be required if aerosols or dust is produced during this task. PPE (gloves, protective clothing, face shield) is required when handling b.p. due to its irritant properties.

Human exposure assessment for professional users

Professional exposure is considered for operators working in metalworking premises. Worker exposure to treated MWF may arise from the following activities:

- 1 Tank-side addition of Biphenyl-2-ol to preserve metalworking fluid requires the biocidal product to be dissolved in a suitable solvent, e.g., propylene glycol, prior to being dosed into the functional fluid. For tank-side addition batches of pre-mixed biocide (containing max. 6%w/w of biocide in a suitable solvent) are typically prepared once per week. On the other hand, biocide concentrates (maximum content 6% Biphenyl-2-ol) may also be added to MWF as shock doses. The total active substance concentration of Biphenyl-2-ol in the system should not exceed 3,000 ppm in the final use dilution. The biocidal product Preventol ON Extra Preservative Solution is added directly to the circulating emulsion during its use (tank-side addition) in a closed system.
- 2 Application phase which involves the end-use of metalworking fluid. The machining operations comprises activities such as metalworking on turning machine and tool settling and dismantling which are performed using 0.3% Biphenyl-2-ol maximum content in MWF. Exposure is estimated according to HEEG opinion on Human exposure assessment to metalworking fluids (Metal working fluid model 2, (TNsG) and BEAT's worked example for PT 13 (Machining of metal parts). Exposure duration is up to 60 min per working day.
- 3 The post application phase refers to other task involved in metalworking procedures such as maintenance, monitoring, cleaning disposal and transfer. Exposure is estimated according to Metal working fluid model 2, (TNsG) and Cleaning of Spray Equipment from BEAT database (TNsG 2007). Exposure duration is assumed 7 hours per working day

The secondary exposure scenario associated with professionals working in the metal working shop is due to transfer of machined metal from lather to storage area. Exposure is assumed similar to the post application phase described.

Human exposure assessment for non professional users

General public is not exposed to metalworking preserved fluids MWF containing Biphenyl-2-ol.

Human exposure assessment from indirect exposure as a result of use

The secondary exposure scenarios identified are: dermal exposure of operators to metal parts contaminated with preserved MWF (see post application phase) and dermal exposure to contaminated coveralls at laundering.

2.2.1.4. Risk characterisation

Summary of risk assessment for industrial use

The exposure for industrial users is considered to be within the acceptable range if adequate PPE is worn. Handling of biocidal product requires the use of PPE due to its classification as irritant. Exposure via inhalation route should be reduced as much as possible due to the hazard properties of solid Biphenyl-2-ol. The use of RPE is considered to reduce inhalation exposure by a factor of 10 (equivalent to the efficacy of technical measures reducing 10-fold airborne levels of Biphenyl-2-ol). RPE would be required if aerosols or dust is produced during this task. It must be noted that appropriate PPE to handle corrosive chemicals must be worn during use of Preventol ON Extra Preservative Solution.

Summary of risk assessment for professional use

The exposure for professional users is considered to be within the acceptable range if adequate

PPE is worn. Tables above show that total exposure is high. Exposure may be overestimated: OPP concentrations may vary among the different activities and professionals are involved in other tasks that do not require exposure to a.s.

When considering the aggregate exposure for professionals working in the premises, the use of risk management measures (RMM) is recommended to reduce exposure via the dermal route. These RMM should be focused in decreasing the residues of preserved MWF on worked metal pieces or machinery. Rinsing of the drilled/cut work pieces with an appropriate solvent after metal working processes is recommended prior to manual handling. In addition, before any maintenance or repair work is carried out the machineries should be flushed. Other risk mitigation measures required to reduce exposure via the dermal route include reducing contact with contaminated surfaces/oily rags, and washing hands regularly.

Assuming a 10-fold reduction in dermal exposure through the use of adequate RMM, no risk is envisaged for the aggregate exposures of operators in metalworking facilities.

Midterm Exposure Scenario	Exposure Adults (mg/kg bw/[d])	AEL (mg/kg bw/[d])	Exposure % AEL
Professional weighting solid OPP to prepare MWF concentrate (6%OPP) and pouring, 20 min/week, Tier 2			
Inhalation*	2.68E-03	0.4	0.6
Dermal**	0.221833	0.4	54.4
Total	0.224516	0.4	55
*.LEV (10 fold reduction) during handling solid b.p. RPE would be required if aerosols or dust are produced **PPE is required to handle solid b.p.			
Chronic Exposure Scenario	Exposure Adults (mg/kg bw/[d])	AEL (mg/kg bw/[d])	Exposure % AEL
Professional (1) metalworking activities, 0.3%OPP, 1 hour daily, Tier 1			
Inhalation	2.06E-05	0.4	0.01
Dermal*	0.37668	0.4	94.17
Total	0.376701	0.4	94.18
*no gloves			
Chronic Exposure Scenario	Exposure Adults (mg/kg bw/[d])	AEL (mg/kg bw/[d])	Exposure % AEL
Professional (1) metalworking activities, 0.3%OPP, 1 hour daily, Tier 2			
Inhalation	2.06E-05	0.4	0.01
Dermal*	0.263934	0.4	65.98
Total	0.263955	0.4	65.99
*impermeable coverall, no gloves			
Chronic Exposure Scenario	Exposure Adults (mg/kg bw/[d])	AEL (mg/kg bw/[d])	Exposure % AEL
Professional (2) other tasks in the workshop, 0.3%OPP, 7 hour daily, Tier 1			
Inhalation	1.44E-04	0.4	0.04
Dermal	0.49665	0.4	124.16
Total	0.496794	0.4	124.20
Chronic Exposure Scenario	Exposure Adults (mg/kg bw/[d])	AEL (mg/kg bw/[d])	Exposure % AEL
Professional (2) other tasks in the workshop, 0.3%OPP, 7 hour daily, Tier 2			
Inhalation	1.44E-04	0.4	0.04
Dermal*	0.040996	0.4	10.25
Total	0.041141	0.4	10.29
*impermeable coverall, gloves			
Chronic Exposure Scenario	Exposure Adults	AEL	Exposure

Scenario	(mg/kg bw/[d])	(mg/kg bw/[d])	% AEL
Professional (2) other tasks in the workshop, 0.3%OPP, 7 hour daily, Tier 2			
Inhalation	1.44E-04	0.4	0,04
Dermal*	0.331943	0.4	82,99
Total	0.332087	0.4	83,02

*impermeable coverall, no gloves

Summary of risk assessment for non-professional use

Not applicable.

Summary of risk assessment from indirect exposure as a result of use

The results indicate an acceptable risk for the professional (chronic) indirect exposure to metal working fluids containing OPP.

Midterm Exposure Scenario	Exposure Adults (mg/kg bw/[d])	AEL (mg/kg bw/[d])	Exposure % AEL
Professional: home laundering of work clothes ¹			
Dermal	4.86E-03	0.4	1.22
Total	4.86E-03	0.4	1.22

¹(40.752 mg OPP deposit on a workday * 5 days / 22,700 cm² coverall) *hands area * 30% transference

2.2.2. Environmental Risk Assessment

2.2.2.1. Fate and distribution in the environment

Considering the hydrolytic stability determined under stringent temperature conditions and at different pH values, it is not expected that hydrolytic processes will contribute to the degradation of OPP in the aquatic systems (estimated DT₅₀ > 1 year).

OPP is rapidly photodegraded in sterile aqueous 0.01 M phosphate buffer (experimental DT₅₀ = 0.3 days). Diketohydroxy-compound (maximum 13.6% AR) and benzoic acid (maximum 7.9% AR) were identified as the major transformation products, other 3 unidentified compounds were found to have a maximum between 1% and 10% of the AR. Innumerable minor phototransformation products (each < 1% AR) were formed. All transformation products occurred transiently and decreased to amounts of < 5% AR at the end of the study. In all cases the QSAR estimates were indicative of a significant potential for rapid degradation in the environment.

The tropospheric half-life of OPP was estimated using the AOPWIN program (v. 1.91, 2000). Using a mean daily OH concentration in air of 0.5 x 10⁶ OH radicals per cm³, a half-life in air of 0.59 days was assessed - corresponding to a chemical life-time in air of about 0.85 days - due to indirect photodegradation. It is not to be expected that it can be carried in the gaseous phase over long distances or can accumulate in air. Furthermore, OPP has a low vapour pressure.

OPP is concluded to be readily biodegradable (71-76% after 28 days and 100% after 16 days, respectively). Moreover, high overall removal rates in activated sludge wastewater treatment plants of 99 to 100% (complete mineralization) were observed in a monitoring study conducted by Körner *et al.* (2000) in a municipal sewage plant Steinhäule located on the Danube River in southern Germany. Several studies in different municipal sewage plants presented by the applicant (Ternes *et al.* (1998), and Lee *et al.* (2005)) confirm the data from

Körner, and a value of 99% elimination efficiency is used in the Tier 2 approach for the risk assessment.

The simple first order DT₅₀ value of ortho-Phenylphenol in the test soil was 1 day (DT₅₀ 2.7 hours) providing an appropriate margin of safety. The DT₅₀ has been re-calculated considering a biphasic approach. A DT₅₀ default value in soil of 30 days (according to the TGD for Risk Assessment Chapter 3, Table 8) is considered to be as worst case for the risk assessment and a DT₅₀ of 15.08 days as a refinement.

Based on two reliable adsorption/desorption studies and the results obtained in the soil degradation study, no potential for translocation into deeper soil layers or even ground water is given. K_{oc} values were 346.7 in the HPLC screening test and 252-392 in the adsorption/desorption (batch equilibrium) study. Based on a classification K_{oc} value of 347 L·kg⁻¹, OPP can be classified as a moderately mobile substance.

Although a log Pow of 3.18 was determined, no indication for a possible bioaccumulative potential of OPP is given due to a calculated steady-state bioconcentration factor (BCF) of 21.7 (wet weight), 114-115 (lipid content). Taking into consideration these low bioconcentration factors and the low computed concentrations in surface water, a significant food chain concern does not exist.

2.2.2.2. Effects assessment

STP compartment

According to TGD for Risk Assessment (EC, 2003), and taking into account the test available with aquatic micro-organisms (according to OECD 209 with activated sludge, EC₅₀ = 56 mg OPP·L⁻¹), an assessment factor of 100 can be applied. Thus, a PNEC_{microorganisms} of 0.56 mg a.i./L is derived.

Surface water compartment

The toxicity of OPP to aquatic organisms is well documented by acute and long-term studies. Three chronic NOEC values for the three trophic levels of the base set (fish, *Daphnia*, algae) are available for the aquatic compartment resulting in NOECs of 0.036 mg a.i./L (*Pimephales promelas*), 0.006 mg a.i./L (*Daphnia magna*) and 0.468 mg a.i./L (*Pseudokirchneriella subcapitata*). A sediment-water chironomid toxicity test using spiked water is available with *Chironomus riparius* with a NOEC of 1.85 mg a.i./L. Since concentrations declined during the test (34-55% present in the water phase after 7 days), initial concentrations in water are not adequate to express the NOEC.

The lowest NOEC value (*Daphnia magna*) of 0.006 mg a.s./L is considered for the PNEC calculation. Since long-term NOECs are available for all three trophic levels, an assessment factor of 10 was applied to the lowest long-term NOEC value. The PNEC_{water} was thus calculated to be 0.0006 mg a.i./L.

Sediment

In two preliminary range finding test (non-GLP) with spiked sediment and spiked water, it was found that the test organisms exposed to spiked water were affected at considerably lower concentrations than the larvae exposed to spiked sediment, with a NOEC of 1.85 mg/L expressed as a concentration in water.

However, it is not agreed to use the NOEC for *C. riparius* because this NOEC is expressed on the basis of initial concentrations in the water phase and, actual concentrations during the 28-days were much lower because of distribution to sediment. For this reason, the equilibrium partitioning on the PNEC_{water} has been used. For this, the Foc in suspended matter (0.1) should

be used instead of the Foc sediment resulting in a $PNEC_{\text{sediment}}$ of 0.0049 mg/kg_{wwt} (0.02254 mg/kg_{dwt}).

$$\begin{aligned}
 PNEC_{\text{sed}} &= (K_{\text{susp-water}}/RHO_{\text{susp}}) * PNEC_{\text{water}} * 1000 && \text{(page 113 of TGD)} \\
 K_{\text{susp-water}} &= F_{\text{water}_{\text{susp}}} + (F_{\text{solid}_{\text{susp}}} * (K_{\text{p}_{\text{susp}}}/1000) * RHO_{\text{solid}}) && \text{(page 47 of TGD)} \\
 &= 0.9 + (0.1 * (34.7/1000) * 2500) = 9.575 \text{ m}^3/\text{m}^3 \\
 PNEC_{\text{sed}} &= (9.575/1150) * 0.0006 * 1000 = 0.0049 \text{ mg/kg} \\
 PNEC_{\text{sed}} &= 0.0049 \text{ mg/kg OPP/kg wet sediment}
 \end{aligned}$$

Terrestrial compartment

For the effects assessment of the soil, compartment tests are available for three trophic levels (terrestrial microorganisms, earthworms, and plants):

- Terrestrial microorganisms (C- and N-cycle):

$$EC_{50} \text{ (28 days)} = 633.5 \text{ mg a.s.} \cdot \text{kg}_{\text{dw}}^{-1} \text{ soil}$$

- Earthworms (*Eisenia fetida*):

$$LC_{50} \text{ (14 days)} = 198.2 \text{ mg a.i.} \cdot \text{kg}^{-1} \text{ soil}$$

$$NOEC \text{ (14 days)} = 125 \text{ mg a.i.} \cdot \text{kg}_{\text{dw}}^{-1} \text{ soil}$$

- Terrestrial plants (*Avena sativa*):

$$LC_{50} \text{ (14 days)} = 53.9 \text{ mg a.i.} \cdot \text{kg}^{-1} \text{ soil}$$

$$NOEC \text{ (14 days)} = 12.5 \text{ mg a.i.} \cdot \text{kg}_{\text{dw}}^{-1} \text{ soil}$$

The lowest result was obtained in the study with plants. A $PNEC_{\text{soil}}$ was calculated on basis of the lowest LC_{50} of three trophic levels using an assessment factor of 1000 (TGD, Table 20).

$$\begin{aligned}
 PNEC_{\text{soil}} &= 53.9 \text{ mg OPP} \cdot \text{kg}^{-1} \text{ dry weight soil} \cdot 10^{-3} \\
 &= 0.054 \text{ mg OPP} \cdot \text{kg}^{-1} \text{ dry weight soil} \\
 &= 0.054 * 1.13 \\
 PNEC_{\text{soil}} &= 0.061 \text{ mg OPP} \cdot \text{kg}^{-1} \text{ wet weight soil}
 \end{aligned}$$

Non-compartment specific effects relevant to the food chain(secondary poisoning)

A flow-through study was conducted to evaluate the bioconcentration of OPP in zebra fish (*Danio rerio*). The arithmetic means of five consecutive steady-state BCF were 21.7 (wet weight), 114-115 (lipid content), indicating a negligible potential of the test substance to bioaccumulate. The achievement of steady-state conditions during the uptake (53 h exposure) phase as well as the consecutive depuration (19 h) were rapid processes.

A risk due to the proposed uses of OPP can be ruled out, since these data show that OPP does not accumulate in the environment. There is no need to assess this exposure route further.

The summary of ecotoxicity data used for the risk assessment are summarised in the Table 2.2.2.2-1.

Table 2.2.2.2-1: Summary of toxicity data used for the risk assessment

Species	Endpoint /Type of test	Results [mg a.i./L]
<i>Oncorhynchus mykiss</i>	Fish acute 96 h - LC ₅₀ Mortality	1
<i>Daphnia magna</i>	Aquatic invertebrates acute 48 h - LC ₅₀ Mortality	2.7
<i>Pseudo-kirchneriella subcapitata</i>	Algae growth inhibition 72 h - NOEC Growth inhibition	0.468
Activated sludge	Microorganisms 3 h - respiration inhibition	56
<i>Pimephales promelas</i> (Fathead minnow)	Fish chronic 21 d - NOEC Reproduction (Egg hatch F1) 21 d - LOEC Reproduction (Egg hatch F1)	36 293
<i>Daphnia magna</i>	Aquatic invertebrates chronic 21 d - NOEC Reproduction	0.006
<i>Avena sativa</i>	14 d - EC ₅₀ Germination rate, mortality and phytotoxicity	53.9
<i>Eisenia fetida</i>	Earthworms 14 d -LC ₅₀ Mortality, weight, abnormal behaviour	198.2
Soil microorganisms	28 d - EC ₅₀ nitrification	633.5
Mallard duck	Birds 14 d - LC ₅₀	>2250
Mallard duck	Birds 5 d - LD ₅₀	>5620
Rat Fischer 344	Mammals acute LD ₅₀ 1 dose + 2 weeks of observation	2733 mg/kg
Beagle Dogs	Mammals chronic NOAEL 1 year	300 mg/kg/day

2.2.2.3. PBT and POP assessment

Assessment of PBT criteria

OPP can be considered readily biodegradable. Monitoring and laboratory studies have also shown that OPP is easily removed in STP systems. Based on literature studies, OPP is also not persistent water-sediment systems, and a soil biodegradation study also has shown that OPP is removed either by sorption or by biodegradation process. Considering the hydrolytic stability determined under stringent temperature conditions and at different pH values it is not expected that hydrolytic processes will contribute to the degradation of OPP in the aquatic systems (estimated DT₅₀ > 1 year), however, from the photolysis study in water, it has been shown that OPP is photolytically unstable in the aqueous medium. Therefore, it is unlikely that OPP persists in the water, sediment or soil compartments.

The assessment of the (potential for) bioaccumulation in the context of PBT or vPvB evaluation makes use of measured bioconcentration factor. When not available, BCF value may be estimated from the octanol/water partition coefficient (K_{ow}) by using (Q)SAR models. The calculated steady-state bioconcentration factor (BCF) for fish of 21.7 L/kg (wet weight), 114-115 (lipid content), indicates a negligible potential of OPP to bioaccumulate. Therefore, OPP does not fulfil the B criterion since its BCF is under the cut-off values proposed in the TGD (BCF > 2,000 for PBT assessment and > 5,000 for vPvB assessment).

The lowest NOEC obtained for OPP was 0.006 mg/L (*Daphnia magna* test). Since the cut off value given by the TGD corresponds to 0.01 mg/L, the substance meets the T criterion.

Assessment of POPs criteria

The vapour pressure of OPP is 0.906 Pa at 25°C, the half-life in air is of 0.587 days, indicating that the criteria for long-range transport potential (vapour pressure < 1000 Pa and half-life in air > 2 days) is not fulfilled. In soil, biodegradation and sorption study was performed to understand the persistence of OPP in this compartment, indicating that OPP is relatively low mobile in soil, although a biodegradation character can also be attributed.

The calculated steady-state bioconcentration factor (BCF) for fish is 21.7 L/kg (wet weight), 114-115 (lipid content), and hence < 5000. Thus, the bioaccumulation criterion is not fulfilled for OPP.

In conclusion, considering the above rationale, it can be concluded that OPP does not fulfil the POPs criteria.

Conclusion:

OPP must not be regarded as a Persistent or Bioaccumulative, Toxic, POP or ED substance because it does not fulfil the criteria. Therefore, OPP is not PBT/vPvB.

2.2.2.4. Exposure assessment

The biocidal product Preventol O Extra presented is the active substance ($\geq 99.5\%$) but it is not actually the final formulation that will be used. Therefore, data of real biocidal product will need to be demanded at national product actualization.

The biocidal PT 13 product Preventol ON Extra Preservative Solution contains ca. 17.7% OPP in alkaline solution. Here due to the pH of > 12, OPP and its alkali salt are present in equilibrium. Environmental exposure may occur by unintended splashes of the biocidal product itself, cleaning operations, or by releases of a preserved matrix (pH < 9) to the environmental compartments. Due to the buffer capacity of the aquatic compartment and the high dilution effect of any alkali release (starting with the waste water channel and the STP) even the local environmental pH will not be affected significantly and will constantly stay below 9, so that a conversion of any released alkali salt form to the genuine phenolic form instantly takes place (the equilibrium is quantitatively expressed in the Henderson Hasselbach equation). Thus for the environment only the exposure to the free phenolic compound OPP must be assessed.

In the present assessment for Preventol O Extra and Preventol ON Extra Preservative Solution, sewage water treatment plants are regarded as the only pathway of direct OPP emissions after use as metalworking fluid (MWF) preservative (see Doc. II-B).

For the environmental risk assessment, two different emission rates (Scenario 1 and 2) resulting from the exchange of the functional fluid have been assessed.

The emission rates to a standard STP are calculated to be 0.329 kg/d for Scenario 1 and 0.165 kg/d for Scenario 2. Predicted Environmental Concentration (PEC) values were determined for different environmental compartments in Doc. II-B.

2.2.2.5. Risk characterisation

Aquatic compartment (incl. sewage treatment plant)

The following risk quotients were derived for the aquatic compartment from the calculated/measured exposure and effect data for OPP (see Table 2.2.2.5-1).

Table 2.2.2.5-1: PEC/PNEC ratios for OPP (aquatic compartment)

Compartments		PEC values		PEC/PNEC	
		Scenario 1 ¹	Scenario 2 ²	Scenario 1 ¹	Scenario 2 ²
STP effluent [mg/L]	Tier 1	8.09E-03	4.05E-03	0.014	0.007
	Tier 2	6.59E-04	3.29E-04	0.001	0.0006
Local concentration in surface water during emission episode [mg/L]	Tier 1	8.09E-04	4.04E-04	1.348	0.675
	Tier 2	6.58E-05	3.29E-05	0.11	0.055
Sediment [mg/kg]	Tier 1	6.74E-03	3.37E-03	1.375	0.688
	Tier 2	5.48E-04	2.74E-04	0.112	0.056

¹ 0.30% = 3000 mg/kg OPP

² 0.15% = 1500 mg/kg OPP

Tier 1: 12.31% of the influent residues being present in the STP effluent water phase

Tier 2: 1% of the influent residues being present in the STP effluent water phase

Sewage treatment plant: The derived risk quotients are clearly < 1, even using the worst-case assumption (Tier 1 approach) of 12.3% of the influent residues being present in the STP effluent water phase for the calculation. Thus, it is considered that there is no risk for microorganisms in a STP caused by OPP used as metalworking fluid preservative.

Surface water: The PEC/PNEC ratios are above 1 for Tier 1 with an in-use concentration of OPP in emulsifiable MWF of 0.3%. The PEC/PNEC ratios for Tier 2 are below 1 therefore, there is no risk to aquatic organisms in surface waters exposed to OPP used as metalworking fluid preservative when the elimination rate obtained in a STP is > 85% for OPP.

Sediment: The PEC/PNEC ratios are above 1 for Tier 1. The PEC/PNEC ratios are < 1 for Tier 2. Therefore, sediment dwelling organisms are not at risk by the intended uses of OPP used as metalworking fluid preservative when the elimination rate obtained in a STP is > 85% for OPP.

All derived risk quotients are clearly < 1 when the in-use concentration of OPP in emulsifiable MWF amounts was 0.15% = 1500 mg/kg and for the reasonable worst case Tier 2. Thus, it is considered that there is no relevant risk for the aquatic environment caused by OPP used as metalworking fluid preservative.

Terrestrial compartment (soil)

To assess the risk for the environmental compartment soil regarding the exposure via sludge, the PNEC_{soil} is compared with the PEC_{soil} (see Table 2.2.2.5-2).

Table 2.2.2.5-2: PEC/PNEC ratios for OPP (terrestrial compartment)

		PEC _{soil} values Concentration in agricultural soil over 30 days [mg/kg _{wwt}]		PEC/PNEC	
		Scenario 1 ¹	Scenario 2 ²	Scenario 1 ¹	Scenario 2 ²
DT ₅₀ = 30 d	Tier 1 ³	5.47E-03	2.73E-03	0.09	0.045
	Tier 2 ⁴	1.75E-03	8.73E-04	0.029	0.014
DT ₅₀ = 15.08 d	Tier 1 ³	4.11E-03	2.06E-03	0.068	0.034
	Tier 2 ⁴	1.31E-03	6.56E-04	0.022	0.011

¹ 0.30% = 3000 mg/kg OPP² 0.15% = 1500 mg/kg OPP³ Tier 1: 3.13% of the STP influent residues being present in STP sludge⁴ Tier 2: 1% of the STP influent residues being present in STP sludge

As the PEC/PNEC ratios are < 1, no relevant risk for soil organisms is indicated due to the use of OPP as metalworking fluid preservative.

Groundwater compartment

According the EU TGD (European Commission, 2003), the predicted concentration of the active substance in soil pore water is taken as a surrogate estimate of the potential concentration in groundwater. No accepted ecological endpoints have been established to enable characterisation of risk to the groundwater compartment (European Commission, 2003). However, the groundwater directive (Directive 2006/118/EC) stipulates a maximum acceptable concentration for pesticides in groundwater of 0.1 µg·L⁻¹. The PECs values are given in Table 2.2.5.2-3.

Table 2.2.2.5-3: PEC values for OPP (groundwater)

		PEC _{qw} values [mg · L ⁻¹]		PEC _{qw} values [µg · L ⁻¹]	
		Scenario 1 ¹	Scenario 2 ²	Scenario 1 ¹	Scenario 2 ²
DT ₅₀ = 30 d	Tier 1 ³	2.80E-04	1.40E-04	0.28	0.14
	Tier 2 ⁴	8.94E-05	4.47E-05	0.09	0.05
DT ₅₀ = 15.08 d	Tier 1 ³	1.45E-04	7.26E-05	0.15	0.07
	Tier 2 ⁴	4.63E-05	2.32E-05	0.05	0.02

¹ 0.30% = 3000 mg/kg OPP² 0.15% = 1500 mg/kg OPP³ Tier 1: 3.13% of the STP influent residues being present in STP sludge⁴ Tier 2: 1% of the STP influent residues being present in STP sludge

From the values presented above it can be seen that emissions associated with the use of OPP as metalworking fluids result in porewater concentrations exceeding this threshold only for worst case Tier 1 and DT₅₀ = 30 days.

However, simulations with FOCUS PEARL for groundwater prove that PEC_{gw} values (80th percentiles of the annual average concentrations in the percolate at 1 m soil depth) of OPP were of <0.0001 µg/L in all scenarios (see Annex PT 13 and Table 2.2.2.5-4).

Table 2.2.2.5-4: Predicted 80th percentile concentrations for OPP in groundwater with FOCUS PEARL

	FOCUS Scenarios	
	Concentration closest to the 80 th percentile [µg·L ⁻¹]	
	Arable land (Maize)	Grassland (Alfalfa)
Châteaudun	<0.0001	<0.0001
Hamburg	<0.0001	<0.0001
Jokioinen	<0.0001	<0.0001
Kremsmünster	<0.0001	<0.0001
Okehampton	<0.0001	<0.0001
Piacenza	<0.0001	<0.0001
Porto	<0.0001	<0.0001
Sevilla	<0.0001	<0.0001
Thiva	<0.0001	<0.0001

It is therefore concluded that OPP no represent a risk to groundwater following the application of sewage sludge to land even in the worst of the cases.

Non compartment specific effects relevant to the food chain (secondary poisoning)

A flow-through study was conducted to evaluate the bioconcentration of OPP in zebra fish (*Danio rerio*). The arithmetic means of five consecutive steady-state BCF were 21.7 (wet weight), 114-115 (lipid content), indicating a negligible potential of the test substance to bioaccumulate. The achievement of steady-state conditions during the uptake (53 h exposure) phase as well as the consecutive depuration (19 h) were rapid processes.

A risk due to the proposed uses of OPP can be ruled out, since these data show that OPP does not accumulate in the environment. There is no need to assess this exposure route further.

A secondary exposure of OPP to man via the food chain can be excluded due to low tonnage of the biocidal product used in whole Europe, rapid degradation in water and minimum amounts which reach the environmental compartments. A risk due to the proposed uses of OPP can be ruled out, since these data show that OPP does not accumulate in the environment. There is no need to assess this exposure route further.

2.2.2.6. Assessment of endocrine disruptor properties

In relation to the potential of OPP to interfere with the hormone system, OPP is present in one of the documents-lists of the Commission staff working document on implementation of the Community Strategy for Endocrine Disruptors - a range of substances suspected of interfering with the hormone systems of humans and wildlife (COM(2004) 1372), and cited as "candidate substance" for a first-in depth study. No endocrine disruption effect was reported in this document or in the following (COM(2007) 1635).

In addition, the prolonged toxicity of OPP to fathead minnow (*Pimephales promelas*) was

tested in a reproductive performance test by ██████████ (2002). In the test, measures of fecundity were assessed daily. Viability of resultant embryos was assessed in animals held in the same treatment regime to which the adults were exposed. A suite of histological and biological endpoints, that potentially are directly reflective of effects associated with endocrine disrupting chemicals, was also evaluated. The results of the study show that OPP does not indicate any adverse effects on reproductive parameters of pair-breeding fathead minnows up to a nominal test concentration of 50 µg a.i./L. With regard to the induction of the biomarker vitellogenin as an early indicator of possible endocrine modulation, no substance-related effects were noted compared to the positive control 17α-ethynylestradiol.

Result of the first EU evaluation project on potential endocrine substances (EUROPEAN COMMISSION, STUDY ON THE SCIENTIFIC EVALUATION OF 12 SUBSTANCES IN THE CONTEXT OF ENDOCRINE DISRUPTER PRIORITY LIST OF ACTIONS, 2002).

From the summary for humans: "The available data from in vivo studies in laboratory mammals (using oral or dermal exposure routes) indicates that *o*-Phenylphenol does not cause adverse effects on reproductive and developmental endpoints (which may be endocrine mediated) at exposure levels where general systemic toxic effects are observed. The lowest NOEL in the in vivo studies was 250 mg·kg_{bw}⁻¹·day⁻¹ for foetotoxic and developmental effects. Limited exposure data for workers and consumers has been located."

For wildlife: "The available aquatic effects data shows that the threshold exposure concentrations of *o*-Phenylphenol above which reproduction of the invertebrate *Daphnia magna* and fish (fathead minnow) are reduced (NOECs = 0.036 mg·L⁻¹ and 0.009 mg·L⁻¹ respectively) are lower than the threshold levels for general toxic effects (i.e. lethality). The effects observed on reproduction in fish were evidently not oestrogen mediated. However, there is no information on the mechanism of action for the effects on reproduction observed in *Daphnia magna*."

The results of this EU evaluation project were also confirmed in a peer evaluation done by the CSTE (2003)

Thus, it can be stated that, to date, no evidence of endocrine disruption activity can be attributed to OPP.

2.3. Overall conclusions

The outcome of the assessment for Biphenyl-2-ol in Product-type 13 is specified in the BPC opinion following discussions at the BPC-9 meeting of the Biocidal Products Committee (BPC). The BPC opinion is available from the ECHA website.

Data on the ventilation system shall be submitted at the biocidal product registration phase.

Further information STP simulation tests will be required six months before the approval of the active substance to support the degradation rate.

2.4. List of endpoints


The most important endpoints, as identified during the evaluation process, are listed in [Appendix I](#).

Appendix I: List of endpoints

Chapter 1: Identity, Physical and Chemical Properties, Classification and Labelling

Active substance (ISO Name)	2-Phenylphenol
Product-type	Working or cutting fluid preservatives

Identity

Chemical name (IUPAC)	2-Phenylphenol
Chemical name (CA)	[1,1'-Biphenyl]-2-ol
CAS No	90-43-7
EC No	201-993-5
Other substance No.	CIPAC No. 246
Minimum purity of the active substance as manufactured (g/kg or g/l)	≥ 995 g/kg
Identity of relevant impurities and additives (substances of concern) in the active substance as manufactured (g/kg)	None
Molecular formula	C ₁₂ H ₁₀ O
Molecular mass	170.2 g/mol
Structural formula	

Physical and chemical properties

Melting point (state purity)	56.7 °C (purity: 99.9%)
Boiling point (state purity)	287 °C (purity: 99.9%)
Thermal stability / Temperature of decomposition	Exothermal decomposition starts at 290 °C. As no decomposition of the test substance could be observed below 150 °C, Biphenyl-2-ol is considered to be stable at room temperature.
Appearance (state purity)	Colourless solid flakes with slight phenolic odour (purity: 99.9%)
Relative density (state purity)	1.237 at 20 °C (purity: 99.9%)
Surface tension (state temperature and concentration of the test solution)	58.72 mN/m at 20.1 °C (0.558 g/L)
Vapour pressure (in Pa, state temperature)	0.474 Pa at 20 °C, 0.906 Pa at 25 °C
Henry's law constant (Pa m ³ mol ⁻¹)	Ratio between vapour pressure and water solubility: 0.15 Pa×m ³ ×mol ⁻¹ at 20 °C and pH 5 0.14 Pa×m ³ ×mol ⁻¹ at 20 °C and pH 7 0.13 Pa×m ³ ×mol ⁻¹ at 20 °C and pH 9

Solubility in water (g/L or mg/L, state temperature)	<p>Results at pH 5: 0.43 g/L at 10°C 0.53 g/L at 20°C 0.70 g/L at 30°C</p> <p>Results at pH 7: 0.45 g/L at 10°C 0.56 g/L at 20°C 0.73 g/L at 30°C</p> <p>Results at pH 9: 0.52 g/L at 10°C 0.64 g/L at 20°C 0.84 g/L at 30°C</p>
Solubility in organic solvents (in g/L or mg/L, state temperature)	<p>Results at 20 °C: <i>n</i>-heptane: 50.3 g/L acetone, 1,2-dichloroethane, ethyl acetate, methanol, <i>p</i>-xylene: > 250 g/L No significant temperature dependence is expected.</p>
Stability in organic solvents used in biocidal products including relevant breakdown products	<p>Biphenyl-2-ol as manufactured does not include an organic solvent in PT 2, 3, 4, 6, 7, 10 and 13. Therefore a study regarding stability in organic solvents does not apply. The b. p. for PT 1 and 9 contains an organic solvent.</p>
Partition coefficient (log P _{ow}) (state temperature)	<p>Log P_{ow}: 3.18 at 22.51 °C. (more accurate value which is to be used exclusively) "the log P_{ow} of Biphenyl-2-ol is nearly independent from pH value when investigated at pH 5, pH 7 and pH 9."</p>
Dissociation constant	<p>pK = 9.5 at 20 °C</p>
UV/VIS absorption (max.) (if absorption > 290 nm state ε at wavelength)	<p>Molar absorptivity: 12800 at 245 nm 8200 at 267 nm The UV-visible spectrum show a band with a maximum at 285 nm and a bandwidth of 40 nm, therefore a short absorption appears above 290 nm.</p>
Flammability or flash point	<p>Biphenyl-2-ol is not highly flammable, does not liberate gases in hazardous amounts when contact with water, does not deliver indications of pyrophoric properties and does not undergo spontaneous combustion.</p>
Explosive properties	<p>Based on scientific judgement it is certified that due to the structural formula Biphenyl-2-ol contains neither oxidising groups nor other chemically instable functional groups. Thus Biphenyl-2-ol is incapable of rapid decomposition with evolution of gases or release of heat, i.e. the solid material does not present any risk for explosion.</p>

Oxidising properties

Based on scientific judgement it is certified that due to the structural formula Biphenyl-2-ol does not contain oxidising groups in its molecular backbone and thus may not react exothermically with a combustible material. Therefore Biphenyl-2-ol does not have oxidising properties.

Auto-ignition or relative self ignition temperature

Biphenyl-2-ol does not undergo spontaneous combustion

Classification and proposed labelling

with regard to physical hazards

None

with regard to human health hazards

Carc 2: H351; Eye Irrit. 2: H319; Skin Irrit. 2: H315; STOT SE 3: H335

with regard to environmental hazards

Aquatic Acute 1:H400; Aquatic Chronic 1; H410

Chapter 2: Methods of Analysis

Analytical methods for the active substance

Technical active substance (principle of method)

Biphenyl-2-ol is separated by means of gas chromatography using flame ionisation detection. The quantitative evaluation is carried out by area normalisation with consideration of water content and non-volatile components.

Impurities in technical active substance (principle of method)

The analytical method for the determination of impurities in the active substance is confidential. This information is provided separately in the confidential part of the dossier.

Analytical methods for residues

Soil (principle of method and LOQ)

HPLC-MS/MS; LOQ = 5 µg/kg

Air (principle of method and LOQ)

GC-MS; LOQ = 0.35 µg/m³.

Water (principle of method and LOQ)

Surface and drinking water: HPLC-MS/MS; LOQ = 0.1µg/L

Body fluids and tissues (principle of method and LOQ)

Not applicable since Biphenyl-2-ol is not classified as toxic or highly toxic.

Food/feed of plant origin (principle of method and LOQ for methods for monitoring purposes)

Citrus Fruit: GC-MS; LOQ = 0.1 µg/kg
QuEChERS Method: EN155662:2008

Food/feed of animal origin (principle of method and LOQ for methods for monitoring purposes)

Meat: GC-MS/MS; LOQ = 0.01 µg/kg

Chapter 3: Impact on Human Health

Absorption, distribution, metabolism and excretion in mammals

Rate and extent of oral absorption:	100% is assumed
Rate and extent of dermal absorption* :	43.19% is assumed
Distribution:	Extensively metabolized. Poorly distributed.
Potential for accumulation:	Low potential for bioaccumulation.
Rate and extent of excretion:	Quickly excreted (12 - 24 h post-dosing).
Toxicologically significant metabolite(s)	No data

* the dermal absorption value is applicable for the active substance and might not be usable in product authorization

Acute toxicity

Rat LD ₅₀ oral	2730.3 mg/kg bw
Rat LD ₅₀ dermal	> 2000 mg/kg bw
Rat LC ₅₀ inhalation	> 36 mg/m ³ (0.036 mg/L)

Skin corrosion/irritation

Skin Irrit. 2 (H315: Causes skin irritation)

Eye irritation

Eye Irrit. 2 (H319: Causes serious eye irritation)

Respiratory tract irritation

No data

Skin sensitisation (test method used and result)

Non Sensitizer (Buehler test on Guinea pigs; 0/10 Number of animals sensitised/total number of animals)
Non Sensitizer (Magnusson-Kligman test on Guinea pigs; 0/20 Number of animals sensitised/total number of animals)

Respiratory sensitisation (test method used and result)

No data

Repeated dose toxicity

Short term

Species/ target / critical effect

Oral: New Zealand White rabbits / increased mortality (13%), gross pathologic alterations and histopathologic alterations
Dermal: Fischer 344 rats/ no systemic effects in any dose group

Relevant oral NOAEL / LOAEL

NOAEL = 100 mg/kg bw/day (teratogenicity oral study)

LOAEL = 250 mg/kg bw/day (teratogenicity oral study)

Relevant dermal NOAEL / LOAEL

NOAEL = 1000 mg/kg bw/day (21-day dermal study)

Relevant inhalation NOAEL / LOAEL

No data

Relevant NOAEC (local effects)

7.5%

Subchronic

Species/ target / critical effect

Rats /urinary bladder/ increased incidence of simple urinary bladder hyperplasia in males and the increased incidence of urinary bladder transitional cell carcinoma in males

Relevant oral NOAEL / LOAEL

NOAEL = 39 mg/kg bw/day (2-years oral study)

LOAEL = 250 mg/kg bw/day (2-years oral study)

Relevant dermal NOAEL / LOAEL

No Data

Relevant inhalation NOAEL / LOAEL

No Data

Relevant NOAEC (local effects)

No Data

Long term

Species/ target / critical effect

Rats /urinary bladder/ increased incidence of simple urinary bladder hyperplasia in males and the increased incidence of urinary bladder transitional cell carcinoma in males

Relevant oral NOAEL / LOAEL

NOAEL = 39 mg/kg bw/day (2-years oral study)

LOAEL = 250 mg/kg bw/day (2-years oral study)

Relevant dermal NOAEL / LOAEL

No Data

Relevant inhalation NOAEL / LOAEL

No Data

Relevant NOAEC (local effects)

No Data

Genotoxicity*In vitro*

Biphenyl-2-ol is considered to be nonmutagenic but it was clastogenic in Chinese hamster ovary cells at cytotoxic concentrations

In vivo

Biphenyl-2-ol is not genotoxic or mutagenic in vivo.

Carcinogenicity

Species/type of tumour	Fischer 344 rat/ neoplasia in urinary bladder (male animals only) B6C3F1 mice/ hepatocellular adenomas(male animals only) The tumours found in mice are not predictive of carcinogenicity for humans. The relevance of urinary bladder tumours in male rats cannot be completely excluded
Relevant NOAEL/LOAEL	200 mg/kg body wt/day 500 mg/kgBW/day

Reproductive toxicityDevelopmental toxicity

Species/ Developmental target / critical effect	New Zealand White rabbits/ No recorded effect on development parameters/ No effects on foetal development
Relevant maternal NOAEL	NOAEL = 100 mg/kg/day
Relevant developmental NOAEL	NOAEL = 250 mg/kg/day

Fertility

Species/critical effect	RatCD Sprague-Dawley/ No recorded effect on reproductive parameters/ bladder calculi, urothelial hyperplasia
Relevant parental NOAEL	NOAEL = 35 mg / kg bw / day
Relevant offspring NOAEL	NOAEL = 125 mg / kg bw / day
Relevant fertility NOAEL	NOAEL = 457 mg / kg bw / day

Neurotoxicity

Species/ target/critical effect	No data
---------------------------------	---------

Developmental Neurotoxicity

Species/ target/critical effect	No data
---------------------------------	---------

Immunotoxicity

Species/ target/critical effect	No data
---------------------------------	---------

Developmental Immunotoxicity

Species/ target/critical effect	No data
---------------------------------	---------

Other toxicological studies

Human data: allergic contact dermatitis or contact sensitivity to Biphenyl-2-ol Other/special studies: Biphenyl-2-ol is carcinogenic in urinary bladder in alkaline conditions in rats

Medical data

No data

Summary

	Value	Study	Safety factor
AEL _{long-term}	0.4 mg/kg bw/day	2-years oral study	100
AEL _{medium-term}	0.4 mg/kg bw/day	2-years oral study	100
AEL _{short-term}	1 mg/kg bw/day	teratogenicity oral study in New Zealand White rabbits	100
ADI ²	0.4 mg/kg bw/day	2-years oral study	100
ARfD	No relevant		

MRLs

Relevant commodities

--

Reference value for groundwater

According to BPR Annex VI, point 68

--

Dermal absorption

Study (*in vitro/vivo*), species tested

Formulation (formulation type and including concentration(s) tested, vehicle)

Dermal absorption values used in risk assessment

Dermal absorption, excretion in vivo, humans.
0.4% (w/v) OPP solution in isopropyl alcohol
43% (100% in corrosive products)

²If residues in food or feed.

Acceptable exposure scenarios (including method of calculation)

Formulation of biocidal product	<p>Preventol O Extra contains ca. $\geq 99.5\%$ of the active substance (a.s.) Biphenyl-2-ol. Formulated into a metal working fluid concentrate up to 6% (60,000 ppm a.s.) and added to the system.</p> <p>Premix containing 6% Biphenyl-2-ol administered directly to the sump of a metalworking operation to a final use dilution concentration.</p> <p>Final use-dilution concentration for Biphenyl-2-ol in metalworking fluid is 0.15 to 0.3% (1,500 to 3,000 ppm).</p> <p>Preventol ON Extra Preservative Solution is an alkaline ready-for-use preservative solution (c.a 17.7% Biphenyl-2-ol, w/w) Likely concentration at use is 0.159 – 0.3% w/v</p>
Intended uses	preservative to control the growth of fungi in metalworking fluids
Industrial users	<p>Formulation: Mix and Load model 7, TNsG 2002, pp.141, revised.</p> <p>PPE (RPE if aerosol or dust is produced) required when handling biocidal product.</p> <p>RMMs recommended to reduce airborne levels.</p> <p>No risk.</p>
Professional users	<p>Biocide replenishment (maximum 6% OPP in MWF): Mix and Load model 7, TNsG 2002, pp.141, revised.</p> <p>Application (metal working activities, maximum 0.3%OPP in MWF): Metal Working Fluid model 2 (2002, part 12, pp. 189, TNsG on Human Exposure), BEAT's worked example for PT 13 (Machining of metal parts.</p> <p>Post application (other tasks, maximum 0.3% OPP in MWF): Metal working fluid model 2 (TNsG, 2002, pp.189), Cleaning of Spray Equipment from BEAT database (TNsG 2007).</p> <p>PPE (coverall) required.</p> <p>RMMs recommended to reduce dermal exposure.</p> <p>No risk.</p>
Non professional users	Not applicable
Indirect exposure	<p>Professional user washing dirty coverall at home.</p> <p>No risk.</p>

Exposure via residue in food

Human exposure to residues of Biphenyl-2-ol on food and feedstuffs can be excluded when the product is applied according to the recommended use

Chapter 4: Fate and Behaviour in the Environment

Route and rate of degradation in water

Hydrolysis of active substance and relevant metabolites (DT₅₀) (state pH and temperature)

pH 5: stable at 50 °C
pH 7: stable at 50 °C
pH 9: stable at 50 °C
Estimated t_{1/2} > 1 year

Photolytic / photo-oxidative degradation of active substance and resulting relevant metabolites

Biphenyl-2-ol:
Experimental DT₅₀: 0.3 days (pure water)
Environmental DT₅₀ [Phoenix, AZ, USA]: 1.7 days
Environmental DT₅₀ [Athens, Greece]: 2.6 days
Diketohydroxy-compound (max. 13.6% at day 1, < 5% after 7 days):
Experimental DT₅₀: 1.3 days (pure water)
Environmental DT₅₀ [Phoenix, AZ, USA]: 7.2 days
Environmental DT₅₀ [Athens, Greece]: 11.1 days

Readily biodegradable (yes/no)

Yes;
71-76% biodegradation after 28 d
100% biodegradation after 14 d
100% biodegradation after 10 d (inherent test)

Inherent biodegradable (yes/no)

Biodegradation in freshwater

Biodegradation in seawater

Not relevant since Biphenyl-2-ol is not used or released in the marine environment at considerable amounts. Therefore, a seawater biodegradation test is not required.

Non-extractable residues

Not relevant due to indoor use.

Distribution in water / sediment systems (active substance)

Not relevant due to indoor use.
Estimation from screening experiments: < 14 d

Distribution in water / sediment systems (metabolites)

Not relevant due to indoor use.

Route and rate of degradation in soil

Mineralization (aerobic)

Results are given as mean value of duplicate test of [phenyl-UL-¹⁴C]-labelled Biphenyl-2-ol in % of the applied radioactivity for day 127 of incubation under aerobic conditions:
9.6% (n = 2, 20 ± 1 °C)

Laboratory studies (range or median, with number of measurements, with regression coefficient)

DT_{50 lab} (20 °C, aerobic):
2.7 hours* (n = 1), r² = 0.994

DT_{90lab} (20 °C, aerobic):
8.81 hours* (n = 1), r² = 0.994

degradation in the saturated zone:

Field studies(state location, range or median with number of measurements)

Not relevant due to indoor use

Anaerobic degradation

Not relevant due to indoor use.

Soil photolysis

Not relevant due to indoor use.

Non-extractable residues

77.4% at day 127 (n = 2, 20 ± 1 °C)

Relevant metabolites - name and/or code, % of applied a.i. (range and maximum)

No relevant metabolites

Soil accumulation and plateau concentration

Not relevant due to indoor use

Adsorption/desorption

K_a , K_d

K_{aoc} , K_{doc}

pH dependence (yes / no) (if yes type of dependence)

Adsorption, OECD Guideline 106:

K_f: 7.04 , 7.47, 8.53, 11.66 (n = 4)

K_{oc}: 252, 355, 389, 393 (n = 4, mean: 347)

Desorption 1:

K_{fdes}: 9.36, 16.42, 16.78, 18.62 (n = 4)

K_{ocdes}: 334, 621, 699, 864 (n = 4)

Adsorption, OECD Guideline 121:

estimated mean K_{oc} value: 346.7

K_d was not reported

pH dependence was not apparent

Fate and behaviour in air

Direct photolysis in air

Not relevant because there is no relevant release of the compound to the air compartment

Quantum yield of direct photolysis

Photo-oxidative degradation in air

DT₅₀ = 0.59 days

Volatilization

Not relevant because there is no relevant release of the compound to the air compartment

Reference value for groundwater

According to BPR Annex VI, point 68

Monitoring data, if available

Soil (indicate location and type of study)

No data presented

Surface water (indicate location and type of study)

Municipal sewage plant Steinhäule located on the Danube River in southern Germany. The plant has mechanical purification devices (primary clarification), activated sludge treatment, biological nitrate removal (nitrification/denitrification), biological phosphate removal and final settlement tanks as main cleaning steps. Concentrations of Biphenyl-2-ol in 24 h influent and effluent samples from 10/11 March 1998

Substance (µg/L)	Influent 10/11 March (8 a.m-8a.m)	Effluent 10/11 March (4 p.m-4 p.m)
Biphenyl-2-ol	1.54 ± 0.349	< 0.015

Ground water (indicate location and type of study)

No data presented

Air (indicate location and type of study)

No data presented

Chapter 5: Effects on Non-target Species**Toxicity data for aquatic species (most sensitive species of each group)**

Species	Time-scale	Endpoint	Toxicity
Fish			
<i>Oncorhynchus mykiss</i>	96 hours	Mortality	LC ₅₀ = 4.0 mg/L Dill <i>et al.</i> (1985)
<i>Pimephales promelas</i>	21 days	Reproduction	NOEC = 0.036 mg/L ████████████████████ (2002)
Invertebrates			
<i>Daphnia magna</i>	48 hours	Mortality	LC ₅₀ = 2.7 mg/L Dill <i>et al.</i> (1985)
<i>Daphnia magna</i>	21 days	Survival & reproduction	NOEC = 0.006 mg/L Bruns (2001)
Algae			

<i>Pseudokirchneriella subcapitata</i>	72 hours	Growth inhibition	$E_rC_{50} = 3.57 \text{ mg/L}$ $E_bC_{50} = 1.35 \text{ mg/L}$ $NOEC = 0.468 \text{ mg/L}$ Hicks (2001)
Microorganisms			
Activated sludge	3 hours	Inhibition of respiratory rate	$EC_{50} = 56 \text{ mg/L}$ Klecka, Landi, and Bodner (1985)

Effects on earthworms or other soil non-target organisms

Acute toxicity to earthworms ..

LC_{50} (14 days) = 198.2 mg/kg
Moser & Scheffczyk (2004)

Reproductive toxicity to earthworms

No study available

Effects on soil micro-organisms

Nitrogen mineralization

EC_{50} (28 days) = 633.5 mg a.s./kg d.wt. soil
Schulz.L (2012)

Carbon mineralization

Effects on terrestrial vertebrates

Acute toxicity to mammals

$LD_{50} = 2733 \text{ mg/kg bw}$ (♂+♀)
[redacted] (1994)

Chronic toxicity to mammals
(Annex IIA, point VI.6.5)

NOAEL = 300 mg/kg diet (1 year)
Cosse *et al.* (1990)

Acute toxicity to birds

$LC_{50} > 2250 \text{ mg/kg bw}$
[redacted] (1986)

Dietary toxicity to birds

$LD_{50} > 5620 \text{ mg/kg diet}$
[redacted] (1986)

Reproductive toxicity to birds

No study available

Effects on honeybees

Acute oral toxicity

No study available

Acute contact toxicity

No study available

Effects on other beneficial arthropods

Acute oral toxicity

No study available

Acute contact toxicity

No study available

Acute toxicity to

No study available

Bioconcentration

Bioconcentration factor (BCF)

BCF = 21.7 (whole fish), 114-115 (lipid content)
Caspers (1999)Depuration time(DT₅₀)

< 1 h (5 µg/L) / < 19 h (50 µg/L)

Depuration time(DT₉₀)

2 h (5 µg/L) / < 6 h (50 µg/L)

Level of metabolites (%) in organisms accounting for > 10 % of residues

No metabolites identified

Chapter 6:Other End Points

Appendix II: List of Intended Uses

Object and/or situation	Product name	Organisms controlled	Formulation		Application			Applied amount per treatment			Re marks:
			Type (d-f)	Conc. of a.s.(i)	method kind (f-h)	number min max	interval between applications (min)	g a.s./L min max	water L/m ² min max	g a.s./m ² min max	
Preservation of metal-working fluids PT 13	Preventol O Extra	fungi and yeasts The test period was 5 weeks	AL	177 g/kg	addition	-	?	0.5-3 g/L	-	-	-
Preservation of metal-working fluids PT 13	Preventol ON Extra Preservative Solution	fungi and yeasts The test period was 5 weeks	AL	177 g/kg	addition	-	?	0.5-3 g/L	-	-	-

Appendix III: List of studies

Data protection is claimed by the applicant in accordance with Article 60 of Regulation (EU) No 528/2012.

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A2.6(01) IIA, II 2.6	Stroech, K.D.	1991	Preventol O Extra (2-Phenylphenol) Synthesis. Date: 1991-02-19 CONFIDENTIAL	Bayer AG, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A2.7(01) IIA, II 2.7	Anonymous	2000	Preventol O Extra in flakes. Date: 2000-02-11	BU, Material Protection Products, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A2.7(01) IIA, II 2.7 also filed: A2.8(01)	Erstling, K.	2005	Determination of main and minor components in Preventol O Extra, 5-batch analysis. Date: 2005-02-16 CONFIDENTIAL	Bayer Industry Services GmbH & Co. OHG, BIS- SUA-Analytics, Leverkusen, Germany	Study No.: G 05/0009/00 LEV	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A2.7(02) IIA, II 2.7	Stroeck, K.	2014	Quality Control Data from the production plant covering approximately 68 months (Jan. 2009 to Sept. 2014) to derive a specification limit for OPP. CONFIDENTIAL	LANXESS Deutschland GmbH Köln, Germany	--	Yes	--	--	LANXESS Deutschland GmbH
A2.8(02) IIA, II 2.8	Feldhues, E.	2006	Additional information on study report No. 2005/0009/00, Determination of main and minor components in Preventol O extra 5-Batch-Analysis. Date: 2006-05-12 CONFIDENTIAL	Bayer Industry Services GmbH & Co KG, BIS-SUA-PUA I, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.1.1(01) IIA, III 3.1 also filed: A3.1.2(01) also filed: A3.1.3(01) also filed A3.10(01)	Erstling, K.	2001 a	Physicochemical properties. Date: 2001-09-13 Amended: 2004-12-02, 2006-03-02, 2006-04-24, 2007-06-26	Bayer AG, Leverkusen, Germany	A 00/0068/01 LEV	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A3.1.3(02) IIA, III 3.1	Erstling, K.	2007	Physicochemical properties of Preventol O Extra	Bayer Industry Services, Leverkusen, Germany	2007/0045/02	Yes	No	Yes	LANXESS Deutschland GmbH
A3.2(01) IIA, III 3.2	Olf, G.	2003	Vapour pressure, Physical-Chemical properties. Date: 2003-02-11 Amended: 2003-02-24 2007-06-29	Bayer AG, Leverkusen, Germany	03/003/01	Yes	No	Yes	LANXESS Deutschland GmbH
A3.2(02) IIA, III 3.2 also filed: A7.3.1(01)	Beiell, U.	2004	Preventol O Extra (<i>o</i> -Phenylphenol) Calculation of Henry's Law Constant and Photodegradation. Date: 2004-09-27	Dr. Knoell Consult GmbH, Mannheim, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.3(01) IIA, III 3.3	Stroech, K.	2006	<i>o</i> -Phenylphenol / Appearance. Date: 2006-04-11	LANXESS Deutschland GmbH, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.4(01) IIA, III 3.4	Erstling, K.	2004	Spectral Data of Preventol O Extra. Date: 2004-07-16 Amended: 2004-12-01	Bayer Industry Services, Leverkusen, Germany	A 02/0162/03 LEV	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A3.5(01) IIA, III 3.5	Erstling, K.	2002	Water solubility. Date: 2002-02-15	Bayer AG, Leverkusen, Germany	A 00/0068/02 LEV	Yes	No	Yes	LANXESS Deutschland GmbH
A3.6(01) – also filed: A3.9(01)	Kausler	1991	Partition coefficient, dissociation constant, pH value. Date: 1991-01-09 Amended: 2005-02-03 2007-06-26	Bayer AG, Leverkusen, Germany	A 89/0062/06 LEV	Yes	No	Yes	LANXESS Deutschland GmbH
A3.6(02) – also filed: A3.9(02)	Erstling, K.	2001 b	Partition coefficient (<i>n</i> -octanol/water) / Dissociation constant. Date: 2001-10-23 Amended: 2001-11-14, 2004-12-03 and 2005-01-14 2007-06-28	Bayer AG, Leverkusen, Germany	A 00/0068/03 LEV	Yes	No	Yes	LANXESS Deutschland GmbH
A3.7(01) IIIA, III.1	Jungheim, R.	2004	Solubility of Preventol O Extra in organic solvents. Date: 2004-07-26	Bayer Industry Services, Leverkusen, Germany	A 02/0162/04 LEV	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A3.7(02) IIIA, III.1	Feldhues, E.	2006 a	Statement Solubility of Preventol O Extra in organic solvents, Temperature dependence. Date: 2006-11-20	Bayer Industry Services, BIS-SUA-PUA I, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.9(03) IIA, III 3.6	Feldhues, E.	2006 b	Statement Partition coefficient n-octanol/water of Preventol O Extra, Temperature and pH dependence. Date: 2006-11-20	Bayer Industry Services, BIS-SUA-PUA I, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.11(01) IIA, III 3.8	Heinz, U.	2004	Determination of safety relevant data of Preventol O Extra. Date: 2004-07-12 Amended: 2005-01-14	Bayer Industry Services, Leverkusen, Germany	04/00223	Yes	No	Yes	LANXESS Deutschland GmbH
A3.13(01) IIA, III 3.10	Olf, G.	2004	Surface tension of Preventol O Extra. Date: 2004-09-16	Bayer Technology Services, Leverkusen, Germany	04006/03	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A3.15(01) IIA, III 3.11	Stroech, K.	2004 a	<i>o</i> -Phenylphenol / Explosive properties. Date: 2004-07-29	Bayer Chemicals AG, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.16(01) IIA, III 3.12	Stroech, K.	2004 b	<i>o</i> -Phenylphenol / Oxidising properties. Date: 2004-07-29	Bayer Chemicals AG, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A3.17(01) IIA, III 3.13 also filed A8.1(02)	Kraus, H.	2006	<i>o</i> -Phenylphenol (OPP) / Reactivity towards container material. Date: 2006-05-30	LANXESS Deutschland GmbH, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A4.1(01) IIA, IV 4.1	Feldhues, E.	2005	Validation of analytical methods for the determination of main and minor components in Preventol O Extra. Date: 2005-02-04 Amended: 2006-04-24 CONFIDENTIAL	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	A 02/0162/08 LEV	Yes	No	Yes	LANXESS Deutschland GmbH
A4.1(02) IIA, IV 4.1	Dick, W.	1990 a	Water – Volumetric method. Date: 1990-12-18 CONFIDENTIAL	ZF-DZA/Analytik LEV/OAL, Leverkusen, Germany	2011- 0131301-90	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A4.1(03) IIA, IV 4.1	Dick, W.	1990 b	Karl Fischer titrant (KF-T) – Equivalent water concentration-Volumetric method. Date: 1990-12-18 CONFIDENTIAL	ZF-DZA/Analytik LEV/OAL, Leverkusen, Germany	2011-0131401-90	No	No	Yes	LANXESS Deutschland GmbH
A4.2(01) IIA, IV 4.2	Brumhard, B.	2004	Method 00829 for the determination of residues of Preventol O Extra in soil by HPLC-MS/MS. Date: 2004-01-05	Bayer Crop Science AG, Monheim am Rhein, Germany	Bayer Method No.: 00829; Report No.: MR- 107/03	Yes	No	Yes	LANXESS Deutschland GmbH
A4.2(02) IIA, IV 4.2	Feldhues, E.	2005 b	Validation of an analytical method for the determination of Preventol O Extra in air samples. Date: 2005-02-21 Amended: 2007-06-20 2010-01-22	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	A 02/0162/05 LEV	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A4.2(03) IIA, IV 4.2	Königer, A.	2010	Validation of a GC method for the determination of Preventol O Extra in air. Date: 2010-01-22	CURRENTA GmbH &Co. OHG Services Analytik Leverkusen Germany	2009/0013/01	Yes	--	--	LANXESS Deutschland GmbH
A4.2(04) IIA, IV 4.2	Brumhard, B.	2003	Enforcement method 00828 (MR-100/03) for the determination of Preventol O Extra in surface and drinking water by HPLC-MS/MS. Date: 2003-12-17 Amended: 2005-03-14 2007-07-02	Bayer Crop Science AG, Monheim am Rhein, Germany	Report No.: MR-100/03; Method No.: 00828	Yes	No	Yes	LANXESS Deutschland GmbH
A4.3(01) IIA, IV 4.3	Stroeck, K.	2014	Residue determination of 2-phenylphenol in meat via GC/MS/MS measurement. 2014-06-16, amended 2014-10-23	Lanxess Deutschland GmbH, Köln, Germany		No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A4.3(02) IIA, IV 4.3	Semrau, J	2011	Determination of residues of orthophenylphenol (OPP) and phenylhydroquinone (PHQ) and their conjugates after a single postharvest application of AGF/1-04 in oranges, Southern Europe 2011.	Eurofins Agrosience Services GmbH, Stade, Germany, (), 2011-12-12	Report No.: S11-01940	Yes	No	Yes	Agrupost, Valencia, Spain
A5 IIA 5.4	Russell, A.D., Hugo, W.B. and Ayliffe, G.A.J.	1990	Principles and practice of disinfection, preservation and sterilisation.	--	--	--	Yes	No	Second Edition, Blackwell Scientific Public
A5.3.1(01) IIA, V 5.3	Bombliès, L. and Wedde, A.	2000	Preventol O Extra (active substance. Determination of the "Minimal Inhibitory Concentration (MIC) against various test microorganisms. Date: 2000-09-16	Labor L+S, Bad-Bocklet-Großenbrach, Germany	01020940	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A5.3.1(02) IIA, V 5.3	Exner, O.	1997	Preventol O Extra: Determination of bactericidal effectiveness in a qualitative suspension disinfection test in accordance with German Society of Hygiene and Microbiology (DGHM) guidelines. Date: 1997-11-28	Bayer AG, Material Protection Business Unit, Krefeld, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A6.1.1(01) IIA, VI 6.1.1	██████████ and ██████████ ██████████	1994	Dowicide™ 1 Antimicrobial: Acute Oral Toxicity Study in Fischer 344 Rats. Date: 1994-07-29	Dow Chemical Company	K-001024-057A	Yes	No	Yes	Dow Chemical Company
A6.1.2(01) IIA, VI 6.1.2	██████████	1991	Preventol O Extra (Schuppen) – Acute Dermal Toxicity Study in Male and Female Wistar Rats. Date: 1991-01-09	Bayer AG	19831	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.1.3(01) IIA, VI 6.1.3	██████████ ██████████ ██████████ and ██████████	1992	<i>ortho</i> -Phenylphenol: Acute Aerosol Inhalation Toxicity Study in Fischer 344 Rats. Date: 1992-02-24	Dow Chemical Company	K-001024-049	Yes	No	Yes	Dow Chemical Company
A6.1.3(01)	Marple et al.	1978	A Dust Generator for Laboratory Use.	--	<i>Am. Ind. Hyg. Assoc. J.</i> 39 : 26-32	--	--	--	--
A6.1.4(01) IIA, VI 6.1.4	██████████	1994 a	Dowicide™ 1 Antimicrobial: Primary Dermal Irritation Study in New Zealand White Rabbits. Date: 1994-07-29	Dow Chemical Company	K-001024-057B	Yes	No	Yes	Dow Chemical Company
A6.1.4(02) IIA, VI 6.1.4	██████████ ██████████	1981 b	Report on the test of Preventol O Extra for irritation of the mucous membrane. Date: 1981-11-04	Fraunhofer-Institut für Toxikologie und Aerosolforschung, Schmallenberg, Germany	T2004666	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.1.5(01) IIA; VI 6.1.5	[REDACTED]	1994 b	Dowicide™ 1 Antimicrobial: Dermal Sensitization Potential in the Hartley Albino Guinea Pig. Date: 1994-07-29	Dow Chemical Company	K-001024-057E	Yes	No	Yes	Dow Chemical Company
A6.1.5(02) IIA; VI 6.1.5	Andersen, K.E. and Hamann, K.	1984	The Sensitizing Potential of Metalworking Fluid Biocides (Phenolic and Thiazole Compounds) in the Guinea-Pig Maximization Test in Relation to Patch-Test Reactivity in Eczema Patients.	Department of Dermatology, Gentofte Hospital, Hellerup, Denmark	<i>Fd. Chem Toxic.</i> 22 (8), pp. 655-660	No	Yes	No	--
A6.2(01) IIA, VI 6.2	Bartels, M.J., Brzak, K.A., McNett, D. and Shabrang, S.N.	1997	<i>ortho</i> -Phenylphenol (OPP): Limited Metabolism Study in Human. Date: 1997-02-03	Dow Chemical Company	HET K-001024-059	Yes	No	Yes	Dow Chemical Company
A6.2(02) IIA, VI 6.2	[REDACTED] [REDACTED] [REDACTED] and [REDACTED] [REDACTED]	1997	<i>ortho</i> -Phenylphenol (OPP): Metabolism of ¹⁴ C-Labelled OPP in B ₆ B ₃ F ₁ Mice and Fischer 344 Rats. Date: 1997-02-06	Dow Chemical Company	HET K-001024-060	Yes	No	Yes	Dow Chemical Company

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.2(03) IIA, VI 6.2	Selim, S.	1996	A Single Open Dose Label Study to Investigate the Absorption and Excretion of ¹⁴ C/ ¹³ C-Labeled <i>ortho</i> -Phenylphenol Formulation after Dermal Application to Healthy Volunteers. Date: 1996-09-19	Bayer AG	P0995002	Yes (GCP)	No	Yes	LANXESS Deutschland GmbH
A6.3.1(01) IIA, VI 6.3.1 also filed: A6.5(02)	██████████ ██████████ ██████████ ██████████ ██████████ and ██████████	1990	<i>ortho</i> -Phenylphenol: Palatability/Probe, Four-Week and One-Year Oral Toxicity Studies in Beagle Dogs. Date: 1990-09-24	Dow Chemical Company	K-001024-039	Yes	No	Yes	Dow Chemical Company
A6.3.2(01) IIA, VI 6.3.2	██████████ and ██████████ ██████████	1993	<i>ortho</i> -Phenylphenol: 21-Day Repeated Dermal Dose Study of Systemic Toxicity in Fischer 344 Rats. Date: 1993-03-03	Dow Chemical Company	K-001024-056	Yes	No	Yes	Dow Chemical Company

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.4.1(01) IIA, VI 6.4	██████████ ██████████ ██████████ and ██████████	1996 a	Technical Grade <i>ortho</i> -Phenylphenol: A Special Subchronic Dietary Study to Examine the Mechanism of Urinary Bladder Carcinogenesis in the Male Rat. Date: 1996-11-11	Bayer AG	92-972-MS	No	No	Yes	LANXESS Deutschland GmbH
A6.5(01) IIA, VI 6.5 also filed: A6.7(01)	██████████ and ██████████ ██████████	1996	Technical Grade <i>ortho</i> -Phenylphenol: A Combined Chronic Toxicity / Oncogenicity Testing Study in the Rat. Date: 1996-02-23, Amended: 1999	Bayer AG	92-272-SC	Yes	No	Yes	LANXESS Deutschland GmbH
A6.6.1(01) IIA, VI 6.6.1	San, R.H.C. and Springfield, K.A.	1989	Salmonella/Mammalian-Microsome Plate Incorporation Mutagenicity Assay (Ames Test). Date: 1989-12-22	Bayer AG	C141.501017	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.6.1(01)	Ames et al.	1975	Methods for detecting carcinogens and mutagens with salmonella-mammalian-microsome mutagenicity test	--	Mutation Res. 31 , 347-363	--	--	--	--
A6.6.1(01)	Maron & Ames	1983	Revised methods for the salmonella mutagenicity test	--	Mutation Res. 113 , 173-215	--	--	--	--
A6.6.2(01) IIA, VI 6.6.2	Tayama, S., Kamiya, N. and Nakagawa, Y.	1989	Genotoxic effects of o-Phenylphenol metabolites in CHO-K1 cells.	Dept. of Toxicology, Tokyo Metropolitan Research Laboratory of Public Health, Tokyo, Japan	<i>Mutat. Res.</i> 223 , pp. 23-33	No	Yes	No	--
A6.6.3(01) IIA, VI 6.6.3	Brendler, S.	1992	Preventol O Extra – Mutagenicity Study for the Detection of Induced Forward Mutations in the CHO-HGPRT Assay In Vitro. Date: 1992-04-09	Bayer AG	21278	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.6.5(01) IIA, VI 6.6.5	██████████ ██████████	2000	Preventol O Extra – Comet Assay In Vivo in Mouse Liver and Kidney. Date: 2000-08-08	Bayer AG	PH 30130	Yes	No	Yes	LANXESS Deutschland GmbH
A6.8.1(02) IIA, VI 6.8.1	██████████ ██████████ ██████ and ██████████	1991	<i>ortho</i> -Phenylphenol (OPP): Gavage Teratology Study in New Zealand White Rabbits. Date: 1991-04-23	Dow Chemical Company	K-001024-045	Yes	No	Yes	Dow Chemical Company
A6.8.1(01) IIA, VI 6.8.1	Kaneda, M., Teramoto, S., Shingu, A. and Yasuhiko, S.	1978	Teratogenicity and Dominant-Lethal Studies with <i>o</i> -Phenylphenol.	Toxicology Division, Institute of Environmental Toxicology, Kodaira, Tokyo, Japan	<i>J. Pesticide Sci.</i> 3 , pp. 365-370	No	Yes	No	--
A6.8.2(01) IIA, VI 6.8.2	██████████ ██████ and ██████████	1995	A Two-Generation Dietary Reproduction Study in Sprague-Dawley Rats Using Technical Grade <i>ortho</i> -Phenylphenol. Date: 1995-09-28	Bayer AG	93-672-VX	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.8.2(02) IIA, VI 6.8.2	██████████ ██████	1990	Two-Generation Dietary Reproduction Study in Rats Using <i>ortho</i> -Phenylphenol. Date: 1990-09-17 (revised report, original report date: 1989-01-13)		85-671-02	Yes	No	Yes	LANXESS Deutschland GmbH
A6.10(01)	Fukushima, S., Kurata, Y., Shibata, M., Ikawa, E. and Ito, N.	1983	Promoting Effect of Sodium <i>o</i> -Phenylphenate and <i>o</i> -Phenylphenol on Two-Stage Urinary Bladder Carcinogenesis.	First Department of Pathology, Nagoya City University Medical School, Nagoya, Japan	<i>Gann.</i> , 74 , pp. 625-632	No	Yes	No	--
A6.10(02)	Fujii, T., Nakamura, K. and Hiraga, K.	1987	Effects of pH on the Carcinogenicity of <i>o</i> -Phenylphenol and Sodium <i>o</i> -Phenylphenate in the Rat Urinary Bladder.,	Dept. of Toxicology, Tokyo Metropolitan Research Laboratory of Public Health, Tokyo, Japan	<i>Fd. Chem. Toxic.</i> 25 (5), pp. 359-362	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.10(03)	██████████ ██████	1994	<i>o</i> -Phenylphenol – Interactions of <i>o</i> -Phenylphenol (OPP) and its metabolites with microsomal prostaglandin-H-synthase: possible implications for OPP-induced tumour formation in the rat urinary bladder. Date: 1994-01-12	Bayer AG	22788	No	No	Yes	LANXESS Deutschland GmbH
A6.12.1(01) IIA, VI 6.12.1	Heyne, R. and Attig, G.	2004	Occupational Medical Experiences with <i>o</i> -Phenylphenol. Date: 2004-12-06	Bayer Industry Services, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
A6.12.6(01) IIA, VI 6.9.6	Adams, R.M.	1981	Allergic contact dermatitis due to <i>o</i> -Phenylphenol.	Palo Alto Medical Clinic, Palo Alto, CA, USA	<i>Contact Dermatitis</i> 7 , p. 332	No	Yes	No	--
A6.12.6(02) IIA, VI 6.9.6	van Hecke, E.	1986	Contact sensitivity to <i>o</i> -Phenylphenol in a coolant.	Dept. of Dermatology, University Hospital, Gent, Belgium	<i>Contact Dermatitis</i> 15 (1), p. 46	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.12.6(03) IIA, VI 6.9.6	Schnuch, A., Geier, J., Uter, W. and Frosch, P.J.	1998	Patch testing with preservatives, antimicrobials and industrial biocides. Results from a multicentre study.	Information Network of Dermatological Clinics in Germany (IVDK)	<i>Br. J. Dermatology</i> 138 , pp. 467-476	No	Yes	No	--
A6.12.6(04) IIA, VI 6.9.6	Geier, J., Kleinhans, D. and Peters, K.-P.	1996	Kontaktallergien durch industriell verwendete Biozide – Ergebnisse des Informationsverbunds Dermatologischer Kliniken (IVDK) und der Deutschen Kontaktallergiegruppe. (Contact Allergy Due to Industrial Biocides– Results of the IVDK and the German Dermatitis Research Group.)	Information Network of Departments of Dermatology in Germany (IVDK)	<i>Dermatosen / Occup. Environ.</i> 44 , pp. 154-159	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A6.12.6(05) IIA, VI 6.12.6	Brasch, J., Henseler, T. and Frosch, P.	1993	Patch Test Reactions to a Preliminary Preservative Series – A retrospective study based on data collected by the “Information Network of Dermatological Clinics” (IVDK) in Germany.	Information Network of Departments of Dermatology in Germany (IVDK)	<i>Dermatosen</i> 41 (2), pp. 71-76	No	Yes	No	--
A6.15(01) IIIA, VI 4	Stroech, K.D.	2013	Residue determination of 4-chloro-3-methylphenol and 2-phenylphenol in edible tissues of 15 broiler chicken that were reared on an area disinfected with the LCB trial product "CMK/OPP 32". date: 2013-01-22	LANXESS Deutschland GmbH,	--	No	No	Yes	LANXESS Deutschland GmbH,
A7.1.1.1.1(01) IIA, VII.7.6.2.1	Reusche, W.	1991	Hydrolysis study of 2-phenylphenol according to OECD guideline 111. Date: 1991-01-02, amended: 2004-12-02	Bayer AG, Leverkusen, Germany	G 89/0056/02 LEV	Yes	No	Yes	Bayer Crop Science AG

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.1.1.1.2(01) IIA, VII.7.6.2.2	Heinemann, O.	2005	[Phenyl-UL- ¹⁴ C]-2-phenylphenol: Phototransformation in Water. Date: 2005-03-15.	Bayer CropScience AG, Monheim, Germany	MEF-05/018	Yes	No	Yes	Bayer Crop Science AG
A7.1.1.1.2(02) IIA, VII.7.6.2.2	Wick, L.Y. and Gschwend, P.M.	1998	Source and chemodynamic behaviour of diphenyl sulfone and <i>ortho</i> - and <i>para</i> -hydroxybiphenyl in a small lake receiving discharges from an adjacent superfund site.	Ralph M. Parsons laboratory, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA	<i>Environ. Sci. Technol.</i> 32 , pp. 1319-1328.	No	Yes	No	--
A7.1.1.1.2(02)	Haag, W. and Hoigné J.	1986	Singlet oxygen in surface waters .3. Photochemical formation and steady-state concentrations in various types of waters	--	<i>Environ. Sci. Technol.</i> , 20 , pp. 341-348	--	Yes	No	--
A7.1.1.1.2(02)	Leifer, A.	1988	The Kinetics of Environmental Aquatic Photochemistry.	--	American Chemical Society, Washington, DC, USA	--	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.1.1.2.1(01) IIA, VII.7.6.1.1	Gonsior, S.J. and Tryska, T.J.	1997	Evaluation of the Ready Biodegradability of o-Phenylphenol. Date: 1997-08-01	Environmental Chemistry Research Laboratory, The Dow Chemical Company, Midland, Michigan	971080	Yes	No	Yes	The DOW Chemical Company
A7.1.1.2.1(02) IIA, VII.7.6.1.1	Kanne, R.	1989	Preventol O Extra. Biodegradation. Date: 1989-07-24	Bayer AG, Institut für Umweltanalyse und Bewertungen, Leverkusen, Germany	51A/88/I	Yes	No	Yes	Bayer AG
A7.1.1.2.1(03)	Painter H.A. and King E.F.	1985	Ring test programme 1983-84. Assessment of biodegradability of chemicals in water by manometric respirometry	Ring test, monitored by the Water Research Centre, Elder Way, UK - Stevenage Herts	EUR 9962 EN	No	No	No	Commission of the EC: Environment and Quality of life

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.1.1.2.1(04)	Kanne, R.	1989b	Preventol O Extra. Biodegradation in Rhine River Water. Date: 1989-07-24	Bayer AG, Institut für Umweltanalyse und Bewertungen, Leverkusen, Germany	Report-No. 51A/88/II	Yes	No	Yes	Bayer AG
A7.1.1.2.2(01) IIA, VII.7.6.1.2	Wellens, H.	1990	Zur biologischen Abbaubarkeit mono- und disubstituierter Benzolderivate.	Abwasserbiologische Laboratorien der HOECHST AG, Frankfurt, Germany	Z. Wasser-Abwasser-Forsch. 23, 85-98	No	Yes	No	--
A7.1.2.1.1(01) IIIA, XII.2.1	Körner W., Bolz U., Süßmuth W., Hiller G., Schuller W., Hanf V. & Hagenmaier H.	2000	Input/Output Balance of Estrogenic Active compounds in a Major Municipal Sewage Plant in Germany.	Institute of Organic Chemistry, University of Tübingen, Germany	<i>Chemosphere</i> 40 , 1131-1142.	No	Yes	No	--
A7.1.2.1.1(01) IIIA, XII.2.1	Bolz, U., Körner, W., Hagenmeier, H.	2000	Development and validation of a GC/MS method for determination of phenolic xenoestrogens in aquatic samples.	Institute of Organic Chemistry, University of Tübingen, Germany	<i>Chemosphere</i> 40 , 929-935.	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.1.2.1.1(02) IIIA, XII.2.1	Ternes, T., Stumpf, M., Schuppert, B., Haberer, K.	1998	Simultaneous Determination of Antiseptics and Acidic Drugs in Sewage and River Water.	ESWE-Institute for Water Research and Water Technology, Wiesbaden, Germany	Vom Wasser, 90, 295-309.	No	Yes	No	--
A7.1.2.1.1(03) IIIA, XII.2.1	Lee, H.-B., Peart, T.E., Svoboda, M.L.	2005	Determination of endocrine-disrupting phenols, acidic pharmaceuticals, and personal-care products in sewage by solid-phase extraction and gas chromatography-mass spectrometry.	Aquatic Ecosystem Protection Research Branch, National Water Research Institute, Environment Canada, Ontario, Canada.	Journal of Chromatography A, 1094, 122-129.	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.1.2.2.2(01) IIIA, XII 2.1	Bruns, E.	2005	Preventol O Extra (<i>ortho</i> -Phenylphenol). Summary of screening experiments concerning the behaviour of <i>ortho</i> -Phenylphenol (OPP) in a "water-sediment system". Date: 2005-03-29	Bayer Industry Services GmbH & Co. OHG, Leverkusen, Germany	--	Yes	No	Yes	Bayer Crop Science AG
A7.1.3(01) IIA, VII 7.7	Erstling, K.	2001	Preventol O Extra in Schuppen – Adsorption/Desorption, during the period June to September 2001. Date: 2001-09-17	Bayer AG, Zentrale Analytik, Leverkusen, Germany	A 0/0068/04 LEV	Yes	No	Yes	LANXESS Deutschland GmbH
A7.2.1(01) IIIA, VII 4, XII 1.1	Fliege, R.	2005	[phenyl-UL- ¹⁴ C]- <i>ortho</i> -Phenylphenol: Aerobic soil metabolism in one European soil. Date: 2005-03-23	Bayer CropScience AG, Development, Metabolism / Environmental fate, Germany	MEF-05/072	Yes	No	Yes	Bayer Crop Science AG

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.2.2.1(02)	Nitsche, M.	2011	Biodegradation of Preventol® O Extra (2-phenylphenol) in soil under aerobic conditions	Lanxess Deutschland GmbH, Leverkusen, Germany	-	No	No	Yes	Lanxess Deutschland GmbH
A7.2.2.1 (02)	Loehr, Raymond C. and Matthews, John E.	1992	Loss of organic chemicals in soil: Pure compound treatability studies	<i>Journal of Soil Contamination</i> 1(4) 339-360	--	--	--	--	--
A7.2.3.1(01) IIIA, XII.1.2	Oddy, A. and Jacob, O.	2005	[¹⁴ C]-2-Phenylphenol: Adsorption to and Desorption from four soils. Date: 2005-03-16	Battelle AgriFood Ltd., Essex, UK	CX/04/019	Yes	No	Yes	LANXESS Deutschland GmbH
A7.3.2 IIIA 12.3	Wasser, C.	2014	Residues of the Combustion of OPP20, Residues in fumes and gases.	Anadiag Laboratories, France 67500 Haguenau	R B4256	No	No	Yes	LANXESS Deutschland GmbH
A7.4.1.1(01) IIA, VII.7.1	██████████	1990	Acute Fish Toxicity of Preventol O Extra. Date: 1990-04-10	Bayer AG, Institut für Umweltanalysen und Bewertungen, Leverkusen, Germany	51 A/88 F	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.4.1.1(02)	[REDACTED]	1991	<i>o</i> -Phenylphenol Toxicity to Fish <i>Chinook salmon</i> (<i>Oncorhynchus tshawytscha</i>). Date: 1991-10-22	British Columbia Research Corp., Vancouver, Canada	2-11-200-222-91001	No	No	Yes	LANXESS Deutschland GmbH
A7.4.1.2(01) IIA, VII.7.2	[REDACTED] [REDACTED] [REDACTED] and [REDACTED] [REDACTED]	1985	Evaluation of the toxicity of Dowicide 1 Antimicrobial, Technical <i>o</i> -Phenylphenol to representative aquatic organisms. Date: 1985-12-12	Mammalian and Environmental Toxicology, Health & Environmental Sciences, Midland, Michigan, USA	ES-811	No	No	Yes	Dow Chemical Company
A7.4.1.2(02)	Kühn, R., Pattard, M., Pernak, K.- D. Winter,	1988	Harmful effects of chemicals in the <i>Daphnia</i> reproduction test as a basis for assessing their environmental hazard in aquatic systems. March 1988	Institute for Water, Land and Air Hygiene of the Federal German Health Office	10603052	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.4.1.3(01) IIA, VII.7.3	Hicks, S.	2002	<i>ortho</i> -Phenylphenol: Growth Inhibition Test with the Green Alga, <i>Selenastrum capricornutum</i> . Date: 2002-03-12	ABC Laboratories, Inc., Missouri, USA	ABC Study No. 46980, Dow Study No. 010167	Yes	No	Yes	Dow Chemical Company
A7.4.1.3(02)	Caspers, N.	1989	Cellular proliferation inhibitory test: <i>Scenedesmus subspicatus</i> CHODAT (green alga). Date: 1989-07-04	Bayer AG	No. 51 A/88	No	No	Yes	LANXESS Deutschland GmbH
A7.4.1.4(01) IIA, VII.7.4	Mueller, G.	1990	Preventol O Extra, 2-phenylphenol, Toxicity to Bacteria. Date: 1990-08-08	Bayer AG, Institute of Environmental Analysis, Leverkusen, Germany	51 A/88B	Yes	No	Yes	LANXESS Deutschland GmbH
A7.4.1.4(01) IIA, VII.7.4	Weyers, A.	2006	Preventol O Extra, Toxicity to Bacteria. Re-Evaluation based on Study Report No. 51 A/88 B, corresponding raw data and additional information provided by the sponsor. Date: 2006-09-05	Bayer Industry Services, Leverkusen, Germany	--	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.4.1.4(02)	Klecka, G.M., Landi, L.P. and Rodner, K.M.	1985	Evaluation of the OECD Activated Sludge, Respiration Inhibition Test	--	<i>Chemosphere</i> 14 , pp. 1239-1251	No	Yes	No	--
A7.4.2(01) IIA, VII.7.5	Fàbregas, E.	2007	<i>o</i> -Phenylphenol - Calculation of the Bioconcentration Factor (BCF). Date: 2007-06-05	Dr. Knoell Consult GmbH, Leverkusen, Germany	Report-No. KC-BCF-08/07	No	No	Yes	LANXESS Deutschland GmbH
A7.4.3.2(01) IIIA, XIII 2.2	Caunter J.E. and [REDACTED]	2002	Preventol O Extra: Determination of Effects on the Reproduction of Fathead minnow (<i>Pimephales promelas</i>). Date: 2002-03-25	Brixham Environmental Laboratory, AstraZeneca UK Limited, Brixham, UK	BL7213/B	Yes	No	Yes	LANXESS Deutschland GmbH
A7.4.3.3.1(01) IIIA, XIII.2.3	Caspers, N.	1999	Investigation of the Ecological Properties of Preventol O Extra, Test on Bioaccumulation. Date: 1999-05-27	Bayer AG, Leverkusen, Germany	793 A/98	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.4.3.4(01) IIIA, XIII 2.4	Bruns, E.	2001	Preventol O Extra, <i>Daphnia magna</i> Reproduction Test. Date: 2001-12-13	Bayer AG, WD-UWS, Institute of Environmental Analysis and Evaluation, Leverkusen	1092 A/01 DL	Yes	No	Yes	LANXESS Deutschland GmbH
7.4.3.4/02	Caspers, N.	1989	Life cycle test with water fleas - <i>Daphnia magna</i> - EC ₅₀ immobilisation and EC ₅₀ reproduction. Date: 1989-10-13	Bayer AG	No. 51 A/88	No	No	Yes	LANXESS Deutschland GmbH
A7.4.3.5.1(01) IIIA, XIII 2.4	Egeler, P. and Gilberg, D.	2005	Preventol O Extra: A study on the toxicity to the sediment dweller <i>Chironomus riparius</i> . Date: 2005-02-28	ETC Oekotoxikologie GmbH, Germany	AI1ME	Yes	No	Yes	LANXESS Deutschland GmbH
A7.5.1.1/01	Reis, K-H.	2007	Effects of 2-Phenylphenol (Preventol O Extra) on the Activity of the Soil Microflora in the Laboratory. Date: 2007-06-21	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany	35591080	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.5.1.1(02)	Schulz, L.	2012	Effects on the activity of soil microflora (Nitrogen transformation test) Date: 2012-02-10	BioChem agrar, Labor für biologische und chemische Analytik GmbH 04827 Gerichshain, Germany	Project-No. 12 10 48 003 N	No	No	Yes	LANXESS Deutschland GmbH
A7.5.1.2(01) IIIA, XIII 3.2	Moser, Th. and Scheffczyk, A.	2004	Preventol O Extra: Acute toxicity to the earthworm <i>Eisenia fetida</i> in an artificial soil test. Date: 2004-12-08	ETC Oekotoxikologie GmbH, Flörsheim, Germany	AI1RA	Yes	No	Yes	LANXESS Deutschland GmbH
A7.5.1.3	Bützler, R., Meinerling, M.	2008	Effects of 2-Phenylphenol (Preventol O Extra) on Terrestrial (Non-Target) Plants: Seedling Emergence and Seedling Growth Test. Date: 2008-10-17	IBACON GmbH, Rossdorf, Germany,	Report No. 35594084	Yes	No	Yes	LANXESS Deutschland GmbH
A7.5.3.1.1(01) IIIA, XIII 1.1	██████████	1986 a	<i>ortho</i> -Phenylphenol Technical: An Acute Oral Toxicity Study with the Mallard. Date: 1986-06-06	Wildlife International Ltd., St. Michaels, Maryland, USA	ES-874 (103-248)	Yes	No	Yes	Dow Chemical Company

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
A7.5.3.1.2(01) IIIA, XIII 1.2	██████████	1986 b	<i>ortho</i> -Phenylphenol Technical: A Dietary LC ₅₀ Study with the Bobwhite. Date: 1986-06-06	Wildlife International Ltd., St. Michaels, Maryland, USA	ES-873 (103-246)	Yes	No	Yes	Dow Chemical Company
A7.5.3.1.2(02) IIIA, XIII 1.2	██████████	1986 c	<i>ortho</i> -Phenylphenol Technical: A Dietary LC ₅₀ Study with the Mallard. Date: 1986-06-06	Wildlife International Ltd., St. Michaels, Maryland, USA	ES-875 (103-247)	Yes	No	Yes	Dow Chemical Company
A7.5.5.1(01) IIIA, 13.3	Fàbregas, E.	2007	<i>o</i> -Phenylphenol - Calculation of the Bioconcentration Factor in Earthworms (BCFearthworm). Date: 2007-06-05	Dr. Knoell Consult GmbH, Leverkusen, Germany	Report-No. KC-BCF-09/07	No	No	Yes	LANXESS Deutschland GmbH
A8.1(01) IIA, VIII 8.1 also filed: A8.2(01) also filed: A8.3(01) also filed: A8.4(01) also filed: A8.5(01)	Anonymous	2004	Safety Data Sheet Preventol O Extra. Date: 2004-03-10	LANXESS Deutschland GmbH, Leverkusen, Germany	011472/23	No	No	--	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B2.2(01) IIB, I 2.2	Bayer MaterialScience AG (Ed.)	2011a	Safety Data Sheet CAUSTIC SODA SOLUTION (50%). Date: 2011-11-04 CONFIDENTIAL	Bayer MaterialScience AG, Leverkusen, Germany	Version 3.1	No	No	No	Bayer MaterialScience AG
B2.2(02) IIB, I 2.2	Bayer MaterialScience AG (Ed.)	2011b	Technical information Caustic Soda Solution 50%. Date: 2011-03-11 CONFIDENTIAL	Bayer MaterialScience AG, Leverkusen, Germany	Specification No: 05452627-01- 11	No	No	No	Bayer MaterialScience AG
B2.3(01) IIB, I 2.3 also filed B3.1(01)	Stroech, K.	2006	<i>o</i> -Phenylphenol / Appearance. Date: 2006-04-11	LANXESS Deutschland GmbH, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
B3.1(01)	Stroech, K.D.	2012a	Preventol® ON Extra Preservative Solution / Appearance Properties. Date: 2012-01-31	LANXESS Deutschland GmbH, Leverkusen, Germany	212951ni-sxx	No	No	Yes	LANXESS Deutschland GmbH
B3.2(01) IIB, III 3.2	Stroech, K.	2004a	<i>o</i> -Phenylphenol / Explosive properties. Date: 2004-07-29	Bayer Chemicals AG, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
B3.2(01)	Stroech, K.D.	2012b	Preventol® ON Extra Preservative Solution / Explosive Properties. Date: 2012-01-31	LANXESS Deutschland GmbH, Leverkusen, Germany	212954ni-sxx	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B3.3(01) IIB, III 3.3	Stroech, K.	2004b	o-Phenylphenol / Oxidising properties. Date: 2004-07-29	Bayer Chemicals AG, Leverkusen, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
B3.3(01)	Stroech, K.D.	2012c	Preventol® ON Extra Preservative Solution / Oxidising Properties. Date: 2012-01-31	LANXESS Deutschland GmbH, Leverkusen, Germany	212955ni-sx	No	No	Yes	LANXESS Deutschland GmbH
B3.4(01) IIB, III 3.4	Heinz, U.	2004	Determination of safety relevant data of Preventol O Extra. Date: 2004-07-12 Amended: 2005-01-14	Bayer Industry Services, Leverkusen, Germany	04/00223	Yes	No	Yes	LANXESS Deutschland GmbH
B3.4(01)	Stroech, K.D.	2012d	Preventol® ON Extra Preservative Solution / Flammability or Spontaneous Ignition Properties. Date: 2012-01-31	LANXESS Deutschland GmbH, Leverkusen, Germany	212952ni-sxx	No	No	Yes	LANXESS Deutschland GmbH
B3 IIB, III 3.4	Heinz, U.	2004	Determination of safety relevant data of Preventol O Extra. Date: 2004-07-12 Amended: 2005-01-14	Bayer Industry Services, Leverkusen, Germany	04/00223	Yes	No	Yes	LANXESS Deutschland GmbH
B3.4(01) B3.6(01) B3.10(01)	Krasemann, R.	2006	Safety-related Data / Product description of Preventol ON Extra Preservative Solution. Date: 2006-10-27	Bayer Industry Services GmbH & Co. OHG, Leverkusen, Germany	2006/01781	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B3.5(01) IIB, III 3.4	Erstling, K.	2007	Determination of acidity/alkalinity	Bayer Industry Services, Leverkusen, Germany	2007/0045/02	Yes	No	Yes	LANXESS Deutschland GmbH
B3.5(01)	Nitsche, M.	2012a	Preventol® ON Extra Preservative Solution Determination of pH - Value. Date: 2012-04-16	LANXESS Deutschland GmbH, Leverkusen, Germany	None	No	No	Yes	LANXESS Deutschland GmbH
B3.5(02)	Nitsche, M.	2012b	Determination of the Alkalinity of Preventol® ON Extra Preservative Solution. Date: 2012-04-13	LANXESS Deutschland GmbH, Leverkusen, Germany	None	No	No	Yes	LANXESS Deutschland GmbH
B3.6(01) IIB, III 3.6	Erstling, K.	2001	Physicochemical properties. Date: 2001-09-13 Amended: 2004-12-02, 2006-03-02 and 2006-04-24	Bayer AG, Leverkusen, Germany	A 00/0068/01 LEV	Yes	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B3.7(01) IIB, III 3.7	European Commission (Ed.)	2006	Content of the product dossier accompanying the active substance for Annex I inclusion. Date: 2006-09-14	European Commission, Directorate-General-JRC, Institute for Health and Consumer Protection, Unit: Toxicology and Chemical Substances, European Chemicals Bureau	--	No	Yes	No	European Commission, European Chemicals Bureau
B3.7(01)	Nitsche, M.	2012c	Storage Stability of the Formulation Preventol ON Extra Preservative Solution at Accelerated Temperature of 40 °C. Date: 2012-05-08	LANXESS Deutschland GmbH, Leverkusen, Germany	None	No	No	Yes	LANXESS Deutschland GmbH
B3.7(02)	Nitsche, M.	2012d	Storage Stability at 0 °C of the Formulation Preventol ON Extra Preservative Solution. Date: 2012-02-17	LANXESS Deutschland GmbH, Leverkusen, Germany	None	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B3.7(03)	Nitsche, M.	2012e	Storage Stability of the Formulation Preventol ON Extra Preservative Solution at Ambient Temperature. (6 months interim report) Date: 2012-08-03	LANXESS Deutschland GmbH, Leverkusen, Germany	None	No	No	Yes	LANXESS Deutschland GmbH
B3.8(01) IIB, III 3.8	Erstling, K.	2007	Physicochemical properties of Preventol O Extra	Bayer Industry Services, Leverkusen, Germany	2007/0045/02	Yes	No	Yes	LANXESS Deutschland GmbH
B3.10(01) -	Olf, G.	2004	Surface tension of Preventol O Extra. Date: 2004-09-16	Bayer Technology Services, Leverkusen, Germany	04006/03	Yes	No	Yes	LANXESS Deutschland GmbH
B3.10(01)	Keldenich, H.-P. and Kokott, H.	2011	Determination of Surface Tension of Preventol ON Extra preservative solution. Date: 2011-02-28	Bayer Technology Services GmbH, Operation Support & Safety, Process and Plant Safety Laboratory, Leverkusen, Germany	Study-No.: 2011/00296e	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B4.1(01) IIA, IV 4.1	Feldhues, E.	2005a	Validation of analytical methods for the determination of main and minor components in Preventol O Extra. Date: 2005-02-04 Amended: 2006-04-24	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	A 02/0162/08 LEV	Yes	No	Yes	LANXESS Deutschland GmbH
B4.1(01)	Nitsche, M.	2012f	Validation of the Test Method AFAM 2301-0272501-99E for Determination of Content of <i>o</i> -Phenylphenol in Preventol® ON Extra Preservative Solution. Date: 2012-04-16	LANXESS Deutschland GmbH, Leverkusen, Germany	None	No	No	Yes	LANXESS Deutschland GmbH
B5.8 IIB, V5.8	Russell, A.D., Hugo, W.B. and Ayliffe, G.A.J.	1990	Principles and practice of disinfection, preservation and sterilisation.	---	Second Edition, Blackwell Scientific Public., London (pages 201 and 204).	No	Yes	No	--

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B5.10(01) IIB, V 5.10	Groetsch, W. and Nothhelfer, B.	2000	SF Preventol OPP. Determination of the bacteriostatic and fungistatic efficacy according to the DGHM-guideline (I/2.1). Date: 2000-06-27	Labor L+S, Bad- Bocklet- Großenbrach, Germany.	Report No. 01020970	No	No	Yes	LANXESS Deutschland GmbH
B5.10(02) IIB, V 5.10	Wachtler, P.	2002	Preventol Preservatives. Report on Preservation Test. Efficacy of Preventol OF 45 in a CaCO ₃ slurry.	Bayer Chemicals, BCH-MPP-TM-IPC, Building R54, Krefeld, Germany Date: 2002-08-05	--	No	No	Yes	LANXESS Deutschland GmbH
B5.10(03) IIB, V5.10	Herbertz, T.	2012	Efficacy study submitted for the registration of Preventol® ON Extra Preservative Solution. The results support the antimicrobial efficacy for Product- type 07 according to the BPD 98/8/EC. Date: 2012-03-08	Lanxess Deutschland GmbH, Leverkusen, Germany	---	No	No	Yes	LANXESS Deutschland GmbH
B5.10(04)	Herbertz, T.	2014	Efficacy of Preventol O Extra in a cooling lubricant.	Lanxess Deutschland GmbH, Leverkusen, Germany	-	-	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B6.1.1 IIB, VI 6.1.1	██████████	1998a	Single Dose Oral Toxicity in Rats/LD 50 in Rats. Date: 1998-12-01	MB Research Laboratories, Spinnerstown, PA, USA	Project No.: MB 98-7078.01	Yes	No	Yes	LANXESS Deutschland GmbH
B6.1.2 IIB, VI 6.1.2	██████████	1998b	Acute Dermal Toxicity in Rabbits/LD 50 in Rabbits. Date: 1998-11-24	MB Research Laboratories, Spinnerstown, PA, USA	Project No.: MB 98-7078.02	Yes	No	Yes	LANXESS Deutschland GmbH
B6.2(1) IIB, VI 6.2	██████████	1998c	Primary Dermal Irritation in Rabbits. Date: 1998-11-24	MB Research Laboratories, Spinnerstown, PA, USA	Project No.: MB 98-7078.03	Yes	No	Yes	LANXESS Deutschland GmbH
B6.2(2) IIB, VI 6.2	██████████	1998d	Primary Eye Irritation/Corrosion in Rabbits. Date: 1998-12-01	MB Research Laboratories, Spinnerstown, PA, USA	Project No.: MB 98-7078.04	Yes	No	Yes	LANXESS Deutschland GmbH
B6.3 IIB, VI 6.3	██████████	1994	Dowicide™ A Antimicrobial: Dermal Sensitization Potential in the Hartley Albino Guinea Pig. Date: 1994-07-29	The Toxicology Research Laboratory, Health and Environmental Science, Dow Chemical Company, Midland, MI, USA	Report No.: K-001025-014E	Yes	No	Yes	Dow Chemical Company

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B6.4 IIB, VI 6.4	Selim, S.	1996	A Single Open Dose Label Study to Investigate the Absorption and Excretion of ¹⁴ C/ ¹³ C-Labeled <i>ortho</i> -Phenylphenol Formulation after Dermal Application to Healthy Volunteers. Date: 1996-09-19	Bayer AG	P0995002	Yes (GCP)	No	Yes	LANXESS Deutschland GmbH
B6.6 IIB, VI 6.6	Oswald, D. & Zürcher, W.	2003	Raumluftmessung im Werk Gummern vom 07. - 09. Januar 2003 [Indoor Air Measurements in the Gummern Plant between 7 th and 9 th January, 2003]. Date: 2003-01-16	Omya AG, Analytical Laboratory, Oftringen, Switzerland	58005.03	No	No	Yes	LANXESS Deutschland GmbH

(Sub)Section / Annex point	Authors (s)	Year	Title	Testing Company	Report No.	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)	Data Owner
B6.6 IIB, VI 6.6	Wachtler, P. & Kretschmer, F.	2002	Preventol VP SP 80005 - Stationäre and personenbezogene Messungen im Bereich der Bahnverladung (Gleis 3+5) Firma Omya AG in Gummern (Österreich) [Preventol VP SP 80005 - Stationary and Personal Measurements in the Train Charging Area of Omya AG in Gummern (Austria)]. Date: 2002-06-18	Bayer AG Werk Uerdingen, Krefeld, Germany	--	No	No	Yes	LANXESS Deutschland GmbH
B8(01) IIB, VIII 8	LANXESS Deutschland GmbH (Ed.)	No data	Safety Data Sheet PREVENTOL ON EXTRA PRESERVATIVE SOLUTION.	LANXESS Deutschland GmbH, Leverkusen, Germany	81238457C	No	No	No	LANXESS Deutschland GmbH