

# Summary of product characteristics for a biocidal product

**Product name:** Udder star

**Product type(s):** PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

**Authorisation number:** EU-0020541-0000

**R4BP 3 asset reference number:** EU-0020541-0004

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	3
4. Authorised use(s)	3
5. General directions for use	9
5.1. Instructions for use	9
5.2. Risk mitigation measures	9
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	10
5.4. Instructions for safe disposal of the product and its packaging	10
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	10
6. Other information	10

## Administrative information

### 1.1. Trade names of the product

Udder Star
Udder Star Max
Udder S Max
Udderdine Star
Uddine Star
Udderdine ST
Uddine ST

### 1.2. Authorisation holder

**Name and address of the authorisation holder**

Name	Boumatic
Address	31, Rue Jules Melotte 4350 Remicourt Belgium

**Authorisation number**

EU-0020541-0000 1-3
---------------------

**R4BP 3 asset reference number**

EU-0020541-0004
-----------------

**Date of the authorisation**

18/11/2019
------------

**Expiry date of the authorisation**

31/10/2029
------------

### 1.3. Manufacturer(s) of the biocidal products

<b>Name of the manufacturer</b>	Christeyns n.v.
<b>Address of the manufacturer</b>	Afrikalaan 182 9000 Ghent Belgium
<b>Location of manufacturing sites</b>	Afrikalaan 182 9000 Ghent Belgium

#### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	1319 - Iodine
<b>Name of the manufacturer</b>	Christeyns n.v.
<b>Address of the manufacturer</b>	Afrikalaan 182 9000 Ghent Belgium
<b>Location of manufacturing sites</b>	Cosayach Nitratos S.AO, Oficina Cala Cala S/N Pozo Almonte Iquique Chile

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Iodine		Active Substance	7553-56-2	231-442-4	0,5
Fatty alcohol ethoxylate			69011-36-5	500-241-6	2,5

### 2.2. Type of formulation

AL - Any other liquid
-----------------------

### 3. Hazard and precautionary statements

#### Hazard statements

Causes serious eye irritation.  
Harmful to aquatic life with long lasting effects.

#### Precautionary statements

Wash hands thoroughly after handling.  
Wear eye protection.  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
If eye irritation persists: Get medical advice.  
Avoid release to the environment.

### 4. Authorised use(s)

#### 4.1 Use description

##### Use 1 - Use # 3.1 – Manual dip treatment, post milking teat disinfectants

#### Product type

PT03 - Veterinary hygiene (Disinfectants)

#### Where relevant, an exact description of the authorised use

-

#### Target organism(s) (including development stage)

Scientific name: Bacteria  
Common name: Bacteria  
Development stage:

Scientific name: yeasts  
Common name: Yeasts  
Development stage:

#### Field(s) of use

Indoor  
Teat disinfection for post –milking, manual dip treatment of cow teats

#### Application method(s)

Open system: dip treatment -  
For post-milking min. 5 minutes contact time

**Application rate(s) and frequencies**

Apply an amount of product that is sufficient to cover the entire teat, at the maximum 10 ml per treatment - Product is ready to use - 2 times per day

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

20kg, 60kg, 200kg, 210kg HDPE drum and 1000kg HDPE IBC

**4.1.1 Use-specific instructions for use**

Product for disinfection of cow teat by dipping treatment, after milking.

Fill the dipping cup with the indicated quantity of undiluted product straight from the original packaging. With every treatment, care should be taken to the volume of treatment product in the dipping cup. To reduce dermal exposure, the use of a dosing pump for filling the product into the application equipment is recommended.

**4.1.2 Use-specific risk mitigation measures**

Wear protective chemical resistant gloves (glove material to be specified by the authorisation holder within the product information) and eye protection.

**4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

See general directions for use of the meta SPC 3

**4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See general directions for use of the meta SPC 3

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of the meta SPC 3

#### 4.2 Use description

##### Use 2 - Use # 3.2 – Manual spraying treatment, post milking teat disinfectants

**Product type**

PT03 - Veterinary hygiene (Disinfectants)

**Where relevant, an exact description of the authorised use**

-

**Target organism(s) (including development stage)**

Scientific name: Bacteria  
Common name: Bacteria  
Development stage:

Scientific name: yeasts  
Common name: Yeasts  
Development stage:

**Field(s) of use**

Indoor

Teat disinfection for post – milking, manual spray treatment of cow teats

**Application method(s)**

Open system: spray treatment -  
For post-milking min. 5 minutes contact time

**Application rate(s) and frequencies**

Apply an amount of product that is sufficient to cover the entire teat, at the maximum 15ml per spray treatment - Product is ready to use - 2 times per day

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

20kg, 60kg, 200kg, 210kg HDPE drum and 1000kg HDPE IBC

#### 4.2.1 Use-specific instructions for use

Product for disinfection of cow teat by spraying, after milking.

Apply an amount of product that is sufficient to cover the entire teat, at the maximum 15 ml per spray treatment

Fill the spray flask with the indicated quantity of undiluted product straight from the original packaging. To reduce dermal exposure, the use of a dosing pump for filling the product into the application equipment is recommended.

#### 4.2.2 Use-specific risk mitigation measures

Wear protective chemical resistant gloves (glove material to be specified by the authorisation holder within the product information), protective clothing, chemical resistant boots, and eye protection.

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of the meta SPC 3

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of the meta SPC 3

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of the meta SPC 3

### 4.3 Use description

#### Use 3 - Use # 3.3 – Automated dip treatment, post-milking teat disinfectants

**Product type**

PT03 - Veterinary hygiene (Disinfectants)

**Where relevant, an exact description of the authorised use**

-

**Target organism(s) (including development stage)**

Scientific name: Bacteria  
Common name: Bacteria  
Development stage:

Scientific name: yeasts  
Common name: Yeasts  
Development stage:

**Field(s) of use**

Indoor

Teat disinfection for post – milking, automated dip treatment of cow teats



<b>Application method(s)</b>	Open system: dip treatment - For post-milking min. 5 minutes contact time.
<b>Application rate(s) and frequencies</b>	Apply an amount of product that is sufficient to cover the entire teat, at the maximum 10ml per dip treatment - Product is ready to use. - 3 times per day
<b>Category(ies) of users</b>	Professional
<b>Pack sizes and packaging material</b>	20kg, 60kg, 200kg, 210kg HDPE drum and 1000kg HDPE IBC

#### 4.3.1 Use-specific instructions for use

Product for disinfection of cow teat by dipping, after milking.

Apply an amount of product that is sufficient to cover the entire teat, at the maximum 10 ml per dip treatment.

Use the undiluted product straight from the original packaging. To reduce dermal exposure, the use of a dosing pump for filling the product into the application equipment is recommended.

#### 4.3.2 Use-specific risk mitigation measures

See general directions for use of the meta SPC 3

#### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See general directions for use of the meta SPC 3

#### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See general directions for use of the meta SPC 3

### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See general directions for use of the meta SPC 3

## 4.4 Use description

### Use 4 - Use # 3.4 – Automated spraying treatment, post-milking teat disinfectants

<b>Product type</b>	PT03 - Veterinary hygiene (Disinfectants)
<b>Where relevant, an exact description of the authorised use</b>	-
<b>Target organism(s) (including development stage)</b>	Scientific name: Bacteria Common name: Bacteria Development stage:  Scientific name: yeasts Common name: Yeasts Development stage:
<b>Field(s) of use</b>	Indoor  Teat disinfection for post – milking, automated spray treatment of cow teats
<b>Application method(s)</b>	Open system: spray treatment - For post-milking min. 5 minutes contact time.
<b>Application rate(s) and frequencies</b>	15 ml per spray treatment - Product is ready to use - 3 times per day
<b>Category(ies) of users</b>	Professional
<b>Pack sizes and packaging material</b>	20kg, 60kg, 200kg, 210kg HDPE drum and 1000kg HDPE IBC

### 4.4.1 Use-specific instructions for use

Product for disinfection of cow teat by spraying, after milking.

Apply an amount of product that is sufficient to cover the entire teat, at the maximum of 15 ml per spray treatment.

Use the undiluted product straight from the original packaging. To reduce dermal exposure, the use of a dosing pump for filling the product into the application equipment is recommended.

#### **4.4.2 Use-specific risk mitigation measures**

See general directions for use of the meta SPC 3

#### **4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

See general directions for use of the meta SPC 3

#### **4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See general directions for use of the meta SPC 3

#### **4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

See general directions for use of the meta SPC 3

### **5. General directions for use**

#### **5.1. Instructions for use**

The products must be brought to temperatures above 20°C before use.

For the post-milking treatment: The teats should be treated as soon as possible after milking in such a way that the teats are fully covered. Let the teats air dry. The cows should be kept in a standing position until the product is fully dried (at least 5 minutes).

#### **5.2. Risk mitigation measures**

In case a combination of pre- and post-milking disinfection is necessary, using another biocidal product not containing iodine has to be considered for pre- milking disinfection.

Keep out of reach of children.

### **5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

First aid: general information: immediately remove any clothing soiled by the product

After inhalation: Supply fresh air; consult doctor in case of complaints.

After skin contact: Wash skin thoroughly. Generally the product does not irritate the skin.

After eye contact: Rinse opened eye for several minutes under running water. Remove contact lenses, if present and easy to do. Continue rinsing.

After swallowing: Rinse out mouth and then drink water. If you feel unwell, get medical advice/attention.

If medical advice is needed, have product container or label at hand.

Do not allow product to enter sewers/ surface or ground water.

### **5.4. Instructions for safe disposal of the product and its packaging**

At the end of the treatment, dispose unused product and the packaging in accordance with local requirements. Used product can be flushed to the municipal sewer or disposed to the manure deposit depending on local requirements. Avoid release to an individual waste water treatment plant.

### **5.5. Conditions of storage and shelf-life of the product under normal conditions of storage**

Protect from frost. Store in a cool, dry place, away from direct sunlight and out of reach from children.

Shelf life: 2 years.

### **6. Other information**