


Regulation (EU) No 528/2012 concerning the making available on the market and use of biocidal products

**PRODUCT ASSESSMENT REPORT OF A BIOCIDAL  
PRODUCT FOR THE MAJOR CHANGE AND  
RENEWAL OF A NATIONAL AUTHORISATION**



Product identifier in R4BP	Agrorat BD-3 pellet
Product type(s):	14 (Rodenticide)
Active ingredient(s):	Bromadiolone
Case No. in R4BP	BC-VG014447-35 (NA-RNL) BC-SF030766-34 (NA-MAC)
Asset No. in R4BP	ES-0003184-0000
Evaluating Competent Authority	Spain
Internal registration/file no	ES/APP(NA)-2018-14-00111
Date	February 2018

## Table of content

<b>1</b>	<b>Conclusion .....</b>	<b>3</b>
<b>2</b>	<b>Summary of the product assessment .....</b>	<b>5</b>
2.1	Administrative information .....	5
2.2	Composition and formulation.....	6
2.3	Classification and Labelling according to the Regulation (EC) No 1272/2008.....	7
2.4	Use(s) appropriate for further authorisation .....	8
<b>2.5</b>	<b>General directions for use.....</b>	<b>22</b>
<b>3</b>	<b>Assessment of the product .....</b>	<b>29</b>
<b>3.1</b>	<b>Use(s) considered appropriate for authorisation after former assessment (uses evaluated by Spain).....</b>	<b>29</b>
3.2	Physical, chemical and technical properties.....	33
3.4	Methods for detection and identification.....	35
<b>3.5</b>	<b>Efficacy against target organisms.....</b>	<b>35</b>
3.6	Risk assessment for human health .....	40
3.7	Risk assessment for animal health.....	60
3.8	Risk assessment for the environment .....	61
3.9	Assessment of a combination of biocidal products .....	78
3.10	Comparative assessment .....	78
<b>4.</b>	<b>Confidential annex (Access level: “Restricted” to applicant and authority).....</b>	<b>79</b>
4.1	Full composition of the product .....	79
4.2	List of studies for the biocidal product (Confidential data) .....	81
4.2	List of studies for the biocidal product (Confidential data).....	84

# 1 Conclusion

The assessment presented in this report includes the major change submitted by the applicant according to Implementing Regulation 354/2013 in order to decrease the content of bromadiolone active substance at a level of 0.0029% w/w due to laid down in Commission Regulation (EU) 2016/1179 of 19 July 2016 amending, for the purposes of its adaptation to technical and scientific progress, Regulation (EC) No 1272/2008 of the European Parliament and of the Council. In addition, this report also includes the conditions for the renewal of the active substance, according Commission Regulation (EU) 2017/1380 of 25 July 2017.

The initial evaluation of the biocidal product AGRORAT BD-5 PELLETT containing of bromadiolone active substance at a level of 0.005% w/w should be taken into account. As the name of the product refers to the content in active substance of the product, the Spanish Competent Authority requested to the applicant changed the product name in order not to mislead the user and for enforcement tasks.

It is concluded after evaluation of new data submitted that the ready-to-use product, AGRORAT BD-3 PELLETT, with the active substance bromadiolone, at a level of 0.0029% w/w, may be authorised for use as a rodenticide (product-type 14). Some of conclusions to the initial assessment remains valid and the new information provided by the applicant to support the decrease of active substance allow granting the authorisation.

Physical, chemical and technical properties remain valid to the initial evaluation other than the stability test. A new accelerated stability test has been submitted. However, no long-term stability test has been submitted, therefore a post-authorisation requirement should be included in the authorisation certificate.

The conclusions about physical hazards and methods for detection and identification remain valid to the initial evaluation and no new information has been submitted.

New efficacy data, semi-field and field trials, have confirmed that AGRORAT BD-3 PELLETT is effective in the proposed areas of use, at the recommended dose rate.

According to Commission Regulation (EU) 2016/1179 the product AGRORAT BD-3 PELLETT, with the active substance bromadiolone, at a level of 0.0029% w/w is classified as SPECIFIC TARGET ORGAN

TOXICITY AFTER REPEATED EXPOSURE. CATEGORY 2 (STOT RE 2); H373 May cause damage to organs (blood) through prolonged or repeated exposure.

Risk assessment has been done with the new content of active substance.

The risk assessment for the environment has been performed for the intended uses indoors, outdoors around buildings and outdoor in open areas and waste dumps. Since the concentration of the active substance has been reduced, the new evaluation shows that the conclusions for the first evaluation remain valid

Therefore, AGRORAT BD-3 PELLETT can be authorised as a rodenticide product against house mice (*Mus musculus*) and brown rats (*Rattus norvegicus*). It is to be used indoors, outdoors around buildings and outdoor in open areas and waste dumps. The users can be general public, professional and trained professional. It is a ready to used pellet bait to be used in tamper-resistant bait stations. The specific intended uses of the product are in section 2.4. of this assessment report.

Please, note that this assessment report includes all the uses requested by the applicant and assessed, only as information for the concerned Member States.

Spanish CA only grants the use of AGRORAT BD-3 PELLETT according to the table 5 included in this assessment report due to our national risk mitigation measures.

## 2 Summary of the product assessment

### 2.1 Administrative information

#### 2.1.1 Identifier in R4BP

AGRORAT BD-3 PELLET
---------------------

#### 2.1.2 Manufacturer(s) of the product

<b>Name of manufacturer</b>	LABORATORIOS AGROCHEM S.L.
<b>Address of manufacturer</b>	C/ Tres Rieres, 10 08292 - Esparreguera (Barcelona) SPAIN
<b>Location of manufacturing sites</b>	C/ Tres Rieres, 10 08292 - Esparreguera (Barcelona) SPAIN

#### 2.1.3 Manufacturer(s) of the active substance(s)

<b>Active substance</b>	Bromadiolone
<b>Name of manufacturer</b>	LABORATORIOS AGROCHEM S.L.
<b>Address of manufacturer</b>	C/ Tres Rieres, 10 08292 - Esparreguera (Barcelona) SPAIN
<b>Location of manufacturing sites</b>	C/ Tres Rieres, 10 08292 - Esparreguera (Barcelona) SPAIN

## 2.2 Composition and formulation

### 2.2.1 Qualitative and quantitative information on the composition

Table 1

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[(1RS,3RS;1RS,3SR)-3-(4'-bromobiphenyl-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxycoumarin	Active substance	28772-56-7	249-205-9	0.0029
-	-	Non-active substances	-	-	-

- The product contains a bittering agent and a dye.
  - Information on the full composition is provided in the confidential annex (see chapter 4).
- According to the information provided the product contains no nanomaterial as defined in Article 3 paragraph 1 (z) of Regulation No. 528/2012

### 2.2.2 Information on the substance(s) of concern

No substance of concern was identified upon initial assessment (the application for authorisation was submitted and the assessment took place before the Biocidal Products Regulation 528/2012 entered into force).

### 2.2.3 Candidate(s) for substitution

No candidate for substitution was identified upon initial assessment (the application for authorisation was submitted and the assessment took place before the Biocidal Products Regulation 528/2012 entered into force).

Now that the Biocidal Products Regulation 528/2012 entered into force, the following substance(s) was/were identified as candidate(s) for substitution upon this renewal:

Bromadiolone does meet the exclusion criteria according to Article 5(1) BPR. Because the following exclusion criteria are met:

- toxic for reproduction category 1B
- persistent, bioaccumulative and toxic

And therefore, Bromadiolone does meet the conditions laid down in Article 10 BPR, and is consequently a candidate for substitution.

## 2.2.4 Type of formulation


Ready-to-use bait: Pellet

## 2.3 Classification and Labelling according to the Regulation (EC) No 1272/2008

Table 2

Classification	
Hazard classes, Hazard categories	Hazard statements
Specific target organ toxicity after repeated exposure. Category 2. (STOT RE 2)	H373 May cause damage to organs (blood) through prolonged or repeated exposure

Table 3

Labelling		
	Code	Pictogram / Wording
Pictograms	GHS08	
Signal word		WARNING
Hazard statements	H373	May cause damage to organs (blood) through prolonged or repeated exposure
Supplemental hazard information	-	
Supplemental label elements	-	
Precautionary statements	P102	Keep out of reach of children
	P103	Read label before use.
	P260	Do not breathe dust.
	P280	Wear protective gloves.
	P314	Get medical advice/attention if you feel unwell.
	P501	Dispose of contents and/ or container as a hazardous waste to a registered establishment or undertaking, in accordance with current regulations.
Note	-	

## 2.4 Use(s) appropriate for further authorisation

In order to make proper use of the standard sentences for SPCs for rodenticides it is considered necessary to split the uses currently evaluated in Spain further down:

**Table 4**

Use(s) considered appropriate for authorisation after former assessment (uses currently <u>evaluated in SPAIN</u> )		Use(s) appropriate for further authorisation	
1	House mice and/or brown rats – general public– in door	1	House mice and brown rats – general public - indoor
		2	Brown Rats – general public – outdoor around buildings
2	House mice and/or brown rats – professionals – indoor	3	House mice – professionals - indoor
		4	Brown Rats – professionals - indoor
		5	House mice and/or brown rats – Professionals – outdoor around buildings
3	House mice and/or brown rats – trained professionals – indoor	6	House mice and/or brown rats – trained professionals - indoor
		7	House mice and/or brown rats – trained professionals – outdoor around buildings
		8	Brown Rats – trained professionals – outdoor open areas & waste dumps

### Uses authorized in Spain according national Risk Mitigation Measures

**Table 5**

Use(s) considered appropriate for authorisation after former assessment (uses currently <u>under authorisation in Spain</u> )	Use(s) appropriate for authorisation in Spain according national Risk Mitigation Measures.
House mice and/or brown rats – general public– in and around buildings	House mice and brown rats – general public - indoor
	Brown Rats – general public – outdoor around buildings
House mice and/or brown rats – professional– in and around buildings	House mice – professionals - indoor
	Brown Rats – professionals - indoor
	Brown Rats – Professionals – outdoor around buildings
House mice and/or brown rats – trained professional– in and around buildings, open areas and waste dumps	House mice and/or brown rats – trained professionals - indoor
	Brown Rats – trained professionals – outdoor around buildings.



### 2.4.1 Use 1– House mice and brown rats– general public – indoor

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Indoor.
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	<b>Mice:</b> bait boxes with 60g of product each 5-10m 60g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10m (5m in case of strong infestation and 10m in case of weak infestation). <b>Rats:</b> bait boxes with 100g of product each 5-10m 100g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters (5m in case of strong infestation and 10m in case of weak infestation).
Category(ies) of users	General public
Pack sizes and packaging material	<b>Maximum pack size of 150g.</b> Number of packed bags per packaging: up to 150g Grams/kg of bait per packed bag: individual sachets of 10 to 100 g Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.

#### 2.4.1.1 Use-specific instructions for use

. The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

#### 2.4.1.2 Use-specific risk mitigation measures

- See section 2.5.2

**2.4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- See section 2.5.3.

**2.4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

-See section 2.5.4

**2.4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

-See section 2.5.5

**2.4.2 Use 2– Brown Rats – general public – Outdoor around buildings**

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Outdoor around buildings
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	<b>Rats:</b> bait boxes with 100g of product each 5-10m  100g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters (5m in case of strong infestation and 10m in case of weak infestation)
Category(ies) of users	General public
Pack sizes and packaging material	<b>Maximum pack size of 150g.</b>  Number of packed bags per packaging: up to 150g Grams/kg of bait per packed bag: individual sachets of 10, 15, 20, 25, 30, 50, 60 and 100 g.

Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.

#### **2.4.2.1 Use-specific instructions for use**

- Place the bait stations in areas not liable to flooding.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

#### **2.4.2.2 Use-specific risk mitigation measures**

- See section 2.5.2

#### **2.4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- See section 2.5.3.

#### **2.4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

- See section 2.5.4

#### **2.4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

- See section 2.5.5

### 2.4.3 Use 3- House mice – professionals – indoor

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice)
Field(s) of use	Indoor.
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations, in sachets or as loose pellet
Application rate(s) and frequency	<b>Mice:</b> bait boxes with 60g of product each 5-10m  60g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters (5m in case of strong infestation and 10m in case of weak infestation).
Category(ies) of users	Professionals
Pack sizes and packaging material	<b>Minimum pack size of 3 kg.</b>  Number of packed bags per packaging: up to 10 kg. Grams/kg of bait per packed bag: individual sachets of 10 to 30g.  Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.  Furthermore, the product can be supplied as loose pellet directly inside the secondary packaging mentioned above.

#### 2.4.3.1 Use-specific instructions for use

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Follow any additional instructions provided by the relevant code of best practice.

### 2.4.3.2 Use-specific risk mitigation measures

-See section 2.5.2

### 2.4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

### 2.4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-See section 2.5.4

### 2.4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-See section 2.5.5

## 2.4.4 Use 4 – Brown Rats – professionals – indoor

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Indoor.
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations, in sachets or as loose pellet.
Application rate(s) and frequency	<b>Rat:</b> bait boxes with 100g of product each 5-10m  100g of bait per bait station. If more than one bait station is needed,

	the minimum distance between bait stations should be of 5-10 meters (5m in case of strong infestation and 10m in case of weak infestation).
Category(ies) of users	Professionals
Pack sizes and packaging material	<p><b>Minimum pack size of 3 kg.</b></p> <p>Number of packed bags per packaging: up to 10 kg. Grams/kg of bait per packed bag: individual sachets of 10 to 100 g</p> <p>Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.</p> <p>Furthermore, the product can be supplied as loose pellet directly inside the secondary packaging mentioned above.</p>

#### 2.4.4.1 Use-specific instructions for use

- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Follow any additional instructions provided by the relevant code of best practice.

#### 2.4.4.2 Use-specific risk mitigation measures

- See section 2.5.2

#### 2.4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 2.4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-See section 2.5.4

#### 2.4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-See section 2.5.5

#### 2.4.5 Use 5 – House mice and/or brown rats – professionals – outdoor around buildings

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Outdoor around buildings
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations, in sachets or as loose pellet
Application rate(s) and frequency	<b>Rats:</b> bait boxes with 100 g of product each 5-10m  100g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters (5m in case of strong infestation and 10m in case of weak infestation).  <b>Mice:</b> bait boxes with 60g of product each 5-10m.  60g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters (5m in case of strong infestation and 10m in case of weak infestation).
Category(ies) of users	Professionals
Pack sizes and packaging material	<b>Minimum pack size of 3 kg.</b>  Number of packed bags per packaging: up to 10 kg. Grams/kg of bait per packed bag: individual sachets 10 to 100g  Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE

or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.

Furthermore, the product can be supplied as loose pellet directly inside the secondary packaging mentioned above.

#### **2.4.5.1 Use-specific instructions for use**

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [*for mice* - at least every 2 to 3 days at] [*for rats* - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- Follow any additional instructions provided by the relevant code of best practice.

#### **2.4.5.2 Use-specific risk mitigation measures**

- Do not apply this product directly in the burrows

#### **2.4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### **2.4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

- See section 2.5.4



### 2.4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-See section 2.5.5

### 2.4.6 Use 6 - House mice and/or brown rats – trained professionals – indoor

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Indoor
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations, in sachets or as loose pellet
Application rate(s) and frequency	<b>Rats:</b> bait boxes with 100-200 g per baiting point <b>Mice:</b> bait boxes with 60-100 g per baiting point
Category(ies) of users	Trained professionals
Pack sizes and packaging material	<b>Minimum pack size of 3 kg.</b> Number of packed bags per packaging: up to 10 kg. Grams/kg of bait per packed bag: individual sachets of 10 to 100 g Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC. Furthermore, the product can be supplied as loose pellet directly inside the secondary packaging mentioned above.

#### 2.4.6.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.
- Follow any additional instructions provided by the relevant code of best practice.

#### 2.4.6.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated

area and their surroundings) about the rodent control campaign

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use the product in pulsed baiting treatments.
- This product shall only be used indoors and in places that are not accessible to children or non-target animals.

**2.4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

**2.4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

-See section 2.5.4.

**2.4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

-See section 2.5.5.

**2.4.7 Use 7 – House mice and/or brown rats – trained professionals – outdoor around buildings**

Product Type(s)	14
Where relevant, an exact	Not relevant for rodenticides

description of the use	
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Outdoor around buildings
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations, in sachets or as loose pellet
Application rate(s) and frequency	<b>Rats:</b> bait boxes with 100-200 g per baiting point <b>Mice:</b> bait boxes with 60-100 g per baiting point
Category(ies) of users	Trained professionals
Pack sizes and packaging material	<b>Minimum pack size of 3 kg.</b>  Number of packed bags per packaging: up to 10 kg. Grams/kg of bait per packed bag: individual sachets of 10 to 100g  Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.  Furthermore, the product can be supplied as loose pellet directly inside the secondary packaging mentioned above.

#### 2.4.7.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
  - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- Follow any additional instructions provided by the relevant code of best practice.

#### 2.4.7.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.

- Do not apply this product directly in the burrows.

### 2.4.7.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

### 2.4.7.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See section 2.5.4

### 2.4.7.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See section 2.5.5

## 2.4.8 Use 8 – Brown Rats – trained professionals – Outdoor open areas & waste dumps

Product Type(s)	14
Where relevant, an exact description of the use	Not relevant for rodenticides
Target organism(s) (including development stage)	<i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Outdoor open areas Outdoor waste dumps
Application method(s)	Ready-to-use bait to be used in tamper-resistant bait stations, in sachets or as loose pellet
Application rate(s) and frequency	<b>Rats:</b> bait boxes with 100-200 g per baiting point.
Category(ies) of users	Trained professionals
Pack sizes and packaging material	<b>Minimum pack size of 3 kg.</b>

	<p>Number of packed bags per packaging: up to 10 kg.          Grams/kg of bait per packed bag: individual sachets of 10 to 100 g</p> <p>Packaging material: Bags, sacks, Buckets, Tubes, Bottles and Sachets. Material: Carton + PET OR LDPE, PE or PP or PET or LDPE or PET / PET MET / PE or PET / ALU / PE or PET / PE or PA / PE, HDPE, PVC.</p> <p>Furthermore, the product can be supplied as loose pellet directly inside the secondary packaging mentioned above.</p>
--	---

#### 2.4.8.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period
- Follow any additional instructions provided by the relevant code of best practice

#### 2.4.8.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.

#### 2.4.8.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 2.4.8.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-See section 2.5.4

#### 2.4.8.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-See section 2.5.5

### 2.5 General directions for use

#### 2.5.1 Instructions for use

##### General Public:

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- *[Do not open the sachets containing the bait - where relevant for the bait formulation in the product].*
- Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.
- Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Do not place bait stations near water drainage systems where they can come into contact with water.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

- Remove the remaining bait or the bait stations at the end of the treatment period.

**Professionals:**

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (*see section 5.3 for the information to be shown on the label*).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after

using the product.

- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely those there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.
- Bait in sachets: Do not open the sachets containing the bait.
- Loose grain: Place the bait in the baiting point by using a dosage devise. Specify the methods to minimise dust (e.g. wet wiping).

### **Trained professionals:**

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (*see section 5.3 for the information to be shown on the label*).

When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant



as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Bait in sachets: Do not open the sachets containing the bait
- Loose Grain: Place the bait in the baiting point by using a dosage devise. Specify the methods to minimise dust (e.g. wet wiping).

### **2.5.2 Risk mitigation measures:**

#### **General Public:**

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation)

or to detect rodent activity).

- The product information (i.e. label and/or leaflet) shall clearly show that:
- The product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- Users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.
- Dispose dead rodents in accordance with local requirements [*The method of disposal shall be described specifically in the national SPC and be reflected on the product label*].

### **Professionals:**

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
  - .the product shall not be supplied to the general public (e.g. "for professionals only").
  - .the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- Users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations")

- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements [*The method of disposal shall be described specifically in the national SPC and be reflected on the product label*]

### **Trained Professionals:**

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements [*The method of disposal shall be described specifically in the national SPC and be reflected on the product label*].

### **2.5.3 Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:
- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, always check for and remove contact lenses, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label *[insert country specific information]*. Contact a veterinary surgeon in case of ingestion by a pet *[insert country specific information]*
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre *[insert national phone number]*"
- Hazardous to wildlife.

#### **2.5.4 Instructions for safe disposal of the product and its packaging**

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements *[The method of disposal shall be described specifically in the national SPC and be reflected on the product label]*.
- Use of gloves is recommended.

#### **2.5.5 Conditions of storage and shelf-life of the product under normal conditions of storage**

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: two years

#### **2.5.6 Other information**

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use

tools such as tongs when disposing them.

- This product contains a bittering agent and a dye.

**Post-authorisation requirements:**

- Long-term stability test within 2 years.

### 3 Assessment of the product

#### 3.1 Use(s) considered appropriate for authorisation after former assessment (uses evaluated by Spain)

##### 3.1.1 Use 1 – House mice and/or brown rats – general public– indoor

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	Indoors
Application method(s)	The biocidal product is a ready to use pellet bait (in sachets or loose pellet)
Application rate(s) and frequency	For <b>rats</b> , each bait point usually contains up to 100g of bait and it should be placed in 10 m <sup>2</sup> depending on the level of infestation. For <b>mice</b> each bait point usually contains up to 50g of bait and it should be placed 10 m <sup>2</sup> depending on the level of infestation.
Category(ies) of users	General public
Pack sizes and packaging material	<ul style="list-style-type: none"> <li>· <b>Loose pellet:</b></li> <li>· <u>Bag or sack (Sachet):</u> Description: Prefabricated bags or serial production bags, both thermal-welded Material: PE or PP or PET or LDPE or PET/PET MET/PE or PET/ALU/PE or PET/PE or PA/ PE. Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <u>Plastic bucket/tub</u> Description: Rectangular or conical bucket sealed Material: HDPE or PE or PP Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <u>Bottle</u> Description: Conical bottle sealed</li> </ul>

	<p>Material: HDPE or PE or PP or PET or PVC  Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg  - <u>Carton Sachet</u>:  Description: Prefabricated bags or serial production bags, both thermal-welded  Material: Tektura (kind of carton) + PET or LDPE  Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</p> <p><b>Pellet in sachets</b>: individual polypropylene or polyethylene sachets each containing 10, 20, 25 or 50g.  - <u>Bag or sack (Sachet)</u>:  Description: Prefabricated bags or serial production bags, both thermal-welded  Material: PE or PP or PET or LDPE or PET/PET MET/PE or PET/ALU/PE or PET/PE or PA/ PE .  Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg  - <u>Paper Bag</u>:  Description: Prefabricated bags or serial production bags, both sewn or glued.  Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg  - <u>Cardboard box</u>:  Description: Self-assembly cardboard box with sealed or glued flaps  Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg  - <u>Plastic bucket</u>  Description: Rectangular or conical bucket sealed  Material: HDPE  Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg</p>
--	---

### 3.1.2 Use 2 – House mice and/or brown rats – professional– indoors

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	indoors
Application method(s)	The biocidal product is a ready to use pellet bait (in sachets or loose pellet)
Application rate(s) and frequency	For <b>rats</b> , each bait point usually contains up to 100g of bait and it should be placed in 10 m <sup>2</sup> depending on the level of infestation.

	For <b>mice</b> each bait point usually contains up to 50g of bait and it should be placed 10 m2 depending on the level of infestation.
Category(ies) of users	Professional
Pack sizes and packaging material	<ul style="list-style-type: none"> <li>· <b>Loose pellet:</b></li> <li>· <u>Bag or sack (Sachet):</u> Description: Prefabricated bags or serial production bags, both thermal-welded  Material: PE or PP or PET or LDPE or PET/PET MET/PE or PET/ALU/PE or PET/PE or PA/ PE . Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <u>Plastic bucket/tub</u> Description: Rectangular or conical bucket sealed Material: HDPE or PE or PP Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <u>Bottle</u> Description: Conical bottle sealed Material: HDPE or PE or PP or PET or PVC Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <u>Carton Sachet:</u> Description: Prefabricated bags or serial production bags, both thermal-welded Material: Tektura (kind of carton) + PET or LDPE Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li><b>Pellet in sachets:</b> individual polypropylene or polyethylene sachets each containing 10, 20, 25 or 50g.</li> <li>· <u>Bag or sack (Sachet):</u> Description: Prefabricated bags or serial production bags, both thermal-welded Material: PE or PP or PET or LDPE or PET/PET MET/PE or PET/ALU/PE or PET/PE or PA/ PE. Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg</li> <li>· <u>Paper Bag:</u> Description: Prefabricated bags or serial production bags, both sewn or glued. Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg</li> <li>· <u>Cardboard box:</u> Description: Self-assembly cardboard box with sealed or glued flaps Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g and 1kg</li> <li>· <u>Plastic bucket</u> Description: Rectangular or conical bucket sealed Material: HDPE Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400 and 500g</li> </ul>

	and 1kg
--	---------

### 3.1.3 Use 3 – House mice and/or brown rats – trained professional–indoors

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	indoors
Application method(s)	The biocidal product is a ready to use pellet bait (in sachets or loose pellet)
Application rate(s) and frequency	For <b>rats</b> , each bait point usually contains up to 100g of bait and it should be placed in 10 m2 depending on the level of infestation.  For <b>mice</b> each bait point usually contains up to 50g of bait and it should be placed 10 m2 depending on the level of infestation.
Category(ies) of users	Trained Professional
Pack sizes and packaging material	<p><b><u>Loose pellet:</u></b> The maximum capacity of the packages is restricted to 10kg. When greater capacity packaging marketed, the content of those have to be divided into bags weighing 10kg or less each.</p> <ul style="list-style-type: none"> <li>· <b><u>Bag or sack (Sachet):</u></b> Description: Prefabricated bags or serial production bags, both thermal-welded Material: PE or PP or PET or LDPE or PET/PET MET/PE or PET/ALU/PE or PET/PE or PA/ PE. Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <b><u>Plastic bucket/tub</u></b> Description: Rectangular or conical bucket sealed Material: HDPE or PE or PP Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <b><u>Bottle</u></b> Description: Conical bottle sealed Material: HDPE or PE or PP or PET or PVC Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> <li>· <b><u>Carton Sachet:</u></b> Description: Prefabricated bags or serial production bags, both thermal-welded Material: Tektura (kind of carton) + PET or LDPE Contents (Net weight): 50, 90, 100, 150, 200, 250, 300, 400, 500 g and 1, 2, 2.5, 3, 5, 10, 15, 20, 25 and 30kg</li> </ul>



	<ul style="list-style-type: none"> <li>· <b>Pellet in sachets:</b> individual polypropylene or polyethylene sachets each containing 10, 20, 25 or 50g.</li> <li>· <b>Bag or sack (Sachet):</b> Description: Prefabricated bags or serial production bags, both thermal-welded Material: PE or PP or PET or LDPE or PET/PET MET/PE or PET/ALU/PE or PET/PE or PA/ PE . Contents (Net weight): 500g and 1, 2, 3, 5, 10, 15, 20, 25 and 30kg.</li> <li>· <b>Paper Bag:</b> Description: Prefabricated bags or serial production bags, both sewn or glued. Contents (Net weight): 500g and 1, 2, 3, 5, 10, 15, 20, 25 and 30kg.</li> <li>· <b>Cardboard box:</b> Description: Self-assembly cardboard box with sealed or glued flaps Contents (Net weight): 500g and 1, 2, 3, 5, 10, 15, 20, 25 and 30kg.</li> <li>· <b>Plastic bucket</b> Description: Rectangular or conical bucket sealed Material: HDPE Contents (Net weight): 500g and 1, 2, 3, 5, 10, 15, 20 and 30kg.</li> </ul>
--	---

### 3.2 Physical, chemical and technical properties

Property	Guideline and Method	Purity of the test substance (% (w/w))	Results	Reference												
Storage stability test – accelerated storage	CIPAC MT46.3	0.0027	<p>Bromadiolone active ingredient initial content: 0.0027 ± 0.0001% w/w</p> <p>Bromadiolone active ingredient final content: 0.0025 ± 0.0001% w/w</p> <p>Δ[C] = -7.41%. The result complies with the tolerance value (-10%).</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Test</th> <th>Initial value</th> <th>Final value</th> </tr> </thead> <tbody> <tr> <td>Relative density (20°C)</td> <td>1.2369 g/mL</td> <td>1.2348 g/mL</td> </tr> </tbody> </table> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Test</th> <th>Initial value</th> <th>Final value</th> </tr> </thead> <tbody> <tr> <td>pH (1% aqueous)</td> <td>8.4</td> <td>8.6</td> </tr> </tbody> </table>	Test	Initial value	Final value	Relative density (20°C)	1.2369 g/mL	1.2348 g/mL	Test	Initial value	Final value	pH (1% aqueous)	8.4	8.6	IUCLID 3.4.1
Test	Initial value	Final value														
Relative density (20°C)	1.2369 g/mL	1.2348 g/mL														
Test	Initial value	Final value														
pH (1% aqueous)	8.4	8.6														

Property	Guideline and Method	Purity of the test substance (% (w/w))	Results			Reference									
			<table border="1"> <tr> <td data-bbox="676 416 839 472">dilution)</td> <td data-bbox="839 416 1066 472"></td> <td data-bbox="1066 416 1238 472"></td> </tr> <tr> <td data-bbox="676 506 839 562"><b>Test</b></td> <td data-bbox="839 506 1066 562"><b>Initial value</b></td> <td data-bbox="1066 506 1238 562"><b>Final value</b></td> </tr> <tr> <td data-bbox="676 573 839 707">Dustiness</td> <td data-bbox="839 573 1066 707">3.0 mg (nearly dust-free)</td> <td data-bbox="1066 573 1238 707">4.4 mg (nearly dust-free)</td> </tr> </table> <p data-bbox="676 730 1270 1178"><b>Conclusion:</b> No change in the sample appearance, colour, odour, weight variation, relative density, dustiness and in the pH value was found for the test item stored in plastic bags after 12 weeks of storage at 35°C, and no variation was found in colour or in either the internal or external configuration, or loss of sample or evident corrosion phenomena of packaging. Therefore, it can be concluded that the sample of Bromadiolone 0.0027% w/w pellet bait formulation is stable in its commercial packaging under the accelerated storage conditions.</p>			dilution)			<b>Test</b>	<b>Initial value</b>	<b>Final value</b>	Dustiness	3.0 mg (nearly dust-free)	4.4 mg (nearly dust-free)	
dilution)															
<b>Test</b>	<b>Initial value</b>	<b>Final value</b>													
Dustiness	3.0 mg (nearly dust-free)	4.4 mg (nearly dust-free)													
Storage stability test – <b>long term storage at ambient temperature</b>	Guidance on Data Requirements for Active Substances and Biocidal Products	0.0027	Study ongoing Final results: May 2019			IUCLID 3.4.1									

Apart from the properties mentioned above, neither new data was not provided nor had new guidance to be taken into account for re-assessment.

Accordingly, the conclusion from the former assessment regarding those physical, chemical and technical properties not provided remains valid.

The renewal is conditioned to the presentation of the long term stability test; therefore, a post-authorisation condition should be showed in the authorisation certificate.

### **3.3 Physical hazards and respective characteristics**

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding physical hazards and respective characteristics remains valid.

### **3.4 Methods for detection and identification**

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding methods for detection and identification remains valid.

### **3.5 Efficacy against target organisms**

AGRORAT BD-3 PELLETT is renewed with a decrease of the active substance concentration from 50 ppm to 29 ppm (major change) and a biocidal product name change (previously AGRORAT BD-5 PELLETT) and is used against Brown rat (*Rattus norvegicus*) and House mouse (*Mus musculus*).

Taking into account that a complete efficacy data package with 0.005% w/w bromadiolone was submitted, and that the change in the formulation is basically in the content of active substance, it is assumed that the level of palatability remains the same with the new composition being at least 20% of palatability in laboratory tests.

The applicant has submitted new studies in order to support the efficacy of the new formulation of the product AGRORAT BD-3 PELLETT against *Rattus norvegicus* and *Mus musculus*: two semi-field trials and two field trials. These studies were carried out at a concentration of 0.0027% w/w bromadiolone, which is considered a worse case, and thus demonstrating the efficacy of the biocidal product with the new concentration of 0.0029% w/w bromadiolone given that the change in the content of co-formulants it is considered minimum and therefore it is not affected to the efficacy of the product. Please, see the summary of the semi-field and field trial submitted by the applicant.

In conclusion, according to the test provided, ES CA consider that the biocidal product with 0.0029% w/w bromadiolone is effective against rats and mice indoor and outdoor.

Experimental data on the efficacy of the biocidal product against target organism(s)							
Function	Field of use envisaged	Test substance	Test organism(s)	Test method	Test system / concentrations applied / exposure time	Test results: effects	Reference
Rodenticide	Semi-field test	<b>Bromadiolone 0.0027% w/w pellet bait.</b>	Brown rat ( <i>Rattus norvegicus</i> )  5 females 5 males  Weight between 205 and 306g.	Semi-field test: Mortality and palatability  According to TNG for PT 14 and Transitional Guidance for PT14	Rats placed by sex in a circular conditioned space with three rectangular surfaces at 20.15-23.8°C of temperature with an air exchange of 20-35 rph and a relative humidity between 42% and 64%. The total area of the habitat was 2.7414 m <sup>2</sup> (0.548m <sup>2</sup> / rat).  Photoperiod: 12 h light/12 h dark  Food, drink and test item were placed in vessels ad libitum.  Acclimation period (3 days), Pre-feeding period (4 days) and Administration period (Bromadiolone pellet bait vs. EPA STANDARD, 4 days) and Observation period.	Mean consumption test item: 45.14% (482.3 g)  Average mortality occurrence: 100% at day 5.3 after the introduction of the test item.  Palatability: Acceptable (≥20%) Mortality: Acceptable (≥90%)	IUCLID 6.7
Rodenticide	Field test (Indoor/Outdoor)	<b>Bromadiolone 0.0027% w/w pellet bait.</b>	Brown rat ( <i>Rattus norvegicus</i> Berk)	Field test. According to Transitional Guidance for PT14 and	The trial was set up in an agricultural habitat (breeding stables for cows, fodder and equipment warehouses).  -Pre-treatment census (5 days):	-Pre-treatment: consumption (on the last 4 days): 669.5 g/day and average tracking score values of 15-24.	IUCLID 6.7

Experimental data on the efficacy of the biocidal product against target organism(s)							
Function	Field of use envisaged	Test substance	Test organism(s)	Test method	Test system / concentrations applied / exposure time	Test results: effects	Reference
				OEPP/EP PO principles: PP 1/114(2)	census bait stations (100 g) and tracking patches -Lag phase: 4 days -Treatment (16 days) : 100 g of poisoned test bait were daily put down in each station -Lag phase: 5 days -Post-treatment census (5 days): census bait stations (100 g) and tracking patches.  *Each bait station will be spaced out 5-10 m from each other (5 m in case of strong infestation; 10 m in case of weak infestation).	Estimate of a population size of a minimum of 30-40 rats. -Post-treatment: no bait takes was recorded. Tracking patches score= 0  Efficacy = 100 %  Percentage of bait consumed after the control operation compared to the amount of bait consumed before the control operation is ≤10% (according TNG for PT 14)	
Rodenticide	Semi-field test	<b>Bromadiolone 0.0027% w/w pellet bait.</b>	House mouse ( <i>Mus musculus</i> )  5 females	Semi-field test: Mortality and	Mice placed by sex in a circular conditioned space with two rectangular surfaces at 18.42-22.26°C of temperature with an air	Mean consumption test item: 41.56 % (84 g)	IUCLID 6.7

Experimental data on the efficacy of the biocidal product against target organism(s)							
Function	Field of use envisaged	Test substance	Test organism(s)	Test method	Test system / concentrations applied / exposure time	Test results: effects	Reference
			5 males Weight between 21 and 33g	palatability According to TNG for PT 14 and Transitional Guidance for PT14	exchange of 20-35 rph and a relative humidity between 55% and 78%. The total area of the habitat was 1.8145m <sup>2</sup> (0.0.363m <sup>2</sup> /mouse). Photoperiod: 12 h light/12 h dark Food, drink and test item were placed in vessels ad libitum. Acclimation period (3 days), Pre-feeding period (4 days) and Administration period (Bromadiolone pellet bait vs. EPA STANDARD, 4 days) and Observation period	Average mortality occurrence: 100% at day 6.2 after the introduction of the test item.  Palatability: Acceptable (≥20%) Mortality: Acceptable (≥90%)	
Rodenticide	Field test (Indoor)	<b>Bromadiolone 0.0027% w/w fresh pellet bait.</b>	House mouse ( <i>Mus musculus</i> L.)	Field test. According to Transitional Guidance for PT14 and OEPP/EPPO principles: PP 1/114(2)	The trial was set up in an agricultural habitat (breeding stables for cows and hens, fodder and equipment warehouses). -Pre-treatment census (5 days): census bait stations (60 g) and tracking patches -Lag phase: 4 days -Treatment (13 days) : 60 g of poisoned test bait were daily put	-Pre-treatment: consumption (on the last 4 days) of 294.3 g/day and average tracking score values of 12-21. Estimate of a population size of a minimum of 75-85 rats. -Post-treatment:	IUCLID 6.7

Experimental data on the efficacy of the biocidal product against target organism(s)							
Function	Field of use envisaged	Test substance	Test organism(s)	Test method	Test system / concentrations applied / exposure time	Test results: effects	Reference
					<p>down in each station</p> <p>-Lag phase: 4 days</p> <p>-Post-treatment census (5 days): census bait stations (60 g) and tracking patches.</p> <p>*Each bait station will be spaced out 5-10 m from each other (5 m in case of strong infestation; 10 m in case of weak infestation).</p>	<p>no bait takes was recorded.</p> <p>Tracking patches score= 0</p> <p>Efficacy = 100 %</p> <p>Percentage of bait consumed after the control operation compared to the amount of bait consumed before the control operation is <math>\leq 10\%</math> (according TNG for PT 14)</p>	

## **3.6 Risk assessment for human health**

### **3.6.1 Assessment of effects of the active substance on human health**

Neither new data was not provided nor had new guidance to be taken into account for re-assessment.

Accordingly, the conclusion from the former assessment regarding effects of the active substance on human health remains valid.

### **3.6.2 Assessment of effects of the product on human health**

Neither new data was not provided nor had new guidance to be taken into account for re-assessment.

Accordingly, the conclusion from the former assessment regarding effects of the product on human health remains valid.

### **3.6.3 Exposure assessment**

Bromadiolone 0.0029% w/w pellet bait is red ready-to-use cereal/pellet bait impregnated containing 0.0029% w/w bromadiolone and packaged in sachets.

Bromadiolone is not volatile. The risk of inhalation exposure to bromadiolone is considered to be potentially significant only during decanting of loose grain bait prior to loading of bait points. On the other hand, dermal exposure of users is likely to be limited to the hands. Exposure of other parts of the body can be discounted as negligible.

Regarding human exposure no studies have been submitted; therefore, the exposure assessment has been performed using the paper "HEEG opinion on a harmonised approach for the assessment of rodenticides (anticoagulants)" agreed at TMII 2011. This paper was based on an operator exposure study conducted by CEFIC/EBPF Rodenticides Data Development Group (Chambers et al. (2004)) and the number of manipulations agreed at TMII 2010.

This opinion was revised by Ad hoc Working Group on Human Exposure in September 2016, including the sentence "*For package sizes ≤ 10kg, loose grains have to be placed on the bait point by using a dosage device (decanting is to be avoided)*". Since AGRORAT BD-3 is put on the market in packs up to 10kg, "mixing & loading (decanting of grain bait)" scenario has not been done, and therefore inhalation exposure is assessed as negligible.



**Identification of main paths of human exposure towards active substance and substances of concern from its use in biocidal product**

Summary table: relevant paths of human exposure							
Exposure path	Primary (direct) exposure			Secondary (indirect) exposure			
	Trained professional use	Professional use	General public (Non-professional)	Trained professional use	Professional use	General public	Via food
Inhalation	No	No	No	No	No	No	No
Dermal	Yes	Yes	Yes	Yes	Yes	Yes	n.a.
Oral	n.a.	n.a.	n.a.	No	No	Yes	n.a.

**List of scenarios**

Summary table: scenarios			
Scenario number	Scenario	Primary or secondary exposure Description of scenario	Exposed group
1.	Application (deploying bait stations)	<p>Primary exposure during the deploying the product or loading and placing the bait boxes.</p> <p>This scenario is taken in accordance to the HEEG Opinion (2012) where only potential dermal exposure is foreseeable, while inhalation exposure is assessed as negligible.</p> <p>On the other hand, following HEEG Opinion (2010) 63 loading manipulations are assumed for professional trained operator while 5 manipulations are considered for non-trained professional and non-professional</p>	Trained professional, professional and general public (non-professionals)

Summary table: scenarios			
Scenario number	Scenario	Primary or secondary exposure Description of scenario	Exposed group
2	Post-application (Cleaning) (refillable and sealed bait stations)	<p>Primary exposure during cleaning/disposal of bait boxes.</p> <p>The operator emptied a loaded bait station containing pellet baits.</p> <p>As in the exposure scenario before, only potential dermal exposure is foreseeable, while inhalation exposure is assessed as negligible.</p> <p>On the other hand, following HEEG Opinion (2010) 16 cleaning manipulations are assumed for professional operator and 5 for non-trained professional and non-professional.</p>	Trained professional, professional and general public (non-professionals)
3	Touching dead rodents	<p>Secondary exposure: dermal contact with dead rodents.</p> <p>Adults may touch dead rodents and dispose them from the treated area. The contact with the products applied in bait stations or outdoors is considered an incidental exposure and therefore it will be very limited.</p>	Bystanders (Adults)
.4	Ingestion of bait by children	<p>Secondary exposure: It is assumed that a child might ingest 5 g of the bait (a general assumption of poison centre specialists, what children would ingest, sees CAR Difethialone, Norway 2007).</p>	Bystanders (Children)

## Professional exposure

### Trained professional (Pest Control Operator)

#### *Scenario [1] – Deploying bait station (Application phase)*

#### **Description of Scenario [1] – Trained Professional user (Pest Control Operator)**

In this scenario the operator may be in contact with the bait when the bait is deployed, loaded and/or placed. Professional operator is bounded to use PPE during the development of the different tasks of his work. Inhalation exposure is considered as negligible during this scenario.

Total systemic exposure has been assessed with (Tier 2) and without PPE (Tier 1).

	Parameters	Value
Tier 1	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg
	Dermal exposure data	2.04 mg <sub>product</sub> /manipulation (75 <sup>th</sup> percentile)
	Number of manipulations during loading	63
Tier 2	PPE (gloves)	10% of permeability (90% of protection)

#### Calculations for Scenario [1]

Summary table: estimated exposure from Trained professional users					
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake
Scenario [1]	Tier 1 / No PPE	-	2.23 x 10 <sup>-7</sup> mg/kg bw/day	-	2.23 x 10 <sup>-7</sup> mg/kg bw/day
Scenario [1]	Tier 2 / PPE (gloves)	-	2.23 x 10 <sup>-8</sup> mg/kg bw/day	-	2.23 x 10 <sup>-8</sup> mg/kg bw/day

#### Scenario [2] – Cleaning/disposal phase

Description of Scenario [2] – Trained Professional user (Pest Control Operator)
<p>In this scenario the operator may be in contact with the bait when the bait is cleaned and/or disposed. Professional operator is bounded to use PPE during the development of the different tasks of his work. Inhalation exposure is considered as negligible during this scenario.</p> <p>Total systemic exposure has been assessed with (Tier 2) and without PPE (Tier 1).</p>

	Parameters	Value
Tier 1	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg
	Dermal exposure data	3.79 mg <sub>product</sub> /manipulation (75 <sup>th</sup> percentile)
	Number of manipulations	16
Tier 2	PPE (gloves)	10% of permeability (90% of protection)

### Calculations for Scenario [2]

Summary table: estimated exposure from trained professional users					
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake
Scenario [2]	Tier 1 / No PPE	-	$1.055 \times 10^{-7}$ mg/kg bw/day	-	$1.055 \times 10^{-7}$ mg/kg bw/day
Scenario [2]	Tier 2 / PPE (gloves)	-	$1.055 \times 10^{-8}$ mg/kg bw/day	-	$1.055 \times 10^{-8}$ mg/kg bw/day

### Combined scenarios

Summary table: combined systemic exposure from trained professional users					
Scenarios combined	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake	
Scenarios [1 + 2] – Tier 1	$1.45 \times 10^{-6}$ mg/kg bw/day	$(2.23 \times 10^{-7} + 1.055 \times 10^{-7})$ mg/kg bw/day	-	$3.29 \times 10^{-7}$ mg/kg bw/day	

Summary table: combined systemic exposure from trained professional users						
Scenarios combined	Estimated inhalation uptake	Estimated uptake	dermal	Estimated uptake	oral	Estimated total uptake
Scenarios [1 + 2] – Tier 2 (PPE)	$1.45 \times 10^{-6}$ mg/kg bw/day	$(2.23 \times 10^{-8} + 1.055 \times 10^{-8})$ mg/kg bw/day		-		$3.29 \times 10^{-8}$ mg/kg bw/day

### Non-trained Professional user

#### Scenario [1] – Deploying bait station (Application phase)

Description of Scenario [1] –professional user		
<p>In this scenario the user may be in contact with the bait when the bait is deployed, loaded and/or placed. Professional user is bounded to use PPE during the development of the different tasks of his work. Inhalation exposure is considered as negligible during this scenario.</p> <p>Total systemic exposure has been assessed with (Tier 2) and without PPE (Tier 1).</p>		
	Parameters	Value
Tier 1	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg
	Dermal exposure data	2.04 mg <sub>product</sub> /manipulation (75 <sup>th</sup> percentile)
	Number of manipulations during loading	5
Tier 2	PPE (gloves)	10% of permeability (90% of protection)

## Calculations for Scenario [1]

Summary table: estimated exposure from professional users						
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake	
Scenario [1]	Tier 1 / No PPE	-	1.77x 10 <sup>-8</sup> mg/kg bw/day		1.77 x 10 <sup>-8</sup> mg/kg bw/day	
Scenario [1]	Tier 2 / PPE (gloves)	-	1.77 x 10 <sup>-9</sup> mg/kg bw/day	-	1.77 x 10 <sup>-9</sup> mg/kg bw/day	

Scenario [2] – Cleaning/disposal phase

Description of Scenario [2] –professional user		
<p>In this scenario the user may be in contact with the bait when the bait is cleaned and/or disposed. Professional operator is bounded to use PPE during the development of the different tasks of his work. Inhalation exposure is considered as negligible during this scenario.</p> <p>Total systemic exposure has been assessed with (Tier 2) and without PPE (Tier 1).</p>		
	Parameters	Value
Tier 1	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg
	Dermal exposure data	3.79 mg <sub>product</sub> /manipulation (75 <sup>th</sup> percentile)
	Number of manipulations	5
Tier 2	PPE (gloves)	10% of permeability (90% of protection)

**Calculations for Scenario [3]**

Summary table: estimated exposure from professional users						
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated uptake	total uptake
Scenario [2]	Tier 1 / No PPE	-	$3.29 \times 10^{-8}$ mg/kg bw/day	-	-	$3.29 \times 10^{-8}$ mg/kg bw/day
Scenario [2]	Tier 2 / PPE (gloves)	-	$3.29 \times 10^{-9}$ mg/kg bw/day	-	-	$3.29 \times 10^{-9}$ mg/kg bw/day

Combined scenarios

Summary table: combined systemic exposure from professional users						
Scenarios combined	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated uptake	Estimated uptake	total uptake
Scenarios [1 + 2] – Tier 1	-	$(1.77 \times 10^{-8} + 3.29 \times 10^{-8})$ mg/kg bw/day	-	-	-	$5.07 \times 10^{-8}$ mg/kg bw/day
Scenarios [1 + 2] – Tier 2 (PPE)	-	$(1.77 \times 10^{-9} + 3.29 \times 10^{-9})$ mg/kg bw/day	-	-	-	$5.07 \times 10^{-9}$ mg/kg bw/day

**General public (Non-professional) exposure**

Although non-professional users are untrained and cannot be expected to wear protective clothing, the application pattern of Bromadiolone 0.0029% w/w pellet bait by the general public is similar to non-trained professional users. The use is occasional, for a short time in a single day and unlikely to be repeated more than once a week. However, in accordance with the CARs on various Rodenticides and proposed by HEEG (2010), fewer manipulations as compared to professionals are considered. Hence, 5 deploying and 5 cleaning manipulations are assumed for a non-professional user.

After use the product is likely to be collected and disposed of in a controlled way (as directed by product labels).

Scenario [1] – Deploying bait station (Application phase)

<b>Description of Scenario [1] – General public (Non-professional)</b>		
<p>In this scenario the user may be in contact with the bait when the bait is loaded and placed. General public (non-professional user) is not bounded to use PPE during the development of the different tasks of product's application although its use is recommended in the product's label. Inhalation exposure is considered as negligible during this scenario.</p>		
	<b>Parameters</b>	<b>Value</b>
Tier 1	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg
	Dermal exposure data	2.04 mg <sub>product</sub> /manipulation (75 <sup>th</sup> percentile)
	Number of manipulations	5

#### Calculations for Scenario [1]

<b>Summary table: systemic exposure from general public (non-professional users)</b>						
<b>Exposure scenario</b>	<b>Tier/PPE</b>	<b>Estimated inhalation uptake</b>	<b>Estimated dermal uptake</b>	<b>Estimated oral uptake</b>	<b>Estimated uptake</b>	<b>total</b>
Scenario [1]	Tier 1 /No PPE	-	1.77 x 10 <sup>-8</sup> mg/kg bw/day	-	1.77 x 10 <sup>-8</sup> mg/kg bw/day	

#### Scenario [2] – Cleaning/disposal phase



### Description of Scenario [2] – General public (non-professional user)

During the process of cleaning the bait boxes, non-professional users are expected to collect and dispose of unused or part-used products.

After use the product is likely to be collected and disposed of in a controlled way (as directed by product labels).

Bait stations for use by the non-professional user (general public) may be supplied as lockable, tamper-proof units that may be refilled by the user.

	Parameters	Value
Tier 1	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg
	Dermal exposure data	5.7 mg <sub>product</sub> /manipulation (75 <sup>th</sup> percentile)
	Number of manipulations	5

### Calculations for Scenario [2]

#### Summary table: estimated exposure from general public (non-professional uses)

Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake
Scenario [2]	Tier 1 / No PPE	-	3.29 x 10 <sup>-8</sup> mg/kg bw/day	-	3.29 x 10 <sup>-8</sup> mg/kg bw/day

Combined scenarios

Summary table: combined systemic exposure from general public (non-professional uses)						
Scenarios combined	Estimated inhalation uptake	Estimated uptake	dermal	Estimated uptake	oral	Estimated total uptake
Scenarios [1 + 2]	-	5.07 x 10 <sup>-8</sup> mg/kg bw/day		-		5.07 x 10 <sup>-8</sup> mg/kg bw/day

**Exposure of the general public**

Adults or children/infants may be present following application and may be incidentally exposed by touching unprotected bait and/or dead rodents. For products applied in bait stations incidental exposure will be very limited.

During application of Bromadiolone 0.0029% w/w pellet bait in rodent control, secondary exposure to the rodenticidal baits may occur. Two scenarios are considered, dermal contact with dead rodents by adults and incidental ingestion of baits by children.

Scenario [3] - Dermal contact with dead rodents

Based on these scenarios as listed in Appendix 7.2.1 of TNsG (2002) Part 3, calculation of secondary exposure is performed:

Description of Scenario [3]		
It is assumed that adults may come into contact with 1 g of the bait on the exterior fur of dead rodents. However, since for hygiene reasons and prevention of diseases dead rodents should not be touched without gloves, a protection factor of 10 is considered appropriate. This assumption was considered as a real case, although it is known that PPE should only be considered for professional users.		
	Parameters	Value
Tier 1	Dermal contact	1 g
	A.S. content of BP	0.0029%
	Dermal absorption:	0.36%
	Operator body weight:	60 kg

Description of Scenario [3]		
Tier 2	PPE (gloves to remove the death rodents)	90%

Calculations for Scenario [3]

Summary table: systemic exposure from secondary exposure of general public						
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated uptake	total
Scenario [3]	Tier 1 / no PPE	-	$1.74 \times 10^{-6}$ mg/kg bw/d	-	$1.74 \times 10^{-6}$ mg/kg bw/d	
Scenario [3]	Tier 2 / with PPE	-	$1.74 \times 10^{-7}$ mg/kg bw/d	-	$1.74 \times 10^{-7}$ mg/kg bw/d	

Further information and considerations on scenario [3]

As it was mentioned before, Tier 2 was developed as a realistic case, considering that no one remove death rodents with their nude hands.

Scenario [4] - Ingestion of bait by children

Description of Scenario [4]		
As a general assumption of poison center specialists, it is assumed that children ingest 5 g of the bait. However, ingestion of 5 g represents a high overestimate of exposure, since baits contain a repellent (denatonium benzoate as bitter agent), which will most likely urge the children to spit the bait. Hence, applying the general assumption of ingestion of 10 mg of bait (TNsG default for a bait with repellent), a second assessment as Tier 2 was performed.		
	Parameters	Value
Tier 1	Amount of BP ingested	5 g

Description of Scenario [4]		
	Oral absorption	100%
	A.S. content of BP	0.0029%
	Children body weight:	10 kg
Tier 2	Amount of BP ingested, considering the presence of a bittering agent	10 mg

Calculations for Scenario [4]

Summary table: systemic exposure from secondary exposure of general public						
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake	
Scenario [4]	Tier 1 / no bittering agent	-	-	0.0145 mg/kg bw/d	0.0145 mg/kg bw/d	
Scenario [4]	Tier 2 / with bittering agent	-	-	$2.9 \times 10^{-5}$ mg/kg bw/d.	$2.9 \times 10^{-5}$ mg/kg bw/d.	

#### Further information and considerations on scenario [5]

As it was mentioned before, Tier 2 was developed as a realistic case, considering that all rodenticide products (as Bromadiolone 0.0029% w/w pellet bait) contains a bittering agent.

#### Monitoring data

No monitoring studies have been submitted; therefore, the exposure assessment has been performed using the paper "HEEG opinion on a harmonised approach for the assessment of rodenticides (anticoagulants)" agreed at TMII 2011. This paper was based on an operator exposure study conducted by CEFIC/EBPF Rodenticides Data Development Group (Chambers et al. (2004)) and the number of manipulations agreed at TMII 2010.

**Dietary exposure**

Not applicable: non exposure is foreseen because the bait boxes with the product must not be placed where food, feeding stuffs, drinking water and surfaces where food is prepared an become contaminated.

**Exposure associated with production, formulation and disposal of the biocidal product**

Please see scenario [2] for professional exposure which is related with disposal of the biocidal product.

**Aggregated exposure**

No aggregated exposure is foreseeable since the product is not intended to be used under another biocidal product type.

- **Summary of exposure assessment**

Scenarios and values to be used in risk assessment			
Scenario number	Exposed group (e.g. professionals, non-professionals, bystanders)	Tier/PPE	Estimated total uptake
1.	Trained professional	Tier 1/ no PPE (unrealistic)	$2.23 \times 10^{-7}$ mg/kg bw/day
1.	Trained professional	Tier 2/ PPE	$2.23 \times 10^{-8}$ mg/kg bw/day
1.	Professional	Tier 1/ no PPE	$1.77 \times 10^{-8}$ mg/kg bw/day
1.	Professional	Tier 2/ PPE	$1.77 \times 10^{-9}$ mg/kg bw/day
1.	Public general (non-professional)	No PPE	$1.77 \times 10^{-8}$ mg/kg bw/day
2.	Trained professional	Tier 1/ no PPE (unrealistic)	$1.05 \times 10^{-7}$ mg/kg bw/day
2.	Trained professional)	Tier 2/ PPE	$1.05 \times 10^{-8}$ mg/kg bw/day
2.	Professional	Tier 1/ no PPE	$3.29 \times 10^{-8}$ mg/kg bw/day
2.	Professional	Tier 2/ PPE	$3.29 \times 10^{-9}$ mg/kg bw/day

Scenarios and values to be used in risk assessment			
Scenario number	Exposed group (e.g. professionals, non-professionals, bystanders)	Tier/PPE	Estimated total uptake
2.	General public (non-professional)	No PPE	$3.29 \times 10^{-8}$ mg/kg bw/day
3.	General public (bystander)	Tier 1/ no PPE (unlikely)	$1.74 \times 10^{-6}$ mg/kg bw/day
3.	General public (bystander)	Tier 2/ PPE (gloves)	$1.74 \times 10^{-7}$ mg/kg bw/day
4.	General public (Children)	Tier 1 (without efficient bitter agent)	0.0145 mg/kg bw/day
4.	General public (Children)	Tier 2 (with bitter agent)	$3.9 \times 10^{-5}$ mg/kg bw/day

### 3.6.4 Risk characterisation for human health

#### Reference values to be used in Risk Characterisation

Reference	Study	NOAEL (LOAEL) ( $\mu\text{g}/\text{kg}$ bw/day)	AF	Correction for oral absorption	Value ( $\mu\text{g}/\text{kg}$ bw/day)
AELshort-term	Teratogenicity study in rabbit	2(*)	600	70%	0.0023
AELmedium-term	Subchronic study in rabbit	0.5	300	70%	0.0012
AELlong-term	-	-	-	-	-
ARfD	Not required	-	Not required	-	Not required
ADI	Not required	-	Not required	-	Not required

(\*) LOAEL

#### Maximum residue limits or equivalent

Exposure to residues in food is not assessed because no contamination on food or feedingstuff is foreseen.

### **Risk for industrial users**

According to industrial exposure measures, where most of processes are performed on closed-systems, non risk is foreseeable for industrial users when the product is manufactured or formulated.

### **Risk for professional users**

### **Risk for Trained professional users (Pest Control Operators)**

### **Systemic effects**

Task/ Scenario	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
Deploying bait station/ Scenario [1]	Tier 1	$5 \times 10^{-4}$	$1.2 \times 10^{-6}$	$2.23 \times 10^{-7}$	18	Yes
	Tier 2			$2.23 \times 10^{-8}$	1	Yes
Cleaning Scenario [2]	Tier 1			$1.05 \times 10^{-7}$	8.8	Yes
	Tier 2			$1.05 \times 10^{-8}$	0.8	Yes

### **Combined scenarios**

Scenarios combined	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
[1] + [2]	Tier 1	$5 \times 10^{-4}$	$1.2 \times 10^{-6}$	$3.29 \times 10^{-7}$	27	Yes
	Tier 2			$3.29 \times 10^{-8}$ $1.52 \times 10^{-6}$	2.7	Yes

## Risk for non-trained professional users

### Systemic effects

Task/ Scenario	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
Deploying bait station/ Scenario [1]	Tier 1	$5 \times 10^{-4}$	$1.2 \times 10^{-6}$	$1.77 \times 10^{-8}$	1.5	Yes
	Tier 2			$1.77 \times 10^{-9}$	0.15	Yes
Cleaning Scenario [2]	Tier 1			$3.29 \times 10^{-8}$	2.75	Yes
	Tier 2			$3.29 \times 10^{-9}$	0.27	Yes

### Combined scenarios

Scenarios combined	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
[1] + [2]	Tier 1	$5 \times 10^{-4}$	$1.2 \times 10^{-6}$	$5.07 \times 10^{-8}$	4.2	Yes
	Tier 2			$5.07 \times 10^{-9}$	0.42	Yes

### Local effects

There is no need to consider local effects separately.

### Conclusion

Exposure for professional users applying 'Bromadiolone 0.0029% w/w pellet bait' for control of rats and mice is acceptable either with or without the use of PPE under the assumption of 5 manipulations per day.



**Risk for General public (non- professional users)****Systemic effects**

Task/ Scenario	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
Deploying bait station/ Scenario [1]	Tier 1	5 x 10 <sup>-4</sup>	2.3 x 10 <sup>-6</sup>	1.77 x 10 <sup>-8</sup>	0.7	Yes
Cleaning / Scenario [2]	Tier 1			3.29 x 10 <sup>-8</sup>	1.4	Yes

**Combined scenarios**

Scenarios combined	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
[1] + [2]	Tier 1	5 x 10 <sup>-4</sup>	2.3 x 10 <sup>-6</sup>	5.07 x 10 <sup>-8</sup>	2.2	Yes

**Local effects**

There is no need to consider local effects separately.

**Conclusion**

Exposure for general public (non-professional) applying 'Bromadiolone 0.0029% w/w pellet bait' for control of rats and mice is acceptable without the use of PPE under the assumption of 5 manipulations per day.

**Risk for the general public**

During application of Bromadiolone 0.0029% w/w pellet bait in rodent control, secondary exposure to the rodenticidal baits may occur. Two scenarios are considered, dermal contact with dead rodents by adults (Scenario [3]) and incidental ingestion of pellets by children (Scenario [4]). Scenarios are described in Appendix 7.2.1 of the TNsG Part 3 for wax baits and have been adopted for grain baits.

In Scenario [3], it is assumed that adults would come into contact with 1 g of the bait on the exterior fur of dead rodents (Tier 1). However, since for hygiene reasons and prevention of diseases dead rodents

should not be touched without gloves. Hence, a protection factor of 10 is considered appropriate for this task (Tier 2).

Children are potentially the group most at risk as they may play inside or around buildings where baits have been placed. Infants could be exposed orally by chewing bait or touching their mouth with contaminated fingers. Two Tiers have been developed: Tier 1, considering that 5 g of the bait is ingested by children and Tier 2 (more realistic) where the presence of the biter agent in the product is taken in account and hence, 10 mg of the product is assumed as ingested by children.

Neither of these scenarios (3 and 4 ) is considered to result in long-term exposure, whereas acute exposure may occur.

### Systemic effects

Task/ Scenario	Tier	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
Dermal contact with dead rodents / Scenario [3]	Tier 1 /no PPE	$2 \times 10^{-3}$	$2.3 \times 10^{-6}$	$1.74 \times 10^{-6}$	75.65	Yes
	Tier 2 /(gloves as PPE)			$1.74 \times 10^{-7}$	7.56	Yes
Ingestion of bait / Scenario [4]	Tier 1 /no biter agent			0.0145	$6.30 \times 10^5$	No
	Tier 2 /biter agent			$2.9 \times 10^{-5}$	1260.8	No

### Local effects

There is no need to consider local effects separately.

### Conclusion

Estimation of secondary exposure scenarios demonstrates that there is no undue risk for adults when touching dead rodents with unprotected hands.

These estimations of secondary exposure scenarios demonstrate that children are at risk by ingesting 5 g or 10 mg of pellets according to the estimations.

However, calculations are based on conservative assumptions which will likely overestimate actual exposure levels. Furthermore, baits are placed according to the risk mitigation measures proposed for anticoagulant rodenticides usually out of the reach of children in tamper-resistant bait stations or in

covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations for trained professional users.

Moreover, Bromadiolone 0.0029% w/w pellet bait baits contain a highly efficient bittering agent to prevent ingestion by children.

#### Risk for consumers via residues in food

Neither new data was not provided nor had new guidance to be taken into account for re-assessment.

Accordingly, the conclusion from the former assessment regarding risks for consumers via residues in food remains valid.

#### Risk characterisation from combined exposure to several active substances or substances of concern within a biocidal product<sup>1</sup>

There is no risk derived from a combined exposure because indirect exposure via the environment is considered negligible, the product is not intended to be mixed with other biocidal or non biocidal products and the product does not contain any other active substance of concern.

#### Summary of risk characterisation

Scenario number	Exposed group (e.g. trained professionals, professionals, general public)	Tier/PPE	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/AEL (%)	Acceptable (yes/no)
1.	Trained professional user	Tier 1	$1.2 \times 10^{-6}$	$2.23 \times 10^{-7}$	18	Yes
1.	Trained professional user	Tier 2	$1.2 \times 10^{-6}$	$2.23 \times 10^{-8}$	1	yes
1.	professional user	Tier 1	$1.2 \times 10^{-6}$	$1.77 \times 10^{-8}$	1.5	Yes
1.	professional user	Tier 2	$1.2 \times 10^{-6}$	$1.77 \times 10^{-9}$	0.15	Yes

Scenario number	Exposed group (e.g. trained professionals, professionals, general public)	Tier/PPE	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/AEL (%)	Acceptable (yes/no)
1.	General public (non- professional)	Tier 1	$2.3 \times 10^{-6}$	$1.77 \times 10^{-8}$	0.7	Yes
2	Trained professional user	Tier 1	$1.2 \times 10^{-6}$	$1.05 \times 10^{-7}$	8.8	Yes
2	Trained professional user	Tier 2	$1.2 \times 10^{-6}$	$1.05 \times 10^{-8}$	0.8	Yes
2	Professional user	Tier 1	$1.2 \times 10^{-6}$	$3.29 \times 10^{-8}$	2.75	Yes
2	Professional user	Tier 2	$1.2 \times 10^{-6}$	$3.29 \times 10^{-9}$	0.27	Yes
2.	General public (Non-professional)	No PPE	$2.3 \times 10^{-6}$	$3.29 \times 10^{-8}$	1.4	Yes
3.	General public (bystander)	Tier 1/ no PPE (unlikely)	$2.3 \times 10^{-6}$	$1.74 \times 10^{-6}$	76	Yes
3.	General public (bystander)	Tier 2/ PPE (gloves)	$2.3 \times 10^{-6}$	$1.74 \times 10^{-7}$	8	Yes
4.	General public (Children)	Tier 1 (without efficient bitter agent)	$2.3 \times 10^{-6}$	0.0145	$6.30 \times 10^5$	No
4.	General public (Children)	Tier 2 (with bitter agent)	$2.3 \times 10^{-6}$	$2.9 \times 10^{-5}$	1261	No

### 3.7 Risk assessment for animal health

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding animal health remains valid.

### 3.8 Risk assessment for the environment

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding the environment remains valid.

#### 3.8.1 Exposure assessment

##### General information

Assessed PT	PT 14
Assessed scenarios	Scenario 1: in and around buildings application, against brown rat. Scenario 2: waste dumps/landfills, against brown rat. Scenario 3: open areas
ESD(s) used	EUBEES 2 Emission Scenario Document for rodenticides.
Approach	A consumption based approach has been used as a suitable protective measure at the local level.
Distribution in the environment	
Groundwater simulation	No
Confidential Annexes	No
Life cycle steps assessed	
Remarks	It has been only evaluate the use of this product against rats since it is the worst case.

##### Emission estimation

###### Scenario [1]: in and around buildings

The worst-case application is for the rat. The scenario is for eradication on a farm. The scenario indicates 2 – 3 applications per year. Bait points for rats are set 5 – 10 m apart. For the purposes of aligning the scenario with human exposure, the scenario assesses exposure from use of 250 g of bait in each of the 10 bait points. The bait points are replenished 5 times in a 21-day programme. There is 1 % direct release of the bait to soil. The scenario presented by the applicant differs from the ESD worst case scenario only regarding the amount of bait in each station, i.e. 200 g instead of 250 g.

ESD worst case:

Input parameters for calculating the local emission
---

Input	Value	Unit	Remarks
Scenario: use in bait points, in and around buildings			
Amount of product used at each refill/application	250	g	
Fraction of active substance in Product	2.9E-03	%	
Area directly exposed to active Substance	0.09	m <sup>2</sup>	
Area indirectly exposed to active substance	550	m <sup>2</sup>	
Number of emission days per Year	21	days	
Number of application sites	10	-	
Number of refills per site	5	-	
Fraction of active substance released directly to soil	0.01	-	
Depth of exposed soil	10	cm	
Fraction of active substance metabolised	21	%	
Bulk density of soil	1.7E03	Kg <sub>wwt</sub> /m <sup>3</sup>	

Applicant's worst case:

Input parameters for calculating the local emission			
Input	Value	Unit	Remarks
Scenario: use in bait points, in and around buildings			
Amount of product used at each refill/application	200	g	
Fraction of active substance in Product	2.9E-03	%	
Area directly exposed to active Substance	0.09	m <sup>2</sup>	
Area indirectly exposed to active substance	550	m <sup>2</sup>	
Number of emission days per Year	21	days	
Number of application sites	10	-	
Number of refills per site	5	-	
Fraction of active substance released directly to soil	0.01	-	

Depth of exposed soil	10	cm	
Fraction of active substance metabolised	21	%	
Bulk density of soil	1.7E03	Kg <sub>wwt</sub> /m <sup>3</sup>	

#### Calculations for Scenario [1]

Calculations have been performed according to EUBEES, Emission document for biocides used as rodenticides

Direct release in the realistic worst case farm scenario based on bait in bait boxes has been calculated as following (equation 2 ESD):

ESD worst case

Parameter	Definition	Units	Value
Amount of product used at each refill/application	Q <sub>prod</sub>	g	250
Fraction of active substance in product	F <sub>C<sub>prod</sub></sub>	-	0,000029
Number of application sites	N <sub>sites</sub>	-	10
Number of refills per site	N <sub>refil</sub>	-	5
Fraction of active substance released directly to soil	F <sub>release, soil</sub>	-	0,01
<b>Local direct emission rate of active substance to soil from a campaign</b>	<b>E<sub>local</sub><sub>soil-campaing</sub> = (Q<sub>prod</sub> × F<sub>C<sub>prod</sub></sub> × N<sub>sites</sub> × F<sub>release, soil</sub>) (2)</b>	<b>g</b>	<b>0.0036</b>

Applicant's worst case

Parameter	Definition	Units	Value
Amount of product used at each refill/application	Q <sub>prod</sub>	g	200

Fraction of active substance in product	$F_{C_{prod}}$	-	0,000029
Number of application sites	$N_{sites}$	-	10
Number of refills per site	$N_{refil}$	-	5
Fraction of active substance released directly to soil	$F_{release, soil}$	-	0,01
<b>Local direct emission rate of active substance to soil from a campaign</b>	<b><math>E_{local_{soil-campaing}} = (Q_{prod} \times F_{C_{prod}} \times N_{sites} \times F_{release, soil})</math> (2)</b>	<b>g</b>	<b>0,0029</b>

The concentration in the soil around each bait box after direct release can be estimated by the equation (3) of the ESD for PT14:

ESD worst case

Parameter	Definition	Units	Value
Local direct emission rate of active substance to soil from a campaign	$E_{soil, D-campaing}$ (2)	g	0.0036
Area directly exposed to active substance	$AREA_{exposed-D}$	m <sup>2</sup>	0.09
Depth of exposed soil	$DEPTH_{SOIL}$	m	0.1
Number of application sites	$N_{sites}$	-	10
Density of exposed soil	$RHO_{soil}$	kg/m <sup>3</sup>	1700
<b>Local concentration in soil due to direct release after a campaign [mg/kg]</b>	<b><math>C_{local_{soil-D}} = (E_{local_{soil-D-campaign}} \times 10E3) / (AREA_{exposed-D} \times DEPTH_{soil} \times RHO_{soil} \times N_{sites})</math> (3)</b>	<b>mg/kg</b>	<b>0.024</b>

Applicant's worst case

Parameter	Definition	Units	Value
-----------	------------	-------	-------



Parameter	Definition	Units	Value
Local direct emission rate of active substance to soil from a campaign	$E_{\text{soil, D-campaing}}$ (2)	g	0.0029
Area directly exposed to active substance	$AREA_{\text{exposed-D}}$	m <sup>2</sup>	0.09
Depth of exposed soil	$DEPTH_{\text{SOIL}}$	m	0.1
Number of application sites	$N_{\text{sites}}$	-	10
Density of exposed soil	$RHO_{\text{soil}}$	kg/m <sup>3</sup>	1700
<b>Local concentration in soil due to direct release after a campaign [mg/kg]</b>	<b><math>C_{\text{local, soil-D}} = (E_{\text{local, soil-D-campaign}} \times 10E3) / (AREA_{\text{exposed-D}} \times DEPTH_{\text{soil}} \times RHO_{\text{soil}} \times N_{\text{sites}})</math> (3)</b>	<b>mg/kg</b>	<b>0.019</b>

The concentration in the soil around the bait box taking into account only disperses release can be estimated by the equation:

ESD worst case

Parameter	Definition	Units	Value
Amount of product used at each refill/application	$Q_{\text{prod}}$	g	250
Fraction of active substance in product	$F_{C_{\text{prod}}}$	-	0.000029
Number of application sites	$N_{\text{sites}}$	-	10
Number of refills per site	$N_{\text{refil}}$	-	5
Fraction released indirectly to soil	$F_{\text{release-ID, soil}}$		0.73
Fraction released directly to soil	$F_{\text{release, soil}}$		0.01
Area indirectly exposed to	$AREA_{\text{exposed-ID}}$	m <sup>2</sup>	550

Parameter	Definition	Units	Value
rodenticide			
Depth of exposed soil	DEPTH <sub>SOIL</sub>	m	0.1
Density of exposed soil	RHO <sub>soil</sub>	kg/m <sup>3</sup>	1700
<b>Concentration in soil due to indirect (disperse) release after a campaign</b>	<b>Clocal<sub>soil-ID</sub> = ((Q<sub>prod</sub> x F<sub>cprod</sub> x N<sub>sites</sub> x N<sub>refil</sub> x 10<sup>3</sup> x F<sub>release,ID soil</sub> x (1-F<sub>release,D soil</sub>)) / (AREA<sub>exposed-ID</sub> x DEPTH<sub>soil</sub> x RHO<sub>soil</sub> x N<sub>sites</sub>)) (4)</b>	<b>mg/kg</b>	<b>0.0028</b>

## Applicant's worst case

Parameter	Definition	Units	Value
Amount of product used at each refill/application	Q <sub>prod</sub>	g	200
Fraction of active substance in the product	F <sub>cprod</sub>	-	0.000029
Number of application sites	N <sub>sites</sub>	-	10
Number of refills per site	N <sub>refil</sub>	-	5
Fraction released indirectly to soil	F <sub>release-ID, soil</sub>		0.73
Fraction released directly to soil	F <sub>release, soil</sub>		0.01
Area indirectly exposed to rodenticide	AREA <sub>exposed-ID</sub>	m <sup>2</sup>	550
Depth of exposed soil	DEPTH <sub>SOIL</sub>	m	0.1
Density of exposed soil	RHO <sub>soil</sub>	kg/m <sup>3</sup>	1700
<b>Concentration in soil due to indirect (disperse) release after a campaign</b>	<b>Clocal<sub>soil-ID</sub> = ((Q<sub>prod</sub> x F<sub>cprod</sub> x N<sub>sites</sub> x N<sub>refil</sub> x 10<sup>3</sup> x F<sub>release,ID soil</sub> x (1-F<sub>release,D soil</sub>)) / (AREA<sub>exposed-ID</sub> x DEPTH<sub>soil</sub> x</b>	<b>mg/kg</b>	<b>0.00224</b>

	<b>RHOsoil x Nsites) (4)</b>		
--	------------------------------	--	--

Total soil concentrations around the bait boxes are the sum of the soil concentrations caused by direct and indirect pollution of the soil:

ESD worst case

<b>Total concentration immediately direct to the bait</b>	$C_{\text{local soil}} = C_{\text{local soil-D}} + C_{\text{local soil-ID}}$	<b>mg/kg</b>	<b>0.0265</b>
---	--	--------------	---------------

Applicant's worst case

<b>Total concentration immediately direct to the bait</b>	$C_{\text{local soil}} = C_{\text{local soil-D}} + C_{\text{local soil-ID}}$	<b>mg/kg</b>	<b>0.0212</b>
---	--	--------------	---------------

### Scenario [2]: waste dumps

This scenario covers control of rats and disposal of rats in waste dumps and landfills where the exposure is assumed to be higher than that described in the open area scenario. In some instances, applications of rodenticides to refuse dumps take place. Mostly the use is limited to occasions of population outbreaks of rats. Often the rodenticides are deployed around the perimeter of the dump, more than in the disposal area itself. The bait may be placed at regular places in special feeding stations in order to prevent other animals from eating the bait.

The worst-case application is for the rat. The scenario is for eradication on an open dump. The scenario indicates 7 applications per year, with 40 kg product per application. There is 90% release of the bait to soil and 365 emission days.

<b>Input parameters for calculating the local emission</b>			
<b>Input</b>	<b>Value</b>	<b>Unit</b>	<b>Remarks</b>
Scenario: use in landfills and dumps			
Amount of product used at each refill/application	40	Kg	
Fraction of active substance in product	29E-03	%	
Number of emission days for control at waste dumps	365	days	

Number of application	7	-	
Fraction of active substance released to soil	0.73	-	
Area exposed to rodenticide	10000	m <sup>2</sup>	
Depth of exposed soil	10	cm	
Bulk density of soil	1.7E03	Kg <sub>wwt</sub> /m <sup>3</sup>	

### Calculations for Scenario [2]

Calculation of E<sub>local soil</sub> (equation 17, ESD PT14)

Parameter	Definition	Units	Value
Amount of product used per application	Q <sub>prod</sub>	g	40
Fraction of active substance in product	F <sub>C<sub>prod</sub></sub>	-	0.000029
Number of application sites	N <sub>sites</sub>	-	7
Fraction of active substance released directly to soil	F <sub>release, soil</sub>	-	0.73
<b>Local direct emission of active substance to soil from a campaign</b>	<b>E<sub>local soil-campaing</sub> = Q<sub>prod</sub> x F<sub>C<sub>prod</sub></sub> x N<sub>sites</sub> x F<sub>release, soil</sub> (17)</b>	<b>kg</b>	<b>5.93E-03</b>

Calculation of C local soil (equation 18, ESD PT14)

Parameter	Definition	Units	Value
Local direct emission of active substance to soil from a campaign	E <sub>local soil, campaing</sub> (2)	kg/m <sup>3</sup>	5.93E-03
Area directly exposed to active substance	AREA <sub>exposed-D</sub>	m <sup>2</sup>	10000
Depth of exposed soil	DEPTH <sub>SOIL</sub>	M	0.1

Density of exposed soil	$RHO_{soil}$	kg/m <sup>3</sup>	1700
Local concentration in soil due to direct release after a campaign [mg/kg]	$C_{local\ soil-D} = (E_{local\ soil-D-campaign} \times 10E3) / (AREA_{exposed-D} \times DEPTH_{soil} \times RHO_{soil} \times N_{sites}) (18)$	mg/kg	0.0035

### Scenario 3: open areas

This scenario covers control of rats and water voles in open areas such as around farmland, parks and golf courses where the aim is to prevent “nuisance” from burrows or “soil heaps” or due to public hygiene reasons. Rodenticides are also used to reduce impacts on game rearing or outside food stores (potato/sugar beet clams).

The main release to the environment is expected when impregnated grain is applied into rat holes. By a spoon or a small shovel, the product is normally poured approximately 30 cm into the rat holes, depending on the slope and general accessibility of the hole. The treated holes are closed by a stone, a piece of board or similar immediately after the application to prevent unintended exposure of children or non-target organisms (e.g. birds, cats and dogs).

A typical initial dose for a rat hole is 100-200 g grain.hole-1; and normally application is repeated twice with an interval of 5-6 days. Inspection of the holes to assess the effect of the control action is usually carried out some 5-6 days after application of the poison and again with similar intervals if repeated applications are necessary.

Input parameters for calculating the local emission			
Input	Value	Unit	Remarks
Scenario: use in landfills and dumps			
Amount of product used at each Refilling in the control operation	200	Kg	
Fraction of active substance in product	2.9E-03	%	
Number of emission days for control at open areas	6	days	
Number of application	2	-	
Fraction of product released to soil during application	0.05	-	
Fraction of product	0.20	-	

released to soil during use			
Soil volume exposed soil around the hole	0.0085	m <sup>3</sup>	
Bulk density of soil	1.7E03	Kg <sub>wwt</sub> /m <sup>3</sup>	

### Calculations for Scenario [3]

#### Calculation of Elocal<sub>soil-campaign</sub> (equation 9, ESD PT14)

Parameter	Definition	Units	Value
Amount of product used at each refilling in the control operation	Q <sub>prod</sub>	g	200
Fraction of active substance in product	F <sub>cprod</sub>	-	0.000029
Number of application sites	N <sub>sites</sub>	-	1
Number of refills per site	N <sub>refil</sub>	-	2
Fraction of the product released to soil during application	F <sub>release, soil, appl</sub>	-	0.05
Fraction of product released to soil during use	F <sub>release, soil, use</sub>		0.2
<b>Local emission of active substance to soil during a campaign</b>	<b>Elocal<sub>soil-campaign</sub> = (Q<sub>prod</sub> X F<sub>cprod</sub> X N<sub>sites</sub> X N<sub>refil</sub> x (F<sub>release, soil, appl</sub> + F<sub>release, soil</sub>)) (9)</b>	<b>g</b>	<b>2.90E-03</b>

#### Calculation of Clocal<sub>soil-campaign</sub> (equation 10, ESD PT14)

Parameter	Definition	Units	Value
Local emission to soil from the episode	Elocal <sub>soil-campaign</sub>	g	5.00E-03
Soil volume exposed to rodenticide	Vsoil <sub>exposed</sub> (eq. 9a ESD)	m <sup>3</sup>	8.50E-03

Parameter	Definition	Units	Value
Density of wet exposed soil	$RHO_{soil}$	$kg/m^3$	1700
Local concentration in soil after a campaign	$C_{local} = \frac{E_{local\text{-campaign}}}{V_{soil\text{-exposed}} \times RHO_{soil}} \times 10^3$ (10)	mg/kg	2.01E-01

### Fate and distribution in exposed environmental compartments

Identification of relevant receiving compartments based on the exposure pathway									
	Fresh-water	Freshwater sediment	Sea-water	Seawater sediment	STP	Air	Soil	Ground-water	Other
Scenario 1	No	No	No	No	No	No	Yes	Yes	
Scenario 2	No	No	No	No	No	No	Yes	Yes	
Scenario 3	No	No	No	No	No	No	Yes	Yes	

### Calculated PEC values

The Predicted Environmental Concentrations for this emission scenario are calculated according TGD II.

Summary table on calculated PEC values <sup>1</sup>								
	PEC <sub>s</sub> TP	PEC <sub>water</sub>	PEC <sub>sed</sub>	PEC <sub>seawater</sub>	PEC <sub>seased</sub>	PEC <sub>soil</sub>	PEC <sub>GW</sub> <sup>2</sup>	PEC <sub>air</sub>
	[mg/l]	[mg/l]	[mg/kg <sub>ww</sub> ↓]	[mg/l]	[mg/kg <sub>wwt</sub> ]	[mg/kg]	[µg/l]	[mg/m <sup>3</sup> ]
Scenario 1	-	-	-			0.026	1.01x10 <sup>-1</sup>	
Scenario 2	-	-	-			0.0035	1.33x10 <sup>-2</sup>	
Scenario 3						0.201	0.7	

### Primary and secondary poisoning

Non-target vertebrates may be exposed to bromadiolone either directly by ingestion of exposed product (primary poisoning) or indirectly by ingestion of the carcasses of target rodents that contain residues of bromadiolone (secondary poisoning).

Assessment of secondary poisoning through the aquatic food chain is not performed for the following reasons: the risk assessment for the aquatic compartment indicates that there will be very low concentrations of bromadiolone in the aquatic compartment, and there was no risk identified of bromadiolone for surface water or sediment dwelling organisms. The justification for not performing an assessment of secondary poisoning via the terrestrial food chain is that secondary poisoning will be limited due to the small area that potentially is contaminated by bromadiolone around buildings and the limited number of earthworms inhabiting this area.

Due to the highly toxic nature of bromadiolone, primary and secondary poisoning presents a hazard to non target mammals and birds following use in and around buildings. The risk assessment of bromadiolone used in and around buildings is summarised by presenting PEC/PNEC ratios for long-term primary and secondary poisoning. The risks posed by use in open areas and on waste dumps can be considered as adequately covered by the same assessment.

For the acute situation, as was agreed at TMIII-06, PNEC derivation for birds and mammals will only apply to long-term effects and acute effects will only be evaluated on a qualitative basis. It is important to stress that this qualitative assessment is not intended to be used for the risk characterisation of primary and secondary poisoning of rodenticides and shall not be used for a comparative assessment. This comparison should only give a first indication of the acute toxicity of the substance.

#### Primary poisoning

##### Tier 1

	<b>PEC (conc. in food, mg/kg)</b>
Birds	29
Mammals	29

**Tier 2** (for bait containing bromadiolone in and around buildings, step 2 (realistic worst case).

<b>Non-target animal</b>	<b>PEC<sub>oral</sub> = ETE, conc. of bromadiolone after one meal (mg/kg)</b>	<b>LD<sub>50</sub> dose (mg/kg bw/d)</b>	<b>PEC<sub>oral</sub> higher than LD<sub>50</sub> (y/n)</b>
Dog	0.92	1.3	n (TF)
Pig	0.16	1.3	n (TF)
Pig, young	0.50	1.3	n (TF)
Tree sparrow	7.25	134	n (TF)
Chaffinch	6.26	134	n (TF)
Wood pigeon	2.26	134	n (TF)
Pheasant	2.25	134	n (TF)



This comparison indicates that birds, pigs and mammals are not at risk for acute primary poisoning.

**Tier 2 long-term risk assessment** for bait containing bromadiolone in and around buildings. Very high risks for long-term primary poisoning of both mammals and birds are identified. However, long-term consumption of these quantities of bromadiolone bait is generally not realistic and should be regarded strictly as worst case.

Non-target animal	PEC = EC, concentration of bromadiolone after one day of elimination (mg/kg)
Dog	1.15
Pig	0.19
Pig, young	0.60
Tree sparrow	8.71
Chaffinch	7.56
Wood pigeon	2.73
Pheasant	2.72

#### Secondary poisoning

**The tier 1 qualitative acute risk assessment** of secondary poisoning based on measured residue levels (presented by the applicant) in target rodents indicates no risk for birds or mammals. However, this qualitative assessment is only an indication and is not intended to be used for the risk characterisation of secondary poisoning of rodenticides.

**The tier 1 long-term risk assessment** based on default (Task Force) residue levels in target rodents results in very high PEC/PNEC values for predatory birds and mammals.

	PNEC <sub>oral</sub> (conc. in food)	PEC <sub>oral</sub> Bromadiolone conc. in target rodent (mg/kg bw), ESD default values
Birds	0.0087 mg/L	13.9
Mammals	0.00019 mg/kg	13.9

### 3.8.2 Risk characterisation

#### **Atmosphere**

Emission to the atmosphere from this use is considered negligible.

**Terrestrial compartment**

Calculated PEC/PNEC values	
	PEC/PNEC <sub>soil</sub>
Scenario 1	0.26
Scenario 2	0.035
Scenario 3	2.03

Conclusion:

Scenario 1 and 2: present ratios of PEC/ PNEC less than 1 so, an acceptable level of risk to soil are predicted from those scenarios. For open areas, the PEC/PNEC ratio is above 1.0 indicating that there are unacceptable risks to the terrestrial compartment when this product is used in the tunnels of open areas. However, the PEC/PNEC ratios calculated indicate a marginal risk based on the PEC that represents a localised “hotspot” of contamination near the entrance of each baited tunnel. According to the EUBEES 2 scenario, the use near the openings of the tunnels is covered by the assessment of the scenario “in and around buildings” with bait box. So, there is no unacceptable risk for the terrestrial compartment (including groundwater) when this product is used near the openings of the tunnels of the target rodents.

**Groundwater**

For scenario open areas the value obtained for ground water is higher than 0.1 µg/L but, considering the localised treated area in the tunnels, the risk for groundwater was not considered relevant

**Primary and secondary poisoning**

Non-target vertebrates may be exposed to bromadiolone either directly by ingestion of exposed product (primary poisoning) or indirectly by ingestion of the carcasses of target rodents that contain residues of bromadiolone (secondary poisoning).

Assessment of secondary poisoning through the aquatic food chain is not performed for the following reasons: the risk assessment for the aquatic compartment indicates that there will be very low concentrations of bromadiolone in the aquatic compartment, and there was no risk identified of bromadiolone for surface water or sediment dwelling organisms. The justification for not performing an assessment of secondary poisoning via the terrestrial food chain is that secondary poisoning will be limited due to the small area that potentially is contaminated by bromadiolone around buildings and the limited number of earthworms inhabiting this area.

Due to the highly toxic nature of bromadiolone, primary and secondary poisoning presents a hazard to non target mammals and birds following use in and around buildings. The risk assessment of bromadiolone used in and around buildings is summarised by presenting PEC/PNEC ratios for long-term primary and secondary poisoning. The risks posed by use in open areas and on waste dumps can be considered as adequately covered by the same assessment.

For the acute situation, as was agreed at TMIII-06, PNEC derivation for birds and mammals will only apply to long-term effects and acute effects will only be evaluated on a qualitative basis. It is important to stress that this qualitative assessment is not intended to be used for the risk characterisation of primary and secondary poisoning of rodenticides and shall not be used for a comparative assessment. This comparison should only give a first indication of the acute toxicity of the substance.

### **Primary poisoning**

In the **Tier 1 assessment** of primary poisoning it is assumed that the whole day's food requirement is satisfied by consumption of wax blocks, and therefore the concentration in food will be the same as the concentration of a.s. in the bait, 50 mg/kg. This is then compared to the long-term PNECs for birds and mammals. The resulting PEC/PNEC ratios in the table below reveal a high risk for both birds and mammals of long-term primary poisoning.

**Table 2.8.4.5-1 PEC/PNEC ratios for primary poisoning – Tier 1 assessment**

	<b>PEC (conc. in food, mg/kg)</b>	<b>PNEC (conc. in food)</b>	<b>PEC/PNEC</b>
Birds	29	0.0087 mg/l	3333
Mammals	29	0.00019 mg/kg	152632

**Tier 2 acute qualitative risk assessment** for bait containing bromadiolone in and around buildings, step 2 (realistic worst case).

Table 2.8.4.5-2 PEC values calculated for birds and mammals

Non-target animal	PEC <sub>oral</sub> = ETE, conc. of bromadiolone after one meal (mg/kg)	LD <sub>50</sub> dose (mg/kg bw/d)	PEC <sub>oral</sub> higher than LD <sub>50</sub> (y/n)
Dog	0.95	1.3	N
Pig	0.16	1.3	N
Pig, young	0.50	1.3	N
Tree sparrow	7.25	134	N
Chaffinch	6.264	134	N
Wood pigeon	2.26	134	N
Pheasant	2.25	134	N

This comparison indicates that birds and mammals are not at risk for acute primary poisoning.

**Tier 2 long-term risk assessment** for bait containing bromadiolone in and around buildings. Very high risks for long-term primary poisoning of both mammals and birds are identified. However, long-term consumption of these quantities of bromadiolone bait is generally not realistic and should be regarded strictly as worst case.

Table 2.8.4.5-3 PEC/PNEC ratios for primary poisoning - Tier 2 assessment long term

Non-target animal	PEC = EC, concentration of bromadiolone after one day of elimination (mg/kg)	PNEC dose (mg/kg bw/day)	PEC/PNEC
Dog	0.95	0.0000056	169643
Pig	0.16	0.0000056	28571
Pig, young	0.50	0.0000056	89286
Tree sparrow	7.25	0.0013	557692
Chaffinch	6.264	0.0013	4818
Wood pigeon	2.26	0.0013	1738
Pheasant	2.25	0.0013	1731

### Secondary poisoning

The **tier 1 qualitative acute risk assessment** of secondary poisoning based on measured residue levels (presented by the applicant) in target rodents indicates no risk for birds or mammals. However,

this qualitative assessment is only an indication and is not intended to be used for the risk characterisation of secondary poisoning of rodenticides.

The **tier 1 long-term risk assessment** based on default residue levels in target rodents results in very high PEC/PNEC values for predatory birds and mammals.

**Table 2.8.4.5-4 PEC/PNEC ratios for secondary poisoning – Tier 1 assessment**

	<b>PNEC<sub>oral</sub></b> <b>(conc. In food)</b>	<b>PEC<sub>oral</sub></b> <b>Bromadiolone conc. in target rodent</b> <b>(mg/kg bw), ESD default values</b>	<b>PEC/PNEC</b>
Birds	0.0087 mg/l	13.9	1600
Mammals	0.00019 mg/kg	13.9	73200

### **Conclusion**

Although the quantity of active substance has been reduced the quantitative risk assessments is that there are still, in some cases, very high unacceptable risks to non-target vertebrates via primary and secondary poisoning.

To minimise the likelihood of target rodents developing resistance to second-generation anticoagulant rodenticides, long-term deployment of baits as a preventative control measure is not recommended. Product labels additionally instruct users to retrieve and securely dispose of all unconsumed baits at the end of control programmes. Both these factors limit the opportunity for exposure and reduce the primary poisoning risk to small non-target animals. Provided that baits are deployed in accordance with the product labelling and other approved guidance on good practice, the primary poisoning risk to non-target mammals may be considered to be negligible.

The risk of secondary poisoning of bromadiolone to birds and small mammals is expected to be significantly reduced by restricting its use to treatment campaigns of limited duration, limiting access of non-target animals to the blocks and removing dead and moribund rodents during a baiting campaign to minimise the opportunity secondary exposure. These mitigation measures are described in good practice guidance documents, in training material for pest control professionals and on the labels of the products. Also, with the aim of harmonising the assessments of second generation anticoagulant rodenticides, a common approach to the use of risk mitigation measures has been agreed at the CA meeting in Nov 2016.

### **3.9 Assessment of a combination of biocidal products**

A use with other biocidal products is not intended.

### **3.10 Comparative assessment**

As bromadiolone is a Candidate for Substitution, a comparative assessment must be carried out as part of the evaluation process.

The Biocidal Products Committee of the European Chemicals Agency published its Opinion on Questions regarding the comparative assessment of anticoagulant rodenticides on 02 March 2017 (Document no. ECHA/BPC/145/2017).

The Opinion states that:

- In the absence of anticoagulant rodenticides, the use of rodenticide biocidal products containing other active substances would lead to an inadequate chemical diversity to minimize the occurrence of resistance in the target harmful organisms. These products also show some significant practical or economical disadvantages for the relevant uses.
- There is insufficient scientific evidence to prove that non-chemical alternative methods of rodent control are sufficiently effective according to the criteria established in agreed Union guidance with a view to prohibit or restrict the authorised uses of anticoagulant rodenticides.

The Opinion forms the basis of the COMMISSION IMPLEMENTING DECISION (EU) 2017/1532 of 7 September 2017 addressing questions regarding the comparative assessment of anticoagulant rodenticides in accordance with Article 23(5) of Regulation (EU) No 528/2012 of the European Parliament and of the Council.

On the basis of this comparative assessment, the authorisation of rodenticide products containing bromadiolone is justified.