

EN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

NEO-ACTIBLOCK BROM

Product type(s)

PT14: Rodenticides

Authorisation number: 2021-10-28-B02

R4BP asset number: MT-0027305-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	NEO-ACTIBLOCK BROM
---------------	--------------------

1.2. Authorisation holder

Name and address of the authorisation holder	Name	Activa s.r.l.
	Address	via Feltre, 32 20132 Milano Italy
Authorisation number	2021-10-28-B02	
<i>R4BP asset number</i>	MT-0027305-0000	
Date of the authorisation	14/12/2012	
Expiry date of the authorisation	31/12/2025	

1.3. Manufacturer(s) of the product

Name of manufacturer	ACTIVA S.r.l.
Address of manufacturer	Via Feltre 32 20132 Milano Italy
Location of manufacturing sites	I.N.D.I.A. Industrie Chimiche S.p.a. - Nona Strada 57 35020 Padova Italy Vebi Istituto Biochimico S.r.l. - Via Desman 43 35010 Borgoricco (PD) Italy

1.4. Manufacturer(s) of the active substance(s)

Active substance	Bromadiolone
Name of manufacturer	Activa S.r.l.
Address of manufacturer	Via Feltre 32 20132 Milano Italy
Location of manufacturing sites	Dr. Tezza s.r.l. - Via Tre Ponti 22 37059 S.Maria di Zevio (VR) Italy

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[3-(4'-Bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	active substance	28772-56-7	249-205-9	0,0027
Denatonium benzoate	benzyl-[2-(2,6-dimethylanilino)-2-oxoethyl]-diethylazanium; benzoate	Non-active substance	3734-33-6	223-095-2	0,001
Calcium hydroxide	calcium; dihydroxide	Non-active substance	1305-62-0	215-137-3	0,375

2.2. Type(s) of formulation

RB Bait (ready for use)

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H373: May cause damage to organs (blood) through prolonged or repeated exposure { 1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard: }.
Precautionary statements	P102: Keep out of reach of children. P308+P313: IF exposed or concerned: Get medical advice. P308+P313: IF exposed or concerned: Get medical attention. P501: Dispose of contents to national and international regulations. P501: Dispose of container to national and international regulations.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. House mice – professionals – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: -
Field(s) of use	indoor use -
Application method(s)	Method: Ready-to-use bait Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 60 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): 0 Number and timing of application: -
Category(ies) of users	professional
Pack sizes and packaging material	Labelled or printed bag/sack in PE/Aluminium/Polyester (thermal welded): from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg. Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed HDPE bucket (sealed): from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg. Bag/sack in PE/Aluminium/Polyester (thermal welded) in printed carton sachet or in printed cardboard: from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate-packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg.

4.1.1. Use-specific instructions

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.1.2. Use-specific risk mitigation measures

4.1.3. Use-specific risk mitigation measures

See “General directions for use (professionals)”.

4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See “General directions for use (professionals)”.

4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See “General directions for use (professionals)”.

4.2. Use description

Table 2. Rats – professionals – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.

Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: -
Field(s) of use	indoor use -
Application method(s)	Method: Ready-to-use bait Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): - Number and timing of application: -
Category(ies) of users	professional
Pack sizes and packaging material	Labelled or printed bag/sack in PE/Aluminium/Polyester (thermal welded): from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg. Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed HDPE bucket (sealed): from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg. Bag/sack in PE/Aluminium/Polyester (thermal welded) in printed carton sachet or in printed cardboard: from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate-packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg.

4.2.1. Use-specific instructions

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

See “General directions for use (professionals)”.

4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See “General directions for use (professionals)”.

4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See “General directions for use (professionals)”.

4.3. Use description

Table 3. House mice and/or rats – professionals – outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: - Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: -
Field(s) of use	outdoor use Outdoor around buildings
Application method(s)	Method: Ready-to-use bait Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: House mouse: 60 g of bait per bait station. Brown rat: 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.

	<p>Dilution (%): -</p> <p>Number and timing of application: -</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>Labelled or printed bag/sack in PE/Aluminium/Polyester (thermal welded): from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg.</p> <p>Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed HDPE bucket (sealed): from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg.</p> <p>Bag/sack in PE/Aluminium/Polyester (thermal welded) in printed carton sachet or in printed cardboard: from 1.5 kg up to 25 kg (containing rodenticide baits of 10-20 g in wax blocks). Packaging is restricted to separate packed bags in PE/Aluminium/Polyester with a maximum packaging size of 10 kg.</p>

4.3.1. Use-specific instructions

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or drainage systems, ensure that bait contact with water is avoided.

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See “General directions for use (professionals)”.

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See “General directions for use (professionals)”.

4.4. Use description

Table 4. House mice – general public – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: -
Field(s) of use	indoor use -
Application method(s)	Method: Ready-to-use bait Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 60 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): - Number and timing of application: -
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Labelled or printed bag/sack in PE/Aluminium/Polyester (thermal welded): from 10 g up to 100 g (containing rodenticide baits of 10-20 g in wax blocks).

	<p>Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed HDPE bucket (sealed): 10 g up to 100 g (containing rodenticide baits of 10-20 g in wax blocks).</p> <p>Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed carton sachet or in labelled or printed cardboard: from 10 g up to 100 g (containing rodenticide baits of 10-20 g in wax blocks).</p>
--	---

4.4.1. Use-specific instructions

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.4.2. Use-specific risk mitigation measures

4.4.3. Use-specific risk mitigation measures

See “General directions for use (general public)”.

4.4.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See “General directions for use (general public)”.

4.4.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See “General directions for use (general public)”.

4.4.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See “General directions for use (general public)”.

4.5. Use description

Table 5. Rats – general public – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: -
Field(s) of use	indoor use -
Application method(s)	Method: Ready-to-use bait Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): - Number and timing of application: -
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Labelled or printed bag/sack in PE/Aluminium/Polyester (thermal welded): from 100 g up to 300 g (containing rodenticide baits of 10-20 g in wax blocks). Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed HDPE bucket (sealed): from 100 g up to 300 g (containing rodenticide baits of 10-20 g in wax blocks). Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed carton sachet or in labelled or printed cardboard: from 100 g up to 300 g (containing rodenticide baits of 10-20 g in wax blocks).

4.5.1. Use-specific instructions

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.5.2. Use-specific risk mitigation measures

4.5.3. Use-specific risk mitigation measures

See “General directions for use (general public)”.

4.5.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See “General directions for use (general public)”.

4.5.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See “General directions for use (general public)”.

4.5.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See “General directions for use (general public)”.

4.6. Use description

Table 6. Rats - general public - outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: -

Field(s) of use	<p>outdoor use</p> <p>Outdoor around buildings</p>
Application method(s)	<p>Method: Ready-to-use bait</p> <p>Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.</p>
Application rate(s) and frequency	<p>Application rate: 100 g wax block bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters</p> <p>Dilution (%): -</p> <p>Number and timing of application: -</p>
Category(ies) of users	<p>general public (non-professional)</p>
Pack sizes and packaging material	<p>Labelled or printed bag/sack in PE/Aluminium/Polyester (thermal welded): from 100 g up to 300 g (containing rodenticide baits of 10-20 g in wax blocks).</p> <p>Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed HDPE bucket (sealed): from 100 g up to 300 g (containing rodenticide baits of 10-20 g in wax blocks).</p> <p>Bag/sack in PE/Aluminium/Polyester (thermal welded) in labelled or printed carton sachet or in labelled or printed cardboard: from 100 g up to 300 g (containing rodenticide baits of 10-20 g in wax blocks).</p>

4.6.1. Use-specific instructions

- Place the bait stations in areas not liable to flooding.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.6.2. Use-specific risk mitigation measures

4.6.3. Use-specific risk mitigation measures

See “General directions for use (general public)”.

4.6.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See “General directions for use (general public)”.

4.6.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See “General directions for use (general public)”.

4.6.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See “General directions for use (general public)”.

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

Professionals

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section “Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment” for the information to be shown on the label).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves (rubber) during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.

General Public

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.
- Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Do not place bait stations near water drainage systems where they can come into contact with water.

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

-
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
 - Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

Professionals

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
 - the product shall not be supplied to the general public (e.g. "for professionals only").
 - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

General Public

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).
- The product information (i.e. label and/or leaflet) shall clearly show that:
 - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.
- Dispose dead rodents in accordance with local requirements.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Professionals

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.

Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre or 118.

- Hazardous to wildlife.

General Public

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:

Dermal exposure, wash skin with water and then with water and soap.

Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label [insert country specific information]. Contact a veterinary surgeon in case of ingestion by a pet [insert country specific information].

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [insert national phone number]".

- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

Professionals

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.

General Public

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Professionals

- Store in a dry, cool and well ventilated place. Do not store at temperature higher than 35°C. Keep the container closed and away from direct sunlight.

- Store in places prevented from the access of children, birds, pets and farm animals.

- Shelf life: 2 years.

General Public

- Store in a dry, cool and well ventilated place. Do not store at temperature higher than 35°C. Keep the container closed and away from direct sunlight.

- Store in places prevented from the access of children, birds, pets and farm animals.

- Shelf life: 2 years.

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.