BORASOL 3% lotion 100 g

- Borasol lotion contains boric acid which is used for treatment of skin inflammation and exhibits antibacterial and antifungal properties. Borasol also has astringent and drying effect.

Composition:
100 g of the lotion contains 3 g of boric acid.

Action:
The preparation has bacteriostatic and fungistatic properties. It has mild antiseptic effects and exhibits astringent and drying properties. It absorbs through the mucous membranes and damaged skin.

Indications:
Inflammation of the skin and external urogenital organs, eczema, burns, bruising, swelling, superficial epidermal injuries.

Contraindications:
Hypersensitivity to boric acid. Do not use in infants and children younger than 11. Do not apply to extensive wounds and open cuts. Avoid long-term use. Do not combine with colloidal silver solutions, methenamine or balsam of Peru.

Adverse effects:
Long-term use on damaged skin or mucous membranes may cause a rash, skin reddening, epidermal desquamation and toxic systemic effects (particularly in infants and young children), the symptoms of which are: vomiting, diarrhoea, convulsions and damage to the internal organs (liver, kidneys).

Interactions:
It shows physicochemical incompatibility with colloidal silver solutions, methenamine or balsam of Peru.

Dosage:
Apply externally; adults: wash or rinse the skin 2–3 times daily or use as a gauze swab dripped with the solution.