

Summary of product characteristics for a biocidal product

Product name: GMP 48

Product type(s): PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

Authorisation number: 2019-07-31-B03

R4BP 3 asset reference number: MT-0013584-0008

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	2
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	3
3. Hazard and precautionary statements	3
4. Authorised use(s)	3
5. General directions for use	8
5.1. Instructions for use	8
5.2. Risk mitigation measures	9
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	9
5.4. Instructions for safe disposal of the product and its packaging	9
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	9
6. Other information	9

Administrative information

1.1. Trade names of the product

Iobac Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc
Dipal Conc

1.2. Authorisation holder

**Name and address of the
authorisation holder**

Name	DeLaval NV
Address	Industriepark-Drongen 10 9031 Gent Belgium

Authorisation number

2019-07-31-B03 1-5

**R4BP 3 asset reference
number**

MT-0013584-0008

Date of the authorisation

02/08/2019

Expiry date of the authorisation

01/08/2029

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer

DeLaval NV

Address of the manufacturer

Industriepark-Drongen 10 9031 Gent Belgium

Location of manufacturing sites

Industriepark-Drongen 10 9031 Gent Belgium

1.4. Manufacturer(s) of the active substance(s)

Active substance

1319 - Iodine

Name of the manufacturer

Sociedad Quimica y Minera (SQM) S.A.

Address of the manufacturer

Supplier EU: SQM Europe NV - Sint-Pietersvliet 7 bus 8 2000 Antwerpen Belgium

Location of manufacturing sites

Los Militares 4290 Piso 4 8320000 Las Condes (Santiago) Chile

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Iodine		Active Substance	7553-56-2	231-442-4	0,75
Alcohol ethoxylate		Non-active substance	68439-46-3		6,34
Sodium iodide		Non-active substance	7681-82-5	231-679-3	0,33

2.2. Type of formulation

SL - Soluble concentrate

3. Hazard and precautionary statements

Hazard statements

Causes serious eye irritation.
Harmful to aquatic life with long lasting effects.
May cause damage to organs thyroid through prolonged or repeated exposure oral route.

Precautionary statements

Keep out of reach of children.
Avoid release to the environment.
Wear eye protection.
Wear face protection.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Wash hands thoroughly after handling.
If eye irritation persists: Get medical advice.
If eye irritation persists: Get medical attention.

4. Authorised use(s)

4.1 Use description

Use 1 - Teat disinfection; concentrated dip products applied manually

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

Teat disinfectant

Target organism(s) (including development stage)

Scientific name:
Common name: Bacteria
Development stage:

	<p>Scientific name: Common name: Yeasts Development stage:</p> <p>Scientific name: Common name: Viruses Development stage:</p>
Field(s) of use	<p>Indoor</p> <p>The product is intended for post-milking teat disinfection. The product is applied directly after milking on the teats of lactating animals.</p>
Application method(s)	<p>Method: Post-milking disinfection of udder teats with diluted concentrate Detailed description: The product is applied manually after milking on the teats by manual dipping using a dip cup</p>
Application rate(s) and frequencies	<p>Application Rate: ca. 5 ml per cow; ca. 2.5 ml per sheep or goat Dilution (%): 20%: Use concentration: 0.15% w/w iodine Number and timing of application: To be applied after milking, 2 times per day. The products can be used all year during the lactation period.</p>
Category(ies) of users	<p>Professional</p>
Pack sizes and packaging material	<p>Can, Plastic: HDPE , 5L Can, Plastic: HDPE , 10L Can, Plastic: HDPE , 20L Can, Plastic: HDPE , 60L Drum, Plastic: HDPE , 200L IBC, Plastic: HDPE , 1000L Bottle, Plastic, HDPE, 1L</p>

4.1.1 Use-specific instructions for use

<p>Directions for use: It is recommended to wear gloves during milking and application to protect the skin and ensure hygienic milking condition. A 20% dilution should be made with clean water. Take 200 ml of the concentrated product and make up to 1L with water to obtain an end concentration of 0.15% iodine. The use solution must be brought to a temperature above 20°C before use. It is possible that foaming occurs while preparing the ready-to-use solution. In that case it is strongly recommended to only start the treatment after the foam level has sufficiently decreased. For manual application: use a clean dip cup. Ensure that teats are clean before use. Apply the diluted product on each teat immediately after milking: DIP: dip every teat in the product. Ensure that the teat is covered to at least three quarters length and replenish the dip cup as necessary; Allow teats to dry after application during at least 5 min and keep the animals standing during this period.</p>
--

4.1.2 Use-specific risk mitigation measures

Use eye/face protection during post-milking teat disinfection by manual dipping.
See 5.2 for general instructions.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

There are no use specific instructions. See 5.3 for general instructions.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

There are no use specific instructions. See 5.4 for general instructions.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.2 Use description

Use 2 - Teat disinfection; concentrated spray products applied manually

Product type

PT03 - Veterinary hygiene (Disinfectants)

Where relevant, an exact description of the authorised use

Teat disinfectant

Target organism(s) (including development stage)

Scientific name:
Common name: Bacteria
Development stage:

Scientific name:
Common name: Yeasts
Development stage:

Scientific name:
Common name: Viruses
Development stage:

Field(s) of use

Indoor

The product is intended for post-milking teat disinfection. The product is applied directly after milking on the teats of lactating animals.

Application method(s)

Method: Post-milking disinfection of udder teats with diluted concentrate
Detailed description:

	The product is applied after manually milking on the teats by spraying using a spray container.
Application rate(s) and frequencies	Application Rate: ca 5 ml per cow; ca. 2.5 ml per sheep or goat Dilution (%): 20%: Use concentration: 0.15% w/w iodine Number and timing of application: To be applied after milking 2 times per day. The product can be used all year during the lactation period.
Category(ies) of users	Professional
Pack sizes and packaging material	Can, Plastic: HDPE, 5L Can, Plastic: HDPE, 10L Can, Plastic: HDPE, 20L Can, Plastic: HDPE, 60L Drum, Plastic: HDPE, 200L IBC, Plastic: HDPE, 1000L Bottle, Plastic: HDPE, 1L

4.2.1 Use-specific instructions for use

<p>Directions for use: A 20% dilution should be made with clean water. Take 200 ml of the concentrated product and make up to 1L with water to obtain an end concentration of 0.15% iodine. The use solution must be brought to a temperature above 20°C before use.</p> <p>It is possible that foaming occurs while preparing the ready-to-use solution. In that case it is strongly recommended to only start the treatment after the foam level has sufficiently decreased. For manual application: use a clean applicator. Ensure that teats are clean before use. Apply the diluted product on each teat immediately after milking: 1) MANUAL TRIGGER SPRAY: spray every teat 1 sec. with the product. Ensure that the teat is covered to at least three quarters length and replenish the spray container as necessary; 2) AUTOMATIC SPRAYER: spray every teat 1 sec. with the product. Allow teats to dry after application during at least 5 min and keep the animals standing during this period.</p>
--

4.2.2 Use-specific risk mitigation measures

Use chemical resistant gloves (glove material to be specified by the authorisation holder within the product information) and eye/face protection during post-milking teat disinfection by manual spraying. See 5.2 for general instructions.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

There are no use specific instructions. See 5.3 for general instructions.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

There are no use specific instructions. See 5.4 for general instructions.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.3 Use description

Use 3 - Teat disinfection, concentrated dip or spray products applied automatically

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	Teat disinfectant
Target organism(s) (including development stage)	Scientific name: Common name: Bacteria Development stage: Scientific name: Common name: Yeasts Development stage: Scientific name: Common name: Viruses Development stage:
Field(s) of use	Indoor The product is intended for post-milking teat disinfection. The product is applied directly after milking on the teats of lactating animals.
Application method(s)	Method: Post-milking disinfection of udder teats with diluted concentrate Detailed description: The product is applied automatically by milking machines after milking on the teats by integrated dipping or spraying device.
Application rate(s) and frequencies	Application Rate: ca. 5 ml per cow; ca. 2.5 ml per sheep or goat Dilution (%): 20%: Use concentration: 0.15% w/w iodine Number and timing of application: To be applied after milking, 3 times per day. The products can be used all year during the lactation period.
Category(ies) of users	Professional
Pack sizes and packaging material	Can, Plastic: HDPE , 5L Can, Plastic: HDPE , 10L Can, Plastic: HDPE , 20L Can, Plastic: HDPE , 60L Drum, Plastic: HDPE , 200L

IBC, Plastic: HDPE , 1000L
Bottle, Plastic, HDPE, 1L

4.3.1 Use-specific instructions for use

Directions for use:

A 20% dilution should be made with clean water. Take 200 ml of the concentrated product and make up to 1L with water to obtain an end concentration of 0.15% iodine.

The use solution must be brought to a temperature above 20°C before use.

It is possible that foaming occurs while preparing the ready-to-use solution. In that case it is strongly recommended to only start the treatment after the foam level has sufficiently decreased.

4.3.2 Use-specific risk mitigation measures

Use eye/face protection during post-milking teat disinfection by automated dipping or spraying.
See 5.2 for general instructions.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

There are no use specific instructions. See 5.3 for general instructions.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

There are no use specific instructions. See 5.4 for general instructions.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. General directions for use

5.1. Instructions for use

There are no general instructions for use. See section 4 for the use specific instructions for use.

5.2. Risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

DIRECT/INDIRECT EFFECTS

According to our experience and to the information provided to us, the product does not have any adverse effects if it is used and handled as specified.

FIRST AID MEASURES

Eye contact: Rinse thoroughly with plenty of water for at least 15 minutes. If eye irritation persists: Get medical advice/attention.

Skin contact: Wash off immediately with soap and plenty of water removing all contaminated clothes and shoes.

Ingestion: Wash out mouth with water. Seek medical advice immediately if symptoms occur and/or large quantities have been ingested.

Inhalation: Move to fresh air.

EMERGENCY MEASURES TO PROTECT THE ENVIRONMENT

Prevent further leakage or spillage if safe to do so.

Methods and materials for containment and cleanup: Dam up. Soak up with inert absorbent material. Prevent product from entering drains. Keep in suitable and closed containers for disposal.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose unused product and the packaging in accordance with local requirements. Used product can be flushed to the municipal sewer or disposed to the manure deposit depending on local requirements. Avoid release to an individual waste water treatment plant.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Storage: Store upright in the tightly closed original container. Protect from direct light, temperatures above 30°C and frost.

Shelf-life: 2 years

6. Other information