ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

NYNA D+ BLOC P

Product type(s)

PT14: Rodenticides

Authorisation number: FR-2013-0105

R4BP asset number: FR-0002632-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	NYNA D+ BLOC P
	SUPER FULGURAT B
	BLOC SD 50
	DIFEBLOC RAT 50
	DIFEBLOC RONGEUR 50
	STOP DIFEBLOC 50
	SUPP' DIFEBLOC 50
	RÉGAL
	RÉGAL'EXKI
	FUGIS
	FUGIS XP
	I

1.2. Authorisation holder

	Name	TRIPLAN SA
Name and address of the authorisation holder	Address	BP 258 LA POSTE FRANCAISE AD500 ANDORRA LA VELLA France
Authorisation number		FR-2013-0105
R4BP asset number		FR-0002632-0000
Date of the authorisation		10/10/2013
Expiry date of the authorisation		31/12/2026

1.3. Manufacturer(s) of the product

Name of manufacturer	AEDES PROTECTA
	75 RUE D'ORGEMONT 95210 SAINT- GRATIEN France
Location of manufacturing sites	LIEU DIT DOUILLAC 81310 PARISOT France

Name of manufacturer	HDA
Address of manufacturer	ZA LA CHARME MENETROL 63200 RIOM France
Location of manufacturing sites	ZA LA CHARME MENETROL 63200 RIOM France

Name of manufacturer	EDIALUX
Address of manufacturer	ZA MACON EST 01750 REPLONGES France
1	EDIALUX BRETAGNE - ZA DE KERAMPAOU 29140 MELGVEN France

Name of manufacturer	INDUSTRIALCHIMICA SRL
Address of manufacturer	VIA SORGAGLIA 40 35020 ARRE (PD) Italy
Location of manufacturing sites	VIA SORGAGLIA 40 35020 ARRE (PD) Italy

me of manufacturer	RATOUCY SAS
--------------------	-------------

	29 RUE DE LA FORET LOOZE –BP145 89303 JOIGNY France
Location of manufacturing sites	29 RUE DE LA FORET 89300 LOOZE France

Name of manufacturer	LARC
Address of manufacturer	ZA DE KERAMPAOU 29140 MELGVEN France
Location of manufacturing sites	ZA DE KERAMPAOU 29140 MELGVEN France

Name of manufacturer	NOXIMA
	CARREFOUR JEAN MONNET – LA CROIX SAINT OUEN 60201 COMPIEGNE France
6	CARREFOUR JEAN MONNET – LA CROIX SAINT OUEN 60201 COMPIEGNE France

Name of manufacturer	FARMAVIT OOD
	BUL TSAR BORIS III, N°63, OFFICE N°1 1612 SOFIA Bulgaria
Location of manufacturing sites	INDUSTRIALNA STR 2PLEVEN DISTRICT 5960 GULIANTSI Bulgaria

Name of manufacturer	AGGRESS
Address of manufacturer	EL. VENIZELOU 158 A 16341 ATHENS Greece
Location of manufacturing sites	ARMA THEBES 32200 VOIOTIA Greece

Name of manufacturer	SOFAR France
Address of manufacturer	ZA DU DREVERS BP 02 29190 PLEYBEN France
Location of manufacturing sites	ZA DU DREVERS BP 02 29190 PLEYBEN France

1.4. Manufacturer(s) of the active substance(s)

Active substance	Difenacoum
Name of manufacturer	Activa S.r.l.
Address of manufacturer	Via Feltre, 32 20132 Milano Italy
	Dr. Tezza S.r.l. Via Tre Ponti, 22 37050 S. Maria di Zevio (VR) Italy

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Difenacoum	3-(3-biphenyl-4- yl-1,2,3,4- tetrahydro-1- naphthyl)-4- hydroxycoumarin	active substance	56073-07-5	259-978-4	0,005

2.2. Type(s) of formulation

Ready-to-use bait : block

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H360D: May damage the unborn child.
	H373: May cause damage to organs blood through prolonged or repeated exposure {1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard:}.
Precautionary statements	P102: Keep out of reach of children.
	P202: Do not handle until all safety precautions have been read and understood.
	P280: Wear protective gloves.
	P308+P313: IF exposed or concerned: Get medical attention.
	P501: Dispose of {0::contents container} to {1::}.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. House mice – professionals (not relevant in France) - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: House mice Development stage: Juveniles and Adults
Field(s) of use	indoor use Indoor
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: 40 g of bait per bait point Dilution (%): 0 Number and timing of application: - 40 g of bait per bait point. If more than one bait point is needed, the minimum distance between bait point should be of 5 meters.
Category(ies) of users	professional
Pack sizes and packaging material	Bait product of 10 or 20 gr units Minimum pack size of 1.5 kg up to 25 kg.
	Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg. Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.1.1. Use-specific instructions

The bait station should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait station are intact and to remove rodent bodies. Re-fill bait when necessary.

4.1.2. Use-specific risk mitigation measures

4.1.3. Use-specific risk mitigation measures

Use-specific risk mitigation measures are not necessary when risk mitigation measures, as detailed in the section "General directions for use (professionals)", are implemented.

4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

Use specific instruction for safe disposal of the product and its packaging are not necessary when instruction for safe disposal of the product and its packaging, as detailed in the section "General directions for use (professionals)", are implemented.

4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (professionals)", are implemented.

4.2. Use description

Table 2. Rats – professionals (not relevant in France) - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles and adults Scientific name: Rattus rattus Common name: Black or roof rat Development stage: Juveniles and Adults
Field(s) of use	indoor use Indoor

Application method(s)	Method: bait application
	Detailed description: Ready-to-use bait to be used in tamper- resistant bait stations
Application rate(s) and frequency	Application rate: 60-100 g of bait per bait point
	Dilution (%): 0
	Number and timing of application: 60-100 g of bait per bait point. If more than one bait point is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	professional
Pack sizes and packaging material	Bait product of 10 or 20 gr units
	Minimum pack size of 1.5 kg up to 25 kg.
	Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.
	Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.2.1. Use-specific instructions

The bait station should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait station are intact and to remove rodent bodies. Re-fill bait when necessary.

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

Use-specific risk mitigation measures are not necessary when risk mitigation measures, as detailed in the section "General directions for use (professionals)", are implemented.

4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

Use specific instruction for safe disposal of the product and its packaging are not necessary when instruction for safe disposal of the product and its packaging, as detailed in the section "General directions for use (professionals)", are implemented.

4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (professionals)", are implemented.

4.3. Use description

Table 3. House mice and rats - professionals (not relevant in France) - outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: house mouse Development stage: Juveniles and Adults Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles and adults Scientific name: Rattus rattus Common name: Black or roof rat Development stage: Juveniles and adults
Field(s) of use	outdoor use Outdoor around building
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: - 40 g of bait per bait point for mice 60-100 g of bait per bait point for rats. Dilution (%): 0 Number and timing of application: - 40 g of bait per bait point for mice.

	- 60-100 g of bait per bait point for rats.
	If more than one bait point is needed, the minimum distance between bait points should be of 5 meters.
Category(ies) of users	professional
Pack sizes and packaging material	Bait product of 10 or 20 gr units
	Minimum pack size of 1.5 kg up to 25 kg.
	Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.
	Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.3.1. Use-specific instructions

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait station should be visited for mice at least every 2 to 3 days at the beginning of the treatment and for rats only 5 to 7 days after the beginning of the treatment and at least weekly afterwards (for mice and rats), in order to check whether the bait is accepted, the bait station are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in bait station in which bait has been damaged by water or contaminated by dirt.

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or drainage systems, ensure that bait contact with water is avoided.

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

Use specific instruction for safe disposal of the product and its packaging are not necessary when instruction for safe disposal of the product and its packaging, as detailed in the section "General directions for use (professionals)", are implemented.

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (professionals)", are implemented.

4.4. Use description

Table 4. House mice and rats - trained professionals - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development	_
stage)	Common name: brown rat
	Development stage: Juveniles and Adults
	Scientific name: Rattus rattus
	Common name: Black or roof rat
	Development stage: Juveniles and Adults
	Scientific name: Mus musculus
	Common name: house mouse
	Development stage: Juveniles ans Adults
Field(s) of use	indoor use
	Indoor
Application method(s)	Method: bait application
	Detailed description: Bait formulations: - ready-to-use bait to
	be used in tamper-resistant bait stations - ready-to-use bait may
	be used in covered and protected baiting points as long as they
	provide the same level of protection for non-target species and
	humans as tamper-resistant bait station
Application rate(s) and frequency	Application rate: Rats - High infestation: 100 g of bait per baiting
	point Low infestation: 60 g of bait per baiting point. Mice - 40 g
	of bait per baiting point.
	Dilution (%): 0
	Number and timing of application:

	Rats: - High infestation: 100 g of bait per baiting point Low infestation: 60 g of bait per baiting point. Mice: - 40 g of bait per baiting point. Permanent baiting, were applicable.
Category(ies) of users	trained professional
Pack sizes and packaging material	Bait product of 10 or 20 gr units Minimum pack size of 1.5 kg up to 25 kg.
	Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.
	Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.4.1. Use-specific instructions

- Remove the remaining product at the end of treatment period.

4.4.2. Use-specific risk mitigation measures

4.4.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.

- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use the product in pulsed baiting treatments.
- Do not apply this product directly in the burrows

4.4.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.4.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

Use specific instruction for safe disposal of the product and its packaging are not necessary when instruction for safe disposal of the product and its packaging, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.4.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.5. Use description

Table 5. Mice and rats - trained professionals - outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: house mouse Development stage: Juveniles and Adults
	Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles and Adults
	Scientific name: Rattus rattus

	Common name: Black or roof rat Development stage: Juveniles and Adults
Field(s) of use	outdoor use
	Outdoor around buildings
Application method(s)	Method: bait application
	Detailed description: Bait formulations: - ready-to-use bait to be used in tamper-resistant bait stations - ready-to-use bait may be used in covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait station
Application rate(s) and frequency	Application rate: Rats - High infestation: 100 g of bait per baiting point Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point.
	Dilution (%): 0
	Number and timing of application: Bait products
	Rats: - High infestation: 100 g of bait per baiting point Low infestation: 60 g of bait per baiting point.
	Mice: - 40 g of bait per baiting point.
	- Permanent baiting were applicable.
Category(ies) of users	trained professional
Pack sizes and packaging material	Bait product of 10 or 20 gr units
	Minimum pack size of 1.5 kg up to 25 kg.
	Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.
	Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.5.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.

- Remove the remaining product at the end of treatment period.

4.5.2. Use-specific risk mitigation measures

4.5.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows

4.5.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.5.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

Use specific instruction for safe disposal of the product and its packaging are not necessary when instruction for safe disposal of the product and its packaging, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.5.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.6. Use description

Table 6. Rats - trained professionals - Outdoor open areas

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles and Adults
	Scientific name: Rattus rattus Common name: Black or roof rat Development stage: Juveniles and adults
Field(s) of use	outdoor use Outdoor open areas.
Application method(s)	Method: bait application Detailed description: Bait formulations: - ready-to-use bait to be used in tamper-resistant bait stations or in covered and protected baiting points
Application rate(s) and frequency	Application rate: - High infestation: 100 g of bait per baiting point Low infestation: 60 g of bait per baiting point. Dilution (%): 0 Number and timing of application: Bait products - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point.
	- Permanent baiting, were applicable.
Category(ies) of users	trained professional

Pack sizes and packaging material	Bait product of 10 or 20 gr units
	Minimum pack size of 1.5 kg up to 25 kg.
	Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.
	Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.6.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- -Remove the remaining product at the end of treatment period.

4.6.2. Use-specific risk mitigation measures

4.6.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows

4.6.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.6.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

Use specific instruction for safe disposal of the product and its packaging are not necessary when instruction for safe disposal of the product and its packaging, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.6.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.7. Use description

Table 7. Brown rats - trained professionals - Sewer

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles and adults
Field(s) of use	indoor use Sewers
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water.
Application rate(s) and frequency	Application rate: 60-100 g of bait per baiting point Dilution (%): 0 Number and timing of application: Bait products 60-100 g of bait per baiting point
Category(ies) of users	trained professional
Pack sizes and packaging material	Bait product of 10 or 20 gr units

Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg. Packaging material: plastic composite-triple layer (polyester/ PET met/polyethylene)

4.7.1. **Use-specific instructions**

Baits must be applied in a way so that they do not come into contact with water and are not washed away.

4.7.2. **Use-specific risk mitigation measures**

4.7.3. Use-specific risk mitigation measures

The product must be used only in sewers connected to a sewage treatment plant. Do not use in rainwater systems with direct releases to the aquatic compartment.

Do not use this product in pulsed baiting treatments.

- 4.7.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment
- 4.7.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section "Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment" for the information to be shown on the label).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves (rubber) during product handling phase.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
- the product shall not be supplied to the general public (e.g. "for professionals only").
- the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

- users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre or or emergency telephone number.
- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 2 years.

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.