

CLH report

Proposal for Harmonised Classification and Labelling

Based on Regulation (EC) No 1272/2008 (CLP Regulation),
Annex VI, Part 2

Substance Name: 4 vinylcyclohexene (VCH)

EC Number: 202-848-9

CAS Number: 100-40-3

Index Number:

Contact details for dossier submitter:

ANSES

253, av. du général Leclerc

94701 Maisons-Alfort Cedex

reach@anses.fr

00 (33)1 56 29 19 30

Version number: 3

Date: May 2011

CONTENTS

Part A.

1	PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING	4
1.1	SUBSTANCE	4
1.2	HARMONISED CLASSIFICATION AND LABELLING PROPOSAL	4
1.3	PROPOSED HARMONISED CLASSIFICATION AND LABELLING BASED ON CLP REGULATION AND/OR DSD CRITERIA	6
2	BACKGROUND TO THE CLH PROPOSAL	9
2.1	HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING	9
2.2	SHORT SUMMARY OF THE SCIENTIFIC JUSTIFICATION FOR THE CLH PROPOSAL	9
2.3	CURRENT HARMONISED CLASSIFICATION AND LABELLING	9
2.4	CURRENT SELF-CLASSIFICATION AND LABELLING	9
3	JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL	9

Part B.

	SCIENTIFIC EVALUATION OF THE DATA	10
1	IDENTITY OF THE SUBSTANCE	10
1.1	NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE	10
1.2	COMPOSITION OF THE SUBSTANCE	11
	Current Annex VI entry: no harmonised classification	11
1.2.1	<i>Composition of test material</i>	14
1.3	PHYSICO-CHEMICAL PROPERTIES	15
2	MANUFACTURE AND USES	17
2.1	MANUFACTURE	17
2.2	IDENTIFIED USES	17
3	CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES	17
4	HUMAN HEALTH HAZARD ASSESSMENT	17
4.1	TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)	17
4.1.1	<i>Non-human information</i>	17
4.1.2	<i>Human information</i>	22
4.1.3	<i>Summary and discussion on toxicokinetics</i>	22
4.2	ACUTE TOXICITY	25
4.3	SPECIFIC TARGET ORGAN TOXICITY – SINGLE EXPOSURE (STOT SE)	25
4.4	IRRITATION	25
4.5	CORROSIVITY	25
4.6	SENSITISATION	25
4.7	REPEATED DOSE TOXICITY	25
4.7.1	<i>Non-human information</i>	29
4.7.1.1	Repeated dose toxicity: oral	29
4.7.1.2	Repeated dose toxicity: inhalation	29
4.7.1.3	Repeated dose toxicity: dermal	29
4.7.1.4	Repeated dose toxicity: other routes	29
4.7.1.5	Human information	29
4.7.1.6	Other relevant information	29
4.7.1.7	Summary and discussion of repeated dose toxicity	31
4.8	SPECIFIC TARGET ORGAN TOXICITY (CLP REGULATION) – REPEATED EXPOSURE (STOT RE)	32
4.9	GERM CELL MUTAGENICITY (MUTAGENICITY)	33

4.9.1	<i>Non-human information</i>	34
4.9.1.1	<i>In vitro data</i>	34
4.9.1.2	<i>In vivo data</i>	35
4.9.2	<i>Human information</i>	35
4.9.3	<i>Other relevant information</i>	35
4.9.4	<i>Summary and discussion of mutagenicity</i>	36
4.9.5	<i>Comparison with criteria</i>	37
4.9.6	<i>Conclusions on classification and labelling</i>	37
4.10	CARCINOGENICITY	38
4.10.1	<i>Non-human information</i>	41
4.10.1.1	<i>Carcinogenicity: oral</i>	41
4.10.1.2	<i>Carcinogenicity: inhalation</i>	41
4.10.1.3	<i>Carcinogenicity: dermal</i>	41
4.10.2	<i>Human information</i>	41
4.10.3	<i>Other relevant information</i>	41
4.10.4	<i>Summary and discussion of carcinogenicity</i>	43
4.10.5	<i>Comparison with criteria</i>	45
4.10.6	<i>Conclusions on classification and labelling</i>	46
4.11	TOXICITY FOR REPRODUCTION	46
4.11.1	<i>Effects on fertility</i>	51
4.11.1.1	<i>Non-human information</i>	51
4.11.1.2	<i>Human information</i>	51
4.11.2	<i>Developmental toxicity</i>	51
4.11.2.1	<i>Non-human information</i>	51
4.11.2.2	<i>Human information</i>	51
4.11.3	<i>Other relevant information</i>	51
4.12	OTHER EFFECTS	52
5	ENVIRONMENTAL HAZARD ASSESSMENT	52
6	OTHER INFORMATION	52
7	REFERENCES	52
8	ANNEXES	54

Part A.

1 PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING

1.1 Substance

Table 1: Substance identity

Substance name:	<i>4-vinylcyclohexene</i>
EC number:	<i>202-848-9</i>
CAS number:	<i>100-40-3</i>
Annex VI Index number:	
Degree of purity:	<i>95% min (for technical product) and 99% (research)</i>
Impurities:	<i>1,5-cyclooctadiene tert-butylcatechol para-tert-butylcatechol 1,5,9-cyclododecatriene 1,2-divinylcyclobutane water</i>

1.2 Harmonised classification and labelling proposal

Table 2: The current Annex VI entry and the proposed harmonised classification

	CLP Regulation	Directive 67/548/EEC (Dangerous Substances Directive; DSD)
Current entry in Annex VI, CLP Regulation	-	-
Current proposal for consideration by RAC	Carc. 1B – H 350	Carc. Cat. 2; R45
Resulting harmonised classification (future entry in Annex VI, CLP)	Carc. 1B – H 350	Carc. Cat. 2; R45

CLH REPORT FOR [4 VINYL CYCLOHEXENE]

Regulation)		
--------------------	--	--

1.3 Proposed harmonised classification and labelling based on CLP Regulation and/or DSD criteria

Table 3: Proposed classification according to the CLP Regulation

CLP Annex I ref	Hazard class	Proposed classification	Proposed SCLs and/or M-factors	Current classification ¹⁾	Reason for no classification ²⁾
2.1.	Explosives	None		None	Not evaluated
2.2.	Flammable gases	None		None	Not evaluated
2.3.	Flammable aerosols	None		None	Not evaluated
2.4.	Oxidising gases	None		None	Not evaluated
2.5.	Gases under pressure	None		None	Not evaluated
2.6.	Flammable liquids	None		None	Not evaluated
2.7.	Flammable solids	None		None	Not evaluated
2.8.	Self-reactive substances and mixtures	None		None	Not evaluated
2.9.	Pyrophoric liquids	None		None	Not evaluated
2.10.	Pyrophoric solids	None		None	Not evaluated
2.11.	Self-heating substances and mixtures	None		None	Not evaluated
2.12.	Substances and mixtures which in contact with water emit flammable gases	None		None	Not evaluated
2.13.	Oxidising liquids	None		None	Not evaluated
2.14.	Oxidising solids	None		None	Not evaluated
2.15.	Organic peroxides	None		None	Not evaluated
2.16.	Substance and mixtures corrosive to metals	None		None	Not evaluated
3.1.	Acute toxicity - oral	None		None	Not evaluated
	Acute toxicity - dermal	None		None	Not evaluated
	Acute toxicity - inhalation	None		None	Not evaluated
3.2.	Skin corrosion / irritation	None		None	Not evaluated
3.3.	Serious eye damage / eye irritation	None		None	Not evaluated
3.4.	Respiratory sensitisation	None		None	Not evaluated
3.4.	Skin sensitisation	None		None	Not evaluated
3.5.	Germ cell mutagenicity	None		None	Not evaluated
3.6.	Carcinogenicity	H350: May cause cancer	None	None	
3.7.	Reproductive toxicity	None		None	Not evaluated
3.8.	Specific target organ toxicity –single exposure	None		None	Not evaluated
3.9.	Specific target organ toxicity – repeated exposure	None		None	Not evaluated

CLH REPORT FOR [4 VINYL CYCLOHEXENE]

3.10.	Aspiration hazard	None		None	Not evaluated
4.1.	Hazardous to the aquatic environment	None		None	Not evaluated
5.1.	Hazardous to the ozone layer	None		None	Not evaluated

¹⁾Including specific concentration limits (SCLs) and M-factors

²⁾Data lacking, inconclusive, or conclusive but not sufficient for classification

Labelling: Signal word: Danger
 Hazard statements: H350
 Precautionary statements: not harmonised

Proposed notes assigned to an entry:

Table 4: Proposed classification according to DSD

Hazardous property	Proposed classification	Proposed SCLs	Current classification ¹⁾	Reason for no classification ²⁾
Explosiveness	None		None	Not evaluated
Oxidising properties	None		None	Not evaluated
Flammability	None		None	Not evaluated
Other physico-chemical properties <i>[Add rows when relevant]</i>	None		None	Not evaluated
Thermal stability	None		None	Not evaluated
Acute toxicity	None		None	Not evaluated
Acute toxicity – irreversible damage after single exposure	None		None	Not evaluated
Repeated dose toxicity	None		None	Not evaluated
Irritation / Corrosion	None		None	Not evaluated
Sensitisation	None		None	Not evaluated
Carcinogenicity	Carc. Cat. 2; R45 May cause cancer	None	None	
Mutagenicity – Genetic toxicity	None		None	Not evaluated
Toxicity to reproduction – fertility	None		None	Not evaluated
Toxicity to reproduction – development	None		None	Not evaluated
Toxicity to reproduction – breastfed babies. Effects on or via lactation	None		None	Not evaluated
Environment	None		None	Not evaluated

¹⁾ Including SCLs

²⁾ Data lacking, inconclusive, or conclusive but not sufficient for classification

Labelling: Indication of danger: T
 R-phrases: R45
 S-phrases: S53

2 BACKGROUND TO THE CLH PROPOSAL

2.1 History of the previous classification and labelling

VCH was not listed in Annex I of the 67/548/EC Directive.

2.2 Short summary of the scientific justification for the CLH proposal

VCH is currently not classified according to Annex VI of CLP. However, IARC classifies it as Group 2B: The agent (mixture) is possibly carcinogenic to humans since 1994. Based on the increased incidence of ovarian tumors in female mice exposed orally by gavage to VCH for two years, and considering the fact that the mode of action described for this effect is plausible in human, a classification as Carc. Cat. 2; R45; May cause cancer, or H350: May cause cancer, is warranted.

2.3 Current harmonised classification and labelling

No current harmonised classification in Annex VI of CLP.

2.4 Current self-classification and labelling

No data available.

3 JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL

VCH has a CMR property (carcinogenicity). Harmonised classification and labelling for CMR and respiratory sensitisation is a Community-wide action under article 115 of REACH and article 36(1) of CLP. VCH is currently not classified according to Annex VI of CLP.

Repeated-dose toxicity and genotoxicity data are presented for information as they may provide relevant data for assessment of carcinogenicity but no classification is discussed and proposed for these endpoints.

Part B.

SCIENTIFIC EVALUATION OF THE DATA


1 IDENTITY OF THE SUBSTANCE

1.1 Name and other identifiers of the substance

Table 5: Substance identity

EC number:	202-848-9
EC name:	4-vinylcyclohexene
CAS number (EC inventory):	100-40-3
CAS number:	
CAS name:	Cyclohexene, 4-ethenyl-
IUPAC name:	4-vinylcyclohexene
CLP Annex VI Index number:	
Molecular formula:	C ₈ H ₁₂
Molecular weight range:	108,18 g/mol

Structural formula:



1.2 Composition of the substance

Table 6: Constituents (non-confidential information)

Constituent	Typical concentration	Concentration range	Remarks
4-vinylcyclohexene		95% - 99%	

Current Annex VI entry: no harmonised classification

Table 7: Impurities (non-confidential information)

Impurity	Typical concentration	Concentration range	Remarks
cycloocta-1,5-diene		3% max	
water		200 ppm max	
tert-butylpyrocatechol		25-200 ppm	
para-tert-butylpyrocatechol	50 ppm		
1,5,9-cyclododecatriene		traces	
1,2-divinylcyclobutane		traces	

Chemical Name: 1,5-cyclooctadiene


EC Number: 203-907-1

CAS Number: 111-78-4

IUPAC Name: cycloocta-1,5-diene

Molecular Formula: C₈H₁₂

Structural Formula:



Molecular Weight: 108.18 g/mol

Typical concentration (% w/w):

Concentration range (% w/w): 3 % max

Classification: No harmonised classification

Chemical Name: water

EC Number: 231-791-2

CAS Number: 7732-18-5

IUPAC Name: water

Molecular Formula: H₂O

Structural Formula: H-O-H

Molecular Weight: 18.02 g/mol

Typical concentration (% w/w):

Concentration range (% w/w): 200ppm max

Classification: No harmonised classification

Chemical Name: tert-butylpyrocatechol


EC Number: 248-325-9

CAS Number: 27213-78-1

IUPAC Name: tert- Butylbenzene-1,2-diol

Molecular Formula: C₁₀H₁₄O₂

Structural Formula:



Molecular Weight: 166.22 g/mol

Typical concentration (% w/w):

Concentration range (% w/w): 25-200 ppm

Classification: No harmonised classification

Chemical Name: para-tert-butylpyrocatechol


EC Number: 202-653-9

CAS Number: 98-29-3

IUPAC Name: 4-tert-Butylbenzene-1,2-diol

Molecular Formula: C₁₀H₁₄O₂

Structural Formula:



Molecular Weight: 166.22 g/mol

Typical concentration (% w/w): 50 ppm

Concentration range (% w/w): Information not available

Classification: No harmonised classification

Chemical Name: 1,5,9-cyclododecatriene


EC Number: 220-437-2

CAS Number: 2765-29-9

IUPAC Name: trans,cis,cis-1,5,9-cyclododecatriene

Molecular Formula: C₁₂H₁₈

Structural Formula:



Molecular Weight: 162.27g/mol

Typical concentration (% w/w): traces

Concentration range (% w/w): Information not available

Classification: No harmonised classification

Chemical Name: 1,2-divinylcyclobutane


EC Number:

CAS Number: 2422-85-7

IUPAC Name: 1,2-divinylcyclobutane

Molecular Formula: C₈H₁₂

Structural Formula:



Molecular Weight: 108.18g/mol

Typical concentration (% w/w): traces

Concentration range (% w/w): Information not available

Classification : No harmonised classification

Table 8: Additives (non-confidential information)

Additive	Function	Typical concentration	Concentration range	Remarks
No data concerning the additives of VCH are available.				

1.2.1 Composition of test material

VCH used in the rat and mouse carcinogenicity studies from NTP had a purity of 98%. Impurities in two lots of test chemical included 0.01 % butylated hydroxytoluene in one and 0.005% tert-butylcatechol in the other, which had been added as inhibitors of peroxide formation

1.3 Physico-chemical properties

Table 9: Summary of physico - chemical properties

Property	Value	Reference	Comment (e.g. measured or estimated)
State of the substance at 20°C and 101,3 kPa	Colourless liquid	NTP Technical Report on the toxicology and carcinogenesis studies of 4-VCH in F344/N rats and B6C3F1 mice, US National Toxicology Program, 1986	
Melting/freezing point	-108,9°C	Handbook of chemistry and physics 2005-2006 National Library of Medicine HSDB Database	
Boiling point	128°C	Handbook of chemistry and physics 2005-2006 National Library of Medicine HSDB Database	
Relative density	0,8299 at 20°C	Handbook of chemistry and physics 2005-2006 National Library of Medicine HSDB Database	
Vapour pressure	2,09 kPa at 25°C	NLM - ChemIDplus Lite National Library of Medicine HSDB Database	
Surface tension	Not available		
Water solubility	50mg/L at 25°C Not water soluble	NLM - ChemIDplus Lite National Library of Medicine HSDB Database Handbook of chemistry and physics 2005-2006	
Partition coefficient n-	Log Pow = 3,93	International	

CLH REPORT FOR [4 VINYLCHYCLOHEXENE]

octanol/water		Chemical Safety Card, NIOSH, 1995 National Library of Medicine HSDB Database	
Flash point	15.9°C (open cup) 16°C (closed cup)	IUCLID Data Set 2006 National Library of Medicine HSDB Database	
Flammability	Flammable, dangerous fire risk. Dangerous fire hazard when exposed to heat or flame OSHA flammability Class: IB	National Library of Medicine HSDB Database MSDS OHS30250, 2009	
Explosive properties	Vapor/air mixtures are explosive	International Chemical Safety Card, NIOSH, 1995	
Self-ignition temperature	269°C	International Chemical Safety Card, NIOSH, 1995 National Library of Medicine HSDB Database	
Oxidising properties	Can react with oxidizers Upon contact with oxygen, VCH undergoes auto-oxidation to produce vinylcyclohexene hydroperoxide	National Library of Medicine HSDB Database	
Granulometry	Non applicable		
Stability in organic solvents and identity of relevant degradation products	Miscible with methanol Soluble in ether, benzene, petroleum ether	National Library of Medicine HSDB Database	
Dissociation constant	Not available		
Viscosity	Not available		
Reactivity towards container material	Incompatible materials: halogens, oxidizing materials Closed containers may rupture violently. Containers may rupture or explode if exposed to heat.	MSDS OHS30250, 2009	
Thermal stability	May polymerise at temperatures above	IARC monographs	

	26,6°C and prolonged exposure to oxygen	volume 60	
Stability/shelf life	Oxidizes in air to form hydroperoxide	National Library of Medicine HSDB Database	
Henry's law constant	0.045 atm.m ³ /mol at 25°C	NLM - ChemIDplus Lite	

2 MANUFACTURE AND USES

2.1 Manufacture

Not relevant for this dossier

2.2 Identified uses

VCH is found in industrial processes involving 1,3-butadiene, including the manufacture of lauric acid. VCH formed may be sold as-is or converted to 4-vinylcyclohexene diepoxide. It has been used as a chemical intermediate for production of flame retardants, flavours and fragrances, in the manufacture of polyolefins, as a solvent and in the manufacture of its diepoxide. Low levels of occupational exposure have been measured during the production and use of 1,3-butadiene.

3 CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES

Not evaluated in this dossier

4 HUMAN HEALTH HAZARD ASSESSMENT

Publications reported in this proposal and not reviewed in the IARC monograph on VCH (IARC, 1994) are indicated in the section: 7 References.

4.1 Toxicokinetics (absorption, metabolism, distribution and elimination)

4.1.1 Non-human information

4-vinylcyclohexene (VCH) has numerous data regarding its metabolism and its interspecies difference. Data are summarised below and grouped by paragraphs.

Smith and co-workers compared the disposition and the *in vitro* metabolism of VCH in female mice and rats (Smith *et al.*, 1990a). Female B6C3F1 mice and Fisher 344 rats were exposed to a single

dose of 400 mg/kg bw [^{14}C]VCH in corn oil by gavage. By 24h, mice and rats eliminated 97 and 88% of the dose, respectively. In both species, urine was the major route of excretion with 50-60% of the administered dose eliminated by this route. The second major route of elimination was expired air, as expired organics (1/3 of the absorbed dose in mice and rats). There were no differences between the rat and the mouse in the disposition of [^{14}C]VCH equivalents in the ovary. No evidence was obtained to indicate that the ovaries of either species retained VCH as a parent compound or as radioactive equivalents. The tissue with the highest concentration of VCH in both species was adipose tissue. The highest concentration of VCH in the adipose tissue of mice was found between 1 and 2 h after VCH treatment, whereas rat adipose tissue continued to accumulate VCH until at least 8 h after dosing. Tissue concentrations in other tissues were slightly higher in the tissues of rats than the corresponding values in mouse tissues over the times studied (6 h in mice, and 8 h in rats).

After an i.p. administration of VCH 800 mg/kg bw, the highest concentration of VCH-1,2-epoxide (41 nmol/mL) was found in blood of mice, 2 h after dosing. VCH-7,8-epoxide was not detected in mice. These monoepoxides were not detected in rats up to 6 h after dosing (less than 2.5 nmol/mL).

The rate of epoxidation of VCH to VCH-1,2-epoxide is 6.5-fold greater in female mice liver microsomes compared to those of female rats. The blood concentration of VCH-1,2-epoxide was present in the blood of mice with the highest concentration at 2 h (41 nmol/mL) whereas the blood concentration of this monoepoxide in rats was < 2.5 nmol/mL up to 8 h after exposure. VCH-7,8-epoxide was not present in the blood of either species at the level of detection (level of sensitivity for VCH epoxide = 2.5 nmol/mL). Mouse hepatic microsomes formed 4 times more of VCH-1,2-epoxide.

Giannarini and co-workers administered 500 mg/kg VCH or VCH-1,2-epoxide or VCH diepoxide i.p. in corn oil to male albino Swiss mice (Giannarini *et al.*, 1981). VCH and VCH-1,2-epoxide were able to enhance the activities of certain xenobiotic transforming enzymes. CYP, cytochrome *b5*, NADPH-cytochrome *c* reductase, aminopyrine-*N*-demethylase and epoxide hydrolase were induced by VCH or VCH-1,2-epoxide (results from four separate tissue pools, each pool consisting of livers from 5-8 mice). On the contrary, *p*-nitroanisole-*O*-demethylase activity was not significantly induced. Hepatic glutathione levels were rapidly depleted after exposure to VCH, VCH-1,2-epoxide or VCH diepoxide, suggesting that glutathione is probably involved in the metabolism of VCH. VCH-1,2-epoxide and VCH diepoxide were more active than VCH, maybe because of the conjugation of the monoepoxide and the diepoxide with GSH.

Rats accumulate more VCH (or equivalent) than mice but mice metabolises VCH to VCH-1,2-epoxide or VCH-7,8-epoxide more rapidly and more efficiently than rats.

Cytochrome involvement during VCH metabolism

Fontaine and co-workers investigated the induction of CYP involved in VCH bioactivation in mice and rats after repeated exposure (Fontaine *et al.*, 2001). B6C3F1 mice and Fischer-344 rats (28-38 days of age) were dosed with either VCH (7.5 mmol/kg i.p. for 10 days), VCH-1,2-epoxide (1.75 mmol/kg/i.p. for 10 days), vinylcyclohexene diepoxide (VCD)¹ (0.4 mmol/kg i.p. for 10 days), or phenobarbital (80 mg/kg i.p. for 5 days). In each treatment group, microsomes were prepared from four individual rats or were pooled from four mice per group (16 mice total). Hepatic microsomes

¹ Vinylcyclohexene diepoxide (VCD) is the ultimate ovotoxic metabolite of VCH (Doerr-Stevens *et al.*, 1999). Its toxicity is described in sections 4.7.1.6 and 4.10.3.

prepared from mice or rats treated repeatedly with VCH demonstrated significantly increased VCH bioactivation *in vitro*, as assessed by increased formation of VCH-1,2-epoxide (3.8- and 2.0-fold in mice and rats, respectively) and VCH-7,8-epoxide, the highest amounts being formed in mice. Although incubations were conducted for up to 60 min to allow possible VCD formation, VCD was only detected in mice pre-treated with either phenobarbital or VCH, but not in control mice or in control or pre-treated rats.

Mice and rats were then dosed with VCH, VCH-1,2-epoxide, or VCD for 10 days and measured for increases in hepatic microsomal CYP levels or activities. Total hepatic CYP levels were elevated only in microsomes from mice pretreated with VCH or VCH-1,2-epoxide. Immunoblotting analysis of microsomes from VCH-treated rodents revealed elevated levels of CYP2A and CYP2B in mice but not in rats. VCH-1,2-epoxide pretreatment also increased CYP2B levels in the mouse. Activities toward specific substrates for CYP2A and CYP2B (coumarin and pentoxyresorufin, respectively) confirmed that VCH and VCH-1,2-epoxide pretreatments resulted in increased catalytic activities of CYP2A and CYP2B in the mouse but not the rat. Pretreatment with phenobarbital, a known inducer of CYP2A and CYP2B, increased VCH bioactivation in both species. Interestingly, metabolism studies with human CYP "Supersomes" (human CYP + P450 reductase + cytochrome b5) reveal that, of eight isoforms tested (CYP1A2, CYP2A6, CYP2B6, CYP2C6, CYP2E1, CYP3A4, CYP4A11 and aromatase), only human CYP2E1 and CYP2B6 were capable of significantly catalyzing VCH epoxidation, whereas CYP2B6, CYP2A6, CYP2E1, and CYP3A4 were capable of catalyzing the epoxidation of the monoepoxides.

Smith and co-workers investigated the biochemical basis for the different susceptibility to 4-vinylcyclohexene -induced ovarian toxicity and carcinogenesis in mice and rats (Smith *et al.*, 1990c). Cytochromes P450 (CYP) involved in the epoxidation of 4-vinylcyclohexene (VCH) in liver microsomes from female B6C3F1 and 129/J (deficient in constitutive expression of CYP2B forms) mice and F344 rats were determined.

In mice liver microsomes, mouse CYP2A forms were involved in VCH epoxidation. No protein immunochemically related to mouse CYP2A was detected in female rat hepatic microsomes. However, VCH epoxidation was catalysed by CYP2B1 in microsomes from both species. But the level of constitutive expression of CYP2B forms is lower in rats compared with mice (0.07 nmol/min/mg protein in rats vs 0.26 nmol/min/mg protein in mice). A high rate of VCH epoxidation was observed in Phenobarbital-treated female rat liver microsomes, suggesting that pre-treatment with Phenobarbital, which induces mainly CYP2B isoforms, could increase the blood level of VCH-1,2-epoxide in VCH-treated rats and increase susceptibility to VCH-induced ovarian carcinogenicity. The constitutive expression in mouse liver of CYP2A and 2B forms, which catalyse ca. 80% of VCH epoxidation by hepatic microsomes from B6C3F1 mice, partially explains the susceptibility of mice to VCH-induced ovarian carcinogenicity. Although CYP2B forms of female Fischer 344 rats can also catalyze VCH epoxidation into VCH-1,2-epoxide, the expression of these isozymes in untreated animals probably does not occur at levels comparable to mice. Other CYPs are involved in VCH epoxidation in female rat hepatic microsomes, but they are probably expressed at very low levels and / or have poor catalytic activity toward VCH.

The effect of repeated exposure of VCH (7.5 mmol/kg/day) on mouse liver microsomal activities and VCH epoxidation was determined by Doerr-Stevens and co-workers (Doerr-Stevens *et al.*, 1999). CYP2B and CYP2A, principle isoforms involved in the bioactivation of VCH, as well as CYP2E1 and CYP3A were evaluated. VCH exposure increased total CYP content (35-83% above control levels) after either 5, 10, or 15 days of treatment. Western blot analysis revealed an induction of CYP2A, CYP2B, and CYP2E1 at day 10. Elevated levels of CYP2A and CYP2B

correlated with marker androstenedione and testosterone 16 α - and 16 β -hydroxylase activities. Microsomes prepared from mice pretreated with VCH for 10 days demonstrated an increase (≥ 2 -fold) in the rate of VCH monoepoxide and diepoxide formation. Microsomal VCH epoxidation was increased to a similar extent by phenobarbital, acetone, and dexamethasone treatment. An increase in cytosolic glutathione S-transferase activity was observed after repeated VCH treatment, an enzyme potentially involved in detoxification of the VCH epoxides. Interestingly, preliminary studies indicated that circulating levels of the monoepoxide (VCH-1,2-epoxide) and diepoxide metabolites of VCH were elevated after repeated dosing of VCH. Overall, the results indicate that repeated exposure of VCH in mice induces CYP-dependent activities, and in turn induction of its metabolism.

The data presented above show that, CYP2A and 2B are the main cytochromes involved in VCH metabolism in rodents. VCH promotes its own metabolism by increasing expression of these cytochromes. Inter-rodent difference in VCH metabolism is correlated to the expression of these 2 cytochromes: VCH is poorly metabolised in rats due to the fact that CYP2A is not present and CYP2B is poorly expressed in rats.

Presence of VCH metabolites in various tissues

It has to be noted that VCD is classified in Annex I of the 67/548/EC Directive as Carc. Cat. 3; R40 - T; R23/24/25.

Keller and co-workers studied the metabolism of VCH in microsomes from CrI:CD BR rat and B6C3F1 mouse liver, lung, and ovary (Keller *et al.*, 1997). Tissue samples were incubated with the test chemicals for 15 min at 37°C, and were tested for their ability to catalyze the following reactions:

- 4-VCH to VCH-1,2-epoxide and VCH-7,8-epoxide
- VCH-1,2-epoxide to VCD and VCH 1,2-diol
- VCH-7,8-epoxide to VCD and VCH 7,8 diol
- Hydrolysis of VCD.

3,3,3-trichloropropene oxide was added for reactions in which an epoxide metabolite was expected.

The reaction 4-VCH to VCH-1,2-epoxide was detected in liver and lung of rat and mouse. Mouse liver had a V_{max} value 56-fold higher than that for rat liver. In lung tissue, the reaction proceeded only 2.5- to 3.8 fold faster in mice than in rats.

The reaction 4-VCH to VCH-7,8-epoxide was detected in rat and mouse liver and in mouse lung. The V_{max} was lower in rat liver compared to mouse liver (0.007 nmol/min/mg protein in rats vs 0.91 nmol/min/mg protein in mice), and it was 12-fold lower than that for metabolism to the VCH-1,2-epoxide.

Rat and mouse livers were very similar in their K_m and V_{max} values for the reaction VCH-1,2-epoxide to VCD and also for VCH-7,8-epoxide to VCD.

VCH-1,2-epoxide to VCH 1,2-diol was only detectable in liver tissue from rats and mice and was similar. The hydrolysis of VCH-7,8-epoxide to VCH-7,8 diol was only detectable in rat liver. (V_{max} = 135.8 nmol/min/mg protein). Hydrolysis of VCD was detectable in rat and mouse liver and lung and in rat ovary. Rat liver had a V_{max} nine-fold higher than did the mouse liver. For lung

and ovary, the reactions seemed to be saturated even at the lowest concentrations where substrate is detectable. Maximum activity is approximately 14-fold lower in rat lung than in rat liver, and is approximately equal in mouse liver and lung. The mouse ovary had activity less than or equal to the rat ovary, and in both species the ovarian activity was low (no control).

In conclusion, this study indicates that the mouse has significant greater capacity to metabolize VCH to reactive species than does the rat, and that the mouse hydrolyzes epoxides less efficiently than the rat.

Metabolism of VCH in mice ovaries or ovarian fractions

The expression of CYP2E1, CYP2A, and CYP2B isoforms was investigated in isolated ovarian fractions from mice exposed to VCH or to its ovotoxic metabolite, VCD (Cannady *et al.*, 2003). B6C3F1 mice were administered i.p. daily for 15 days with VCH (7.4 mmol/kg bw/d) or VCD (0.57 mmol/kg bw/d). Ovaries were removed and either isolated into specific ovarian compartments for mRNA analysis, fixed for immunohistochemistry, or prepared for enzymatic assays. mRNA and protein for all isoforms were expressed/distributed in all ovarian fractions from vehicle-treated mice. In small preantral follicles (= F1 follicles) which are specifically destroyed by VCH or VCD, VCH or VCD dosing increased ($p < 0.05$) mRNA encoding CYP2E1 ($645 \pm 14\%$ in VCH above control; $582 \pm 16\%$ in VCD above control), CYP2A ($689 \pm 8\%$ in VCH above control; $730 \pm 22\%$ in VCD above control), and CYP2B ($246 \pm 7\%$ in VCH above control). VCH dosing altered ($p < 0.05$) mRNA encoding CYP2E1 in non targeted F3 follicles ($168 \pm 7\%$) and CYP2A in interstitial cells ($207 \pm 19\%$) above control. Immunohistochemical analysis revealed the greatest staining intensity for all CYP isoforms in the interstitial cells. VCH dosing altered ($p < 0.05$) staining intensity in interstitial cells for CYP2E1 ($19 \pm 2.4\%$ below control) and CYP2A ($39 \pm 5\%$ above control). Staining intensity for CYP2B was increased ($p < 0.05$) above control in granulosa cells of small preantral ($187 \pm 42\%$) and antral ($63 \pm 8\%$) follicles. Catalytic assays in ovarian homogenates revealed that CYP2E1 and CYP2B were functional. Only CYP2E1 activity was increased ($149 \pm 12\%$ above control; $p < 0.05$) by VCH dosing. The results demonstrate that mRNA and protein for CYP isoforms known to bioactivate VCH are expressed in the mouse ovary and are modulated by *in vivo* exposure to VCH and VCD. Interestingly, there is high expression of these isoforms in the interstitial cells. Thus, the ovary may contribute to ovotoxicity by promoting bioactivation of VCH to the toxic metabolite, VCD.

Rajapaksa and coworkers evaluated the role of ovarian CYP2E1 in VCH-induced ovotoxicity (Rajapaksa *et al.*, 2007). Ovaries from B6C3F1 mice, CYP2E1 wild-type (+/+) mice and null (-/-) mice were sampled at postnatal day 4. Then they were cultured for 15 days with VCD (30 μ M), or VCH-1,2-epoxide (125-1000 μ M). Female CYP2E1 +/+ and -/- mice (28 days of age) were administered i.p. daily for 15 days with VCH, VCH-1,2-epoxide, or VCD. Following culture or *in vivo* dosing, ovaries were histologically evaluated. In the *in vitro* study, VCD decreased ($p < 0.05$) primordial and primary follicles in ovaries from all three groups of mice. VCH-1,2-epoxide decreased ($p < 0.05$) primordial follicles in B6C3F1 and CYP2E1 +/+ ovaries, but not in CYP2E1 -/- ovaries in culture. VCH-1,2-epoxide did not affect primary follicles in any group of mouse ovaries. However, after *in vivo* exposure, primordial and primary follicles were reduced ($p < 0.05$) by VCD and VCH-1,2-epoxide in CYP2E1 +/+ and -/-. VCH reduced significantly primordial and primary follicles in CYP2E1 +/+ mice and in CYP2E1 +/+ and -/- mice, respectively. The data demonstrate that, whereas *in vitro* ovarian bioactivation of VCH or VCH-1,2-epoxide requires CYP2E1 enzyme, *in vivo* CYP2E1 plays a minimal role. Thus, the findings support that hepatic

metabolism dominates the contribution made by the ovary in bioactivation of VCH and VCH-1,2-epoxide to the ovotoxic metabolite, VCD.

Although the CYPs known to participate in VCH bioactivation are present in different cell types of the ovaries, potentially contributing to ovotoxicity by promoting bioactivation of VCH to the toxic metabolite, VCD, other studies tend to show that liver is the main organ for metabolism and bioactivation of VCH.

4.1.2 Human information

Smith and Sipes investigated the epoxidation of VCH in human hepatic microsomes (Smith and Sipes, 1991). Microsomes were prepared from livers from organ donors dying in traumatic accidents and from patients undergoing surgical resection of liver tumors (male and female). Twelve human liver microsomal samples (1.0 mg/mL microsomal protein) were incubated with VCH (1 mM) and 3,3,3-trichloropropene oxide, an inhibitor of epoxide hydrolase for 20 min. The major microsomal metabolite of VCH was VCH-1,2-epoxide. The rate of production of this metabolite ranged from 0.13 to 1.25 nmol/mg microsomal protein/min. VCH 7,8 epoxide was formed at rates approximately 6-fold lower than VCH-1,2-epoxide (< 0.01-0.21 nmol/mg microsomal protein/min. No differences between males and females were observed in the rate of hepatic microsomal VCH-1,2-epoxide formation. This was not investigated for the monoepoxide VCH-7,8-epoxide. The rate of VCH epoxidation by humans was lower than that from female mice and comparable to the VCH epoxidation rates with hepatic microsomes obtained from female rats (see results from Smith *et al.*, 1990c). These studies indicate that VCH-1,2-epoxide is the major epoxide of VCH formed by human hepatic microsomes, as in mice and rats. However, the metabolism of VCH proceeds at a slower rate by human microsomes compared with hepatic microsomes obtained from mice or rats. VCH 7,8 epoxide were not quantified in mice and rats in the previous study of Smith and coworkers, probably because of the shorter incubation time (5 min vs 20 min). Epoxide hydrolase activity toward VCH epoxides was present in human liver microsomes since an epoxide hydrolase inhibitor (3,3,3 trichloropropene oxide) was added to the system (also used in mice and rats microsomes in the previous study of Smith). These results should be taken with caution since CYP activities in microsomes prepared from human livers could be modified by possible exposure to drugs or environmental chemicals. Then, the rate of formation of VCH mono- and diepoxides could have been different from that observed in this study.

Additionally, Fontaine and co-workers have demonstrated that human CYP "Supersomes" (human CYP + P450 reductase + cytochrome b5) are able to catalyze VCH epoxidation, resulting in the formation of mono- and diepoxide metabolites of VCH (Fontaine *et al.*, 2001). CYP2E1 and CYP2B6 were capable of significantly converting VCH into VCH monoepoxide; whereas CYP2B6, CYP2A6, CYP2E1, and CYP3A4 were capable of catalyzing the epoxidation of the monoepoxides into the ultimate metabolite VCD.

4.1.3 Summary and discussion on toxicokinetics

The major route of excretion of VCH in mice and rats is urine (50-60% of the dose), the second route being expired air as expired organics. Elimination of VCH is slower in rats than in mice. The tissue with the highest concentration of VCH in both species was adipose tissue. VCH or its metabolites were found similarly in ovary from mice and rats.

The major monoepoxide VCH-1,2-epoxide was found in mice blood but not in rats. This could be explained by a higher rate of formation of this metabolite in mice, and/or by a more efficiently

detoxification by e.g. epoxide hydrolases in rats compared to mice. Monoepoxide metabolites are further epoxidized in VCD, a more potent ovary toxicant. Mono- and/or diepoxide metabolites are probably formed in the liver and are distributed in the ovary via circulating blood to exert their toxicities. However, CYP isoforms present in the mouse ovary are able to bioactivate VCH or the ovotoxic metabolite VCD.

After a repeated exposure, VCH is able to induce CYP involved in its own bioactivation in mice and rats. CYP (particularly, CYP2A and CYP2B) are involved in the epoxidation of VCH in liver microsomes from female mice. VCH epoxidation into VCH-1,2-epoxide can be catalyzed, by CYP2B forms in female Fischer 344 rats, but at a lower degree than in mice. Detoxication of VCH epoxides can be mediated by glutathione conjugation and/or hydrolysis by epoxide hydrolase.

Interestingly, it was demonstrated that human hepatic microsomes and human CYP “supersomes” were able to catalyze VCH epoxidation into monoepoxides and to epoxidize these latter in VCD, the ultimate metabolite, like rodent hepatocytes. CYP2E1 and CYP2B6 were found to significantly catalyze VCH into VCH monoepoxide; epoxidation of the monoepoxides into VCD was mediated by CYP2B6, CYP2A6, CYP2E1, and CYP3A4.

Female human microsomes demonstrated epoxidation of VCH at rates 13-fold and 2-fold less than those in mice and rats, respectively (Smith and Sipes, 1991), and total hepatic CYP per milligram of protein is significantly lower in humans than in rodents (Imaoka *et al.*, 1991; Shimada *et al.*, 1994). Interestingly, out of the eight human hepatic CYP isoforms tested in current studies (CYP1A2, CYP2A6, CYP2B6, CYP2C6, CYP2E1, CYP3A4, CYP4A11 and aromatase), CYP2E1 and CYP2B6, were the only isoforms that significantly catalyzed the epoxidation of VCH.

Previous experiments focused on the role of CYP2E1 in the epoxidation of VCH because it has been reported to metabolize the structurally related compounds styrene and 1,3-butadiene (Lieber, 1997; Niesma *et al.*, 1998; Fontaine *et al.*, 2001). Studies showed that, although hepatic microsomes from mice and rats pretreated with acetone showed increases in VCH-1,2-epoxide formation from VCH, hepatic microsomes from mice or rats pretreated with VCH for 5 or 10 days demonstrated no increases in CYP2E1 protein levels or activity (Fontaine *et al.*, 2001). Those data, combined with the data showing no differences in epoxidation of VCH or its monoepoxides in CYP2E1-deficient mouse hepatic microsomes compared with those of mice that do have CYP2E1, indicated that CYP2E1 is not an important isoform in the species-specific bioactivation of VCH. Current studies reconfirm this conclusion because neither VCH, VCH-1,2-epoxide, nor VCD pretreatment for 10 days affected CYP2E1 levels or activity in mice or rats.

Interestingly, although CYP2B6 and CYP2E1 were the only CYP isoforms that catalyzed VCH epoxidation in humans, CYP2B6, CYP2A6, CYP2E1, and CYP3A4 catalyzed the epoxidation of both monoepoxides to form the diepoxide. Although CYP3A4 is the major hepatic CYP isoform in humans, human liver CYP2A6 expression is relatively low (approximately 4%) (Cheng and Schenkman, 1982; Shimada *et al.*, 1994). However, since the rate of formation of VCH 1,2-epoxide was shown to be significantly limited in humans compared with the mouse or rat (Smith and Sipes, 1991), the possibility of VCH-1,2-epoxide or VCH-7,8-epoxide bioactivation to VCD by these particular enzymes could be low in livers from humans exposed to VCH. However, information regarding the rate of epoxidation of VCH monoepoxides in VCD in human hepatic microsomes is missing.

Nevertheless, comparisons of VCH metabolism and hepatic CYP induction demonstrate that CYP2A and CYP2B are important CYP isoforms in the species-dependent bioactivation in rodents and, therefore, ovotoxicity of VCH. The increased expression of CYP2A and CYP2B seen exclusively in the mouse appears to be due to repeated treatment with VCH or VCH-1,2-epoxide.

This indicates that, with repeated exposure to VCH, the mouse is exposed to a greater concentration of the ovotoxic metabolites via enhanced bioactivation. The rat is resistant to the ovotoxicity of VCH, at least in part, because the increases in CYP levels/activities do not occur following repeated exposure to VCH. It is not known if exposure of humans to VCH would result in elevated levels of CYP isoforms. Perhaps studies with cultures of human hepatocytes could help address this question.

Available literature indicates that ovotoxicity of VCH is due to the formation of the ultimate ovotoxicant metabolite VCD, which is carcinogenic to mice and rats (skin tumors in both species and ovary tumor in mice). Metabolisation of VCH into mono- and di-epoxides metabolites is a critical step in the outcome of ovotoxicity and tumors. As long as human hepatic microsomes and human CYP “supersomes” have been demonstrated to be able to metabolise VCH into VCD and that a range of human CYPs are able to catalyse the epoxidation of VCH, an *in vivo* metabolism (maybe slight) of VCH into VCD in humans cannot be ruled out. Moreover, no data are available to evaluate in which extent human CYPs levels/activities would be sensitive in VCH activation leading to increased VCD formation with chronic exposure to VCH, VCD or any other inducers of the specific CYPs involved in VCH metabolism in human.




Figure 1 . Metabolic pathway for 4-vinylcyclohexene. All of the reactions shown were studied as was the hydrolysis of 4-vinylcyclohexene diepoxide. Thickness of lines indicates the relative velocity of the reactions in liver, compared to other reactions in liver. Solid lines indicates the reaction rate for mouse liver, dashed line indicates the reaction rate for rat liver (Keller *et al.*, 1997)

4.2 Acute toxicity

Not evaluated in this dossier.

4.3 Specific target organ toxicity – single exposure (STOT SE)

Not evaluated in this dossier.

4.4 Irritation

Not evaluated in this dossier.

4.5 Corrosivity

Not evaluated in this dossier.

4.6 Sensitisation

Not evaluated in this dossier.

4.7 Repeated dose toxicity

Table 10: Summary table of relevant repeated dose toxicity studies

Method	Results	Remarks	Reference
Rat (Fischer 344) (28 days old) 10/sex/group 0, 10, 40, 80 mg/kg (in corn oil) VCD 0, 100, 400, 800 mg/kg (in corn oil) VCH 0, 42.5, 170, 340 mg/kg (in corn oil) VCH-1,2-epoxide or VCH-7,8-epoxide 30 days Intra-peritoneal administration	No detectable oocyte loss occurred in rats at the highest dose of 7.4 mmol/kg VCH. In this study, ED50 of VCD and VCH-1,2-epoxide were determined for comparison: ED50 VCD = 0.4 mmol/kg ED50 VCH-1,2-epoxide = 1.4 mmol/kg	VCH did not induce oocyte loss in female rats exposed for 30 days.	Smith <i>et al.</i> , 1990b
Mouse (B6C3F1) (28 days old) 0, 10, 40, 80 mg/kg (in corn oil) VCD 0, 100, 400, 800 mg/kg (in corn oil) VCH 0, 42.5, 170, 340 mg/kg (in corn oil) VCH-1,2-epoxide or VCH-7,8-epoxide	The dose of VCH which reduced the small oocyte count to 50% that of control (=ED50) was 2.7 mmol/kg In this study, ED50 of VCD, VCH-1,2-epoxide and VCH-7,8-epoxide were determined for comparison: ED50 VCD = 0.2 mmol/kg		Smith <i>et al.</i> , 1990b

CLH REPORT FOR [4 VINYLCHYCLOHEXENE]

<p>30 days Intra-peritoneal administration</p>	<p>ED50 VCH-1,2-epoxide = 0.5 mmol/kg ED50 VCH-7,8-epoxide = 0.7 mmol/kg</p>		
<p>Rat (Fischer 344) 10/sex/group 0, 50, 100, 200, 400 or 800 mg/kg bw/d (in corn oil by gavage) 13 weeks, 5 days/week</p>	<p>Reduced final body weights in male (≥ 400 mg/kg bw/d) and in female rats (800 mg/kg bw/d)</p> <p>Histopathologic effects:</p> <ul style="list-style-type: none"> - hyaline droplet degeneration of the proximal convoluted tubules of the kidney in dosed male rats (severity dose related). - Inflammation in the submucosa of the nonglandular portion of the stomach was seen in one male and three females in the 800 mg/kg groups; this acute lesion consisted of focal infiltration of neutrophils and diffuse edema in the gastric submucosa. 	<p>This study is considered as valid, although this is a range-finding study, and no ophthalmological examination, no haematology and no clinical biochemistry were performed</p> <p>NOAEL < 50 mg/kg bw/d (male) NOAEL = 200 mg/kg bw/d (female)</p>	<p>NTP, 1986</p>
<p>Mouse (B6C3F1) 10/sex/group 0, 75, 150, 300, 600, or 1200 mg/kg bw/d (in corn oil by gavage) 13 weeks, 5 days/week</p>	<p>Reduced final body weights in female mice receiving 600 mg/kg bw/d</p> <p>Mortality: 9/10 male mice at 1200 mg/kg bw/d and 2/10 and 4/10 female mice at 300 and 1200 mg/kg bw/d, respectively.</p> <p>Histopathological effects:</p> <ul style="list-style-type: none"> - Reduction (level not specified) in the number of primary follicles and mature graafian follicles in the ovaries of female mice exposed to 1200 mg/kg bw/d (ovaries from the lower dose groups were not similarly examined) - Mild, acute inflammation of the 	<p>This study is considered as valid, although this is a range-finding study, and no ophthalmological examination, no haematology and no clinical biochemistry were performed</p> <p>NOAEL = 150 mg/kg bw/d (female) NOAEL = 600 mg/kg bw/d (male)</p>	<p>NTP, 1986</p>

	<p>stomach was seen in the 1200 mg/kg groups in three males that died before the end of the study and in one female that lived to the end of the study.</p>		
<p>Rat (Sprague-Dawley) (male and female) 0, 250, 1000, and 1500 ppm (= 0, 1.11, 4.42, and 6.63 mg/L) by inhalation 13 weeks, 5 days/week, 6 hr/ day</p>	<p>Results:</p> <ul style="list-style-type: none"> - lethargy (1500 ppm) - decreased body weights (significant) in male rats exposed to 1500 ppm compared to controls - significantly lower body weight gains in male and female rats at 1500 ppm. - increased absolute and/or relative liver weights of male and female rats exposed to 1000 or 1500 ppm - increased relative kidney weight in male rats exposed to 1000 or 1500 ppm - increased accumulation of hyaline droplets in the kidneys of all dosed male rats (although compound- related, the droplets were not accompanied by cytotoxicity) - ovarian atrophy was noted in 2/10 female rats exposed to 1500 ppm VCH. Atrophy in these two rats was morphologically distinct from that seen in mice, in that the primary change was a decrease in the numbers of corpora lutea. Since the decrease on corpora lutea were only observed on 2 of 10 rats and was not observed on previously conducted studies with VCH, the authors considered this effect as spurious and not compound-related. However, in absence of 	<p>This study is considered as valid although there is no post-treatment period.</p> <p>NOAEL = 250 ppm</p> <p>In this inhalation study, ovarian atrophy (decrease in the numbers of corpora lutea) was observed in high-dose female rats.</p>	<p>Bevan, 1996</p>

	<p>historical control incidence and of VCH-induced ovarian toxicity in mice, this effect could not be totally disregarded.</p> <p>No effects on:</p> <ul style="list-style-type: none"> - haematological parameters - clinical chemistry evaluation - urinalysis 		
<p>Mouse (B6C3F1) (male/female) 0, 50, 250, or 1000 ppm (= 0, 0.22, 1.11, and 4.42 mg/L) by inhalation 13 weeks, 5 days/week, 6 hr/ day</p>	<p>Results:</p> <ul style="list-style-type: none"> - lethargy observed in the 1000 ppm VCH-exposed mice. - mortality: all male mice and 5/10 female mice on Test Days 11 or 12, at 1000 ppm (three additional female mice exposed to 1000 ppm VCH died prior to study completion). - ovarian atrophy (level not specified) in females exposed to 1000 ppm (5/10 vs 0/10 in the other groups). The ovarian atrophy was characterized as a severe reduction of all developmental stages of ovarian follicles. - Splenic atrophy in 1/10 females exposed to 50 ppm and 1/10 females exposed to 250 ppm (0/10 in control females) - Testicular atrophy: in 4/10 males exposed to 1000 ppm and 1/10 males exposed to 250 ppm (but the incidence in the control group was 4/10) - Thymic atrophy: in 3/10 males exposed to 1000 ppm and 1/10 males exposed to 250 ppm. This effect was not observed in control and low-dose male groups. This finding 	<p>This study is considered as valid although there is no post-treatment period.</p> <p>NOAEL = 250 ppm</p>	<p>Bevan, 1996</p>

	likely represents a secondary, stress-related effect of exposure to a lethal concentration of VCH.		
--	--	--	--

4.7.1 Non-human information

4.7.1.1 Repeated dose toxicity: oral

In an oral 13-week study in rats and mice, VCH induced decreased final body weights in male and female and hyaline droplet degeneration of the proximal convoluted tubules of the kidney in dosed male rats (severity dose related) (NTP, 1986). Increased incidence of mortality was observed in female and male mice. The number of primary follicles and mature graafian follicles was reduced in the ovaries of high-dose female mice (ovaries from the lower dose groups were not similarly examined). VCH induced inflammation of stomach in rats and mice

4.7.1.2 Repeated dose toxicity: inhalation

After exposure to VCH vapors by inhalation for 13 weeks, ovarian atrophy was observed in female mice (Bevan, 1996). Other effects were lethargy, decreased body weights, and increased incidence of mortality in high-dose mice. Rats were also exposed to VCH vapors. After an exposure period of 13 weeks, lethargy, reduced body weights (in males) and reduced body weight gains (in both sexes), increased liver weight in both male and female rats were observed. Male rats displayed an increased relative kidney weight in high-dose, probably related to an increased accumulation of hyaline droplets in the kidneys seen at all doses. Interestingly, ovarian atrophy was noted in some high-dose female rats. However, atrophy was morphologically distinct from that seen in mice, in that the change was a decrease in the numbers of corpora lutea. Since the decrease on corpora lutea were only observed on 2 of 10 rats and was not observed on previously conducted studies with VCH, the authors considered this effect as spurious and not compound-related. However, in absence of historical control incidence and due to VCH-induced ovarian toxicity in mice, this effect could not be totally disregarded.

4.7.1.3 Repeated dose toxicity: dermal

No dermal data have been reported for VCH.

4.7.1.4 Repeated dose toxicity: other routes

No data

4.7.1.5 Human information

No human data.

4.7.1.6 Other relevant information

Based on information displayed in IR/CSA, Section R.6.2.5.2, the metabolic pathway approach is usually reserved to some toxicological endpoints. Here, data concerning VCD are presented as

supportive evidence. However, the metabolism of VCH is not fast enough to base hazard identification only on studies conducted with that metabolite itself.

Table 11: Summary table of relevant repeated dose toxicity studies for VCD

Method	Results	Remarks	Reference
<p>Rat (Fischer 344) (28 days old)</p> <p>10/sex/group</p> <p>0, 10, 40, 80 mg/kg (in corn oil) VCD</p> <p>0, 100, 400, 800 mg/kg (in corn oil) VCH</p> <p>0, 42.5, 170, 340 mg/kg (in corn oil) VCH-1,2-epoxide or VCH-7,8-epoxide</p> <p>30 days</p> <p>Intra-peritoneal administration</p>	<p>The dose of VCD which reduced the small oocyte count to 50% that of control (=ED50) was 0.4 mmol/kg.</p> <p>In this study, ED50 of VCH and VCH-1,2-epoxide were determined for comparison: ED50 VCH > 7.4 mmol/kg ED50 VCH-1,2-epoxide = 1.4 mmol/kg</p>	<p>VCD is the most potent ovotoxicant in rats after an i.p. administration, compared to VCH and VCH monoepoxides.</p>	<p>Smith <i>et al.</i>, 1990b</p>
<p>Mouse (B6C3F1) (28 days old)</p> <p>0, 10, 40, 80 mg/kg (in corn oil) VCD</p> <p>0, 100, 400, 800 mg/kg (in corn oil) VCH</p> <p>0, 42.5, 170, 340 mg/kg (in corn oil) VCH-1,2-epoxide or VCH-7,8-epoxide</p> <p>30 days</p> <p>Intra-peritoneal administration</p>	<p>The dose of VCD which reduced the small oocyte count to 50% that of control (=ED50) was 0.2 mmol/kg</p> <p>In this study, ED50 of VCH, VCH-1,2-epoxide and VCH-7,8-epoxide were determined for comparison: ED50 VCH = 2.7 mmol/kg ED50 VCH-1,2-epoxide = 0.5 mmol/kg ED50 VCH-7,8-epoxide = 0.7 mmol/kg</p>	<p>VCD is the most potent ovotoxicant in mice after an i.p. administration, compared to VCH and VCH monoepoxides.</p> <p>VCD has a similar ovotoxic potency both in rats and mice.</p>	<p>Smith <i>et al.</i>, 1990b</p>
<p>Rat (Fischer 344)</p> <p>0, 3.75, 7.5, 15, 30, 60 mg/rat (in acetone)</p> <p>13 weeks, 5 days/week</p> <p>Dermal exposure</p>	<p>No mortality</p> <p>Reduced final body weights in the high-dose groups.</p> <p>Clinical signs at high dose: redness, scabs, and ulceration at the application site and burrowing behavior after dermal application.</p> <p>Histopathologic effects:</p> <ul style="list-style-type: none"> - Hyperplasia of the sebaceous glands and acanthosis (hyperplasia) and hyperkeratosis of the squamous epithelium were seen at the site of 	<p>NOAEL_{sys} = 30 mg/rat</p> <p>NOAEL_{local} < 15 mg/rat</p>	<p>NTP, 1989</p>

	application in all treated groups. The severity of the lesions was greatest at 60 mg/rat.		
<p>Mouse (B6C3F1)</p> <p>10/sex/group</p> <p>0, 0.625, 1.25, 2.5, 5, 10 mg/mouse (in acetone)</p> <p>13 weeks, 5 days/week</p> <p>Dermal exposure</p>	<p>No effect on mortality and on body weights.</p> <p>Increased relative liver (in all treated males and in the two highest dose group females) and kidney weights (from 1.25 and 2.5 in males and females, respectively).</p> <p>Histopathological effects:</p> <ul style="list-style-type: none"> - Compound-related lesions of the skin included sebaceous gland hyperplasia and acanthosis (hyperplasia) (8/10 males and 2/10 females that received 10 mg/mouse and 1/10 males that received 5 mg/mouse) and hyperkeratosis of the stratified squamous epithelium at the site of application (8/10 males and 8/10 females that received 10 mg/mouse and 5/10 males and 6/10 females that received 5 mg/ mouse). - Diffuse ovarian atrophy observed in all females that received 10 mg/mouse and in 4/10 females that received 5 mg/mouse. - Uterine atrophy in 2/10 females that received 10 mg/mouse 	<p>NOAEL_{sys} < 0.625 mg/ mouse (M)</p> <p>NOAEL_{sys} = 1.25 mg/ mouse (F)</p> <p>NOAEL_{local} = 2.5 mg/ mouse</p>	<p>NTP, 1989</p>

4.7.1.7 Summary and discussion of repeated dose toxicity

Repeated dose toxicity data are presented for information as they may provide relevant data for assessment of carcinogenicity and no classification is discussed and proposed for this endpoint.

VCH produced a similarity of effects whatever the route of exposure (by oral gavage or by inhalation). Indeed, inflammation of stomach, ovarian atrophy in female mice and hyaline droplet degeneration in male rats were observed via both routes of exposure. Other toxic effects were mortality in mice, and reduced body weight. Inflammation of stomach observed in rats and mice

exposed orally to VCH is probably due to the route of exposure (gavage) since these effects were not observed in the 13-week inhalation study.

It should be noted that ovaries were target organs in both rodent species in the 13-week inhalation study, but not in the 13-week oral toxicity (mice only) and in the oral carcinogenicity (mice only) studies. Histopathological findings were different since in mice severe reduction of all developmental stages of ovarian follicles was observed, whereas only the number of corpora lutea was decreased in female rats.

In the 13-week oral NTP study and the 13-week inhalation study, ovarian toxicity was observed in high-dose female mice only, concomitantly to increased mortality. Since no histopathological examination of ovaries from lower dose groups in the NTP study was performed (since this study was aimed to determine dose range for the carcinogenicity study), no information on VCH-induced ovarian toxicity at non lethal doses is available. Although examination of all treated animals was carried out in the 13-week inhalation, the spacing between mid- and high-dose groups could be too large to observe any effect on ovary. However, in a reproductive study (Grizzle *et al.*, 1994), number of primordial oocytes/follicles in F1 high-dose females was decreased by ca. 33%. No mortality occurred in these animals, but they displayed some toxicity effects: decreased body weight from PND 77-117 (8-9%), increased relative liver weight (ca. 8%) and increased feed consumption were observed in these animals (see section 4.12). But, once again, in this study, only F1 control and high-dose groups were examined.

It should be noted that VCD, the ultimate metabolite of VCH, is a very potent ovotoxicant both in rats and mice after an i.p. administration. It is assumed that the ovotoxicity of VCH is attributed to that of VCD and that the epoxidation of VCH in VCD is required to affect ovary. Moreover, based on the dose which reduced the small oocyte count to 50% that of control (=ED50), the potency of VCD to destroy small oocytes is similar in rats and mice when i.p. administered. Interestingly, VCD is not able to produce ovotoxicity in rats after a 13-week dermal exposure. This could be due to differences in distribution of VCD to the target tissue (i.e. ovary) or absorption/excretion rates compared to an i.p. administration.

4.8 Specific target organ toxicity (CLP Regulation) – repeated exposure (STOT RE)

Not evaluated in this dossier.

4.9 Germ cell mutagenicity (Mutagenicity)

Table 12: Summary table of relevant in vitro and in vivo mutagenicity studies

Method	Results	Remarks	Reference
<p>~ OECD guideline 471</p> <p><i>S. typhimurium</i> TA 1535, TA 1537, TA 98 and TA 100</p> <p>with and without metabolic activation (rat or hamster S9 fractions)</p> <p>Preincubation protocol</p> <p>0, 3.3, 10, 33, 100, 333, 1,000 µg/plate in DMSO</p>	<p>VCH did not produce increase in revertants in strains TA100, TA1535, TA1537, and TA98, with or without metabolic activation (rat or hamster S9), when tested according to the pre-incubation protocol. Doses up to 1000 µg per plate were used for each experiment, except without metabolic activation where the highest dose studied was 333 µg per plate (no explanation given in the NTP study report). Cytotoxicity was observed in TA100 strain incubated with 1000 µg per plate, with rat S9.</p>	<p>This study is considered as valid.</p> <p>VCH was not mutagenic in this bacterial reverse mutation assay.</p>	NTP, 1986
<p>~ OECD guideline 474</p> <p>Crl:CD BR (Sprague-Dawley) rats</p> <p>Male and female</p> <p>5/sex/group</p> <p>Inhalation study (vapors)</p> <p>2-day study: 0, 500, 1000, 2000 ppm VCH;</p> <p>13-week-study: 0, 250, 1000, 1500 ppm VCH</p> <p>2-day study: 6 h/ day on two consecutive days</p> <p>13-week study: 6 h/day, 5 days/week</p>	<p>VCH did not induce micronuclei in rats after a 2-day or a 13-week period exposure.</p> <p>Toxicity:</p> <p>2-day study:</p> <p>Clinical signs of toxicity were noted in rats and included decreased responsiveness to sound stimulus, inactivity, and narcosis/sleep induction during both exposures in each VCH treatment group. Animal arousal occurred within approximately 10 minutes after cessation of exposure. No clinical signs of toxicity were noted in rats prior to each exposure or during the recovery period. In rats, body weights for the 2000 ppm VCH-exposed group were significantly lowered at both the 24- and 48-h post-exposure time points compared to the controls (102 and 70%, respectively) .</p> <p>13-week study:</p> <p>There was no compound-related mortality in rats exposed to VCH. Clinical signs of toxicity were evident in male and female rats in all VCH-exposure groups. The most prevalent signs were lethargy, clear</p>	<p>This study is considered as valid.</p>	Bevan, 2001

	discharge from the mouth, and stained fur. Body weight gain for male rats exposed to 1000 ppm and 1500 ppm VCH were significantly lower from controls (12-15% reduction).		
<p>~ OECD guideline 474 B6C3F1/CrBR mice Male and female 5/sex/group Inhalation study (vapors) 2-day study: 0, 250, 500, 1000 ppm VCH; 13-week-study: 0, 50, 250, 1000 ppm VCH 2-day study: 6 h/ day on two consecutive days 13-week study: 6 h/day, 5 days/week</p>	<p>VCH did not induce micronuclei in mice after a 2-day or a 13-week period exposure.</p> <p>Toxicity: 2-day study: Clinical signs of toxicity were not observed in mice. Body weight gain for the 1000 ppm VCH-exposed male mice were significantly less than controls at the 24-h post-exposure timepoint. I</p> <p>13-week study: There was significant compound-related mortality in mice exposed to VCH. all of the male mice (10/10) and 5/10 female mice exposed to 1000 ppm VCH died on test days 11 or 12. Three additional high-dose females died prior to study completion. No clinical signs of toxicity were observed in the 250 ppm VCH-exposed mice; however, tremors and lethargy were observed in one male and two female 50 ppm VCH-exposed mice. Body weight gain for female mice exposed to 250 ppm VCH were significantly lower from controls (82% reduction)</p>	<p>No concurrent positive control substance administered to mice. No toxicity observed in female mice</p>	<p>Bevan, 2001</p>

4.9.1 Non-human information

4.9.1.1 In vitro data

In a bacterial reverse mutation assay (Ames test), did not produce increase in revertants in strains TA100, TA1535, TA1537, and TA98, with or without metabolic activation (rat or hamster S9), when tested according to the pre-incubation protocol. Doses up to 1000 µg per plate were used for each experiment, except without metabolic activation where the highest dose studied was 333 µg per plate (no explanation given in the NTP study report. Cytotoxicity was observed in TA100 strain incubated with 1000 µg per plate, with rat S9 (NTP, 1986).

4.9.1.2 In vivo data

Micronuclei were assessed in rats and mice exposed by inhalation to VCH vapors for two consecutive days or for 13 weeks (Bevan *et al.*, 1996). VCH concentrations were: 0, 250 (mice only), 500, 1000, 2000 (rats only) ppm VCH in the 2-day study; and 0, 50 (mice only), 250, 1000, 1500 (rats only) ppm VCH in the 13-week-study. VCH did not induce micronuclei in rats and mice. However, the results in mice are of limited value since no concurrent positive controls were used in the mice studies. Signs of toxicity and decreased body weight gain were observed in treated rats. Body weight gain was decreased in high-dose male mice in the 2-day study and increased mortality was observed in the high-dose groups in the 13-week study.

4.9.2 Human information

No human data.

4.9.3 Other relevant information

Table 13 Summary table of relevant mutagenicity studies for VCD

Method	Results	Remarks	Reference
Reverse mutation test <i>Salmonella typhimurium</i> TA100, TA1535, TA1537, TA98 strains	TA100, TA1535 and TA98: positive with/without S9 (from 50, 170 and 500 µg/mL, respectively) TA1537: equivocal without S9 and positive with S9, from 1700 µg/mL	/	NTP, 1989
Gene conversion <i>Saccharomyces cerevisiae</i>	Positive without S9 from 3500 µg/mL	Not tested with S9	IARC, 1994
Mitotic crossing-over <i>Saccharomyces cerevisiae</i>	Positive without S9 from 3500 µg/mL	Not tested with S9	IARC, 1994
Reverse mutation <i>Saccharomyces cerevisiae</i>	Positive without S9 from 3500 µg/mL	Not tested with S9	IARC, 1994
Micronucleus formation <i>Allium cepa</i>	Positive without S9 from 700 µg/mL	Not tested with S9	IARC, 1994
Micronucleus formation <i>Vicia faba</i>	Positive without S9 from 1400 µg/mL	Not tested with S9	IARC, 1994
Gene mutation Chinese hamster V79 lung cells, <i>hprt</i> locus	Positive without S9 from 140 µg/mL	Not tested with S9	IARC, 1994
Gene mutation Chinese hamster V79 lung cells, <i>hprt</i> locus	Positive without S9 from 700 µg/mL	Not tested with S9	IARC, 1994
Mouse L5178Y lymphoma cells	Positive from 25 µg/mL	Not tested with S9	NTP, 1989
Sister chromatid exchanges Chinese hamster ovary cells	Positive from 3.73 µg/mL without S9 and from 37.3 µg/mL with S9		NTP, 1989
Micronucleus formation Chinese hamster V79 lung cells	Negative without S9	Not tested with S9	IARC, 1994
Chromosome aberration assay Chinese hamster ovary cells	Positive from 37.8 µg/mL without S9 and from 447 µg/mL with S9	/	NTP, 1989

4.9.4 Summary and discussion of mutagenicity

The mutagenicity database for VCH is limited. VCH was not mutagenic to four strains of *Salmonella typhimurium* with or without metabolic activation. The S9 used comes from hamster or rats. Metabolism data have shown that rat is not a potent species for transforming VCH into VCD. Therefore, the added value of this test regarding VCH mutagenicity *in vitro* with metabolic activation is questioned. VCH did not increase micronucleus frequency in rats and mice (although

the results in mice are of a limited value). However, the diepoxide metabolite of VCH, VCD, was mutagenic to *Salmonella typhimurium* and to *Saccharomyces cerevisiae* (NTP, 1989; IARC, 1994). It also caused gene conversion and mitotic crossing-over in *S. cerevisiae*. Micronuclei were induced by the compound in cells of two plant species, *Allium cepa* and *Vicia faba*, but not in Chinese hamster V79 lung cells (without S9). VCD induced mutations at both the hprt and tk loci in cultured mammalian cells. In rodent cell lines, it induced sister chromatid exchange and chromosomal aberrations.

4.9.5 Comparison with criteria

4.9.6 Conclusions on classification and labelling

Information regarding mutagenicity are displayed as supporting evidence for the carcinogenicity endpoint due to the positive *in vitro* results of VCD. However, no classification is discussed and proposed for this endpoint for VCH.

4.10 Carcinogenicity

Table 14: Summary table of relevant carcinogenicity studies

Method	Results	Remarks	Reference
<p>Rat (Fischer 344) (7 weeks old)</p> <p>50 /sex/group</p> <p>103 weeks, 5 days per week</p> <p>0, 200 or 400 mg/kg bw/d (gavage, in corn oil)</p>	<p>Results:</p> <p>Mortality:</p> <ul style="list-style-type: none"> - male: control 17/50, low 37/50*, high 45/50* - female: control 10/50; low 22/50; high 36/50* <p>The survivals of the high- and low-dose male rats were significantly lower than that of the vehicle controls after week 5 (43 high-dose rats vs 49 control rats) and week 88 (26 low-dose rats vs 36 control rats), respectively. In female rats, the survivals of the high- and low-dose groups were significantly reduced after week 3 (42 high-dose rats vs 50 control rats) and week 102 (31 low-dose rats vs 41 control rats), respectively.</p> <p>Body weights: no effects, except for high dose males late in the study</p> <p>Neoplastic and non-neoplastic effects:</p> <ul style="list-style-type: none"> - slightly increased incidence of epithelial hyperplasia of the forestomach in males (1/50, 3/50, 5/47¹) - slightly increased incidence squamous-cell papillomas or carcinomas (combined) of the skin in high-dose males (control, 0/50; low-dose, 1/50; high-dose, 4/50*) - marginally increased incidence of adenomas or squamous-cell carcinomas (combined) of the clitoral gland in low dose female 	<p>Only two concentrations used</p> <p>Purity: 98%; impurities in two lots of test chemical included 0.01 % butylated hydroxytoluene in one and 0.005% tert-butylcatechol in the other, which had been added as inhibitors of peroxide formation</p> <p>Due to the high incidence of mortality in low and high-dose-rats, it is not possible to relate the increased incidence of adenomas or squamous-cell carcinomas (combined) of the clitoral gland and of squamous-cell papillomas or carcinomas (combined) of the skin to exposure to VCH. This study is considered as inadequate to evaluate the carcinogenic potential of VCH.</p>	<p>NTP, 1986</p> <p>Collins <i>et al.</i>, 1987</p>

	rats (survival more similar to control) (control, 1/50; low-dose, 5/50*; high-dose, 0/49)		
<p>Mouse (B6C3F1) (7 weeks old)</p> <p>50 /sex/group</p> <p>103 weeks, 5 days per week</p> <p>0, 200 or 400 mg/kg (gavage, in corn oil)</p>	<p>Results:</p> <p>Decreased body weights and mortality in both males (control, 13/50; low-dose, 11/50; high-dose, 43/50*) and females (10/50, 11/50, 33/50*, respectively) in the high-dose groups. Neither gross observations nor histopathologic evaluations revealed a specific cause of death in any of the dosed mouse groups.</p> <p>The survivals of the high dose male and female mice were significantly lower than that of the vehicle controls after week 29 and after week 32, respectively.</p> <p>Non-neoplastic effects:</p> <ul style="list-style-type: none"> - mild, acute inflammatory lesions and epithelial hyperplasia of the forestomach, especially in males (0/47, 7/50, 7/46) - increased incidence of a number of other non-neoplastic lesions: lung congestion, (high dose male and female) splenic red pulp atrophy in high dose males, congestion of the adrenal gland in high dose females, cytologic alteration of the adrenal cortex in low and high dose females <p>Neoplastic effects:</p> <p>Females:</p> <ul style="list-style-type: none"> - significant treatment-related increase in the incidence of: <ul style="list-style-type: none"> • granulosa-cell tumours or carcinomas of the ovary (terminal rates: control, 1/39 (3%); low-dose, 9/38* (24%); high-dose 7/16* 	<p>Only two concentrations used</p> <p>Purity: 98%; impurities in two lots of test chemical included 0.01 % butylated hydroxytoluene in one and 0.005% tert-butylcatechol in the other, which had been added as inhibitors of peroxide formation)</p> <p>Due to the extensive and early incidence of mortality in male mice, it is not possible to evaluate the carcinogenic potential of VCH. Nevertheless, survival is similar in control and treated female mice. Based on the significantly increased incidence of granulosa-cell tumours of the ovary and of mixed tumours composed of epithelial and granulosa cells of the ovary, VCH is considered as carcinogenic to female mice.</p>	<p>NTP, 1986</p> <p>Collins <i>et al.</i>, 1987</p>

	<p>(44%); and overall rates (adjustment for mortality): control, 1/49 (2%); low-dose, 10/48* (21%); high-dose, 13/47* (28%)²</p> <ul style="list-style-type: none"> • and of mixed benign tumours composed of epithelial and granulosa cells of the ovary (terminal rates: control, 0/39, low-dose, 24/38* (63%), high-dose, 4/16* (25%); and overall rates (adjustment for mortality): control, 0/49, low-dose, 25/48* (52%), and high-dose: 11/47* (23%)).² <p>(uncommon ovarian neoplasms: incidence in historical corn oil control female B6C3F1 mice: 1.2% = 12/1028 animals)</p> <p>- slight increase of adrenal gland adenoma in high-dose group (terminal rates: control, 0/40; low-dose; 3/39 (8%); high-dose: 2/17 (12%); overall rates (adjustment for mortality): control, 0/50; low-dose, 3/49 (6%); high-dose; 4/48* (8%))</p> <p>The incidences of granulosa-cell hyperplasia and tubular-cell hyperplasia of the ovary were also increased in treated females.</p> <p>Males :</p> <p>- increased incidences of malignant lymphomas and alveolar/bronchiolar adenomas or carcinomas (combined) of the lung seen in the males surviving to the end of the study (malignant lymphomas: 3/37 (8%); 5/39 (13%); 4/7 (57%); alveolo-bronchiolar adenomas or carcinomas [combined]: 3/37 (8%); 9/39* (23%); 3/7* (43%) (the extensive mortality seen in the high-dose male mice confounded the interpretation of these incidences). After adjustment for mortality (i.e. overall rates), the incidences of</p>		
--	--	--	--

	lymphoma (control, 4/50; low-dose, 7/50; high-dose, 5/50; p = 0.01; incidental tumour trend test; p = 0.001 incidental pairwise test for high dose versus control) and of alveolar-bronchiolar adenoma or carcinoma (control, 4/49; low-dose, 11/50; high-dose, 4/50; p = 0.047; incidental tumour trend test) were slightly increased in treated males		
--	---	--	--

* Statistically significant

1 The forestomachs from 47 high-dose male rats were examined.

2 With regard to terminal rates, they should normally be based on a total of 40 control, 39 low-dose, and 17 high-dose animals. However, ovary tissues from one female rat per group were not examined.

4.10.1 Non-human information

4.10.1.1 Carcinogenicity: oral

Carcinogenic studies were performed with mice and rats (NTP, 1986; Collins, 1987). Due to the poor survival of low and high-dose male, and high-dose female rats, it is not possible to evaluate the carcinogenic potential of VCH in rats. Increased mortality was observed in high-dose female and male mice. The most pronounced effect is the significant treatment-related increased incidence of granulosa-cell tumours of the ovary (overall rates: control, 2.0%; low-dose, 21%*; high-dose 28%*) and of mixed tumours composed of epithelial and granulosa cells of the ovary (0%, 52%*, 23%*, respectively) (uncommon ovarian neoplasms; incidence in NTP historical corn oil control female B6C3F1 mice: 1.2% = 12/1028 animals) in female mice.

4.10.1.2 Carcinogenicity: inhalation

No data available.

4.10.1.3 Carcinogenicity: dermal

No data available.

4.10.2 Human information

No human data.

4.10.3 Other relevant information

Table 15: Summary table of relevant carcinogenicity studies for VCD

Method	Results	Remarks	Reference
Rat (Fischer 344) (7-8 weeks old)	Mortality: - male: control 43/50, low	Only two concentrations used	NTP, 1989

CLH REPORT FOR [4 VINYLCHYCLOHEXENE]

<p>60 /sex/group 105 weeks, 5 days per week 0, 15, 30 mg/rat (in acetone) Dermal application</p>	<p>42/50, high 46/50 - female: control 23/50; low 27/50; high 35/50 Neoplastic and non-neoplastic effects: - increased incidences of acanthosis and sebaceous gland hypertrophy of skin from the scapula or back (M + F) - squamous cell papillomas (M) and squamous cell carcinomas (M + F) (squamous cell carcinomas: male: vehicle control, 0/50; low dose, 33/50; high dose, 36/50; female: 0/50; 16/50; 34/50). - increased incidences of basal cell adenomas or carcinomas (combined) (male: 0/50; 1/50; 6/50; female: 0/50; 3/50; 4/50).</p>		
<p>Mouse (B6C3F1) (8-9 weeks old) 60 /sex/group 103 weeks, 5 days per week 0, 2.5, 5, 10 mg/mouse (in acetone) Dermal application</p>	<p>Mortality: - male: control 12/50, low 15/50, mid 46/50*, high 50/50* (wk 85) - female: control 20/50; low 19/50, mid 35/50, high 50/50* (wk 83) Neoplastic and non-neoplastic effects: - acanthosis, hyperkeratosis, and necrotizing inflammation of the skin over the scapula or back. - squamous cell carcinomas (male: vehicle control, 0/50; low dose, 14/50; mid dose, 39/50; high dose, 42/50; female: 0/50; 6/50; 37/50; 41/50). - increased follicular atrophy and tubular hyperplasia of the ovary (atrophy: 12/50; 43/49; 42/49; 47/50; tubular hyperplasia: 5/50; 35/49; 38/49; 34/50). - benign or malignant granulosa cell tumors (0/50; 0/49; 7/49; 12/50)</p>		<p>NTP, 1989</p>

	<p>and benign mixed tumors (0/50; 0/49; 11/49; 6/50) in mid- and high-dose females.</p> <ul style="list-style-type: none"> - increased combined incidences of luteomas, granulosa cell tumors, benign mixed tumors, or malignant granulosa cell tumors in mid- and high-dose female mice (1/50; 0/49; 17/49; 18/50). - marginally increased incidences of alveolar bronchiolar adenomas or carcinomas (combined) in exposed female mice (4/50; 9/50; 11/50; 7/50). 		
--	--	--	--

4.10.4 Summary and discussion of carcinogenicity

In an oral carcinogenesis study (gavage) in mice and rats, VCH produced a significant treatment-related granulosa-cell and mixed tumors of the ovary and adrenal subcapsular tumors in female mice (after adjustment for mortality, incidences of granulosa-cell tumors or carcinomas were 2%, 21% and 28% (significant), and those of mixed tumors composed of epithelial and granulosa cells were 0%, 52% (significant), and 23% (significant), in control, low- and high-dose female mice, respectively). The ovarian tumors were considered as uncommon ovarian neoplasms since the incidence in historical corn oil control female B6C3F1 mice was 1.2% (= 12/1028 animals). Increased incidence of lymphoma and of lung tumours were observed in male mice, but it was not possible to determine the carcinogenic potential of VCH in male mice due to extensive and early mortality. Increased incidences of squamous-cell tumours of the skin in male rats and of clitoral gland tumours in female rats were found, but the studies were considered as inadequate due to poor survival in treated rats compared to controls. It should be noted that IARC has classified VCH as possibly carcinogenic to humans (Group 2B) (IARC, 1994).

In an oral 13-week study, a reduction in the number of primary follicles and mature graafian follicles in the ovaries was observed in female mice but not in female rats exposed to VCH. In an inhalation 13-week study, female mice exhibited also ovarian atrophy, characterized as a severe reduction of all developmental stages of ovarian follicles. Interestingly, in this inhalation study, ovarian atrophy (decrease in the numbers of corpora lutea) was observed in high-dose female rats. However, this effect was morphologically distinct from that seen in mice, in that the effect was a decrease in the numbers of corpora lutea.

Based on the review of ovarian toxicity and carcinogenicity in eight recent national toxicology program studies, Maronpot demonstrated a relationship between antecedent ovarian hypoplasia, atrophy, and hyperplasia, and subsequent ovarian neoplasia (Maronpot, 1987). In addition, he observed that pathologic changes in other tissues such as the adrenal glands and uterus were associated with the treatment-related ovarian changes (it should be reminded that congestion of the adrenal gland and slight increase of adrenal gland adenoma in high-dose female mice were observed in the oral carcinogenicity study). The findings were interpreted as indicating a relationship between

previous ovarian toxicity and subsequent ovarian neoplasia. Therefore, ovarian atrophy observed in female mice exposed to VCH is considered as an early event in VCH-induced ovarian carcinogenesis in mice.

It should be noted that VCH-diepoxide (VCD), the ultimate metabolite of VCH, is also able to produce ovarian tumors in mice and to cause ovotoxicity in mice (but also in rats, contrary to VCH). Indeed, VCD is classified by IARC as possibly carcinogenic to humans (Group 2B). It induced ovary tumors after dermal exposure in mice, and skin tumors in rats and mice at the site of application. It is genotoxic in many *in vitro* assays: it was mutagenic in *Salmonella typhimurium* (Ames test), in *Saccharomyces cerevisiae* (gene conversion and mitotic crossing-over), cultured mammalian cells (at both the *hprt* and *tk* loci), in rodent cell lines (sister chromatid exchange and chromosomal aberrations), and in two plant species, *Allium cepa* and *Vicia faba* (micronucleus test). VCD has a greater ovotoxicity potency in rodents than VCH after an i.p. administration and is able to destroy small oocytes both in rats and mice when administered by i.p. Interestingly, VCD is not able to produce ovotoxicity in rats after a 13-week or a 105-week dermal exposure. This could be due to differences in distribution of VCD to the target tissue (i.e. ovary) or absorption/excretion rates compared to an i.p. administration. Based on the fact that VCD is ovotoxic in rats by i.p. and that it is a critical step in ovarian tumor induction, VCD could be expected to cause ovary tumors in rats when orally or intraperitoneally administered. However, since no oral or i.p. carcinogenesis study in rats exposed to VCD has been located in the literature, it is not possible to prove this assumption. Moreover, VCD induced skin tumors in rats at the site of application. Since VCH was orally (and not dermally) administered to rats, it is not possible to know if VCH could produce local tumors like VCD.

It is remarkable that VCH and VCD are both ovotoxic (in mice, and in both mice and rats, respectively) and ovarian carcinogens in mice. In rodents, it was demonstrated that VCH is primarily submitted to epoxidation catalyzed by hepatic CYP to produce monoepoxides: the major metabolite being VCH-1,2-epoxide. Then VCH-1,2-epoxide undergoes an epoxidation to produce VCD. It has been demonstrated that VCH-induced ovarian tumors are dependent on the metabolism of VCH to the diepoxide metabolite, VCD, which is responsible for the destruction of oocytes, which is a critical step in the induction of ovary carcinogenesis. Detoxication of VCH epoxides can be mediated by glutathione conjugation and/or hydrolysis by epoxide hydrolase. However, mice produced the epoxide metabolites at a higher rate of formation, and they detoxified less efficiently the epoxides by e.g. epoxide hydrolases compared to rats. This could explain the difference in susceptibility to VCH-mediated ovotoxicity in mice and rats. In the oral carcinogenesis study, rats may be resistant to ovarian tumor induction by VCH because of the amount of VCH converted to epoxides is insufficient to produce oocyte destruction, or perhaps due to the poor survival in these animals, which do not allow to draw a firm conclusion for the carcinogenic effects of VCH in rats.

Since it was demonstrated that human hepatic microsomes and human CYPs are able to catalyse *in vitro* the epoxidation of VCH in mono- and diepoxide (VCD), it cannot be ruled out that this reaction could occur in women exposed to VCH. Although the study available regarding human metabolism of VCH seems to show that human is less potent to transform it into its monoepoxide, VCH-1,2-epoxide, than rat (and subsequently than mouse), some metabolism still occurs. Nevertheless, information about the levels of VCD formed *in vitro* in human hepatocytes is missing. Moreover, no data is available to evaluate in which extent human is sensitive to induction of the CYPs responsible for VCH metabolism, after repeated exposure to VCH. Therefore, we are not able to evaluate the potential chronic effect of VCH exposure in human.

Overall, the only valid study to assess carcinogenic effects of VCH is the oral carcinogenicity study in female mice exposed to VCH. The results observed with male mice and female and male rats could not be used to evaluate the hazard potential of VCH because of the poor survival in these

animals. However, based on the well-described mechanism by which ovarian tumors are produced (metabolisation of VCH in VCD and subsequent destruction of small oocytes), and on the fact that *in vitro* epoxidation of VCH was observed in human hepatocytes, it cannot be ruled out that this mechanism is relevant to human.

4.10.5 Comparison with criteria

Rationale for classification in Carc. 1B:

The CLP criteria for classification in Carc. 1B are as follows:

“Known or presumed human carcinogens

A substance is classified in Category 1 for carcinogenicity on the basis of epidemiological and/or animal data. A substance may be further distinguished as:

Category 1A, known to have carcinogenic potential for humans, classification is largely based on human evidence, or

Category 1B: Category 1B, presumed to have carcinogenic potential for humans, classification is largely based on animal evidence. The classification in Category 1A and 1B is based on strength of evidence together with additional considerations (see section 3.6.2.2). Such evidence may be derived from:

- *human studies that establish a causal relationship between human exposure to a substance and the development of cancer (known human carcinogen); or*
- *animal experiments for which there is sufficient (1) evidence to demonstrate animal carcinogenicity (presumed human carcinogen). In addition, on a case-by-case basis, scientific judgement may warrant a decision of presumed human carcinogenicity derived from studies showing limited evidence of carcinogenicity in humans together with limited evidence of carcinogenicity in experimental animals.”*

In the CLP, sufficient evidence of carcinogenicity is defined as when *“a causal relationship has been established between the agent and an increased incidence of malignant neoplasms or of an appropriate combination of benign and malignant neoplasms in (a) two or more species of animals or (b) two or more independent studies in one species carried out at different times or in different laboratories or under different protocols. An increased incidence of tumours in both sexes of a single species in a well-conducted study, ideally conducted under Good Laboratory Practices, can also provide sufficient evidence. A single study in one species and sex might be considered to provide sufficient evidence of carcinogenicity when malignant neoplasms occur to an unusual degree with regard to incidence, site, type of tumour or age at onset, or when there are strong findings of tumours at multiple sites;”*

Limited evidence of carcinogenicity is defined as when *“the data suggest a carcinogenic effect but are limited for making a definitive evaluation because, e.g. (a) the evidence of carcinogenicity is restricted to a single experiment; (b) there are unresolved questions regarding the adequacy of the design, conduct or interpretation of the studies; (c) the agent increases the incidence only of benign neoplasms or lesions of uncertain neoplastic potential; or (d) the evidence of carcinogenicity is restricted to studies that demonstrate only promoting activity in a narrow range of tissues or organs.”*

Based on experimental studies in rodents, a causal relationship between the oral exposure to VCH and the increased incidence of ovary tumors has been demonstrated in female mice. The incidences of mixed tumours composed of epithelial and granulosa cells is 0%, 52%, and 23%, in control, low- and high-dose female mice, respectively. It should be noted that these types of tumors are uncommon ovarian neoplasms, with an incidence in NTP historical corn oil control female B6C3F1

mice of 1.2%. The mechanism by which VCH induces ovary tumors is well understood. VCH is epoxidized in the ultimate metabolite VCD in rodents (with a greater extent in mice compared to rats). Then VCD exerts its ovotoxicity in destroying small oocytes, which is a critical step in the VCH-induced ovarian tumors.

According to IR/CSA (Section R.6.2.5.2), “data from hazard identification studies conducted with [...] primary metabolite itself can be used to identify hazards for the parent compound.”. Therefore, ovotoxicity and carcinogenicity data of VCD can be used to support the classification of VCH. VCD is a more potent ovotoxic in mice, compared to VCH. VCD is a more potent ovotoxic in mice, compared to VCH, and is ovotoxic in both mice and rats, unlike VCH which is not ovotoxic in rats in a 13-week toxicity study, probably because the amount of VCD formed in rats is not enough to exert ovarian toxicity. Moreover, a dermal carcinogenicity demonstrated that its ovotoxicity leads to ovary carcinogenesis in female mice like VCH. This type of tumors was not observed in rats (only local skin tumors were produced at the site of application). Ovotoxicity was demonstrated as an increase of cysts. However, since VCH is able to destroy small oocytes in female rats by i.p. administration and not after a dermal exposure, it could be expected that VCD produces ovotoxicity (and ovarian carcinogenicity) after i.p. administration but not after dermal exposure, probably because of differences in absorption/excretion/distribution of VCD via these routes. No oral or i.p. carcinogenesis study in rats exposed to VCH is available to us in order to confirm this assumption.

It is concluded that the data provided in the report provide **sufficient evidence** of carcinogenic effects of VCH. Moreover, human hepatocytes and human CYPs are able to epoxidize VCH in mono- and diepoxide metabolites *in vitro*. Therefore, there is no mechanistic evidence that could lead to think that these effects are not relevant for human.

A classification **Carc. 1B –H350** is therefore warranted (Carc. Cat. 2; R45 according to Directive 67/548/EEC). As no data are available by inhalation or dermal route, it is proposed not to specify route of exposure in the hazard statement.

Classification in Carc. 1A is not appropriate as it should be based on human data and no human data specific of VCH are available.

Classification in Carc. 2 is not appropriate as the mechanism of action for inducing ovary tumors is well defined (metabolisation of VCH in VCD, which destroys small oocytes, which is a critical step in the outcome of ovary tumor). Moreover it occurred at a high incidence whereas the historical control incidence is low (1.2%).

4.10.6 Conclusions on classification and labelling

Based on the increased incidence of ovarian tumors in female mice exposed to VCH for two years by gavage, a classification as **Carc. 1B – H350: May cause cancer** (Carc. Cat. 2; R45 according to Directive 67/548/EEC) is proposed for VCH, with no specific route of exposure added.

4.11 Toxicity for reproduction

Although results obtained in the 90-day toxicity study on mice show in females a reduction in the number of primary follicles and mature graafian follicles in the ovaries of female mice exposed to

1200 mg/kg bw/d (ovaries from the lower dose groups were not similarly examined) and that one of the metabolites (VCD) is a known ovotoxic, only one reprotoxicity study (Reproductive assessment by continuous breeding (RACB) protocol) is available in the literature. The RACB protocol is summarized to allow comparison with current OECD test guideline 416 (Figure 1). For the F₀ cohabitation and lactation phases, 100 male and 100 female Swiss (CD-1) mice, 11 weeks of age, were assigned to four dose groups as follows: control group, 40 males and 40 females, and each treatment group, 20 males and 20 females. The doses of VCH were 100, 250, and 500 mg/kg bw/day in corn oil, administered once daily per os. Body weights were taken once weekly for determining dose rate. Feed and water consumption were monitored during treatment weeks 1, 2, 5, 9, 13, and 18 (the last week for females only).

During Week 1 of exposure to VCH, animals were housed two per cage by sex by dose group. During Weeks 2 through 15 of exposure, animals were housed in breeding pairs within dose groups, and newborn litters were euthanized immediately after evaluation. Starting at Week 16 of exposure, the breeding pairs were separated, and F₀ females were allowed to deliver and rear the final litter until PND 21. Because no deleterious reproductive effects were observed during the 14-week cohabitation period, the F₀ crossover mating was not conducted, and a limited F₁ generation fertility assessment was conducted using only control and high-dose F₁ animals. All F₀ males were singly housed beginning at Week 16 and then euthanized without a necropsy during Week 17, and all F₀ females were euthanized without a necropsy shortly after their litters reached PND 21. Only pups from the control and high-dose groups were weaned. Treatments were administered to all F₀ animals until euthanization. Data collected during the F₀ cohabitation were date of delivery of each litter, number, sex and weight of pups per litter, number of litters per breeding pairs, PND 0 dam body weight, and feed, water, and body weight data during Weeks 1, 2, 5, 9, 12, and 18 (females). On PND 0, 4, 7, 14, and 21, surviving pups were counted, sexed and weighed for all dams delivering a litter after Week 16.

During the F₁ fertility assessment, the F₁ generation from the control and high-dose groups were housed two per cage by sex within dose beginning at weaning (PND 21). Oral dosing of VCH in corn oil was initiated on PND 22 (prior to direct oral dosing, possible indirect exposure to VCH may have occurred through the gametes, *in utero*, or during lactation). Dose rate was based upon weekly body weights. At 74 ± 10 days of age, 20 males and 20 females per dose group were housed as nonsibling breeding pairs for 7 days or until a vaginal copulatory plug was observed, whichever was sooner. Litter data resulting from the F₁ cohabitation were collected as described for the F₀ cohabitation. After delivery of the litters, vaginal smears were collected from F₁ females for 12 days. All F₁ parents were euthanized and necropsied. Feed and water consumption were measured during Weeks 1 (breeding), 2, 3, and 4 of the F₁ fertility assessment.

At the necropsy, the body, paired kidney with attached adrenal, and liver weights were collected for both sexes immediately following CO₂ asphyxiation. For males, rights testis, left testis with attached epididymis, right epididymis, prostate, and seminal vesicles with coagulating glands (glandular secretions not removed) were weighed at necropsy. Paired ovaries (with attached oviducts) and uterus (with upper half of vagina) were weighed in females. Evaluation of sperm parameters included right cauda epididymis sperm motility, concentration, and morphology, and homogenization-resistant right testis spermatid concentration. All tissues, except ovaries with attached oviducts, were fixed in 10% neutral buffer formalin. The left testis and epididymis were embedded in glycol methacrylate, sectioned at 2.5- μ m thickness, and stained with the hematoxylin/PAS. Paired ovaries with attached oviducts were fixed in Bouin/s fixative for 24 h and then transferred to 70% ethanol to await embedding into paraffin. Ovaries were serially sectioned at

6- μm thickness and every 20th section was mounted, stained with hematoxylin and eosin, and evaluated for the number of primary, growing, and antral follicles.




Figure 1 – VCH reproductive assessment by continuous breeding flow diagram (Grizzle *et al.*, 1994)

This study is only reported to support ovotoxicity of VCH in mice. Classification regarding reprotoxic effects is not proposed here. Nevertheless, we consider that more data should be gathered before evaluating in depth VCH reprotoxicity. Indeed, the screening study available confirms VCH effect on testicular sperm concentration and oocytes/follicles without apparently impacting fertility.

Table 16: Summary table of relevant reproductive toxicity studies

Method	Results	Remarks	Reference
<p>Mouse (CD-1 (ICR) BR outbred Swiss albino) (11 weeks old)</p> <p>20 /sex/treated group</p> <p>40/sex/control group</p> <p>0, 100, 250 or 500 mg/kg (gavage, in corn oil)</p> <p>Exposure period:</p> <p>F0 males: 14 weeks (from 11 weeks of age)</p> <p>F0 females: 20 weeks (from 11 weeks of age)</p> <p>F1 males: 14 weeks (from 22 days of age)</p> <p>F1 females: 16 weeks (from 22 days of age)</p>	<p>Results:</p> <p>F0:</p> <p>0, 100, 250 or 500 mg/kg:</p> <p>No effect on mortality, feed/water consumption and clinical signs.</p> <p>500 mg/kg</p> <ul style="list-style-type: none"> - Slightly decreased postpartum dam weight (F) <p>F1 :</p> <p>0, 100, 250 or 500 mg/kg: No effect on mortality, clinical signs and water consumption.</p> <p>500 mg/kg</p> <ul style="list-style-type: none"> - Decreased mean body weight from PND77-117 (males: PND 77, 31.51 g vs 34.07 g in controls, and PND 117, 32.79 g vs 35.24 g in controls; females: PND 77, 26.20 g vs 28.40 g in controls, and PND 117, 28.00 g vs 30.60 g in controls) Increased relative liver weight (males: 60.46 vs 55.59 mg/g bw in controls; females: 62.08 vs 57.52 mg/g bw in controls) - Increased feed consumption (M + F) - Increased epididymal sperm motility (but within the historical control range) (85.5% vs 68.9% in control group) - Decreased testicular sperm concentration (11.3x10⁴/mg testis tissue vs 13.6x10⁴/mg testis tissue in control group) - Decreased mean number of primordial oocytes/follicles (140.6 vs 208.9 per ovary), growing follicles (23.2 vs 51.2 per 	<p>Reproductive effects of VCH in mice were assessed via the continuous breeding (RACB) protocol.</p>	<p>Grizzle <i>et al.</i>, 1994</p>

	<p>ovary) and antral follicles (4.95 vs 7.40 per ovary)</p> <ul style="list-style-type: none"> - No effect on oestrus - No effect on mating index, fertility index, number of live pups per litter, number of F2 pups born alive, sex ratio 		
--	---	--	--

4.11.1 Effects on fertility

4.11.1.1 Non-human information

Reproductive effects of VCH were assessed in Swiss mice via the continuous breeding (RACB) protocol (Grizzle *et al.*, 1994). Mice were exposed to 0, 100, 250 or 500 mg/kg VCH in corn oil by gavage. The treatment did not induce changes in mortality, feed/water consumption and clinical signs in treated parental generation, but a slightly decreased postpartum weight was observed in dam treated with 500 mg/kg bw/d VCH. In F1 mice, VCH did not affect mortality, clinical signs and water consumption at any dose. However, the high-dose group displayed a decreased mean body weight from PND77-117, an increased relative liver weight and an increased feed consumption. In the presence of a slight toxicity (decreased body weight of dams (8%) and increased relative liver weight in F1 males and females), sperm count (85.5% vs 68.9% in control group, but within the historical control range) and number of oocytes [decreased mean number of primordial oocytes/follicles (140.6 vs 208.9 per ovary), growing follicles (23.2 vs 51.2 per ovary) and antral follicles (4.95 vs 7.40 per ovary)] were decreased, although reproductive capacity was not altered in F0 and in F1.

4.11.1.2 Human information

No data

4.11.2 Developmental toxicity

4.11.2.1 Non-human information

No data

4.11.2.2 Human information

No data

4.11.3 Other relevant information

No data

4.12 Other effects

Not evaluated in this dossier.

5 ENVIRONMENTAL HAZARD ASSESSMENT

Not evaluated in this dossier.

6 OTHER INFORMATION

Information considered in this report was collected by a literature search performed on PubMed up to July 2010. References regarding toxicity and metabolism of VCH that were not evaluated by IARC for their classification in 1994 are mentioned in the references by an exponent 2 in the list below.

As we have seen that VCH is planned for registration on November 30th, 2010, a consultation was performed by emailing concerned registrants in order to require the existing data they would like to be considered. No response was collected.

7 REFERENCES

4-vinyl-1-cyclohexene – National Library of Medicine HSDB Database, <http://toxnet.nlm.nih.gov/cgi-bin/sis/search/r?dbs+hsdb:@term+@DOCNO+2872>

Bevan C, Keller DA, Panepinto AS, Bentley KS. 2001. Effect of 4-vinylcyclohexene on micronucleus formation in the bone marrow of rats and mice. *Drug Chem Toxicol.* 24 (3): 273-285.²

Bevan C, Stadler JC, Elliott GS, Frame SR, Baldwin JK, Leung HW, Moran E, Panepinto AS. 1996. Subchronic toxicity of 4-vinylcyclohexene in rats and mice by inhalation exposure. *Fundam Appl Toxicol.* 32 (1): 1-10.²

Cannady EA, Dyer CA, Christian PJ, Sipes IG, Hoyer PB. 2003. Expression and activity of cytochromes P450 2E1, 2A, and 2B in the mouse ovary: the effect of 4-vinylcyclohexene and its diepoxide metabolite. *Toxicol Sci.* 73 (2): 423-430.²

ChemADVISOR, Inc. MSDS OHS30250, 2009

ChemIDplus Lite, physical properties, -
<http://chem.sis.nlm.nih.gov/chemidplus/jsp/common/PhysicalProperties.jsp?calledFrom=lite>

Cheng KC and Schenkman JB (1982) Purification and characterization of two constitutive forms of rat liver microsomal cytochrome P-450. *J Biol Chem* **257**:2378–2385.²

Doerr-Stevens JK, Liu J, Stevens GJ, Kraner JC, Fontaine SM, Halpert JR, Sipes IG. 1999. Induction of cytochrome P-450 enzymes after repeated exposure to 4-vinylcyclohexene in B6C3F1 mice. *Drug Metab Dispos.* 27 (2): 281-287.²

² This publication was not reviewed in the IARC monograph (IARC, 1994)

Fontaine SM, Hoyer PB, Halpert JR, Sipes IG. 2001. Role of induction of specific hepatic cytochrome P450 isoforms in epoxidation of 4-vinylcyclohexene. *Drug Metab Dispos.* 29 (9): 1236-1242.²

Giannarini C, Citti L, Gervasi PG, Turchi G. 1981. Effects of 4-vinylcyclohexene and its main oxirane metabolite on mouse hepatic microsomal enzymes and glutathione levels. *Toxicol Lett.* 8 (1-2): 115-121.²

Grizzle TB, George JD, Fail PA, Seely JC, Heindel JJ. 1994. Reproductive effects of 4-vinylcyclohexene in Swiss mice assessed by a continuous breeding protocol. *Fundamental and Applied Toxicology.* 22: 122-129.

Handbook of chemistry and physics 2005-2006

IARC (1994). 4-Vinylcyclohexene. In *IARC Monographs on the Evaluation of Carcinogenic Risks of Chemicals to Humans: Some Industrial Chemicals*, Vol. 60, pp. 347-359. International Agency for Research on Cancer, Lyon, France.

IARC (1994). 4-Vinylcyclohexene diepoxide. In *IARC Monographs on the Evaluation of Carcinogenic Risks of Chemicals to Humans: Some Industrial Chemicals*, Vol. 60, pp. 361-373. International Agency for Research on Cancer, Lyon, France.

Imaoka S, Fujita S and Funae Y. 1991. Age-dependent expression of cytochrome P-450 in rat liver. *Biochim Biophys Acta.* 1097: 187-192.²

International Chemical Safety Card, NIOSH, 1995

IUCLID Dataset, prepared by Experien Health Sciences Inc. 10/07/2006

Keller DA, Carpenter SC, Cagen SZ, Reitman FA. 1997. In vitro metabolism of 4-vinylcyclohexene in rat and mouse liver, lung, and ovary. *Toxicol Appl Pharmacol.* 144 (1): 36-44.²

Lieber CS. 1997. Cytochrome P-4502E1: its physiological and pathological role. *Physiol Rev.* 77: 517-544.²

Maronpot RR. 1987. Ovarian toxicity and carcinogenicity in eight recent National Toxicology Program studies. *Environ Health Perspect.* 73: 125-130.

Nieusma JL, Claffey DJ, Koop DR, Chen W, Peter RM, Nelson SD, Ruth JA and Ross D. 1998. Oxidation of 1,3-butadiene to (R)- and (S)-butadiene monoxide by purified recombinant cytochrome P450 2E1 from rabbit, rat, and human. *Toxicol Lett.* 95: 123-129.²

NTP Technical Report on the Toxicology and carcinogenesis studies of 4-vinylcyclohexene (CAS N°. 100-40-3) in F344/N rats and B6C3F₁ mice (gavage studies), US National Toxicology Program, NTP TR 303, August 1986.

NTP Technical Report on the Toxicology and carcinogenesis studies of 4-vinyl-1-cyclohexene diepoxide (CAS N°. 106-87-6) in F344/N rats and B6C3F₁ mice (dermal studies), US National Toxicology Program, NTP TR 362, November 1989.

Rajapaksa KS, Cannady EA, Sipes IG, Hoyer PB. 2007. Involvement of CYP 2E1 enzyme in ovotoxicity caused by 4-vinylcyclohexene and its metabolites. *Toxicol Appl Pharmacol.* 221 (2): 215-221.²

Shimada T, Yamazaki H, Mimura M, Inui Y and Guengerich FP. 1994. Interindividual variations in human liver cytochrome P450 enzymes involved in the oxidation of drugs, carcinogens, and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther.* 270: 414-423.²

Smith BJ, Carter DE, Sipes IG. 1990a. Comparison of the disposition and *in vitro* metabolism of 4-vinylcyclohexene in the female mouse and rat. *Toxicol Appl Pharmacol.* 105 (3): 364-371.

Smith BJ, Mattison DR and Sipes IG. 1990b. The role of epoxidation in 4-vinylcyclohexene-induced ovarian toxicity. *Toxicol Appl Pharmacol.* 105 (3): 372-381.

Smith BJ, Sipes IG, Stevens JC, Halpert JR. 1990c. The biochemical basis for the species difference in hepatic microsomal 4-vinylcyclohexene epoxidation between female mice and rats. *Carcinogenesis.* 11 (11): 1951-1957.²

Smith BJ, Sipes IG. 1991. Epoxidation of 4-vinylcyclohexene by human hepatic microsomes. *Toxicol Appl Pharmacol.* 109 (2): 367-371.

8 ANNEXES