

CLH report

Proposal for Harmonised Classification and Labelling

**Based on Regulation (EC) No 1272/2008 (CLP Regulation),
Annex VI, Part 2**

Substance Name:

Dodecyl methacrylate

EC Number: 205-570-6

CAS Number: 142-90-5

Index Number: 607-247-00-9

Contact details for dossier submitter:

BAuA
Federal Institute for Occupational Safety and Health
Federal Office for Chemicals
Friedrich-Henkel-Weg 1-25
D-44149 Dortmund, Germany

Dossier prepared by:
Industry in accordance with Article 37(6) of CLP Regulation

Version number: 2.0

Date: October 2016

CONTENTS

Part A

1	PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING	4
1.1	SUBSTANCE	4
1.2	HARMONISED CLASSIFICATION AND LABELLING PROPOSAL	4
1.3	PROPOSED HARMONISED CLASSIFICATION AND LABELLING BASED ON CLP REGULATION	5
1.4	RELATION TO COMMON GROUP ENTRY WITH INDEX NO 607-134-00-4 IN ANNEX VI	6
2	BACKGROUND TO THE CLH PROPOSAL	7
2.1	HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING	7
	SUBSTANCES USED IN ANALOGY WITH DODECYL METHACRYLATE	7
2.2	SHORT SUMMARY OF THE SCIENTIFIC JUSTIFICATION FOR THE CLH PROPOSAL	8
2.3	CURRENT HARMONISED CLASSIFICATION AND LABELLING	8
2.3.1	<i>Current classification and labelling in Annex VI, Table 3.1 in the CLP Regulation</i>	8
2.4	CURRENT SELF-CLASSIFICATION AND LABELLING	8
2.4.1	<i>Current self-classification and labelling based on the CLP Regulation criteria</i>	8
3	JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL	9

Part B

1	IDENTITY OF THE SUBSTANCE	10
1.1	NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE	10
1.2	COMPOSITION OF THE SUBSTANCE	10
1.3	PHYSICO-CHEMICAL PROPERTIES	11
2	MANUFACTURE AND USES	13
2.1	MANUFACTURE	13
2.2	IDENTIFIED USES	13
3	CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES	14
4	HUMAN HEALTH HAZARD ASSESSMENT	14
4.1	TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)	14
4.1.1	<i>Non-human information</i>	14
4.1.2	<i>Human information</i>	17
4.1.3	<i>Summary and discussion on toxicokinetics</i>	17
4.2	IRRITATION	18
4.2.1	<i>Skin irritation</i>	18
4.2.1.1	Non-human information	20
4.2.1.2	Human information	22
4.2.1.3	Summary and discussion of skin irritation	22
4.2.1.4	Comparison with criteria	22
4.2.1.5	Conclusions on classification and labelling	22
4.2.2	<i>Eye irritation</i>	23
4.2.2.1	Non-human information	25
4.2.2.2	Human Information	25
4.2.2.3	Summary and discussion of eye irritation	25
4.2.2.4	Comparison with criteria	26
4.2.2.5	Conclusion on classification and labelling	26
4.2.3	<i>Respiratory tract irritation</i>	26
4.3	CORROSIVITY	26
4.4	SPECIFIC TARGET ORGAN TOXICITY – SINGLE EXPOSURE (STOT SE)	26
5	ENVIRONMENTAL HAZARD ASSESSMENT	27
5.2	DEGRADATION	27
5.2.1	<i>Stability</i>	27
5.2.2	<i>Biodegradation</i>	27

5.2.2.1	Screening tests	27
5.2.3	<i>Summary and discussion of degradation</i>	28
5.3	BIOACCUMULATION	28
5.3.1	<i>Aquatic Bioaccumulation</i>	28
5.3.2	<i>Summary and discussion of aquatic bioaccumulation</i>	30
5.4	AQUATIC TOXICITY	30
5.4.1	<i>Fish</i>	32
5.4.1.1	Short-term toxicity to fish	32
5.4.1.2	Long-term toxicity to fish	32
5.4.2	<i>Aquatic invertebrates</i>	32
5.4.2.1	Short-term toxicity to aquatic invertebrates	32
5.4.2.2	Long-term toxicity to aquatic invertebrates	33
5.4.3	<i>Algae and aquatic plants</i>	34
5.5	COMPARISON WITH CRITERIA FOR ENVIRONMENTAL HAZARDS (SECTIONS 5.1 – 5.4).....	35
5.6	CONCLUSIONS ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS (SECTIONS 5.1 – 5.4)	36
6	REFERENCES.....	37

Part A.

1 PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING

1.1 Substance

Table 1: Substance identity

Substance name:	Dodecyl methacrylate
EC number:	205-570-6
CAS number:	142-90-5
Annex VI Index number:	607-247-00-9
Degree of purity:	≥ 80 %
Impurities:	
Stabilizer	

1.2 Harmonised classification and labelling proposal

Table 2: The current Annex VI entry and the proposed harmonised classification

	CLP Regulation
Current entry in Annex VI, CLP Regulation	Skin Irrit. 2, H315 Eye Irrit. 2, H319 STOT SE 3, H335 C ≥ 10 % Aquatic Acute 1, H400 Aquatic Chronic 1, H410
Current proposal for consideration by RAC	Deletion of: Skin Irrit. 2, H315 Eye Irrit. 2, H319 STOT SE 3, H335 Aquatic Acute 1, H400 Aquatic Chronic 1, H410
Resulting harmonised classification (future entry in Annex VI, CLP Regulation)	none

Table 3: Proposed modification of Annex VI entry No. 607-134-00-4

Index No. 607-134-00-4	Wording of the international chemical identifier
Current entry in Annex VI, CLP Regulation	monoalkyl or monoaryl or monoalkyaryl esters of methacrylic acid with the exception of those specified elsewhere in this Annex
Proposed modified entry in Annex VI, CLP Regulation	monoalkyl or monoaryl or monoalkyaryl esters of methacrylic acid with the exception of those specified elsewhere in this Annex and dodecyl methacrylate

1.3 Proposed harmonised classification and labelling based on CLP Regulation

Table 4: Proposed classification according to the CLP Regulation

CLP Annex I ref	Hazard class	Proposed classification	Proposed SCLs and/or M-factors	Current classification ¹⁾	Reason for no classification ²⁾
2.1.	Explosives				
2.2.	Flammable gases				
2.3.	Flammable aerosols				
2.4.	Oxidising gases				
2.5.	Gases under pressure				
2.6.	Flammable liquids				
2.7.	Flammable solids				
2.8.	Self-reactive substances and mixtures				
2.9.	Pyrophoric liquids				
2.10.	Pyrophoric solids				
2.11.	Self-heating substances and mixtures				
2.12.	Substances and mixtures which in contact with water emit flammable gases				
2.13.	Oxidising liquids				
2.14.	Oxidising solids				
2.15.	Organic peroxides				
2.16.	Substance and mixtures corrosive to metals				
3.1.	Acute toxicity - oral				
	Acute toxicity - dermal				
	Acute toxicity - inhalation				
3.2.	Skin corrosion / irritation	none		Skin Irrit. 2 H315	
3.3.	Serious eye damage / eye irritation	none		Eye Irrit. 2 H319	
3.4.	Respiratory sensitisation				
3.4.	Skin sensitisation				
3.5.	Germ cell mutagenicity				
3.6.	Carcinogenicity				
3.7.	Reproductive toxicity				

CLP Annex I ref	Hazard class	Proposed classification	Proposed SCLs and/or M-factors	Current classification ¹⁾	Reason for no classification ²⁾
3.8.	Specific target organ toxicity –single exposure	none		STOT SE 3 H335; C ≥ 10 %	
3.9.	Specific target organ toxicity – repeated exposure				
3.10.	Aspiration hazard				
4.1.	Hazardous to the aquatic environment	none		Aquatic Acute 1 H400 Aquatic Chronic 1 H410	
5.1.	Hazardous to the ozone layer				

¹⁾ Including specific concentration limits (SCLs) and M-factors

²⁾ Data lacking, inconclusive, or conclusive but not sufficient for classification

Labelling: **Signal word:** none

Hazard statements: none

Precautionary statements: none

Proposed notes assigned to an entry: none

1.4 Relation to common group entry with Index No 607-134-00-4 in Annex VI

The common group entry with the Index No. 607-134-00-4 is an umbrella entry for certain esters of methacrylic acid. The international chemical identifier reads as “monoalkyl or monoaryl or monoalkyaryl esters of methacrylic acid with the exception of those specified elsewhere in this Annex” and it classifies all substances encompassed by this definition as Skin Irrit. 2, Eye Irrit 2 and STOT SE 3 H335.

Along with several other esters of methacrylic acid dodecyl methacrylate is exempted from this entry by the fact that it currently has its own entry in Annex VI. With the proposed modification of the current classification of dodecyl methacrylate however the current Annex VI entry would effectively be removed and therefore removing the current exemption at the same time. The deletion of the Annex VI entry for dodecyl methacrylate would therefore only affect the environmental classification of the substance, as the classification for the human health endpoints would be reinstated by the group entry.

It is therefore proposed to simultaneously change the wording of the international chemical identifier of the aforementioned group entry 607-134-00-4 by appending the words “and dodecyl methacrylate” to explicitly exempt the substance from the scope of the group entry.

2 BACKGROUND TO THE CLH PROPOSAL

2.1 History of the previous classification and labelling

Dodecyl methacrylate was primarily classified and labelled by authorities with Xi, R 36/37/38, S26, 28, 60 and adopted of Annex I of Directive 67/548/EEC, Index No. 607-134-00-4 (monoalkyl or monoaryl or monoalkyaryl esters of methacrylic acid with the exception of those specified elsewhere in this Annex). This group classification is not based on data of individual member substances.

In 1995, the Methacrylate Producers Association (MPA), Washington, submitted preliminary results from an algal toxicity study in accordance with TSCA 8e to the coordinator of the Office of Pollution Prevention and Toxics at the Environmental Protection Agency (EPA), Washington DC and submitted in January 1996 the concerning study to EPA.

On this base ECB amended the classification of Dodecyl methacrylate with N, R50/53 which was adopted in 2004 in the 29th ATP to the DSD (Annex I of Directive 67/548/EEC, Index No. 607-247-00-9, R36/37/38, N, R50/53, S26, 28, 60, 61) after the introduction of the environmental endpoints into the classification criteria.

Studies on algal toxicity were repeated and showed that the study, which induced the environmental classification, was invalid.

A first EU classification and labelling dossier was submitted to the German competent authority (BAuA) in 2005. In January 2007 deletion of environmental classification was discussed and approved by the Technical Committee on Classification and Labelling of Dangerous Substances (TC C&L) (ECBI/08/07 Rev. 2), but not implemented.

With implementation of the CLP regulation the substance was classified and labelled as Skin Irrit. 2 (H315), Eye Irrit. 2 (H319), STOT SE 3 (H335), Aquatic Acute 1 (H400) and Aquatic Chronic 1, (H410).

Substances used in analogy with dodecyl methacrylate

Pure dodecyl alcohol (dodecanol, lauryl alcohol) is used only on a small scale to produce dodecyl methacrylate. In the large-scale production of long-chain aliphatic methacrylate esters technical mixtures are used of fatty (long-chain aliphatic) alcohols of natural or synthetic origin. As these substances are of main interest on the market, several toxicological studies are available with mixtures of long-chain methacrylates containing dodecyl methacrylate and were used in this CLH dossier.

Table 5: Physico chemical properties of the substances used in studies in this classification dossier:

Substance name	CAS-No	Molecular formula	MW	Log Pow	Water solubility [mg/l]
2-Ethylhexyl methacrylate	688-84-6	C ₁₂ H ₂₂ O ₂	191	5.59 ^a	3.07 ^a
Dodecyl methacrylate	142-90-5	C ₁₆ H ₃₀ O ₂	254	6.68 ^b	< 0.001 ^a
Tridecyl methacrylate	2495-25-2	C ₁₇ H ₃₂ O ₂	268	7.17 ^b	0.01409 ^c
Isotridecyl methacrylate	94247-05-9	C ₁₇ H ₃₂ O ₂	268	7.09 ^b	0.01628 ^c
Tetradecyl methacrylate	2549-53-3	C ₁₈ H ₃₄ O ₂	282	7.66 ^b	0.004461 ^c
Pentadecyl methacrylate	6140-74-5	C ₁₉ H ₃₆ O ₂	297	8.15 ^b	0.001409 ^c
Hexadecyl methacrylate	2495-27-4	C ₂₀ H ₃₈ O ₂	311	8.64 ^b	0.0004442 ^b
Octadecyl methacrylate	32360-05-7	C ₂₂ H ₄₂ O ₂	339	9.62 ^b	0.0000437 ^b

^a Measured data

^b Calculated data, ^c Calculated data are higher than predicted from experimental data with dodecyl methacrylate

2.2 Short summary of the scientific justification for the CLH proposal

Data from the REACH registration were taken as a basis for this CLH proposal.

Based on the available/presented data the classification/labelling with Skin Irrit. 2 (H315), Eye Irrit. 2 (H319), STOT SE 3 (H335), Aquatic Acute 1 (H400) and Aquatic Chronic 1 (H410) is deemed to be not justified.

2.3 Current harmonised classification and labelling

2.3.1 Current classification and labelling in Annex VI, Table 3.1 in the CLP Regulation

Table 6: Current entry in Annex VI, Table 3.1 in the CLP Regulation (Index-No.: 607-247-00-9)

Classification		Labelling			Specific Conc. Limits, M-factors	Notes
Hazard Class and Category Code(s)	Hazard Statement Code(s)	Pictogram, Signal Word Code(s)	Hazard statement Code(s)	Suppl. Hazard statement Code(s)		
Eye Irrit. 2	H319	GHS07	H319	-	STOT SE 3: C ≥ 10 %	
STOT SE 3	H335	GHS09	H335			
Skin Irrit. 2	H315	Wng	H315			
Aquatic Acute 1	H400		H400			
Aquatic Chronic 1	H410		H410			

2.4 Current self-classification and labelling

2.4.1 Current self-classification and labelling based on the CLP Regulation criteria

The following industry self-classification(s) and labelling are publically available in the ECHA C&L Inventory (October 2016).




Figure 1: C&L notifications submitted to ECHA (October 2016, www.echa.eu)

3 JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL

For Dodecyl methacrylate a harmonised classification had been developed under 67/548/EC. Assessments performed under the OECD chemicals programme and in order to achieve a registration under REACH indicated, that according to new data the existing classification no longer reflects the criteria in Annex 1 of the CLP regulation (Regulation (EC) No 1272/2008). This document represents an update of the harmonised classification according to the currently available and most reliable information following a comprehensive assessment of the key data on behalf of the 2010 registrants under REACH.

Part B.

SCIENTIFIC EVALUATION OF THE DATA


1 IDENTITY OF THE SUBSTANCE

1.1 Name and other identifiers of the substance

Table 7: Substance identity

EC number:	205-570-6
EC name:	Dodecyl methacrylate
CAS number (EC inventory):	142-90-5
CAS number:	142-90-5
CAS name:	2-Propenoic acid, 2-methyl-, dodecyl ester
IUPAC name:	Dodecyl methacrylate
CLP Annex VI Index number:	607-247-00-9
Molecular formula:	C ₁₆ H ₃₀ O ₂
Molecular weight range:	254.42 g/mol

Structural formula:



1.2 Composition of the substance

Table 8: Constituents (non-confidential information)

Constituent	Typical concentration	Concentration range	Remarks
Dodecyl methacrylate	Ca. 99.3 %	95-100 %	

For further information on the composition of the substance refer to the IUCLID file.

1.3 Physico-chemical properties

Table 9: Summary of physico - chemical properties

Property	Value	Reference	Comment (e.g. measured or estimated)
State of the substance at 20°C and 101,3 kPa	liquid		observation
Melting/freezing point	Melting Point: -7 °C (atmospheric pressure (1013 hPa) assumed)	Brandes and Möller (2003),	Measured, handbook data
Boiling point	307 – 318 °C	Brandes , Möller (2003), Nabert, Schön, Redeker (2005)	Measured, handbook data
Relative density	0.87 g/cm ³	Brandes and Möller (2003),	Measured, handbook data
Vapour pressure	0.06 Pa at 20 °C	Rehberg , Fisher (1948),	Measured, dynamic method, extrapolated Clausius Clapeyron equation
Surface tension	waiving		In accordance with column 2 of REACH Annex VII, the surface tension of the substance does not need to be tested because due to its chemical structure, no surface activity is predicted.
Water solubility	< 1 µg/L at 25 °C	Dr. U. Noack Laboratories (2004)	Measured acc.US-EPA OPPTS 830.7860, column elution method
Partition coefficient n-octanol/water	LogPOW 6.68	Syracuse research Corporation (2000)	Calculated, KOWWIN™ v1.67 in EPI web 4.0
Flash point	> 110 °C	Brandes and Möller (2003),	Measured, handbook data
Flammability	waiving	BAM (2013)	Flammability upon ignition (solids, gases): Testing can be waived, substance is a liquid. Flammability in contact with water: The classification procedure needs not to be applied because the substance does not contain metals or metalloids. Pyrophoric properties: The classification procedure needs not to be applied because the substance is known to be stable into contact with air at room temperature for prolonged periods of time (days).
Explosive properties	waiving	BAM (2013)	The classification procedure needs not to be applied because there are no chemical groups associated with explosive properties present in the molecule.

CLH REPORT DODECYL METHACRYLATE

Self-ignition temperature	295 °C @ 1003 hPa	AQura GmbH (2008)	Measured acc. DIN 51794
Oxidising properties	waiving	BAM (2013)	The study does not need to be conducted for flammable liquids.
Granulometry	waiving		The substance is a liquid at 20°C. In accordance with column 2 of REACH Annex VII, the particle size distribution (Granulometry) study does not need to be performed as the substance is marketed or used in a non solid or granular form.
Stability in organic solvents and identity of relevant degradation products	waiving		In accordance with REACH annex XI, the study was not conducted because it is not critical
Dissociation constant	waiving		In accordance with REACH annex XI, the study was not conducted as the test substance does not dissociate based on structural alerts
Viscosity	6.24 mm ² /s @ 20 °C 3.74 mm ² /s @ 40 °C	Evonik RohMax Additives GmbH (2008)	measured acc. DIN 51 562, read across of 2-propenoic acid, 2-methyl-, C12-16-alkyl esters

2 MANUFACTURE AND USES

2.1 Manufacture

The ester is produced either by

- direct esterification of methacrylic acid with the corresponding fatty alcohol (such products may contain up to 1 % methacrylic acid as a low molecular weight impurity)
- or trans-esterification/alcoholysis of methyl methacrylate with the corresponding fatty alcohol (such products may contain up to 1 % methyl methacrylate as a low molecular weight impurity).

Pure dodecyl alcohol (dodecanol, lauryl alcohol) is used only on a small scale to produce the ester. In the large-scale production of long-chain aliphatic methacrylate esters technical mixtures are used of fatty (long-chain aliphatic) alcohols of natural or synthetic origin.

The carbon chain length distribution of the resulting mix of long-chain aliphatic methacrylate esters mirrors the chain length distribution of the alcohol(s) used.

A typical raw material for the production of dodecyl methacrylate is a C12-rich alcohol mixture of natural origin with approx. 65-70 % dodecanol, approx 25 % tetradecanol (lauryl and myristyl alcohol) and approx. 5-10 % of higher alkyl alcohols.

2.2 Identified uses

The esters are monomers for the production of polymers. Typical uses of the polymers are in lubricant additives, paint resins, floor care products, sizing agents for paper, reactive adhesives and reactive coatings.

3 CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES

Not evaluated in this dossier.

4 HUMAN HEALTH HAZARD ASSESSMENT

In this chapter only toxicokinetics and irritation are discussed

4.1 Toxicokinetics (Absorption, Metabolism, Distribution and Elimination)

4.1.1 Non-human information

Physico chemical properties of the substance will enable qualitative judgements of the TK behaviour (Guidance on information requirements and chemical safety assessment Chapter R.7.c, R.7.12 Guidance on Toxicokinetics):

In general with a calculated log Pow of 6.68 of dodecyl methacrylate absorption into the blood from GI absorption, respiratory absorption or skin is not expected. (log Pow values between -1 and 4 are favourable for absorption). With a water solubility of < 1 µg/l the substance is poorly soluble. The molecular weight is 254 g/mol and the substance is not a skin sensitizer.

Experimental *in vitro* studies of the toxicokinetics of dodecyl methacrylate are only available for dermal absorption. Experimental *in vitro* studies with the structurally related substance ethylhexyl methacrylate are used to assess the metabolism of dodecyl methacrylate.

Absorption

GI absorption

No experimental data are available for GI absorption.

Substances with a molecular weight below 500 g/mol, high water solubility and a log Pow between -1 and 4 are favourable for absorption. With log Pow > 4 passive diffusion through membranes is not expected but the substance may form micelles and be absorbed into the lymphatic system. But with a water solubility of < 1 µg/l very low concentrations of the substance are bioavailable so that the substance is poorly absorbed. No signs of systemic toxicity are indicating that absorption has occurred were seen in an acute oral toxicity test up to 5000 mg/kg bw.

GI absorption is not the favoured route of absorption. Only a low amount of the substance may be absorbed by micellular solubilisation due to the very poor water solubility of the substance.

Respiratory absorption – Inhalation

No experimental data are available for respiratory absorption.

The vapour pressure of dodecyl methacrylate is only 0.06 Pa @ 20 °C and therefore the volatility is far too low for inhalation in a gaseous form (substances with low volatility have a vapour pressure of less than 0.5 kPa).

Inhalation is not the favoured route of absorption.

Dermal absorption

Dodecyl methacrylate is a liquid substance with a molecular weight between $MW > 100 < 500$ g/mol which would favour dermal uptake, but with a very low water solubility of 1 µg/l dermal uptake from the stratum corneum into the epidermis is likely to be too low. With $\log Pow > 6$ the rate of transfer between the stratum corneum and the epidermis will be slow and will limit absorption across the skin. Uptake into the stratum corneum itself may be slow.

Although dodecyl methacrylate has a skin binding structure (methacrylate) it was not sensitizing in *in vivo* tests in mice and guinea pigs. The substance is not skin irritating or corrosive, so that the substance itself will not enhance penetration through damaged skin. No signs of systemic toxicity indicating absorption were observed in an acute dermal toxicity study up to 3000 mg/kg bw.

The dermal absorption (steady-state flux) of dodecyl methacrylate has been estimated by calculation using the principles defined in the Potts and Guy prediction model (Heylings JR, 2013).

Table 10: Terms used for categorising absorption of chemicals through human skin:

Kp (cm/h)	Absorption Rate (µg/cm ² /h)	Relative Absorption Rate Category	Predicted Absorption from Normal Exposure
1E-02 – 1E-01	>500	Very fast	Very high
1E-03 – 1E-02	100-500	Rapid - Fast	High
1E-04 – 1E-03	10-50 50-100	Slow - Moderate Moderate - Rapid	Moderate
1E-05 – 1E-04	0.1-10	Very slow - Slow	Low
1E-06 – 1E-05	0.001-0.1	Extremely - Very slow	Minimal
<1E-06	<0.001	Extremely slow	Negligible

Based on a molecular weight of 254.41 g/mol and a $\log Pow$ of 6.68, the predicted flux of Dodecyl methacrylate is 0.003 µg/cm²/h; the relative dermal absorption is minimal.

Metabolism

No data are available of the metabolism of dodecyl methacrylate *in vivo*.

Assumed dodecyl methacrylate will be absorbed the prominent pathway for the metabolism of higher methacrylate esters starts with ester hydrolysis resulting in methacrylic acid and the corresponding alcohol (Jones, 2002), (McCarthy and Witz, 1997). While the acid is further metabolised via the valine pathway of the citric acid cycle (ECETOC, 1996; European Union, 2002) the alcohol may be further metabolised by the two standard metabolic pathways of fatty alcohols (first: oxidation: fatty alcohol → aldehyde → acid, and subsequently CoA-mediated fatty acid metabolism - or secondly : glucuronidation of the alcohol and excretion).

Alkyl esters of methacrylic acid up to C₈ (2-ethylhexyl methacrylate) showed rapid metabolism with half lives in rat blood of less than 30 min (Jones, 2002):

Series of *in vitro* and *in vivo* studies with methacrylates were used to develop PBPK that accurately predict the metabolism and fate of these monomers. The studies confirmed that alkyl methacrylate esters are rapidly hydrolysed by ubiquitous carboxylesterases. First pass (local) hydrolysis of the parent esters has been shown to be significant for all routes of exposure. *In vivo* measurements of rat liver indicated this organ as the greatest esterase activity. Similar measurements for skin microsomes indicated approximately a 20-fold lower activity than for liver. However, this activity was substantial and capable of almost complete first-pass metabolism of the alkyl methacrylates. For example, no parent ester penetrated whole rat skin *in vitro* for n-butyl methacrylate,

octyl methacrylate or dodecyl methacrylate tested experimentally with only methacrylic acid identified in the receiving fluid. In addition, model predictions indicate that esters of ethyl methacrylate or larger would be completely hydrolysed before entering the circulation via skin absorption. This pattern is consistent with a lower rate of absorption for these esters such that the rate is within the metabolic capacity of the skin. Parent ester also was hydrolyzed by S9 fractions from nasal epithelium and was predicted to be effectively hydrolysed following inhalation exposure.

These studies showed that any systematically absorbed parent ester will be effectively removed during the first pass through the liver (CL as % LBF, see Table 11). In addition, removal of methacrylic acid from the blood also occurs rapidly (T₅₀ %; see Table 11).

Table 11: Rate constants for the ester hydrolysis by rat-liver microsomes and predicted systemic fate kinetics from methacrylates following i.v. administration

Ester	V _{max}	K _m	CL (%LBF)	T _{50%} (min)	C _{max} (MAA) (mg L ⁻¹)	T _{max} (MAA) (min)
Methacrylic acid (CAS 79-41-4; MAA)	-	-	51.6%	-	-	-
Methyl methacrylate (CAS 80-62-6; MMA)	445.8	164.3	98.8%	4.4	14.7	1.7
Ethyl methacrylate (CAS 97-63-5; EMA)	699.2	106.2	99.5%	4.5	12.0	1.8
Isobutyl methacrylate (CAS 97-86-9; i-BMA)	832.9	127.4	99.5%	11.6	7.4	1.6
n-Butyl methacrylate (CAS 97-88-1; n-BMA)	875.7	77.3	99.7%	7.8	7.9	1.8
Hexyl methacrylate (CAS 142-09-6; HMA)	376.4	34.4	99.7%	18.5	5.9	1.2
2-Ethylhexyl methacrylate (CAS 688-84-6; 2EHMA)	393.0	17.7	99.9%	23.8	5.0	1.2
Dodecyl methacrylate (OMA)	224.8	11.0	99.9%	27.2	5.0	1.2

V_{max} (nM/min/mg) and K_m (μM) from rat-liver microsome (100 μg/ml) determinations;

CL = clearance as % removed from liver blood flow,

T₅₀ % = Body elimination time (min) for 50 % parent ester,

C_{max} = maximum concentration (mg/L) of MAA in blood,

T_{max} = time (min) to peak MAA concentration in blood from model predictions.

GSH conjugation, the second potential pathway, has only been observed with small alkyl methacrylates (methyl methacrylate/MMA, ethyl methacrylate/EMA) but was no longer measurable with butyl methacrylate. Moreover, GSH conjugation was only detectable with MMA and EMA at high concentrations which are only achievable under laboratory conditions (Elovaara et al. 1983, Mc Carthy et al 1994).

Table 12: Summary of the peak rates of absorption of MAA and alkyl-methacrylate esters through whole rat and human skin.

Ester	Molec. Volume	Rat whole skin				Human whole skin		
		Peak rate of appearance -- Parent Ester --		Peak rate of appearance -- MAA --		Period of peak abs. rate	Absorbed dose	Predicted rate of absorption $\mu\text{g cm}^{-2}\text{h}^{-1}$
		$\mu\text{g cm}^{-2}\text{h}^{-1}$	$\pm\text{SEM}$	$\mu\text{g cm}^{-2}\text{h}^{-1}$	$\pm\text{SEM}$	h	% of applied/over x h	
MAA	78.96*	360	± 20.9	4584**	± 344	5-8	70%/24	327.0**
MMA	93.198	360	± 20.9	108**	± 4.59	2.5-24	11.3%/24	33.4**
EMA	107.436			<i>190**</i>	-			<i>13.6**</i>
iBMA	135.646			<i>56**</i>	-			<i>4.0**</i>
nBMA	135.856			40.9	± 9.4	2-10	0.4%/10	2.9**
6HMA	164.277			<i>20**</i>	-			<i>1.4**</i>
2EHMA	191.66*			<i>9**</i>	-			<i>0.6**</i>
OMA	192.696			10.3	± 0.65	8-24	0.24%/24	0.7**
DMA	249.536			11.8	± 2.11	8-24	0.26%/24	0.8**

The values in normal type were obtained experimentally, whilst those in italics are predicted values.

** Values are predicted rates of appearance of total chemical including parent ester and metabolite

Distribution

As the bioavailability of dodecyl methacrylate is very low that means neither GI- and respiratory absorption nor dermal absorption are expected and complete metabolism is predicted, only a very low amount of the substance comes into consideration for distribution in blood or plasma and accumulation in organs and tissues.

In theory the lipophilic molecule is likely to distribute into cells and then the intracellular concentration may be higher than extracellular concentration particular in fatty tissues, but this is of secondary importance as the bioavailability of the substance is very low.

Accumulation

In case dodecyl methacrylate should be absorbed accumulation in adipose tissue could be expected as the calculated log Pow is 6.68, but before it should be completely metabolized.

Excretion

As absorption is very low respectively not expected and complete metabolism very fast excretion of dodecyl methacrylate is hardly relevant.

4.1.2 Human information

No human information is available

4.1.3 Summary and discussion on toxicokinetics

According to log Pow > 4 bioaccumulation of dodecyl methacrylate is expected. Otherwise with < 1 $\mu\text{g/l}$ the substance is poorly soluble in water. Therefore the bioavailability of the substance is very low. QSAR modelling for dermal skin absorption predicted minimal absorption with a calculated flux of 0.003 $\mu\text{g/cm}^2/\text{h}$ (Heylings, 2013). *In vitro* studies with rat liver showed fast ester hydrolysis with alkyl methacrylates up to C8-methacrylates. The same metabolism is predicted for dodecyl methacrylate particularly as the available concentration in the body will be very low.

4.2 Irritation

4.2.1 Skin irritation

Table 13: Summary table of relevant skin irritation studies

Method	Results	Remarks	Reference
<p>rabbit (New Zealand White)</p> <p>Coverage: semioclusive (shaved)</p> <p>Vehicle: unchanged (no vehicle)</p> <p>OECD Guideline 404 (Acute Dermal Irritation / Corrosion) (adopted 21 May 1981. EEC Directive 84/449/EEC, Part B: methods for the determination of Toxicity, B5. Acute Toxicity. Skin irritation. Official Journal of the European Communities, No L251, pp. 106-108)</p>	<p>not irritating</p> <p>Erythema score:</p> <p>0.66 of max. 4 (animal #1) (Time point: mean 24+48+72 h) (fully reversible within: 72 h) (4-h semioclusive exposure, reevaluated acc. CLP-criteria)</p> <p>0 of max. 4 (animal #2) (Time point: mean 24+48+72 h) (4-h semioclusive exposure, reevaluated acc. CLP-criteria)</p> <p>0.33 of max. 4 (animal #3) (Time point: mean 24+48+72 h) (fully reversible within: 72 h) (4-h semioclusive exposure, reevaluated acc. CLP-criteria)</p> <p>Edema score:</p> <p>1.33 of max. 4 (animal #1) (Time point: mean 24+48+72 h) (fully reversible within: 8 days) (4-h semioclusive exposure, reevaluated acc. CLP-criteria)</p> <p>1.66 of max. 4 (animal #2) (Time point: mean 24+48+72 h) (fully reversible within: 8 days) (4-h semioclusive exposure, reevaluated acc. CLP-criteria)</p> <p>1 of max. 4 (animal #3) (Time point: mean 24+48+72 h) (fully reversible within: 8 days) (4-h semioclusive exposure, reevaluated acc. CLP-criteria)</p>	<p>2 (reliable with restrictions)</p> <p>weight of evidence</p> <p>experimental result</p> <p>Test material (CAS name): Isotridecyl methacrylate</p> <p>Form: liquid</p>	Schreiber (1989)
<p>rabbit (albino rabbits)</p> <p>Coverage: occlusive (shaved and shaved/abraded)</p> <p>Vehicle: unchanged (no vehicle)</p> <p>according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FDA Draize</p>	<p>slightly irritating</p> <p>Erythema score:</p> <p>1.25 of max. 4 (animal: # 1, # 2, #3, #4, #5, #6) (Time point: 24 and 72 h) (not fully reversible within: 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. DSD (overall mean).)</p> <p>2 of max. 4 (animal #1) (Time point: mean 24 + 72 h) (not fully reversible</p>	<p>2 (reliable with restrictions)</p> <p>weight of evidence</p> <p>experimental result</p> <p>Test material (EC name): 2-Propenoic acid, 2-methyl-, C12-16-alkyl esters</p> <p>Methacrylic acid</p>	Sterner and Stigilc (1977)

Method	Results	Remarks	Reference
(1959)	<p>within: 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>1.5 of max. 4 (animal: #2, #3, #4, #6) (Time point: mean 24 + 72 h) (not fully reversible within: 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>1 of max. 4 (animal: #5) (Time point: mean 24 + 72 h) (not fully reversible within: 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>Edema score:</p> <p>0.08 of max. 4 (animal: # 1, # 2, #3, #4, #5, #6) (Time point: 24 and 72 h) (not fully reversible within: 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. DSD (overall mean))</p> <p>0 (animal: #1, #2, #3, #4, #6) (Time point: mean 24 + 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>0.5 of max. 4 (animal #5) (Time point: mean 24 + 72 h) (not fully reversible within: 72 h) (occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p>	<p>ester of an alcohol mixture with a mean C-number of 12,6 = C12.6 methacrylate</p> <p>(65 % dodecyl methacrylate,</p> <p>25 % Tetradecyl methacrylate,</p> <p>10 % higher alkyl methacrylates up to octadecyl methacrylate)</p> <p>Form: liquid</p>	
<p>rabbit (New Zealand White)</p> <p>Coverage: occlusive (shaved and shaved/abraded)</p> <p>Vehicle: unchanged (no vehicle)</p> <p>according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FDA Draize (1959)</p>	<p>slightly irritating</p> <p>Erythema score:</p> <p>1.67 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: 24 and 72 h) (not fully reversible within: 72 h) (Occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. DSD criteria)</p> <p>1.5 of max. 4 (animal: #1, #3, #4, #6) (Time point: mean 24 + 72 hours) (not fully reversible within: 72 h) (Occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>2 of max. 4 (animal: #2, #5) (Time</p>	<p>2 (reliable with restrictions)</p> <p>weight of evidence</p> <p>experimental result</p> <p>Test material (IUPAC name): decyl methacrylate</p> <p>Form: liquid</p>	<p>Sterner W, Chibanguza (1978)</p>

Method	Results	Remarks	Reference
	<p>point: mean 24 + 72 hours) (not fully reversible within: 72 h) (Occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>Edema score:</p> <p>0.92 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: 24 and 72 h) (not fully reversible within: 72 h) (Occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. DSD criteria)</p> <p>1 of max. 4 (animal: #1, #2, #3, #4, #5) (Time point: mean 24 + 72 hours) (not fully reversible within: 72 h) (Occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p> <p>2 of max. 4 (animal: #6) (Time point: mean 24 + 72 hours) (not fully reversible within: 72 h) (Occlusive, exposure time 24 h, observation time 72 h, intact skin, reevaluated acc. CLP criteria)</p>		
<p>rabbit (New Zealand White)</p> <p>Coverage: occlusive</p> <p>Vehicle: unchanged (no vehicle)</p> <p>range finding study</p>	<p>not irritating</p> <p>Erythema score:</p> <p>1 of max. 4 (animal: #1, #2) (Time point: mean 24 + 72 hours) (fully reversible within: 7 days) (Occlusive, exposure time 24 h, observation time 7 days, intact skin, reevaluated acc. CLP criteria)</p> <p>Edema score:</p> <p>0.5 of max. 4 (animal: #1, #2) (Time point: mean 24 + 72 hours) (fully reversible within: 7 days) (Occlusive, exposure time 24 h, observation time 7 days, intact skin, reevaluated acc. CLP criteria)</p>	<p>2 (reliable with restrictions)</p> <p>weight of evidence</p> <p>experimental result</p> <p>Test material: Dodecyl-, Pentadecyl methacrylate</p> <p>Form: liquid</p>	<p>Parsons RD (1981)</p>

4.2.1.1 Non-human information

No study on skin irritation potential is available of the single substance dodecyl methacrylate. The skin irritation was assessed in a weight of evidence approach with four available studies for structurally related long-chain alkyl methacrylates: One study according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FDA Draize (1959) with Methacrylic acid ester of an alcohol mixture with a mean C-number of 12,6, CAS: 90551-76-1 (65 % dodecyl methacrylate,

25 % Tetradecyl methacrylate, 10 % higher alkyl methacrylates up to octadecyl methacrylate), one skin irritation screening test with two animals conducted in 1981 with Dodecyl-, Pentadecyl methacrylate (app. equal parts of C12-, C13, C14-and C15-methacrylates), another FDA Draize study with n-Decyl methacrylate and one study acc. OECD 404 with the structurally related substance isotridecyl methacrylate. Only the data for the shaved, intact skin were used for evaluation. In studies carried out with more than 3 animals both approaches, the overall mean score and the average score determined per animal, were used for evaluation.

C12,6 methacrylate: 6 rabbits were dermally exposed to 0.5 mL of C12,6 methacrylate. Two application sites per animal were treated, one site was left intact, the other site was abraded. Test sites were covered with an occlusive dressing for 24 hours. Animals were observed for 72 hours. Irritation was scored by the method of Draize et al, 1959.

The treated abraded skin sites showed identical effects as the intact sites. For reevaluation only the scores of the intact skin were used.

As the test was performed with 6 animals both, the CLP and DSD approaches for evaluation have to be conducted acc. Guidance on the application of the CLP criteria.

With the CLP approach the response of the individual animal values were averaged over the two observation days (24 hours and 72 hours after application) separate for erythema and edema. The mean erythema values were 1 for one animal, 1.5 for four animals and 2 for one animal. Erythema scores were not fully reversible within 72 hours. All mean scores were below 2.3.

With the DSD approach the average score overall animals was used separate for erythema and edema. The overall mean erythema score was 1.25 and the mean overall edema score 0.08. Both values are below 2.3.

Performance of the study does not comply with requirements of the relevant recent EU and OECD guidelines, where semi-occlusive dressing, an exposure period of 4 hours, treatment of only intact skin and a recovery period of up to 14 days is stipulated. This study is therefore of limited adequacy for C&L purposes due to intensity of the exposure regime and too short recovery period.

Dodecyl-, Pentadecyl methacrylate: In an acute skin irritation range finding study (1981) 2 New Zealand White rabbits were exposed to Dodecyl-, Pentadecyl methacrylate which contains app. equal parts of C12-, C13-, C14 and C15-methacrylates for 24 h under occlusive conditions. Mean erythema score was 1 in both animals, mean edema score was 0.5 in both animals. All signs of irritation were fully reversible within 7 days. According CLP criteria the substance is not irritating in this study.

n-Decyl methacrylate: In a primary dermal irritation study conducted in 1978 New Zealand White rabbits were dermally exposed (intact and scarified skin) under occlusive conditions to 0.5 mL undiluted n-Decyl methacrylate for 24 hours. Animals then were observed for 3 days. Irritation scores for intact skin were reevaluated according to CLP criteria. 2/6 animals reached the maximum irritation score of 2 for erythema and 1/6 animal the maximum irritation score of 2 for edema. Irritations were not fully reversible within the observation time of 72 hours. Otherwise the exposure time was longer than 4 hours.

In this study n-Decyl methacrylate was slightly irritating to skin. According to CLP criteria effects both erythema and oedema effects are < 2.3. With the DSD approach the mean erythema score was 1.67, the mean edema score was 0.92.

Isotridecyl methacrylate was tested in a primary dermal irritation study acc. OECD 404. 3 New Zealand White rabbits were dermally exposed for 4 hours with 0.5 g undiluted test substance under semioclusiv conditions. Animals were observed after 1h, 24h, 48h 72h and after 8 or 9 days. The test was reevaluated acc. CLP criteria. Mean erythema scores (24 +48 +72 h) were 0, 0.33 and 0.66 of max. 4. Mean edema scores (24 +48 +72 h) were 0, 1.33 and 1.66 of max. 4. All erythema scores were fully reversible within 72 h, all edema scores within 8 days. Under CLP criteria Isotridecyl methacrylate is not irritating to skin.

4.2.1.2 Human information

Human information is not available

4.2.1.3 Summary and discussion of skin irritation

By design, the observation period of the two studies with C12,6 methacrylate and n-Decyl methacrylate were too short to observe full recovery of the animals and also the duration of exposure was longer than the current guideline value. But in analogy to isotridecyl methacrylate and Dodecyl-, Pentadecyl methacrylate full recovery after 8/7 days is assumed. In analogy dodecyl methacrylate is considered to be slightly irritating to skin but not a skin irritant according to the CLP criteria.

4.2.1.4 Comparison with criteria

In four studies with structurally related substances to dodecyl methacrylate the criteria for classification acc. CLP criteria were not reached. Mean erythema and oedema scores were < 2.3 in all animals. As two studies were carried out for only 72 h, reversibility was demonstrated with the structurally related substances isotridecyl methacrylate and Dodecyl-, Pentadecyl methacrylate which were fully reversible within 8/7 days.

4.2.1.5 Conclusions on classification and labelling

According to CLP criteria dodecyl methacrylate has not to be classified as irritating to skin. Current classification should be deleted.

4.2.2 Eye irritation

Table 14: Summary table of relevant eye irritation studies

Method	Results	Remarks	Reference
rabbit (New Zealand White) Vehicle: unchanged (no vehicle) OECD Guideline 405 (Acute Eye Irritation / Corrosion)	not irritating (not classified) Cornea score: 0 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) Iris score: 0 of max. 2 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) Conjunctivae score: 0 of max. 3 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) Chemosis score: 0 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h)	2 (reliable with restrictions) weight of evidence experimental result Test material (CAS name): Isotridecyl methacrylate Form: liquid	Schreiber (1989)
rabbit (New Zealand White) Vehicle: unchanged (no vehicle) according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FAD Draize (1959)	not irritating Cornea score: 0 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 hr) Iris score: 0 of max. 2 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 hr) Conjunctivae score: 0 of max. 3 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 hr) Chemosis score: 0 of max. 3 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 hr)	2 (reliable with restrictions) weight of evidence experimental result Test material (EC name): 2-Propenoic acid, 2-methyl-, C12-16-alkyl esters Methacrylic acid ester of an alcohol mixture with a mean C-number of 12,6 = C12.6 methacrylate (65 % dodecyl methacrylate, 25 % Tetradecyl methacrylate, 10 % higher alkyl methacrylates up to octadecyl methacrylate) Form: liquid	Sterner and Chibanguza (1978a)

Method	Results	Remarks	Reference
<p>rabbit (New Zealand White)</p> <p>Vehicle: unchanged (no vehicle)</p> <p>according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FDA Draize (1959)</p>	<p>not irritating</p> <p>Cornea score:</p> <p>0 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p> <p>Iris score:</p> <p>0 of max. 2 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p> <p>Conjunctivae score:</p> <p>0 of max. 3 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p> <p>Chemosis score:</p> <p>0 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p>	<p>2 (reliable with restrictions)</p> <p>weight of evidence</p> <p>experimental result</p> <p>Test material (IUPAC name): decyl methacrylate</p> <p>Form: liquid</p>	<p>Sterner W, Chibanguza G (1978b)</p>
<p>rabbit (Albino Rabbits)</p> <p>Vehicle: unchanged (no vehicle)</p> <p>no data</p>	<p>not irritating</p> <p>Cornea score:</p> <p>0 of max. 4 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p> <p>Iris score:</p> <p>0 of max. 2 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p> <p>Conjunctivae score:</p> <p>0.67 of max. 4 (animal #2) (Time point: mean 24 + 48 + 72 h) (fully reversible (48 h)) (not rinsed)</p> <p>0 of max. 0 (animal: #1, #2, #3, #4, #5, #6) (Time point: mean 24 + 48 + 72 h) (not rinsed)</p>	<p>2 (reliable with restrictions)</p> <p>weight of evidence</p> <p>experimental result</p> <p>Test material: Dodecyl-, Pentadecyl methacrylate</p> <p>Form: liquid</p>	<p>Mastri CW (1975)</p>

4.2.2.1 Non-human information

No study on eye irritation potential is available of the single substance dodecyl methacrylate. The eye irritation was assessed in a weight of evidence approach with four available studies for structurally related long-chain alkyl methacrylates: One study according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FDA Draize (1959) with methacrylic acid ester of an alcohol mixture with a mean C-number of 12,6, CAS: 90551-76-1 (65 % dodecyl methacrylate, 25 % Tetradecyl methacrylate, 10 % higher alkyl methacrylates up to octadecyl methacrylate), one study with Dodecyl-, Pentadecyl methacrylate (app. equal parts of C12-, C13, C14-and C15-methacrylates) (1975), one study with n-decyl methacrylate (1978) and one study according to OECD 405 with the structurally related substance isotridecyl methacrylate.

C12,6 methacrylate: In a study following an FDA guideline (Draize protocol) C12,6-methacrylate (0.1 ml) was instilled into the right eye of six New Zealand White rabbits. The lids were then gently held together for one second. The test eyes were not washed out following the instillation. The left eye remained untreated for control. The eyes were examined at 24, 48 and 72 hours from beginning of test. Eye irritation was scored for signs of corneal damage (density, area), iris reaction and lesions of the conjunctivae (erythema, chemosis and discharge). There were no signs of damage to cornea and iris and no signs of redness and chemosis of the conjunctiva. All irritation scores were 0.

Dodecyl-, Pentadecyl methacrylate: In a primary eye irritation study (1975) with 6 Albino rabbits animals were exposed with 0.1 ml undiluted Dodecyl-, Pentadecyl methacrylate which contains approximately equal parts of C12-, C13-, C14 and C15-methacrylates. Eyes were not washed. Irritation scores were evaluated after 24, 48 and 72 hours. Mean irritation scores for erythema and iris were 0. Maximum mean irritation score of conjunctiva (redness and chemosis) was 0.67. Irritations were fully reversible within 7 days. In this study Dodecyl-, Pentadecyl methacrylate is not irritating to eyes according to the CLP criteria.

n-Decyl methacrylate: In a primary eye irritation study (according to Appraisal of the Safety of Chemicals in foods, drugs and cosmetics, FDA Draize (1959)) 0.1 ml undiluted n-Decyl methacrylate was instilled into the conjunctival sac of the left eye of 6 New Zealand White rabbits, (2.4 -2.6 kg body weight) for 72 hours (not rinsed). Animals were observed for 7 days. Irritation was scored according to Draize scoring system and reevaluated according CLP criteria. Mean irritation scores (24 + 48 + 72 hours) for cornea, iris, conjunctiva and chemosis were 0 for all animals. In this study Decyl methacrylate is not irritating to eyes.

Isotridecyl methacrylate was tested in an eye irritation study according to OECD 405. 0.1 ml test substance was instilled into the right eye of 3 New Zealand White rabbits. The lids were then gently held together for one second. The test eyes were not washed out following the instillation. The left eye remained untreated for control. The examination of the cornea was secured with the aid of fluorescein after recording the observation at 24 hours. The grades of lesions at 24, 48 and 72 hours of the cornea, iris and conjunctiva were examined. There were no signs of damage to cornea and iris and no signs of redness and chemosis of the conjunctiva. All irritation scores were 0.

4.2.2.2 Human Information

No data are available on human information

4.2.2.3 Summary and discussion of eye irritation

No signs of eye irritation were observed in four studies with structurally related long-chain alkyl methacrylates (C12.6 methacrylate, Dodecyl-,Pentadecyl methacrylate, n-Decyl methacrylate and

isotridecyl methacrylate). Maximum irritation score for conjunctiva was 0.67 with Dodecyl-, Pentadecyl methacrylate which was fully reversible within 7 days. Irritation scores in three further studies were 0 for all irritation parameters at every observation time point. In analogy dodecyl methacrylate is considered not to be an eye irritant.

4.2.2.4 Comparison with criteria

The application of Dodecyl methacrylate to rabbit eyes does not induce effects which are relevant for a classification as eye irritant in accordance with the CLP criteria. In four studies with structurally related long chain alkyl methacrylates in only one study the highest induced irritation score for conjunctiva was 0.67 which was fully reversible within 48 h. All other scores were 0 at 24, 48 and 72 h.

4.2.2.5 Conclusion on classification and labelling

According to CLP criteria dodecyl methacrylate has not to be classified as irritating to eyes. Current classification should be deleted.

4.2.3 Respiratory tract irritation

No data are available on respiratory tract irritation. As vapour pressure of dodecyl methacrylate is < 0.1 Pa, inhalation of the gaseous form is not a route of exposure. The physico chemico properties with a very low vapour pressure cannot exclude an exposure to the aerosol form. Since no data on dodecyl methacrylate are available for the aerosol form and the existing classification seems to be based on a group approach, a comparison with criteria is not possible... The lack of irritating properties on the skin and the eye gives supporting evidence that the current classification as STOT SE 3 may not be justified and should be deleted.

4.3 Corrosivity

See irritation

4.4 Specific target organ toxicity – single exposure (STOT SE)

See 4.2.3 Respiratory tract irritation.

5 ENVIRONMENTAL HAZARD ASSESSMENT

5.2 Degradation

Table 15: Summary of relevant information on degradation

Method	Results	Remarks	Reference
Test type: ready biodegradability activated sludge (mixture of 2 storage lakes, 3 municipal sewage plants and 1 industrial sewage plant) equivalent or similar to OECD Guideline 301 C (Ready Biodegradability: Modified MITI Test (I))	readily biodegradable % Degradation of test substance: 88.5 after 2 d (O ₂ consumption) 7.5 after 2 d (O ₂ consumption) 21 after 3 d (O ₂ consumption) 60 after 10 d (O ₂ consumption) 72 after 15 d (O ₂ consumption)	1 (reliable without restriction) Test material: Dodecyl methacrylate	Fraunhofer (1988)
Test type: ready biodegradability activated sludge (mixture of 2 storage lakes, 3 municipal sewage plants and 1 industrial sewage plant) equivalent or similar to OECD Guideline 301 C (Ready Biodegradability: Modified MITI Test (I))	readily biodegradable % Degradation of test substance: 76.6 after 28 d (O ₂ consumption) 12.8 after 5 d (O ₂ consumption) 59.2 after 10 d (O ₂ consumption) 62.2 after 11 d (O ₂ consumption)	2 (reliable with restriction) Read across: Test material: Methacrylic acid ester 13.6 (68 % esters of C-13- alcohols, 32 % esters of C-15- alcohols, containing a total of 35 % esters of branched alcohols)	Fraunhofer (1988b)

5.2.1 Stability

No data are available on hydrolytic stability of dodecyl methacrylate.

5.2.2 Biodegradation

5.2.2.1 Screening tests

Two studies are available on biodegradation of dodecyl methacrylate and a mixture of the structurally related substances C13- and C15- alkyl methacrylates (linear and branched).

The ready biodegradation of dodecyl methacrylate was investigated in a study conducted according to EEC Directive 84/449/EEC, Degradation – biodegradation, Modified MITI Test, published in official Journal of the European Communities No. L251/199) over a period of 28 days using sludge samples from different places like rivers, lakes, municipal and industrial sewage plants as inoculums (30 mg/L) and 100 mg/L test substance. The biodegradation rate was determined by measurement of O₂ consumption. Inoculum blank and procedural/functional control with reference substance aniline was performed.

After 28 days the degradation of dodecyl methacrylate reached 88.5 % (Fraunhofer 1988). 60 % degradation was found after 10 days. The reference substance reached the pass level of 60 % at day 7 (93.8 % after 28 d). This study is regarded as reliable without restriction and satisfies the guideline requirements for ready biodegradation. Dodecyl methacrylate proved to be readily biodegradable.

In a second study (Fraunhofer 1988b) the ready biodegradation of methacrylic acid ester 13.6 (68 % esters of C-13- alcohols, containing a total of 32 % esters of C-15-alcohols, 35 % esters of branched alcohols) was investigated in a study conducted according to EEC Directive 84/449/EEC, Degradation – biodegradation, Modified MITI Test, published in official Journal of the European Communities No. L251/199) over a period of 28 days using sludge samples from different places like rivers, lakes, municipal and industrial sewage plants as inoculums (30 mg/L) and 100 mg/L test substance. The biodegradation rate was determined by measurement of O₂ consumption. Inoculum blank and procedural/functional control was performed with the reference substance aniline.

After 28 days the degradation of methacrylic acid ester 13.6 reached 76.6 %.The reference substance reached 84.0 % after 28 d. This study is regarded as reliable with restriction and satisfies the guideline requirements for ready biodegradation. Methacrylic acid ester 13.6 proved to be readily biodegradable.

5.2.3 Summary and discussion of degradation

Dodecyl methacrylate (Fraunhofer 1988) and a mixture of C13 and C15 alkyl methacrylates (Fraunhofer 1988) were demonstrated to be readily biodegradable in biodegradation tests according to OECD guideline 301 C (modified MITI tests). 88.5 % and 76.6 % biodegradation were achieved within 28 days, respectively. The 10 day window criteria were fulfilled in both tests.

5.3 Bioaccumulation

5.3.1 Aquatic Bioaccumulation

The studies on aquatic bioaccumulation are summarised in the following table:

Table 16: Summary of relevant information on aquatic bioaccumulation

Method	Results	Remarks	Reference
<i>Danio rerio</i> aqueous (freshwater) flow-through Total uptake duration: 56 h Details of method: Calculation of the uptake and depuration rate constants and the BCF: The uptake rate constant (k ₁), the depuration rate constant (k ₂), the kinetic steady state bioconcentration factor (BCF _k) were calculated by linear and nonlinear regression functions using data for concentrations of 2-Ethylhexylmethacrylate in whole fish measured in the extracts. Calculations of means and ranges were done with Excel spreadsheets (Microsoft	BCF: 37 (whole fish) Elimination: yes; DT50: 1.5 h yes; DT95: 6 h	Read across 2 (reliable with restrictions) key study experimental result Test material (EC name): 2-ethylhexyl methacrylate	Schäfers (2006)

Method	Results	Remarks	Reference
<p>Inc.) while linear and non-linear regressions were conducted with the program SigmaStat 2.03 (SPSS Inc. 1997).</p> <p>Calculation of the steady state BCF: The test substance is known to be taken up quickly due to the high partition coefficient and to be rapidly metabolized, leading to a very fast elimination. A non-GLP pre-study showed that the steady state can be expected to be achieved within the first 8-12 h. Two further sampling dates after 32 h and 56 h were included to provide certainty about the BCF. The BCFSS was calculated by dividing the mean of the values for the 2-Ethylhexyl methacrylate concentration in fish which represent the worst case steady state by the mean measured relevant concentrations in the water.</p> <p>Calculation of the depuration rate constants: The depuration rate constant (k₂) was calculated using the measured concentrations in fish during the depuration phase by applying a model regarding fish as one compartment. The model assumes that the concentration of the test substance in the fish (C_f) is decreasing exponentially: $C_f(t) = C_f(t_i) * e^{(-k_2 * t)}$ C_f(t): concentration in fish at sampling time in days (µg/Kg) C_f(t_i): steady state concentration in fish corresponding to the concentration at start of the depuration phase (= 100%) k₂: depuration rate constant k₂ was calculated by linear regression applied to the ln-transformed concentrations in fish.</p> <p>Calculation of the uptake rate constant: The uptake rate constant k₁ was calculated by a non-linear regression of the ratios C_f/C_w against time during the uptake phase and using the depuration rate constant fitted before. The fitted model assumes an attenuation of uptake by simultaneous depuration, increasing with increasing C_f up to an steady state between uptake and depuration. For the one compartment kinetics eq. 3 was fitted: $C_f/C_w = k_1/k_2 * (1 - e^{(-k_2 * t)})$ k₁: uptake rate constant C_f: concentration in fish (µg/kg) C_w: concentration in water (µg/L) k₁ was calculated by non-linear regression using the k₂ values obtained in the depuration phase .</p> <p>Calculation of kinetic BCFk: The kinetic BCF for the one compartment model is given by $BCF_k = k_1/k_2$</p> <p>OECD Guideline 305 (Bioconcentration: Flow-through Fish Test)</p>			

5.3.2 Summary and discussion of aquatic bioaccumulation

No *in vivo* study is available with dodecyl methacrylate but a fish bioaccumulation study (OECD guideline 305) with the structurally related substance ethylhexyl methacrylate. The measured BCF was 37.

Bioavailability of ethylhexyl methacrylate is expected to be higher than dodecyl methacrylate due to the lower log Pow, lower molecular weight and higher water solubility of ethylhexyl methacrylate (EHMA, CAS No: 688-84-6, C₁₂H₂₂O₂: log Pow. 5.59; MW 198.31 g/mol; water solubility: 3.07 mg/l; Dodecyl methacrylate, C₁₆H₃₀O₂: log Pow 6.68, MW 254.42 g/mol, water solubility: < 1 µg/l). Ethylhexyl methacrylate and dodecyl methacrylate are both alkyl methacrylates and the same way of rapid metabolism is expected. Metabolism of ethylhexyl methacrylate is indeed faster than dodecyl methacrylate, but the concentration of dodecyl methacrylate in organisms is much lower than ethylhexyl methacrylate due to the lower bioavailability. Nevertheless, an exact BCF of dodecyl methacrylate cannot be estimated.

However, the calculated log Pow of dodecyl methacrylate is 6.68. According to Guidance on the application of the CLP criteria (Annex III 4.3) read across should only be considered if no experimental BCF or log Pow data or no predicted log Pow data are available. The log Pow is above the CLP cut-off (log Pow ≥ 4) and thus dodecyl methacrylate has potential to bioaccumulate in organisms.

5.4 Aquatic toxicity

Table 17: Summary of relevant information on aquatic toxicity

Method	Results	Remarks	Reference
Fish			
<i>Oncorhynchus mykiss</i> freshwater flow-through OECD Guideline 203 (Fish, Acute Toxicity Test)	LC ₅₀ (96 h): > 62 mg/L act. ingr. (highest test concentration) (meas. (arithm. mean)) based on: mortality (Test solutions were cloudy and grey. The amount of undissolved material increased with increasing test concentration.)	3 (not reliable) weight of evidence experimental result Test material (EC name): dodecyl methacrylate Form: liquid	Springborn laboratories Inc. (1995)
<i>Leuciscus idus</i> freshwater static DIN 38412 part 15	LC ₅₀ (48 h): 1080 mg/L test mat. (test concentration 6 orders of magnitude above the solubility of the test substance) (nominal) based on: mortality	3 (not reliable) weight of evidence experimental result Test material (EC name): dodecyl methacrylate Form: liquid	Institut Fresenius, Chemische und biologische Laboratorien GmbH, 6204 (1988)

CLH REPORT DODECYL METHACRYLATE

Method	Results	Remarks	Reference
Daphnia			
<i>Daphnia magna</i> freshwater flow-through OECD Guideline 202 (Daphnia sp. Acute Immobilisation Test)	EC ₅₀ (48 h): > 2 mg/L test mat. (meas. (arithm. mean)) based on: mobility	3 (not reliable) weight of evidence experimental result Test material (EC name): dodecyl methacrylate	Putt (1995)
<i>Daphnia magna</i> freshwater semi-static OECD Guideline 211 (Daphnia magna Reproduction Test) (adopted September 1998)	NOEC (21 d): >= 5.73 µg/L test mat. (meas. (arithm. mean)) based on: reproduction (and immobilisation) (test concentration above the water solubility)	1 (reliable without restriction) key study experimental result Test material (EC name): dodecyl methacrylate Form: liquid	NOACK (2005)
Algae			
<i>Desmodesmus subspicatus</i> (algae) freshwater static OECD Guideline 201 (Alga, Growth Inhibition Test) (1984)	EC ₅₀ (72 h): > 10 µg/L test mat. (nominal) based on: biomass and growth rate (95 % confidence interval: not applicable) NOEC (72 h): 10 µg/L test mat. (nominal) based on: biomass and growth rate (95 % confidence interval: not applicable) LOEC (72 h): > 10 µg/L test mat. (nominal) based on: biomass and growth rate (95 % confidence interval: not applicable)	1 (reliable without restriction) key study experimental result Test material (EC name): dodecyl methacrylate Form: liquid	NOACK (2005b)
<i>Selenastrum capricornutum</i> (new name: <i>Pseudokirchnerella subcapitata</i>) (algae) freshwater static OECD Guideline 201 (Alga, Growth Inhibition Test)	EC ₅₀ (96 h): > 0.19 mg/L act. ingr. (meas. (initial)) based on: growth rate NOEC (96 h): 0.0062 mg/L act. ingr. (meas. (initial)) based on: growth rate	3 (not reliable) weight of evidence experimental result Test material (EC name): dodecyl methacrylate Form: liquid	Hoberg (1995)

5.4.1 Fish

5.4.1.1 Short-term toxicity to fish

Two studies are available on short term toxicity to fish. The first study according to DIN 38412 part 15 (Fresenius, 1988) with *leuciscus idus* (golden orfe) was conducted with dodecyl methacrylate (97.2 %) at nominal test concentrations of 950, 1000, 1050 and 1100 mg/l. The test solution was prepared using an ultrasonic stirrer.

At 950 mg/l 0/10 fish were dead, 1/10 at 100 mg/l, 2/10 at 1050 mg/l and 10/10 at 1100 mg/l. LC₅₀ was calculated to 1080 mg/l, LCo was 950 mg/l.

As the test concentration was 6 orders of magnitude above the solubility of the test substance, the test is regarded as invalid.

A second test was conducted with a mixture of 69.13 % dodecyl methacrylate and 27.4 % tetradecyl methacrylate acc. OECD guideline 203 with rainbow trout (Springborn 1995) under flow through conditions at nominal concentrations of 13, 21, 35, 58 and 97 mg dodecyl methacrylate/l = active ingredient (a. i.). These concentrations are corresponding with mean measured concentrations of 8.8, 12, 13, 24 and 62 mg a. i. /l (measured by HPLC). Acetone was used as solubilizer in a concentration of 0.167 ml/l. No mortality was observed up to the highest measured concentration of 62 mg/l a. i. LC₅₀ was considered to be > 62 mg/l.

Throughout the study, test solutions were observed to be cloudy and grey and contained a surface film of undissolved test material. As the test was performed with solvent and in a range of concentrations four to five orders of magnitude above the water solubility of the substance (< 1 µg/l) and the substance was introduced into the test medium by rapid stirring, a true solution has not been achieved under the test conditions. Therefore the test is regarded as invalid.

OVERALL COMMENT OF TOXICITY TO FISH: Despite the fact that the available tests are problematic from a technical point of view, it appears that saturated solutions of dodecyl methacrylate are non-toxic to fish, so that there was no necessity to repeat the tests.

5.4.1.2 Long-term toxicity to fish

No data are available on long term toxicity to fish

5.4.2 Aquatic invertebrates

5.4.2.1 Short-term toxicity to aquatic invertebrates

One study is available on the acute toxicity to daphnia. The test was conducted with a mixture of 69.13 % dodecyl methacrylate and 27.4 % tetradecyl methacrylate according to OECD guideline 202 (Putt, 1995) under flow through conditions with *daphnia magna* at nominal concentrations 0.39, 0.65, 1.1, 1.8 and 3.0 mg/l dodecyl methacrylate = active ingredient (a. i.) corresponding with measured concentration of 0.17, 0.30, 0.59, 0.80 and 2.0 mg a. i./l (measured by HPLC). The test material was introduced into the test medium by rapid stirring. Following 48 h of exposure, 35 % immobilization was observed among daphnids exposed to 2.0 mg a. i./l. During the same period 20, 15, 10 and 25 % immobilization was observed among daphnids exposed to the 0.17, 0.30, 0.58 and 0.80 mg a. i./l treatment levels. 5 % immobilization was observed among daphnids exposed to the control solutions. All mobile daphnids exposed to the highest treatment level (2.0 mg a. i./l)

exhibited lethargic behaviour. Sublethal effects were also observed among several organisms exposed to the 0.80 mg a. i./l treatment level. No sublethal effects were observed among daphnids exposed to the remaining concentrations tested.

Based on these results EC_{50} (48 h) was empirically estimated to be > 2.0 mg a. i./l, the highest achievable concentration. The slope of the concentration effect curve at 48 h did not establish a relationship between exposure and effect sufficient to empirically determine a NOEC for dodecyl methacrylate in this test.

In the absence of a clear concentration effect relationship and taking into account that emulsions with concentrations three to four orders of magnitude above the water solubility instead of solutions were tested the test is regarded as invalid.

A 21 d daphnia reproduction test indicates that dodecyl methacrylate is non-toxic to daphnia at a concentration one order of magnitude above the limit of water solubility. Saturated solutions of dodecyl methacrylate are non-toxic to daphnia under acute and chronic conditions.

5.4.2.2 Long-term toxicity to aquatic invertebrates

One study is available on the long term toxicity to aquatic invertebrates (NOACK, 2005). The test was conducted with dodecyl methacrylate, purity 98.34 %. The 21-day-chronic toxicity of dodecyl methacrylate to *Daphnia magna* STRAUSS was studied under semi static conditions according to OECD guideline 211. Daphnids were exposed to dodecyl methacrylate at a limit concentration of 10 $\mu\text{g/L}$ (nominal). This concentration is higher than the solubility in water (< 1 $\mu\text{g/L}$) but has, nevertheless, been chosen with regard to the feasibility of attaining appropriate and analysable test concentrations at 10 $\mu\text{g/L}$.

10 test organisms, individually held were used for the limit concentration and control. At test start they were 2 to 24 h old. The test method was semi-static. Test solutions were renewed daily.

Concentrations of dodecyl methacrylate in the stock solution, limit concentration and control of fresh (0 h) and old (24 h) media were determined via HPLC. Samples were taken and analysed on days 0, 7, 16, 20 (fresh media) and on days 1, 8, 17, 21 (old media). The test item concentrations decreased within 24 h. All effect values were given based on the time weighted mean measured concentration for the limit concentration of 5.73 $\mu\text{g/L}$.

The average number of juveniles per parent in the control group was 85 after 21 days. The reproductive output at the limit concentration was not statistically significant reduced compared to the control. The coefficient of variation around the mean number of living offspring produced per parent in the control group was 5.02 % and shows very small variances between the control replicates.

No winter eggs, males, ephippia, stillborn juveniles and aborted eggs occurred in control or test groups.

The mean day of release of first brood was 9 in the control group and the limit concentration. There was no difference between the two groups. At the limit concentration and the control group 4 broods were released during the test period.

The intrinsic rates of natural increase (IR) of the surviving parent animals accounting for generation time and offspring numbers were used for calculation of population growth. The mean IR of the surviving daphnids of the limit concentration was compared to the control by One Way Analysis of

Variances ($p < 0.05$). There was no statistically significant difference. The intrinsic rate was comparable for the control and limit concentration.

The no observed effect concentration (NOEC) after 21 days based on reproduction capacity is the tested limit concentration of 5.73 $\mu\text{g/L}$. No statistically significant test item related effects were observed at the limit concentration when compared to the control group. No immobilisation of parent animals occurred in the control or test group.

Water quality parameters as pH-value, dissolved oxygen, water hardness and temperature were determined to be within the acceptable limits.

In order to prove the validity of the test system and test conditions at the testing facility, an acute immobilization test according to DIN 38412 L11 was carried out with potassium dichromate as reference item once per month. The EC_{50} of the reference item at 1.84 mg/L after 24 h was within the validity range of 1.0 to 2.5 mg/L according to DIN 38412 L30.

The 21 day $\text{LC}_{50}/\text{EC}_{50}$ based on reproduction/immobilisation was greater than 5.73 $\mu\text{g/L}$ (mean measured concentration). The 21-day NOEC based on reproduction/immobilisation was $\geq 5.73 \mu\text{g/L}$ (mean measured concentration). Production of offspring in the treated groups indicated that Dodecyl methacrylate did not have an effect on the reproduction at concentrations lower or equal than 5.73 $\mu\text{g/L}$.

This study is classified as acceptable and satisfies the guideline requirements for a chronic toxicity study with freshwater invertebrates.

5.4.3 Algae and aquatic plants

Two studies are available on algae toxicity.

The first was conducted with a mixture of 69.13 % dodecyl methacrylate and 27.4 % tetradecyl methacrylate according to OECD guideline 201 with *Selenastrum capricornutum* (new name: *Pseudokirchneriella subcapitata*) under static conditions at nominal concentrations of 0.0063, 0.013, 0.025, 0.050 and 0.10 mg a. i. /l (a. i. = dodecyl methacrylate, corresponding with 0.0091, 0.019, 0.036, 0.072 and 0.145 mg/l test substance (Hoberg, 1995). Acetone was used as solubilizer in a concentration of 0.1 ml/l. Measured concentrations by HPLC were higher than nominal concentrations: 0.068, 0.016, 0.0274, 0.062 and 0.19 mg a. i./l.

EC_{50} value for growth rate ($\text{ECr}_{50}(96 \text{ h}) > 0.19 \text{ mg/l}$) was above the highest nominal concentration tested. NOEC was 0.0062 mg/l (measured concentration).

In the absence of a clear concentration-effect relationship and taking into account that emulsions and not solutions of the material were tested, the NOEC which had been reported in the test report is irrelevant. The study is regarded as invalid.

A second study was conducted with dodecyl methacrylate (purity 98.34 %) for 72 h acc. OECD guideline 201 as limit test under static conditions with *Desmodesmus subspicatus* at test concentrations of 10 $\mu\text{g/l}$ with an initial cell density of nominally $10\text{E}+1 \text{ cell/ml}$ (NOACK, 2005b). Three replicates were tested for the limit concentration and six for the control.

The recovery rate of the limit concentration was 105 % at the test start and 94 % at test end. All effect values are based on nominal test concentrations.

The test concentration of 10 µg/l was higher than the solubility in water (< 1 µg/l) but was, nevertheless, been chosen with regard to the feasibility of attaining an analysable test concentration of 10 µg/l.

EC₅₀ and NOEC based on growth inhibition and biomass production were > 10 µg/l and ≥ 10 µg/l, respectively. The study is acceptable and satisfies the guideline requirements for Algae, Growth Inhibition study. The study is regarded as valid without restrictions.

Validity of ecotoxicity studies with solvent in which the test concentrations exceed the limit of solubility by several orders of magnitude

Several ecotoxicity tests with dodecyl methacrylate have been performed at concentrations orders of magnitude above the limit of solubility. In those cases, rapid stirring or solvent or both have been used to disperse the test material. In those studies, no attempt has been made to determine whether the test material was dissolved and only in one study it was acknowledged that the test material was present in the form of an emulsion. Based on the measured water solubility of dodecyl methacrylate (< 1 µg/l) it can be assumed, that in those cases the test material was present almost entirely as small, undissolved droplets forming an emulsion. That has the consequence that the 'concentration' no longer determines the dose in the test organism but the stochastic, individual contact of the test organism with droplets of the test material and the kinetics of the subsequent absorption of the droplet by the test organism. Test results obtained this way are artefacts and not representative. They cannot be used establishing a concentration-effect relationship. Therefore, test results have only been used when the test was performed without solvent and the nominal test concentration was not higher than approximately ten-fold above the solubility of dodecyl methacrylate in water.

5.5 Comparison with criteria for environmental hazards (sections 5.1 – 5.4)

Table 18: Comparison with criteria for environmental hazards

	Criteria for environmental hazards	Dodecyl methacrylate	Conclusion
Rapid Degradation	Readily biodegradable in a 28-day test for ready biodegradability	88 % after 28 days (O ₂ consumption) 10 day window passed	Rapidly degradable
Bioaccumulation	Log Kow ≥ 4 BCF ≥ 500	Estimated Log Kow = 6.68	Bioaccumulative
Aquatic Toxicity	Acute toxicity data: LC ₅₀ /EC ₅₀ /ErC ₅₀ ≤ 1 mg/L Chronic toxicity data: NOEC ≤ 1mg/L	Algae: ErC ₅₀ 72h > 10 µg/l (test concentration higher than the water solubility of 1 µg/l) NOEC 72 h: ≥ 10 µg/l (test concentration higher than the water solubility of 1 µg/l) Invertebrates: NOEC 21d ≥ 5.73 µg/l (test concentration higher than the water solubility of 1 µg/l) Fish: No valid study	Not acute and chronic toxic up to the water solubility

Criteria for the classification with Aquatic Acute 1 and Aquatic Chronic 1-3:

The values for acute and chronic toxicity are above the water solubility. Therefore, the criteria for the classification of the substance with Aquatic Acute 1 and Aquatic Chronic 1-3 are not fulfilled.

Criteria for the classification with “Aquatic Chronic 4”

- Poorly soluble substance for which no acute toxicity is recorded at levels up to the water solubility
 AND
- which are not rapidly degradable
 AND
- have an experimentally determined $BCF \geq 500$ (or, if absent, a $\text{Log Pow} \geq 4$)

The substance is rapidly degradable. Therefore, this criterion for the classification with Aquatic Chronic 4 is not fulfilled.

**5.6 Conclusions on classification and labelling for environmental hazards
(sections 5.1 – 5.4)**

As dodecyl methacrylate is rapidly degradable and the values for acute and chronic aquatic toxicity are above the water solubility of the substance classification and labelling according to the CLP criteria for environmental hazards is not required. The current classification with Aquatic Acute 1 (H400) and Aquatic Chronic 1 (H410) is not justified and should be deleted.

6 REFERENCES

AQura GmbH (2008), Auto ignition temperature of Isodecyl methacrylate (LIMA) acc DIN N ISO/IEC 17025, unpublished study of Evonik RohMax Additives GmbH, D-64295-Darmstadt, Germany

BAM (2013): Expert judgement by BAM Federal Institute for Materials Research and Testing, Division 2.2, Berlin, Germany.

Brandes E, Möller W (2003), Sicherheitstechnische Kenngrößen Band 1: Brennbare Flüssigkeiten und Gase. Wirtschaftsverlag NW, Verlag für neue Wissenschaft GmbH, Bremerhaven: 346, ISBN 3-89701-745-8

Dr. U. Noack Laboratories (2004) Lauryl methacrylate Water Solubility (Generator Column Method), Unpublished study on behalf of RohMax Additives GmbH, D-64293 Darmstadt, Germany

ECETOC. (1996) Joint Assessment of Commodity Chemicals No. 35, Methacrylic Acid, CAS No. 79-41-4; March 1996

Elovaara E, Kivisto H, Vainio H. (1983) Effects of methyl methacrylate on non-protein thiols and drug metabolizing enzymes in rat liver and kidneys. Arch Toxicol 52(2), 109-121

European Chemicals Bureau (2002). European Union - Risk Assessment Report on Methyl methacrylate. European Union - Risk Assessment Report, Vol. 22.

Evonik RohMax Additives GmbH (2008), Determination of the Kinematic Viscosity of Methacrylic acid ester of aliphatic alcohols C12-C16, Unpublished report on behalf of Evonik RohMax Additives GmbH, D-64297 Darmstadt, Germany

Fraunhofer (1988) Biotic Degradation, The Modified MITI Test, Test Substance: Dodecylmethacrylate CAS-No. 142-90-5. Unpublished report on behalf of Röhm GmbH, D-64293 Darmstadt, Germany. Testing laboratory: Fraunhofer-Institut für Umweltchemie und Ökotoxikologie (FhG), Germany. Owner company: Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany.

Fraunhofer (1988b) Biotic Degradation, The Modified MITI Test, Test Substance: Methacrylic ester CAS-No. 90552-04-8. Unpublished report on behalf of Röhm GmbH, D-64293 Darmstadt, Germany. Testing laboratory: Fraunhofer-Institut für Umweltchemie und Ökotoxikologie (FhG), Germany. Owner company: Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany.

Heylings JR (2013) Higher Methacrylates: In Silico Prediction of Dermal Absorption, Dermal Technology Laboratory Ltd., Med IC4, Keele University Science and Business Park, Keele, Staffordshire, ST5 5NL UK, study on behalf of Higher Methacrylates REACH Task Force.

Hoberg JR (1995) Lauryl methacrylate - Toxicity to the freshwater green alga, *Selenastrum capricornutum*, Springborn Laboratories Inc., Wareham, Massachusetts, USA, Unpublished report on behalf of Methacrylate Producers Association (MPA), Washington DC, USA

Institut Fresenius (1988) Examination of Dodecyl methacrylate, Institut Fresenius, Chemische und biologische Laboratorien GmbH, unpublished report on behalf of Röhm GmbH, Darmstadt

Jones O (2002) Using physiologically based pharmacokinetic modelling to predict the pharmacokinetics and toxicity of methacrylate esters. Thesis, March 21, 2002, Univ. of Manchester

- Mastri CW (1975). Eye irritation Test - Albino Rabbits. Unpublished report on behalf of Rohm & Haas Company. Testing laboratory: Industrial BIO-TEST Laboratorie Inc. Report no.: IBT No.: 601-06217. Owner company: Dow Chemicals. Study number: UNTER 00-095. Report date: 1975-10-17.
- McCarthy,TJ; Witz,G. (1997) Structure-activity relationships in the hydrolysis of acrylate and methacrylate esters by carboxylesterase *in vitro*. Toxicology 116(1-3), 153-158
- McCarthy,TJ et al (1994) The relationship of selected acrylate esters toward Glutathione and Deoxyribonucleosides *in vitro*: Structure-activity relationships, Fundam. Appl. Toxicol. 22: 543 – 548
- Nabert, Schön, Redeker (2005), Sicherheitstechnische Kenngrößen brennbarer Gase und Dämpfe Band II, Deutscher Eichverlag, ISBN 3-8064-9946-2
- NOACK (2005) Lauryl methacrylate: Daphnia magna STRAUS Reproduction Test (Semi-Static, 21 d), Limit Test., Dr. U Noack-Laboratories, D-31157 Sarstedt, Germany, Unpublished report on behalf of Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany
- NOACK (2005b) Lauryl methacrylate: Alga, Growth Inhibition Test with *Desmodesmus subspicatus*, 72 h, Dr. U Noack-Laboratories, D-31157 Sarstedt, Germany, Unpublished report on behalf of Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany
- Parsons RD (1981). Dodecyl-Pentadecyl methacrylate Acute Range finding Skin Irritation. Unpublished study on behalf of Rohm and Haas. Testing laboratory: Rohm and Haas Company Toxicology department. Report no.: 80R 0206. Owner company: Dow Chemicals. Study number: UNTER 00-096. Report date: 1981-03-20.
- Putt AE (1995) Lauryl methacrylate - Acute toxicity to daphnids (*Daphnia magna*) under flow-through-conditions, Springborn Laboratories Inc., Wareham, Massachusetts, USA, Unpublished report on behalf of Methacrylate Producers Association (MPA)
- Rehberg C.E., Fisher C.H. (1948), Properties of Monomeric and Polymeric Alkyl Acrylates and Methacrylates, Ind. Eng. Chem. 40(8): 1429-1433 (1948)
- Schäfers C. (2006) Fish, Bioconcentration of 2-Ethylhexyl methacrylate according to the OECD-Guideline 305, Fraunhofer Institute for Molecular Biology and Applied Ecology (IME) Schmollenberg-Grafschaft, Germany, unpublished report on behalf of Methacrylate Producers Association
- Schreiber G (1989) Test report on Acute Dermal Irritation / Corrosion of Isotridecylmethacrylat, Fraunhofer-Institut für Umweltchemie und Ökotoxikologie (FhG), Schmollenberg, Germany, Unpublished report on behalf of Evonik Röhm GmbH, D-64293 Darmstadt, Germany
- Schreiber G (1989) Test Report on Acute Eye Irritation / Corrosion of Isotridecylmethacrylat, Fraunhofer-Institut für Umweltchemie und Ökotoxikologie (FhG), Schmollenberg, Germany, Unpublished report on behalf of Evonik Röhm GmbH, D-64293 Darmstadt, Germany
- Sterner W, Chibanguza (1978). Prüfung von "Decylmethacrylat, roh" auf primäre Hautreizwirkung beim Kaninchen. [Primary skin irritation test of Decyl methacrylate]. Unpublished report on behalf of Evonik Oil Additives S. A. S., D-64293 Darmstadt, Germany. Testing laboratory: International Bio-Research (IBR), Germany. Report no.: 1-3-170-78. Owner company: Evonik Oil Additives S. A. S., D-64293 Darmstadt, Germany. Study number: UNTER 78-017.

Sterner W, Chibanguza G (1978a). Prüfung von "Methacrylsäureester 12,6, roh" im Augenreiztest am Kaninchen. [Eye irritation test in the rabbit with methacrylic acid ester 12,6, raw]. Unpublished report on behalf of Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany. Testing laboratory: International Bio-Research (IBR). Report no.: 1-3-179-78. Owner company: Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany. Study number: UNTER77-005.

Sterner W, Chibanguza G (1978b). Prüfung von "Decylmethacrylat, roh" im Augenreiztest am Kaninchen. Unpublished report on behalf of Evonik Industries AG, D-64293 Darmstadt, Germany. Testing laboratory: International Bio-Research, Inc. (IBR). Report no.: 1-3-178-78. Owner company: Evonik Industries AG, D-64293 Darmstadt, Germany. Study number: UNTER 78-016.

Sterner W, Stigilc A (1977). Prüfung von "Methacrylsäureester" (12,6) auf primäre Hautreizwirkung beim Kaninchen. [Primary skin irritation test of methacrylic acid ester 12.6]. Unpublished report on behalf of Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany. Testing laboratory: International Bio-Research (IBR), Germany. Report no.: 1-3-438/1-77. Owner company: Evonik RohMax Additives GmbH, D-64293 Darmstadt, Germany. Study number: UNTER 77-006.

Springborn laboratories Inc. (1995) Lauryl methacrylate-Acute toxicity to rainbow trout (*Oncorhynchus mykiss*) under flow through conditions, Springborn laboratories Inc., Environmental Sciences Division, Wareham, Massachusetts, Unpublished report on behalf of Methacrylate Producers Association

Syracuse research Corporation (2000), KOWWIN™ v1.67 in EPI web 4.0, U.S. Environmental Protection Agency