

# Summary of product characteristics for a biocidal product

**Product name:** Harmonix Rodent Paste

**Product type(s):** PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

**Authorisation number:** IE/BPA 70695

**R4BP 3 asset reference number:** IE-0024533-0000

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	2
5. General directions for use	16
5.1. Instructions for use	16
5.2. Risk mitigation measures	16
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	16
5.4. Instructions for safe disposal of the product and its packaging	17
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	17
6. Other information	17

## Administrative information

### 1.1. Trade names of the product

Harmonix Rodent Paste
-----------------------

### 1.2. Authorisation holder

<b>Name and address of the authorisation holder</b>	Name	2022 ENVIRONMENTAL SCIENCE FR SAS
	Address	3, place Giovanni Da Verrazzano 69009 LYON France
<b>Authorisation number</b>		IE/BPA 70695
<b>R4BP 3 asset reference number</b>		IE-0024533-0000
<b>Date of the authorisation</b>		24/07/2020
<b>Expiry date of the authorisation</b>		24/07/2025

### 1.3. Manufacturer(s) of the biocidal products

<b>Name of the manufacturer</b>	Bayer SAS
<b>Address of the manufacturer</b>	16 rue Jean-Marie Leclair, CS 90106 69266 Lyon (Cedex 09) France
<b>Location of manufacturing sites</b>	INDUSTRIALCHIMICA Srl. Via Sorgaglia 40 I-35020 Arre Italy
	Kollant S.r.l., via C. Colombo 7/7 30030 Vigonovo (VE) Italy
	IRIS 1126A, avenue du Moulins - Route de Saint Privat F- 30340 SALINDRES France

#### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	1443 - Cholecalciferol
<b>Name of the manufacturer</b>	Bayer S.A.S
<b>Address of the manufacturer</b>	16 rue Jean-Marie Leclair, CS 90106 69266 Lyon (Cedex 09) France
<b>Location of manufacturing sites</b>	Fermenta Biotech Limited, Village Takoli; P.O. Nagwain District Mandi - 175 121 Himachal Pradesh India
	Fermenta Biotech Limited, Z-109 B & C, SEZ II, Dahej ,Taluka - Vagara District Bharuch - 392130 Gujarat India

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Cholecalciferol		Active Substance	67-97-0	200-673-2	0,077

### 2.2. Type of formulation

RB - Bait (ready for use)
---------------------------

## 3. Hazard and precautionary statements

Hazard statements

Precautionary statements

## 4. Authorised use(s)

## 4.1 Use description

### Use 1 - House mouse, Norway rat, Roof rat – professionals - indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: <i>Rattus norvegicus</i> Common name: Norway Rat Development stage: Adults and juveniles  Scientific name: <i>Rattus rattus</i> Common name: Roof rat Development stage: Adults and juvenils  Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: Adults and juvenils
<b>Field(s) of use</b>	Indoor  Indoor
<b>Application method(s)</b>	Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
<b>Application rate(s) and frequencies</b>	Application Rate: For rats: 100g to 200g of bait per baiting point, For mice: 20g of bait per baiting point. Dilution (%): - Number and timing of application:  For rats:  100-200 g bait per baiting point depending upon infestation level.  If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).  An amount of 200 g of product must be added when bait has been totally consumed within one control interval.  For mice:  20 g bait per baiting point depending upon infestation level.  If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).  An amount of 20 g of product must be added when bait has been totally consumed within one control interval.

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

*Bait in individual tea bag sachet, Long fibre paper, 20g*

*Primary Packaging: Tea bag (cellulose), 20g*

*Secondary Packaging:*

*Type of packaging: Bucket,*

*Size/ Volume of the packaging: up to 10 kg*

*Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket*

*Type of the packaging: Box*

*Size/volume of the packaging: up to 15 kg*

*Material of the packaging: Inner COEX PET/ LDPE plastic sachet in a cardboard box*

*Type of the packaging: Bag-*

*Size/volume of the packaging: up to 5 kg-*

*Material of the packaging: COEX PET/PA/PE Plastic bag, with handle and reclosable zip*

**4.1.1 Use-specific instructions for use**

The bait stations should be visited at least every 1 to 2 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

Bait should be secured so that it cannot be dragged away from the bait station.  
Remove the remaining bait or the bait stations at the end of the treatment period.

#### 4.1.2 Use-specific risk mitigation measures

Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.  
Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.  
Do not wash the bait stations with water between applications.  
Do not use the product in pulsed baiting treatments.

#### 4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Prevent skin contact when disposing remains of baits.

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Keep away from food, drink and animal feeding stuffs.

### 4.2 Use description

#### Use 2 - House mouse, Norway rat , Roof rat – professionals – Outdoor: around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Norway rat Development stage: Adults and juveniles  Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults and juveniles  Scientific name: Mus musculus Common name: House mouse Development stage: Adults and juveniles
<b>Field(s) of use</b>	Outdoor  Outdoor: around buildings

**Application method(s)**

Method: Bait application  
Detailed description:  
Ready-to-use bait to be used in tamper-resistant bait stations

**Application rate(s) and frequencies**

Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point.  
Dilution (%): -  
Number and timing of application:

For rats:

100-200 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).

An amount of 200 g of product must be added when bait has been totally consumed within one control interval.

For mice:

20 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).

An amount of 20 g of product must be added when bait has been totally consumed within one control interval.

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

*Bait in individual tea bag sachet, Long fibre paper, 20g*

*Primary Packaging: Tea bag (cellulose), 20g*

*Secondary Packaging:*

*Type of packaging: Bucket,*  
*Size/ Volume of the packaging: up to 10 kg*  
*Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket*

*Type of the packaging: Box*  
*Size/volume of the packaging: up to 15 kg*  
*Material of the packaging: Inner COEX PET/ LDPE plastic sachet in a cardboard box*



Type of the packaging: Bag-  
Size/volume of the packaging: up to 5 kg-  
Material of the packaging: COEX PET/PA/PE Plastic bag, with handle and reclosable zip

#### 4.2.1 Use-specific instructions for use

Place the bait stations in areas not liable to flooding.

The bait stations should be visited at least every 1 to 2 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

Bait should be secured so that it cannot be dragged away from the bait station.  
The campaign has to be terminated if there is no further consumption of baits.  
Remove the remaining bait or the bait stations at the end of the treatment period.

#### 4.2.2 Use-specific risk mitigation measures

Do not apply this product directly in the burrows.

The product information (i.e. label and/or leaflet) shall clearly show that:

- a) the product shall not be supplied to the general public (e.g. "for professionals only").
- b) the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- c) users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

Do not wash the bait stations with water between applications.

Search for and remove dead rodents during treatment, at least as often as baiting stations are inspected.

Dispose dead rodents in household waste or in a rendering plant. Avoid direct contact.

Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Do not use the product in pulsed baiting treatments.

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Prevent skin contact when disposing remains of baits.

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Keep away from food, drink and animal feeding stuff

### 4.3 Use description

#### Use 3 - House mouse, Norway rat, Roof rat – trained professionals – Indoor

##### Product type

PT14 - Rodenticides (Pest control)

##### Where relevant, an exact description of the authorised use

Not relevant for rodenticides

##### Target organism(s) (including development stage)

Scientific name: *Rattus norvegicus*  
Common name: Norway rat  
Development stage: Adults and juveniles

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: Adults and juveniles

Scientific name: *Mus musculus*  
Common name: House mouse  
Development stage: Adults and juveniles

##### Field(s) of use

Indoor

Indoor

##### Application method(s)

Method: Bait application  
Detailed description:  
Ready-to-use bait to be used in tamper-resistant bait stations

Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations

##### Application rate(s) and frequencies

Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point.

Dilution (%): -  
Number and timing of application:

For rats:

100-200 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).

An amount of 200 g of product must be added when bait has been totally consumed within one control interval.

For mice:

20 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).

An amount of 20 g of product must be added when bait has been totally consumed within one control interval.

Permanent baiting (please see use-specific instructions for use)

**Category(ies) of users**

Trained professional

**Pack sizes and packaging material**

*Bait in individual tea bag sachet, Long fibre paper, 20g*

*Primary Packaging: Tea bag (cellulose), 20g*

*Secondary Packaging:*

*Type of packaging: Bucket,*

*Size/ Volume of the packaging: up to 10 kg*

*Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket*

*Type of the packaging: Box*

*Size/volume of the packaging: up to 15 kg*

*Material of the packaging: Inner COEX PET/LDPE plastic sachet in a cardboard box*

*Type of the packaging: Bag-*

*Size/volume of the packaging: up to 5 kg-*

*Material of the packaging: COEX PET/PA/PE Plastic bag, with handle and reclosable zip*



#### 4.3.1 Use-specific instructions for use

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.  
The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).  
Bait should be secured so that it cannot be dragged away from the bait station.

#### 4.3.2 Use-specific risk mitigation measures

Do not use the product in pulsed baiting treatments.  
The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").  
Do not use in areas where resistance to the active substance can be suspected.  
Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

#### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

see chapter 5.4

#### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

see chapter 5.5

### 4.4 Use description

**Use 4 - House mouse, Norway rat, Roof rat – trained professionals – outdoor: around buildings**

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Norway rat Development stage: Adults and juveniles</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults and juveniles</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: Adults and juveniles</p>
<b>Field(s) of use</b>	<p>Outdoor</p> <p>Outdoor: around buildings</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations</p> <p>Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations</p> <p>Direct application of ready-to-use bait into the burrow.</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point. Dilution (%): - Number and timing of application:</p> <p>For rats:</p> <p>100-200 g bait per baiting point depending upon infestation level.</p> <p>If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).</p> <p>An amount of 200 g of product must be added when bait has been totally consumed within one control interval.</p> <p>For mice:</p> <p>20 g bait per baiting point depending upon infestation level.</p> <p>If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).</p> <p>An amount of 20 g of product must be added when bait has been totally consumed within one control interval.</p>

Permanent baiting (please see use-specific instructions for use)

**Category(ies) of users**

Trained professional

**Pack sizes and packaging material**

*Bait in individual tea bag sachet, Long fibre paper, 20g*

*Primary Packaging: Tea bag (cellulose), 20g*

*Secondary Packaging:*

*Type of packaging: Bucket,*

*Size/ Volume of the packaging: up to 10 kg*

*Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket*

*Type of the packaging: Box*

*Size/volume of the packaging: up to 15 kg*

*Material of the packaging: Inner COEX PET/ LDPE plastic sachet in a cardboard box*

*Type of the packaging: Bag-*

*Size/volume of the packaging: up to 5 kg-*

*Material of the packaging: COEX PET/PA/PE Plastic bag, with handle and reclosable zip*

**4.4.1 Use-specific instructions for use**

Place the baiting points in areas not liable to flooding.

Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Bait should be secured so that it cannot be dragged away from the bait station.

Additional elements to be considered for direct application in burrows:  
 Baits must be placed to minimise the exposure to non-target species and children.  
 Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

#### 4.4.2 Use-specific risk mitigation measures

Do not use this product in pulsed baiting treatments.  
 The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").  
 Do not use in areas where resistance to the active substance can be suspected.  
 Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

#### 4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

see chapter 5.4

#### 4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

see chapter 5.5

### 4.5 Use description

#### Use 5 - House mouse, Norway rat, Roof rat – trained professionals – outdoor open areas & waste dumps

**Product type**

PT14 - Rodenticides (Pest control)

**Where relevant, an exact description of the authorised use**

Not relevant for rodenticides

**Target organism(s) (including development stage)**

Scientific name: *Rattus norvegicus*  
 Common name: Norway rat  
 Development stage: Adults and juveniles

Scientific name: *Rattus rattus*  
 Common name: Roof rat  
 Development stage: Adults and juveniles

Scientific name: *Mus musculus*  
 Common name: House mouse  
 Development stage: Adults and juveniles

<b>Field(s) of use</b>	<p>Outdoor</p> <p>Outdoor: open areas &amp; waste dumps</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations</p> <p>Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations</p> <p>Direct application of ready-to-use bait into the burrow.</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point. Dilution (%): - Number and timing of application:</p> <p>For rats:</p> <p>100-200 g bait per baiting point depending upon infestation level.</p> <p>If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).</p> <p>An amount of 200 g of product must be added when bait has been totally consumed within one control interval.</p> <p>For mice:</p> <p>20 g bait per baiting point depending upon infestation level.</p> <p>If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).</p> <p>An amount of 20 g of product must be added when bait has been totally consumed within one control interval.</p>
<b>Category(ies) of users</b>	<p>Trained professional</p>
<b>Pack sizes and packaging material</b>	<p><i>Bait in individual tea bag sachet, Long fibre paper, 20g</i></p> <p><i>Primary Packaging: Tea bag (cellulose), 20g</i></p> <p><i>Secondary Packaging:</i></p> <p><i>Type of packaging: Bucket,</i> <i>Size/ Volume of the packaging: up to 10 kg</i> <i>Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket</i></p>



*Type of the packaging: Box*  
*Size/volume of the packaging: up to 15 kg*  
*Material of the packaging: Inner COEX PET/LDPE plastic sachet in a cardboard box*

*Type of the packaging: Bag-*  
*Size/volume of the packaging: up to 5 kg-*  
*Material of the packaging: COEX PET/PA/PE Plastic bag, with handle and reclosable zip*

#### **4.5.1 Use-specific instructions for use**

Place the bait stations in areas not liable to flooding.  
Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.  
Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.  
The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).  
Bait should be secured so that it cannot be dragged away from the bait station.

Additional elements to be considered for direct application in burrows:  
Baits must be placed to minimise the exposure to non-target species and children.  
Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

#### **4.5.2 Use-specific risk mitigation measures**

Search for and dispose dead rodents in the infested area at each visit to prevent secondary poisoning.  
At the beginning of the campaign, visit the bait points at the latest after 1 - 2 days and at least on a weekly basis afterwards. The same applies to baiting campaigns that last for more than 35 days.  
Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.  
Do not use this product in pulsed baiting treatments.  
The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").  
Do not use in areas where resistance to the active substance can be suspected.  
Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

#### **4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### **4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

see chapter 5.4

#### **4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

see chapter 5.5

### **5. General directions for use**

#### **5.1. Instructions for use**

- 1) Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- 2) The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- 3) Where possible, bait stations must be fixed to the ground or other structures.
- 4) Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- 5) Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- 6) Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3. for the information to be shown on the label).
- 7) When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- 8) If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- 9) If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a (anticoagulant) rodenticide, or the use of traps as an alternative control measure.
- 10) Bait in sachets: Do not open the sachets containing the bait.

#### **5.2. Risk mitigation measures**

- 1) Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.

#### **5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- 1) In case of:
  - Dermal exposure, wash skin with water and then with water and soap.
  - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
  - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label [insert country specific information]. Contact a veterinary surgeon in case of ingestion by a pet [insert country specific information].
- 2) Hazardous to wildlife.

#### 5.4. Instructions for safe disposal of the product and its packaging

- 1) At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.

#### 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- 1) Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- 2) Store in places prevented from the access of children, birds, pets and farm animals.
- 3) Shelf-life: 36 months

#### 6. Other information

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

The product contains a bittering agent and a dye.  
Please note Ireland applies a minimum Professional pack size of 2.5kg instead of 3.0kg