

# **Summary of product characteristics for a biocidal product**

**Product name:** Klerat Voksblok

**Product type(s):** PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

**Authorisation number:** BPR-reg.nr. 732-1

**R4BP 3 asset reference number:** DK-0009630-0000

## **Table Of Contents**

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	1
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
5. General directions for use	6
5.1. Instructions for use	6
5.2. Risk mitigation measures	7
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	8
5.4. Instructions for safe disposal of the product and its packaging	8
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	8
6. Other information	8

## Administrative information

### 1.1. Trade names of the product

Klerat Voksblok

### 1.2. Authorisation holder

Name and address of the authorisation holder	Name Syngenta Crop Protection AG  Address Rosentalstrasse 67 4058 Basel Switzerland
Authorisation number	BPR-reg.nr. 732-1
R4BP 3 asset reference number	DK-0009630-0000
Date of the authorisation	22/07/2013
Expiry date of the authorisation	01/07/2024

### 1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	Bábolna Bioenvironmental Centre Ltd
Address of the manufacturer	Dr. Köves János u. 3 2943 Bábolna Hungary
Location of manufacturing sites	Dr. Köves János u. 3 2943 Bábolna Hungary

### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	11 - Brodifacoum
<b>Name of the manufacturer</b>	Vertellus Specialties UK Ltd
<b>Address of the manufacturer</b>	Halebank WA8 8NS Widnes United Kingdom
<b>Location of manufacturing sites</b>	Hale Bank WA8 8NS Widnes United Kingdom

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	Active Substance	56073-10-0	259-980-5	0,005

### 2.2. Type of formulation

Ready-to-use-bait: block

## 3. Hazard and precautionary statements

<b>Hazard statements</b>	<p>May damage the unborn child.</p> <p>May cause damage to organs blood through prolonged or repeated exposure .</p>
<b>Precautionary statements</b>	<p>If medical advice is needed, have product container or label at hand.</p> <p>Obtain special instructions before use.</p> <p>Do not handle until all safety precautions have been read and understood.</p> <p>Wear protective gloves.</p> <p>IF exposed or concerned: Get medical attention.</p> <p>Store locked up.</p>

Keep out of reach of children.  
Dispose of contents to as hazardous waste according to local requirements.

## 4. Authorised use(s)

### 4.1 Use description

#### Use 1 - Trained Professionals (R1) - indoor and outdoor around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides.
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: juveniles and adults</p> <p>Scientific name: Rattus rattus Common name: Black or roof rat Development stage: Adults and juveniles</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: juveniles and adults</p>
<b>Field(s) of use</b>	<p>Indoor Outdoor</p> <p>In and around buildings within a maximum distance of 10 meter from the Buildings. The control of rats is however allowed also where an occurrence of rats farther than 10 meters acts as a source of rats in buildings (source control). The product must always be placed in tamper resistant bait stations to prevent poisoning of humans and animals.</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: Rats: 20-60 g. Mice: 20 g Dilution (%): None. Number and timing of application: Rats: 20-60 g per bait station. If more than one bait station is needed, the distance between bait stations should be 5-10 metre.</p> <p>Mice: 20 g per bait station. If more than one bait station is needed, the distance between bait stations should be 2-5-metre (move periodically).</p>

<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	Loose 20g bait within a PP/PE tub or pail: 3, 5 and 10 kg.

#### 4.1.1 Use-specific instructions for use

- The biocidal product may only be used by trained licensed professionals (R1) against the brown rat (*Rattus norvegicus*), the black rat (*Rattus rattus*) and the house mouse (*Mus musculus*) in and around buildings within a maximum distance of 10 meter from the buildings. The control of rats is however allowed also where an occurrence of rats farther than 10 meters acts as a source of rats in buildings (source control).
- In the case of the use of the biocidal product for source control of rats, the conditions in the Danish Environmental Protection Agency's guidance on the assessment and authorization of biocides against rats and mice must be met ("Miljøstyrelsens vejledning om vurdering og godkendelse af kemiske bekæmpelsesmidler til rotter og mus").
- Follow any additional instructions provided by the relevant code of best practice. See the Danish Environmental Protection Agency's guidance on the assessment and authorization of biocides against rats and mice ("Miljøstyrelsens vejledning om vurdering og godkendelse af kemiske bekæmpelsesmidler til rotter og mus")
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Replace eaten bait only after 3 days and then at maximum 7 day intervals.

#### 4.1.2 Use-specific risk mitigation measures

- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals (at least twice a week).
- Do not apply this product directly in the burrows.

#### 4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See other section.

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See other section.

## 4.2 Use description

### Use 2 - Trained Professionals (R1) - sewers

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	-
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: juveniles and adults
<b>Field(s) of use</b>	Other  Sewers
<b>Application method(s)</b>	Method: Bait application - sewers - control rats Detailed description: -
<b>Application rate(s) and frequencies</b>	Application Rate: 20 – 200 grams per manhole Dilution (%): 0 Number and timing of application: 20 – 200 grams per manhole
<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	Loose 20 g bait within a PP/PE tub or pail: 3, 5 and 10 kg.

### 4.2.1 Use-specific instructions for use

- The biocidal product may only be used by trained licensed professionals (R1) against the brown rat (*Rattus norvegicus*) in sewers.
- The product must be secured to structures in the sewer to prevent it being washed away.
- Follow any additional instructions provided by the relevant code of best practice. See the Danish Environmental Protection Agency's guidance on the assessment and authorization of biocides against rats and mice ("Miljøstyrelsens vejledning om vurdering og godkendelse af kemiske bekæmpelsesmidler til rotter og mus").

### 4.2.2 Use-specific risk mitigation measures

See other section.

**4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided

**4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See other section.

**4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

See other section.

**5. General directions for use**

**5.1. Instructions for use**

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product locked up and out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining bait and the bait stations at the end of the treatment period.
- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.

## 5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional (R1) holding certification demonstrating compliance with the applicable training requirements. See section 6 for the exact information that should be included on the label.
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke.
- Wash hands and directly exposed skin after using the product.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.
- Not for use against any other pest than indicated on the label and not for use in higher doses than indicated.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
  - the product shall not be supplied to the general public (e.g. "for trained professionals (R1) only").
  - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
  - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

### **5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
  - Dermal exposure, wash skin with water and then with water and soap.
  - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
  - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.
- Hazardous to wildlife.
- Bait stations must be labelled with a warning that mice or/and rats are being controlled in the area and instructions on how to act in case of poisoning. Bait stations must be labelled with contact information on the pest controllers responsible for the treatment.

### **5.4. Instructions for safe disposal of the product and its packaging**

- At the end of the treatment dispose the uneaten bait and the packaging as hazardous waste in accordance with local requirements.
- The product may not be disposed of in sewers, including rainwater canals .

### **5.5. Conditions of storage and shelf-life of the product under normal conditions of storage**

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store the product locked up and out of the reach of children, birds, pets and farm animals and other non-target animals.
- Store the product away from food, drink and animal feeding stuffs.
- Shelf life: 2 years.

## **6. Other information**

The label of the product should contain the following information (in Danish):

III. Etiketten skal indeholde nedenstående oplysninger. Oplysninger i citationstegn skal angives ordret:

1) I hovedfeltet:  
"Gnavermiddel"

Må kun anvendes til bekæmpelse af brun rotte (*Rattus norvegicus*), husrotte (*Rattus rattus*) og husmus (*Mus musculus*) i og omkring bygninger og kun i en afstand op til 10 meter fra en bygning samt til bekæmpelse af brun rotte i kloakker. Bekæmpelse er dog også tilladt, hvor en forekomst af rotter længere væk end 10 meter fungerer som kilde til rotter i bygninger.

Må kun anvendes af personer, der er autoriserede af Miljøstyrelsen til at bekæmpe rotter med en R1 autorisation."

2) I advarselsfeltet:  
"FARE"

Kan skade det ufødte barn (H360D).

Kan forårsage organskader (blod) ved længerevarende eller gentagen eksponering (H373).

Indhent særlige anvisninger før brug.

Anvend ikke produktet, før alle advarslser er læst og forstået.

Vask huden efter arbejdet.

Der må ikke spises, drikkes eller ryges under brugen af dette produkt.

Bær beskyttelseshandsker.

Brug kemisk resistente beskyttelseshandsker ved brug af produktet." (Handskematerialet skal angives af godkendelsesindehaveren i produktdatabladet).

"Overtrædelse af nedenstående særligt fremhævede forskrifter kan medføre straf:

Må kun anvendes til bekæmpelse af brun rotte (*Rattus norvegicus*), husrotte (*Rattus rattus*) og husmus (*Mus musculus*) i og omkring bygninger og kun i en afstand op til 10 meter fra en bygning samt til bekæmpelse af brun rotte i kloakker. Bekæmpelse er dog også tilladt, hvor en forekomst af rotter længere væk end 10 meter fungerer som kilde til rotter i bygninger.

Må kun anvendes af personer, der er autoriserede af Miljøstyrelsen til at bekæmpe rotter med en R1 autorisation.

Ved kildebekæmpelse skal betingelserne i Miljøstyrelsens vejledning om vurdering og godkendelse af kemiske bekæmpelsesmidler til rotter og mus være opfyldt.

Doseringen af det biocidholdige produkt skal afpasses bekæmpelsen af rotter og mus på den pågældende lokalitet, dog må der højest anvendes 50 g pr. udlægningssted mod rotter og 20 g pr. udlægningssted mod mus af hensyn til faren for forgiftning af mennesker, pattedyr og fugle.

Må ikke anvendes mod andre skadevoldere og ikke i højere doseringer end de i brugsanvisningen nævnte.

For brug i og omkring bygninger og til kildebekæmpelse: Det biocidholdige produkt skal anbringes i aflåste foderstationer, så mennesker og dyr udenfor målgruppen ikke kan komme i kontakt med giften.

For brug i kloakker: Det biocidholdige produkt skal fastgøres i kloakken, så det ikke kan skylles væk.

Døde mus og rotter skal eftersøges to gange ugentligt eller efter behov og bortskaffes som dagrenovation af hensyn til faren for sekundær forgiftning af rovfugle og rovdyr.

Produktet må ikke bruges mere end 35 dage uden en evaluering af angrebets status og effekten af behandlingen.

Giftrester og emballage skal afleveres til den kommunale affaldsordning for farligt affald efter endt bekæmpelse.

Foderstationer skal påføres en tekst, hvoraf det fremgår, at der foregår bekæmpelse af mus og rotter i området, og hvorledes man forholder sig i tilfælde af forgiftning. Foderstationer skal være forsynet med kontaktinformation på dem, som udfører bekæmpelsen.

Må ikke benyttes til permanent bekæmpelse af mus og rotter, såsom forebyggende brug i sikringsordninger.

Ved anbringelse af foderstationer tæt på overfladevand eller afløbssystemer skal det sikres, at det biocidholdige produkt ikke kommer i kontakt med vand.

Må ikke tømmes i kloakafløb.

Opbevares under lås og utilgængeligt for børn.

Må ikke opbevares sammen med fødevarer, drikkevarer og foderstoffer."

#### Farepiktogrammet GHS08

Miljøstyrelsen henleder opmærksomheden på, at godkendelsesindehaver skal anføre oplysninger om førstehjælp, herunder:

"Hvis der er brug for lægehjælp, medbring da beholderen eller etiketten.

VED eksponering eller mistanke om eksponering: Søg lægehjælp.

Dette produkt indeholder et antikoagulerende stof.

Ved indtagelse kan der forekomme forsinkede symptomer, såsom næseblod og blødende tandkød.

I svære tilfælde kan der forekomme blå mærker og blod i afføring eller urin.

Modgift: K1-vitamin; må kun administreres af medicinsk/veterinært personale.

Hvis produktet:

- kommer på huden, vaskes huden med vand og sæbe.
- kommer i øjnene, skyldes øjnene med øjenrensevæske eller vand; hold øjnene åbne i mindst 10 minutter.
- kommer i munden, skyldes munden grundigt med vand.

Giv aldrig en bevidstløs person noget gennem munden.

Fremkald ikke opkastning.

Hvis produktet synkes, søges der straks lægehjælp. Vis produktets beholder eller etiketten til lægen.

Kontakt en dyrlæge, hvis produktet intages af et kåledyr. "

#### 3) I deklarationsfeltet:

a) Teksten "Gnavermiddel BPR-reg. nr. 732-1. Aktivstof og biocidholdigt produkt er godkendt efter biocidforordningen (Forordning (EU) nr. 528/2012)".

b) Oplysning om præparattype: "voksblok" for dette præparat.

c) Indholdet af aktivstof i vægtprocent (% w/w) og g/kg.

d) Udløbsdatoen skal anføres. Denne dato må højst være 2 år efter produktionsdatoen. Etikettens dato kan udformes som en hen-visning til en produktionsdato andetsteds på emballagen.

e) Batchnummer eller – betegnelse skal anføres.

f) Pakningsstørrelse i kg.

g) Godkendelsesindehavers navn og adresse.

#### 4) Brugsanvisningen:

Oplysninger om skadevoldere, anvendelsesområde og doseringer.

Følgende retningslinjer gælder i forhold til bortskaftelse.

Der skal mærkes med sikkerhedssætning P501: "Indholdet/beholderen bortskaffes i overensstemmelse med kommunale regler for affaldshåndtering."

Derudover skal mærkningen ske efter retningslinjerne:

"Tom emballage og rester skal afleveres til den kommunale affaldsordning for farligt affald."

