

Summary of product characteristics for a biocidal product

Product name: CAID BLOCK

Product type(s): PT14 - Rodenticides (Pest control)

Authorisation number: FR-2013-0003

R4BP 3 asset reference number: FR-0000976-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	1
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
5. General directions for use	14
5.1. Instructions for use	14
5.2. Risk mitigation measures	15
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	15
5.4. Instructions for safe disposal of the product and its packaging	16
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	16
6. Other information	16

Administrative information

1.1. Trade names of the product

CAID BLOCK

1.2. Authorisation holder

Name and address of the authorisation holder	Name	LIPHATECH
	Address	Bonnel - CS 10005 47480 Pont Du Casse France
Authorisation number	FR-2013-0003	
R4BP 3 asset reference number	FR-0000976-0000	
Date of the authorisation	04/03/2013	
Expiry date of the authorisation	07/02/2023	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	LIPHATECH SAS
Address of the manufacturer	Bonnel - CS 10005 47480 PONT DU CASSE France
Location of manufacturing sites	Production centre, avenue Jean Serres, ZA Malère 47480 PONT DU CASSE France

1.4. Manufacturer(s) of the active substance(s)

Active substance	14 - Chlorophacinone
Name of the manufacturer	LIPHATECH S.A.S
Address of the manufacturer	Bonnel - CS 10005 47480 PONT DU CASSE France
Location of manufacturing sites	LIPHATECH S.A.S at AlzChem Trostberg GmbH - Chemie Park Trostberg - Dr Albert Frank strasse 32 83308 Trostberg Germany

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Chlorophacinone	Chlorophacinone	Active Substance	3691-35-8	223-003-0	0,005

2.2. Type of formulation

RB - Bait (ready for use)

3. Hazard and precautionary statements

Hazard statements	May damage the unborn child.
Precautionary statements	<p>Obtain special instructions before use.</p> <p>Do not handle until all safety precautions have been read and understood.</p> <p>Wear protective gloves.</p> <p>IF exposed or concerned: Get medical advice.</p> <p>IF exposed or concerned: Get medical attention.</p> <p>Store locked up.</p> <p>Dispose of contents to in accordance with local/regional/national regulation.</p>

4. Authorised use(s)

4.1 Use description

Use 1 - Use # 1 House mice and/or rats – trained professionals – indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: All Scientific name: <i>Rattus norvegicus</i> Common name: Brown rat Development stage: All Scientific name: <i>Rattus rattus</i> Common name: Roof rat Development stage: All
Field(s) of use	Indoor Indoor
Application method(s)	Bait application - Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations - Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations
Application rate(s) and frequencies	- rats: 200 g of bait per bait point. - mice: 100 g of bait per bait point. - 0 - Bait products: - rats: 200 g of bait per bait point. - mice: 100 g of bait per bait point.
Category(ies) of users	Trained professional
Pack sizes and packaging material	Minimum pack size of 3 kg. (In France only: minimum pack size of 5 kg) - Bait formulations: Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton (3 kg-25 kg) ; - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton). Loose bait in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton with integral PE liner (3kg-25 kg) ; - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

4.1.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.
 - *[When available]* Follow any additional instructions provided by the relevant code of best practice.

4.1.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign *[in accordance with the applicable code of good practice, if any]*.
 - Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
 - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
 - Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
 - Do not use the product in pulsed baiting treatments.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.2 Use description

Use 2 - Use # 2 Mice and/or rats – trained professionals – outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Scientific name: *Mus musculus*

Target organism(s) (including development stage)

Common name: House mouse
Development stage: all

Scientific name: Rattus norvegicus
Common name: Brown rat
Development stage: all

Scientific name: Rattus rattus
Common name: Roof rat
Development stage: all

Field(s) of use

Outdoor

Outdoor around buildings

Application method(s)

Bait application -

Bait formulations:

- Ready-to-use bait to be used in tamper-resistant bait stations
- Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations
- Not in France : Direct application into the burrows

Application rate(s) and frequencies

- rats: 200 g of bait per bait point. - mice: 100 g of bait per bait point. - 0 -

Bait products:

- rats: 200 g of bait per bait point.
- mice: 100 g of bait per bait point.

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size of 3 kg.

(In France only: minimum pack size of 5 kg)

- Bait formulations:

- Bait formulations:

Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in:

- PP opaque bucket (3 kg-25 kg) ;
- Opaque cardboard carton (3 kg-25 kg) ;
- Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

Loose bait in:

- PP opaque bucket (3 kg-25 kg) ;
- Opaque cardboard carton with integral PE liner (3kg-25 kg) ;
- Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

4.2.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period [except when applied in burrows].

- [When available] Follow any additional instructions provided by the relevant code of best practice.
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.3 Use description

Use 3 - Use # 3 Mice and/or rats – trained professionals – Outdoor open areas & waste dumps

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mouse
Development stage: Other

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: all

Scientific name: *Rattus rattus*
Common name: Roof rat
Development stage: all

Field(s) of use

Application method(s)	<p>Outdoor</p> <p>Outdoor open areas Outdoor waste dumps (rats only)</p> <p>Bait application - - Ready-to-use bait to be used in tamper-resistant bait stations - Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations - [Not in France: Direct application into the burrows]</p>
Application rate(s) and frequencies	<p>- rats: 200 g of bait per bait point. - mice: 100 g of bait per bait point. - 0 - Bait products: - rats: 200 g of bait per bait point. - mice: 100 g of bait per bait point.</p>
Category(ies) of users	<p>Trained professional</p>
Pack sizes and packaging material	<p>Minimum pack size of 3 kg. (In France only: minimum pack size of 5 kg)</p> <p>- Bait formulations: Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton (3 kg-25 kg) ; - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton). Loose bait in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton with integral PE liner (3kg-25 kg) ; - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).</p>

4.3.1 Use-specific instructions for use

<ul style="list-style-type: none"> - Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding. - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. - Remove the remaining product at the end of treatment period [except when applied in burrows]. - [When available] Follow any additional instructions provided by the relevant code of best practice. - For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
--

4.3.2 Use-specific risk mitigation measures

<ul style="list-style-type: none"> - Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any]. - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
--

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.4 Use description

Use 4 - Use # 4 - Rats - trained professionals - sewers

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: Rattus norvegicus
Common name: Brown rat
Development stage: All

Field(s) of use

Other
Sewers

Application method(s)

Bait application -
- Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water.
- Covered and protected bait points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations

Application rate(s) and frequencies

- 200 grams per manhole - 0 -
Bait products:
- 200 grams per manhole

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size of 3 kg.
(In France only: minimum pack size of 5 kg)

- Bait formulations:
Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in:
 - PP opaque bucket (3 kg-25 kg) ;
 - Opaque cardboard carton (3 kg-25 kg) ;
 - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).
- Loose bait in:
 - PP opaque bucket (3 kg-25 kg) ;
 - Opaque cardboard carton with integral PE liner (3kg-25 kg) ;
 - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

4.4.1 Use-specific instructions for use

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.4.2 Use-specific risk mitigation measures

- [If national policy or legislation requires it] Place baits only in sewer systems which are connected to the sewage treatment plant.
- Do not use this product in pulsed baiting treatments.

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.5 Use description

Use 5 - Use # 5 (not relevant in France) – House mice – professionals – Indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: All
Field(s) of use	Indoor
Application method(s)	Bait application - - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	- 100 g of bait per bait station. - 0 - - 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 1 meter.
Category(ies) of users	Professional
Pack sizes and packaging material	Minimum pack size of 3 kg. - Bait formulations: Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton (3 kg-25 kg) ; - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton). Loose bait in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton with integral PE liner (3kg-25 kg) ; - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

4.5.1 Use-specific instructions for use

- The bait stations should be visited only 2 to 3 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- [When available] Follow any additional instructions provided by the relevant code of best practice

4.5.2 Use-specific risk mitigation measures

-

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.6 Use description

Use 6 - Use # 6 (not relevant in France) – Rats – professionals – indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: All Scientific name: Rattus rattus Common name: Roof rat Development stage: all
Field(s) of use	Indoor Indoor
Application method(s)	Bait application - - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	- 200 g of bait per bait station. - 0 - - 200 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 4 meters.
Category(ies) of users	Professional
Pack sizes and packaging material	Minimum pack size of 3 kg. - Bait formulations: Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in: - PP opaque bucket (3 kg-25 kg) ; - Opaque cardboard carton (3 kg-25 kg) ;

- Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).
- Loose bait in:
- PP opaque bucket (3 kg-25 kg) ;
 - Opaque cardboard carton with integral PE liner (3kg-25 kg) ;
 - Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

4.6.1 Use-specific instructions for use

- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- *[When available]* Follow any additional instructions provided by the relevant code of best practice

4.6.2 Use-specific risk mitigation measures

-

4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.7 Use description

Use 7 - Use # 7 (not relevant in France) - House mice and/or rats – professionals – outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mouse
Development stage: all

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: all

Scientific name: *Rattus rattus*
Common name: Roof rat
Development stage: All

Field(s) of use

Outdoor

Outdoor around buildings

Application method(s)

Bait application -

- Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequencies

Rats: 200 g of bait per bait station. House mice: 100 g of bait per bait station. - 0 -
Rats: 200 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 4 meters.

House mice: 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 1 meters.

Category(ies) of users

Professional

Pack sizes and packaging material

Minimum pack size of 3 kg.

- Bait formulations:

Wrapped blocks (10-140 g) in PP or PE, opaque or transparent sachets in:

- PP opaque bucket (3 kg-25 kg) ;

- Opaque cardboard carton (3 kg-25 kg) ;

- Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

Loose bait in:

- PP opaque bucket (3 kg-25 kg) ;

- Opaque cardboard carton with integral PE liner (3kg-25 kg) ;

- Opaque PE, PP or HDPE prefilled bait stations (up to 60 bait stations in cardboard carton).

4.7.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited *[for mice - at least every 2 to 3 days at]**[for rats - only 5 to 7 days after]* the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- *[When available]* Follow any additional instructions provided by the relevant code of best practice

4.7.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.7.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.7.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.7.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. General directions for use

5.1. Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
 - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
 - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
 - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
 - The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
 - Where possible, bait stations must be fixed to the ground or other structures.
 - Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened.
 - *[If national policy or legislation requires it]* When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
 - Bait should be secured so that it cannot be dragged away from the bait station.
 - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
 - Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
 - Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
 - When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- FOR TRAINED PROFESSIONAL ONLY**
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
 - If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
 - If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- FOR PROFESSIONALS ONLY** Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- FOR PROFESSIONALS ONLY** Remove the remaining bait or the bait stations at the end of the treatment period.
- Do not open the sachets containing the bait.

5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign *[in accordance with the applicable code of good practice, if any]*.
 - The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
 - **FOR TRAINED PROFESSIONAL ONLY** Do not use in areas where resistance to the active substance can be suspected.
 - Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
 - **FOR TRAINED PROFESSIONAL ONLY** Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
 - Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
 - Dispose dead rodents in accordance with local requirements.
- **FOR PROFESSIONAL ONLY** To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
 - **FOR PROFESSIONAL ONLY** Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
 - **FOR PROFESSIONAL ONLY.** The product information (i.e. label and/or leaflet) shall clearly show that:
 - the product shall not be supplied to the general public (e.g. "for professionals only").
 - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
 - **FOR PROFESSIONAL ONLY** Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label [insert country specific information]. Contact a veterinary surgeon in case of ingestion by a pet [insert country specific information]
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [insert national phone number]"
- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months

- In France only : Do not store at temperature above 25°C.

6. Other information

- **(in France only)** : The authorisation holder has to monitor the resistance phenomenon of rodent populations toward the active substance bromadiolone. Results of the resistance monitoring must be submitted at the renewal of the product.
- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.