

# Summary of product characteristics for a biocidal product

**Product name:** Harmonix Rodent Paste

**Product type(s):** PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

**Authorisation number:** 1070-16

**R4BP 3 asset reference number:** DK-0015178-0000

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	2
5. General directions for use	12
5.1. Instructions for use	12
5.2. Risk mitigation measures	13
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	13
5.4. Instructions for safe disposal of the product and its packaging	13
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	13
6. Other information	13

## Administrative information

### 1.1. Trade names of the product

Harmonix Rodent Paste
Harmonix 3D Paste
Harmonix Nager Paste

### 1.2. Authorisation holder

<b>Name and address of the authorisation holder</b>	Name	2022 ENVIRONMENTAL SCIENCE FR SAS
	Address	3, place Giovanni Da Verrazzano 69009 LYON France
<b>Authorisation number</b>	1070-16	
<b>R4BP 3 asset reference number</b>	DK-0015178-0000	
<b>Date of the authorisation</b>	28/07/2020	
<b>Expiry date of the authorisation</b>	24/07/2025	

### 1.3. Manufacturer(s) of the biocidal products

<b>Name of the manufacturer</b>	Bayer SAS
<b>Address of the manufacturer</b>	16 rue Jean-Marie Leclair, CS 90106 69266 Lyon (Cedex 09) France
<b>Location of manufacturing sites</b>	INDUSTRIALCHIMICA Srl. Via Sorgaglia 40 I-35020 Arre Italy
	Kollant S.r.l., via C. Colombo 7/7 30030 Vigonovo (VE) Italy
	IRIS 1126A, avenue du Moulins - Route de Saint Privat F- 30340 SALINDRES France

#### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	1443 - Cholecalciferol
<b>Name of the manufacturer</b>	Bayer S.A.S
<b>Address of the manufacturer</b>	16 rue Jean-Marie Leclair, CS 90106 69266 Lyon (Cedex 09) France
<b>Location of manufacturing sites</b>	Fermenta Biotech Limited, Village Takoli; P.O. Nagwain District Mandi - 175 121 Himachal Pradesh India
	Fermenta Biotech Limited, Z-109 B & C, SEZ II, Dahej ,Taluka - Vagara District Bharuch - 392130 Gujarat India

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Cholecalciferol		Active Substance	67-97-0	200-673-2	0,077

### 2.2. Type of formulation

RB - Bait (ready for use)
---------------------------

## 3. Hazard and precautionary statements

<b>Hazard statements</b>	
<b>Precautionary statements</b>	

## 4. Authorised use(s)

## 4.1 Use description

### Use 1 - House mouse, Norway rat, Roof rat – professionals - indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Norway Rat Development stage: Adults and juveniles  Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults and juvenils  Scientific name: Mus musculus Common name: House mouse Development stage: Adults and juvenils
<b>Field(s) of use</b>	Indoor  Indoor
<b>Application method(s)</b>	Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
<b>Application rate(s) and frequencies</b>	Application Rate: For rats: 100g to 200g of bait per baiting point, For mice: 20g of bait per baiting point. Dilution (%): - Number and timing of application:  For rats:  100-200 g bait per baiting point depending upon infestation level.  If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).  An amount of 200 g of product must be added when bait has been totally consumed within one control interval.  For mice:  20 g bait per baiting point depending upon infestation level.  If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).  An amount of 20 g of product must be added when bait has been totally consumed within one control interval.

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

Bait in individual tea bag sachet, Long fibre paper, 20g

Primary Packaging: Tea bag (cellulose), 20g

Secondary Packaging:

Type of packaging: Bucket,

Size/ Volume of the packaging: up to 10 kg

Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket

Type of the packaging: Box

Size/volume of the packaging: up to 15 kg

Material of the packaging: Inner COEX PET/LDPE plastic sachet in a cardboard box

Type of the packaging: Bag-

Size/volume of the packaging: up to 5 kg

Material of the packaging: Plastic: COEX PET/PA/PE Plastic bag, with handle and reclosable zip

**4.1.1 Use-specific instructions for use**

The bait stations should be visited at least every 1-2 days at the beginning for the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. Carry out pre-baiting survey of the infested area and on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and extent of the infestation.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. Bait should be secured so that it cannot be dragged away from the bait station.

Remove the remaining bait and the bait stations at the end of the treatment period.

**4.1.2 Use-specific risk mitigation measures**

Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. The product information (i.e. label and or leaflet) shall clearly show that:

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek

advice from the product supplier or call a pest control service.  
Do not wash the bait stations with water between applications.  
Do not use product in pulsed baiting treatments.

#### 4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Prevent skin contact when disposing remains of baits.

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Keep away from food, drink and animal feeding stuffs.

### 4.2 Use description

#### Use 2 - House mouse, Norway rat , Roof rat – professionals – Outdoor: around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Norway rat Development stage: Adults and juvenils  Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults and juvenils  Scientific name: Mus musculus Common name: House mouse Development stage: Adults and juvenils
<b>Field(s) of use</b>	Outdoor  Outdoor: around buildings
<b>Application method(s)</b>	Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations (See document CA-Nov16-Doc.4.1c-Final on the concept of tamper-resistant bait stations.)

**Application rate(s) and frequencies**

Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point.  
Dilution (%): -  
Number and timing of application:

For rats:

100-200 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).

An amount of 200 g of product must be added when bait has been totally consumed within one control interval.

For mice:

20 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).

An amount of 20 g of product must be added when bait has been totally consumed within one control interval.

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

Bait in individual tea bag sachet, Long fibre paper, 20g

Primary Packaging: Tea bag (cellulose), 20g

Secondary Packaging:

Type of packaging: Bucket,  
Size/ Volume of the packaging: up to 10 kg  
Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket

Type of the packaging: Box  
Size/volume of the packaging: up to 15 kg  
Material of the packaging: Inner COEX PET/LDPE plastic sachet in a cardboard box

Type of the packaging: Bag-  
Size/volume of the packaging: up to 5 kg  
Material of the packaging: Plastic: COEX PET/PA/PE Plastic bag, with handle and reclosable zip



#### 4.2.1 Use-specific instructions for use

Place the bait stations in areas not liable to flooding.  
The bait stations should be visited at least every 1 to 2 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

Bait should be secured so that it cannot be dragged away from the bait station.  
The campaign has to be terminated if there is no further consumption of baits.  
Remove the remaining bait or the bait stations at the end of the treatment period.

#### 4.2.2 Use-specific risk mitigation measures

Do not apply this product directly in the burrows.

The product information (i.e. label and/or leaflet) shall clearly show that:

- a) the product shall not be supplied to the general public (e.g. "for professionals only").
- b) the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- c) users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

Do not wash the bait stations with water between applications.

Search for and remove dead rodents during treatment, at least as often as baiting stations are inspected.

Dispose dead rodents in household waste or in a rendering plant. Avoid direct contact.

Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Do not use the product in pulsed baiting treatments.

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Prevent skin contact when disposing remains of baits.

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Keep away from food, drink and animal feeding stuffs.

#### 4.3 Use description

**Use 3 - House mouse, Norway rat, Roof rat – trained professionals – Indoor**

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Norway rat Development stage: Adults and juveniles</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults and juveniles</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: Adults and juveniles</p>
<b>Field(s) of use</b>	<p>Indoor</p> <p>Indoor</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point. Dilution (%): - Number and timing of application:</p> <p>For rats:</p> <p>100-200 g bait per baiting point depending upon infestation level.</p> <p>If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).</p> <p>An amount of 200 g of product must be added when bait has been totally consumed within one control interval.</p> <p>For mice:</p> <p>20 g bait per baiting point depending upon infestation level.</p> <p>If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).</p> <p>An amount of 20 g of product must be added when bait has been totally consumed within one control interval.</p> <p>Permanent baiting (please see use-specific instructions for use)</p>
<b>Category(ies) of users</b>	Trained professional

## Pack sizes and packaging material

Bait in individual tea bag sachet, Long fibre paper, 20g

Primary Packaging: Tea bag (cellulose), 20g

Secondary Packaging:

Type of packaging: Bucket,

Size/ Volume of the packaging: up to 10 kg

Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket

Type of the packaging: Box

Size/volume of the packaging: up to 15 kg

Material of the packaging: Inner COEX PET/LDPE plastic sachet in a cardboard box

Type of the packaging: Bag-

Size/volume of the packaging: up to 5 kg

Material of the packaging: Plastic: COEX PET/PA/PE Plastic bag, with handle and reclosable zip

### 4.3.1 Use-specific instructions for use

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Bait should be secured so that it cannot be dragged away from the bait station.

### 4.3.2 Use-specific risk mitigation measures

Do not use the product in pulsed baiting treatments.

The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").

Do not use in areas where resistance to the active substance can be suspected.

Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

see chapter 5.4

#### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

see chapter 5.5

### 4.4 Use description

#### Use 4 - House mouse, Norway rat, Roof rat – trained professionals – outdoor: around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Norway rat Development stage: Adults and juveniles</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults and juveniles</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: Adults and juveniles</p>
<b>Field(s) of use</b>	<p>Outdoor</p> <p>Outdoor: around buildings</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: For rats: 100g to 200g of bait per baiting point; For mice: 20g of bait per baiting point. Dilution (%): - Number and timing of application:</p>

For rats:

100-200 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 10 - 20 linear metres (in case of a low infestation) and 3 – 10 linear meters (in case of a high infestation).

An amount of 200 g of product must be added when bait has been totally consumed within one control interval.

For mice:

20 g bait per baiting point depending upon infestation level.

If more than one bait station is needed, the minimum distance between bait stations should be 5 - 20 linear metres (in case of a low infestation) and 2 – 5 linear meters (in case of a high infestation).

An amount of 20 g of product must be added when bait has been totally consumed within one control interval.

**Category(ies) of users**

Trained professional

**Pack sizes and packaging material**

Bait in individual tea bag sachet, Long fibre paper, 20g

Primary Packaging: Tea bag (cellulose), 20g

Secondary Packaging:

Type of packaging: Bucket,  
 Size/ Volume of the packaging: up to 10 kg  
 Material of the packaging: Inner LDPE plastic sachet in a plastic PP Bucket

Type of the packaging: Box  
 Size/volume of the packaging: up to 15 kg  
 Material of the packaging: Inner COEX PET/ LDPE plastic sachet in a cardboard box

Type of the packaging: Bag-  
 Size/volume of the packaging: up to 5 kg  
 Material of the packaging: Plastic: COEX PET/PA/PE Plastic bag, with handle and reclosable zip

#### 4.4.1 Use-specific instructions for use

Place the baiting points in areas not liable to flooding.  
Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.  
Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.  
The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).  
Bait should be secured so that it cannot be dragged away from the bait station.

#### 4.4.2 Use-specific risk mitigation measures

Do not use this product in pulsed baiting treatments.  
The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").  
Do not use in areas where resistance to the active substance can be suspected.  
Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.

#### 4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

see chapter 5.4

#### 4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

see chapter 5.5

### 5. General directions for use

#### 5.1. Instructions for use

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

Where possible, bait stations must be fixed to the ground or other structures.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3. for the information to be shown on the label).

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation. If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a (anticoagulant) rodenticide, or the use of traps as an alternative control measure.

Bait in sachets: Do not open the sachets containing the bait.

## 5.2. Risk mitigation measures

Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.

## 5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

1) In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by an animal.

2) Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name"; "active substance(s)" and "in case of incident, call a poison centre 82 12 12 12.

3) Hazardous to wildlife.

## 5.4. Instructions for safe disposal of the product and its packaging

- 1) At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.
- 2) Prevent skin contact when disposing remains of baits.

## 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- 1) Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- 2) Store in places prevented from the access of children, birds, pets and farm animals.
- 3) Keep away from food, drink and animal feeding stuffs.
- 4) Shelf-life: 36 months

## 6. Other information

1) Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

2) The product contains a bittering agent and a dye.

[Due to technical deficits of the SPC editor we have to list the following points currently at this point of the SPC:

Cholecalciferol (Vitamin D3, Active substance) Content in BP (w/w%) 0.077 (technical), 0.075 (pure)

Minimum purity of the active substance according to Regulation (EU) 2019/637: 970 g/kg

[(Ph. Eur.\* 7.0; 01/2008:0575 corrected 6.5): min. 970 g/kg (includes also any precholecalciferol present)]

Oplysninger til etiket og brugsanvisning for Harmonix Rodent Paste, BPR-reg. nr. 1070-16

I. Etiketten skal udformes i overensstemmelse med det godkendte resumé af det biocidholdige produkts egenskaber, jf. artikel 69, stk. 1, i biocidforordningen [1] (Forordning (EU) nr. 528/2012), og CLP-Forordningen [2].

II. Det er udelukkende ansøgers/godkendelsesindehaverens ansvar, at etiket, mærkning og pakning lever op til lovens krav, jf. biocidforordningen artikel 69. Etiket og mærkning skal være på dansk.  
Nedenstående tekst i afsnit III er bidrag til overholdelsen af reglerne og således kun en del af de krav, som stilles til blandt andet etiketten.

III. Etiketten skal indeholde nedenstående oplysninger. Oplysninger i citationstegn skal angives ordret:

1) I hovedfeltet:

"Gnavermiddel

Må kun anvendes til bekæmpelse af brun rotte (*Rattus norvegicus*), husrotte (*Rattus rattus*), husmus (*Mus musculus*) i og omkring bygninger og kun i en afstand op til 10 meter fra en bygning. Bekæmpelse er dog også tilladt, hvor en forekomst af rotter længere væk end 10 meter fungerer som kilde til rotter i bygninger.

Må kun anvendes af personer, der er autoriserede af Miljøstyrelsen til at bekæmpe rotter med enten en R1 eller R2 autorisation."

2) I advarselsfeltet:

"**FORSIGTIG**

Vask huden grundigt efter brug.

Overtrædelse af nedenstående særligt fremhævede forskrifter kan medføre straf:

**Må kun anvendes til bekæmpelse af brun rotte (*Rattus norvegicus*), husrotte (*Rattus rattus*), husmus (*Mus musculus*) i og omkring bygninger og kun i en afstand op til 10 meter fra en bygning. Bekæmpelse er dog også tilladt, hvor en forekomst af rotter længere væk end 10 meter fungerer som kilde til rotter i bygninger.**

**Må kun anvendes af personer, der er autoriserede af Miljøstyrelsen til at bekæmpe rotter med enten en R1 eller R2 autorisation.**

Ved kildebekæmpelse skal betingelserne i Miljøstyrelsens vejledning om vurdering og godkendelse af kemiske bekæmpelsesmidler til rotter og mus være opfyldt.

Doseringen af det biocidholdige produkt skal afpasses bekæmpelsen af mus og rotter på den pågældende lokalitet, dog må der højst anvendes 20 g pr. udlægningssted for mus og 200 g pr. udlægningssted for rotter af hensyn til faren for forgiftning af mennesker, pattedyr og fugle.

Må ikke anvendes mod andre skadevoldere og ikke i højere doseringer end de i brugsanvisningen nævnte.

Det biocidholdige produkt skal anbringes i aflåste foderstationer, så mennesker og dyr udenfor målgruppen ikke kan komme i kontakt med giften.

Døde mus og rotter skal eftersøges to gange ugentligt eller efter behov og bortskaffes som dagrenovation af hensyn til faren for sekundær forgiftning af rovfugle og rovdyr.

Produktet må ikke bruges mere end 35 dage uden en evaluering af angrebets status og effekten af behandlingen.

Giftrester og emballage skal afleveres til den kommunale affaldsordning for farligt affald efter endt bekæmpelse.

Foderstationer skal påføres en tekst, hvoraf det fremgår, at der foregår bekæmpelse af mus og rotter i området, og hvorledes man forholder sig i tilfælde af forgiftning. Foderstationer skal være forsynet med kontaktinformation på dem, som udfører bekæmpelsen.

Må ikke benyttes til permanent bekæmpelse af mus og rotter, såsom forebyggende brug i sikringsordninger.

Ved anbringelse af foderstationer tæt på overfladevand eller afløbssystemer skal det sikres, at det biocidholdige produkt ikke kommer i kontakt med vand.

Opbevares utilgængeligt for børn

Må ikke opbevares sammen med fødevarer, drikkevarer og foderstoffer.

Emballagen må ikke genanvendes"

Miljøstyrelsen henleder opmærksomheden på, at godkendelsesindehaver skal anføre oplysninger om førstehjælp, herunder:

"Indeholder cholecalciferol Der findes ingen modgift. Ved forgiftning skal symptombehandles.

Ved indtagelse, skyl munden. Ved indtagelse eller mistanke om forgiftning, ring til en giftinformation eller en læge.



Hvis et dyr uden for målgruppen indtager produktet, ring til en dyrlæge."

3) I deklarationsfeltet:

- a) Teksten "Gnavermiddel BPR-reg. nr. 1070-16. Aktivstof og biocidholdigt produkt er godkendt efter biocidforordningen (Forordning (EU) nr. 528/2012)".
- b) Oplysning om præparattype: "pasta" for dette præparat.
- c) Indholdet af aktivstof i vægtprocent (% w/w) og g/kg.
- e) Udløbsdatoen skal anføres. Denne dato må højst være tre år efter produktionsdatoen. Etikettens dato kan udformes som en hen-visning til en produktionsdato andetsteds på emballagen.
- f) Batchnummer eller – betegnelse skal anføres.
- g) Pakningsstørrelse i kg.
- h) Godkendelsesindehavers navn og adresse.

4) Brugsanvisningen:

Oplysninger om skadevoldere, anvendelsesområde og doseringer.

Følgende retningslinjer gælder i forhold til bortskaffelse.

Der skal mærkes med sikkerhedssætning P501: "Indholdet/holderen bortskaffes i overensstemmelse med kommunale regler for affaldshåndtering."

Derudover skal mærkningen ske efter retningslinjerne:

Tom emballage og rester skal afleveres til den kommunale affaldsordning for farligt affald.