

**EN**

***ANNEX***

**SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT**

Vertox Oktablok

**Product type(s)**

PT14: Rodenticides

**Authorisation number:** IE/BPA 70533

**R4BP asset number:** IE-0000528-0000

---

## 1. ADMINISTRATIVE INFORMATION

### 1.1. Trade name(s) of the product

Trade name(s)	Vertox Oktablok
---------------	-----------------

### 1.2. Authorisation holder

Name and address of the authorisation holder	Name	PelGar International Limited
	Address	18 rue des Remparts d'Ainay 69002 Lyon France
Authorisation number	IE/BPA 70533	
<i>R4BP asset number</i>	IE-0000528-0000	
Date of the authorisation	18/07/2013	
Expiry date of the authorisation	01/07/2026	

### 1.3. Manufacturer(s) of the product

Name of manufacturer	PelGar International Limited
Address of manufacturer	Unit 13 Newman Lane GU34 2QR Alton United Kingdom of Great Britain and Northern Ireland (the)
Location of manufacturing sites	Unit 13 Newman Lane GU34 2QR Alton United Kingdom of Great Britain and Northern Ireland (the)

### 1.4. Manufacturer(s) of the active substance(s)

Active substance	Brodifacoum
Name of manufacturer	PelGar International Limited
Address of manufacturer	Unit 13 Newman Lane GU34 2QR Alton United Kingdom of Great Britain and Northern Ireland (the)
Location of manufacturing sites	Prazska 54 280 02 Kolin Czechia

---

## 2. PRODUCT COMPOSITION AND FORMULATION

### 2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	active substance	56073-10-0	259-980-5	0,005

### 2.2. Type(s) of formulation

RB Bait (ready for use)

### 3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	<p>H373: May cause damage to organs blood through prolonged or repeated exposure { 1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard: }.</p> <p>H360D: May damage the unborn child.</p> <p>EUH401: To avoid risks to human health and the environment, comply with the instructions for use.</p> <p>EUH208: Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction.</p>
Precautionary statements	<p>P201: Obtain special instructions before use.</p> <p>P202: Do not handle until all safety precautions have been read and understood.</p> <p>P280: Wear protective gloves.</p> <p>P308+P313: IF exposed or concerned: Get medical advice/attention.</p> <p>P314: Get medical attention if you feel unwell.</p> <p>P405: Store locked up.</p> <p>P501: Dispose of contents to in accordance with national regulations.</p>

## 4. AUTHORISED USE(S)

### 4.1. Use description

**Table 1. Professional - indoors and outdoors (around buildings)**

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: Adults and juveniles  Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Adults and juveniles  Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: Adults and juveniles
Field(s) of use	indoor use outdoor use  Indoors Outdoors - around buildings
Application method(s)	Method: bait application  Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: RATS: 10 to 60 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be 5 metres. MICE: 5 to 20 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be 2 metres.  Dilution (%): N/A  Number and timing of application:  Mice High infestation – 5 to 20g bait in bait stations every 2 metres Low infestation – 5 to 20g bait in bait stations every 5 metres Rats High infestation – 10 to 60g bait in bait stations every 5 metres Low infestation – 10 to 60g bait in bait stations every 10 metres  Standard baiting program (max 35 days - see Directions for use)..
Category(ies) of users	professional
Pack sizes and packaging material	5g, 10g, 20g, 28g, 50g and 60g blocks loose or wrapped in PP or PE for amateur use and loose 5g, 10g, 20g, 28g, 50g , 60g, 100g

or 200g blocks for professional use, within the following outer packaging:

- PE or PP tub or pail

Minimum pack size 3.0 kg

**-\* Please note IRELAND applies a minimum Professional pack size of 2.5 Kg**

Maximum pack size: 20kg

Loose 5g, 10g, 20g, 28g, 50g, 60g, 100g or 200g blocks either unlined or within a PP or wire-tied PE bag within a double-walled or fibreboard carton

Minimum pack size 3.0 kg

**-\* Please note IRELAND applies a minimum Professional pack size of 2.5 Kg**

Maximum pack size: 20kg

5g, 10g, 20g, 28g, 50g or 60g blocks.

Fibreboard carton 'uni-trays' (moulded styrene tray with pop-out blocks)

**Minimum pack size 3.0 kg**

**-\* Please note IRELAND applies a minimum Professional pack size of 2.5 Kg**

Maximum pack size: 12kg

2, 3 or 4 x 5g blocks packed in a single or multi-use tamper-proof mouse HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper.

Multiple mouse boxes within pack:

10g - 250

15g - 125

20g – 96, 120, 144

30g – 96, 120, 144

40g – 72, 96, 120

60g – 48, 60, 72, 96

80g – 32, 48, 60, 72, 96

1 or 2 x 10g or 1 x 20g blocks packed in a single or multi-use tamper-proof mouse PET, HIPS, PVC, PE, HDPE or PP bait station, all packed in multiples of 1, 2, 3, 4, 5 or 6 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Additional blocks may be included as refills in the outer.

Multiple mouse boxes within pack:

10g - 250

20g – 96, 120, 144

30g – 96, 120, 144

40g – 72, 96, 120

50g - 48, 60, 72, 96

60g – 48, 60, 72, 96

80g – 32, 48, 60, 72, 96

100g - 32, 48, 60, 72, 96

120g - 32, 48, 60, 72, 96

1, 2 or 3 x 20g blocks packed in a single or multi-use tamper-proof rat PET, HIPS, PVC, PE, HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper

Additional blocks may be included as refills in the outer.

Multiple rat boxes within pack:

20g – 96, 120, 144  
 40g – 72, 96, 120  
 60g – 48, 60, 72, 96  
 80g – 32, 48, 60, 72, 96  
 100g - 32, 48, 60, 72, 96  
 120g - 32, 48, 60, 72, 96  
 240g – 16, 32, 48, 60, 72, 96  
 50g blocks packed in a single or multi-use tamper-proof rat HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper  
 Multiple rat boxes within pack:  
 50g – 60, 72, 96  
 80g – 32, 48, 60, 72, 96  
 100g - 32, 48, 60, 72, 96  
 200g – 16, 32, 48, 60, 72, 96  
 1 -20 x 20g in a PE sachet or PE, PVC or HIPS clamshell/blister packed in an outer with a multi-use PE, PP, PET, HIPS, PVC tamperproof rodent bait station.  
 Refill pack - Blister pack, pouch or sachet in a cardboard outer.  
 Multiple units within pack:  
 80g – 32, 48, 60, 72, 96  
 100g - 32, 48, 60, 72, 96  
 200g – 16, 32, 48, 60, 72, 96  
 400g – 8, 16, 32, 48, 60, 72, 96  
 Refill pack - Blister pack, pouch or sachet (paper/PE, AL/PE, paper/Al/PE, PE, PP, PET, HIPS, PVC blister pack, pouch or sachet) in a cardboard outer.  
 Multiple units within pack:  
 80g – 32, 48, 60, 72, 96  
 100g - 32, 48, 60, 72, 96  
 200g – 16, 32, 48, 60, 72, 96  
 400g – 8, 16, 32, 48, 60, 72, 96

#### 4.1.1. Use-specific instructions

##### MICE

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- [When available] Follow any additional instructions provided by the relevant code of best practice.

##### RATS

- The bait stations should be visited at least every 5 to 7 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- [When available] Follow any additional instructions provided by the relevant code of best practice.

##### WHEN USED OUTDOORS

- Protect bait from the atmospheric conditions (eg. rain, snow, etc.). Place the bait stations in areas not liable to flooding.

- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

---

#### 4.1.2. Use-specific risk mitigation measures

#### 4.1.3. Use-specific risk mitigation measures

- Do not apply this product directly in burrows.

#### 4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

INDOOR

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

OUTDOOR

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

N/A

#### 4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

N/A

### 4.2. Use description

**Table 2. Trained professional - indoors and outdoors (around buildings) and in sewers**

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: Adults and juveniles  Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Adults and juveniles  Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: Adults and juveniles
Field(s) of use	indoor use outdoor use  Indoors Outdoors - around buildings In sewers



Application method(s)	<p>Method: bait application</p> <p>Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations Covered and protected baiting points Direct application of ready-to-use bait into the burrow SEWERS- Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water.</p>
Application rate(s) and frequency	<p>Application rate: RATS: 10 to 60 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be 5 metres. MICE: 5 to 20 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be 2 metres.</p> <p>Dilution (%): N/A</p> <p>Number and timing of application:</p> <p>Mice  High infestation – 5 to 20g bait in bait stations every 2 metres  Low infestation – 5 to 20g bait in bait stations every 5 metres  Rats  High infestation – 10 to 60g bait in bait stations every 5 metres  Low infestation – 10 to 60g bait in bait stations every 10 metres</p> <p>Standard baiting program (max 35 days - see Directions for use)..</p> <p>Pulsed baiting</p>
Category(ies) of users	trained professional
Pack sizes and packaging material	<p>5g, 10g, 20g, 28g, 50g and 60g blocks loose or wrapped in PP or PE for amateur use and loose 5g, 10g, 20g, 28g, 50g , 60g, 100g or 200g blocks for professional use, within the following outer packaging:</p> <p>- PE or PP tub or pail  Minimum pack size 3.0 kg  <b>-* Please note IRELAND applies a minimum Professional pack size of 2.5 Kg</b>  Maximum pack size: 20kg  Loose 5g, 10g, 20g, 28g. 50g, 60g, 100g or 200g blocks either unlined or within a PP or wire-tied PE bag within a double-walled or fibreboard carton  Minimum pack size 3.0 kg  <b>-* Please note IRELAND applies a minimum Professional pack size of 2.5 Kg</b>  Maximum pack size: 20kg  5g, 10g, 20g, 28g, 50g or 60g blocks.  Fibreboard carton 'uni-trays' (moulded styrene tray with pop-out blocks)  Minimum pack size 3.0 kg  <b>-* Please note IRELAND applies a minimum Professional pack size of 2.5 Kg</b>  Maximum pack size: 12kg</p>

2, 3 or 4 x 5g blocks packed in a single or multi-use tamper-proof mouse HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper.

Multiple mouse boxes within pack:

10g - 250  
15g - 125  
20g - 96, 120, 144  
30g - 96, 120, 144  
40g - 72, 96, 120  
60g - 48, 60, 72, 96  
80g - 32, 48, 60, 72, 96

1 or 2 x 10g or 1 x 20g blocks packed in a single or multi-use tamper-proof mouse PET, HIPS, PVC, PE, HDPE or PP bait station, all packed in multiples of 1, 2, 3, 4, 5 or 6 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper. Additional blocks may be included as refills in the outer.

Multiple mouse boxes within pack:

10g - 250  
20g - 96, 120, 144  
30g - 96, 120, 144  
40g - 72, 96, 120  
50g - 48, 60, 72, 96  
60g - 48, 60, 72, 96  
80g - 32, 48, 60, 72, 96  
100g - 32, 48, 60, 72, 96  
120g - 32, 48, 60, 72, 96

1, 2 or 3 x 20g blocks packed in a single or multi-use tamper-proof rat PET, HIPS, PVC, PE, HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper.

Additional blocks may be included as refills in the outer.

Multiple rat boxes within pack:

20g - 96, 120, 144  
40g - 72, 96, 120  
60g - 48, 60, 72, 96  
80g - 32, 48, 60, 72, 96  
100g - 32, 48, 60, 72, 96  
120g - 32, 48, 60, 72, 96  
240g - 16, 32, 48, 60, 72, 96

50g blocks packed in a single or multi-use tamper-proof rat HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper.

Multiple rat boxes within pack:

50g - 60, 72, 96  
80g - 32, 48, 60, 72, 96  
100g - 32, 48, 60, 72, 96  
200g - 16, 32, 48, 60, 72, 96

1 -20 x 20g in a PE sachet or PE, PVC or HIPS clamshell/blister packed in an outer with a multi-use PE, PP, PET, HIPS, PVC tamperproof rodent bait station.

Multiple units within pack:

80g - 32, 48, 60, 72, 96  
100g - 32, 48, 60, 72, 96  
200g - 16, 32, 48, 60, 72, 96

---

400g – 8, 16, 32, 48, 60, 72, 96 Refill pack - Blister pack, pouch or sachet (paper/PE, AL/PE, paper/Al/PE, PE, PP, PET, HIPS, PVC blister pack, pouch or sachet) in a cardboard outer. Multiple units within pack: 80g – 32, 48, 60, 72, 96 100g - 32, 48, 60, 72, 96 200g – 16, 32, 48, 60, 72, 96 400g – 8, 16, 32, 48, 60, 72, 96
---

#### **4.2.1. Use-specific instructions**

- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- If used for pulsed baiting: - Replace eaten bait only after 3 days and then at maximum 7 day intervals. Collect any spilled bait and dead rodents.

#### **OUTDOOR**

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period [Not applicable where explicitly authorised for permanent baiting].
- [When available] Follow any additional instructions provided by the relevant code of best practice.

---

## SEWER USE

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.

## COVERED/PROTECTED BAIT POINTS

- [For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species].

[When available] Follow any additional instructions provided by the relevant code of best practice.

## DIRECT APPLICATION IN BURROWS

- Baits must be placed to minimise the exposure to non-target species and children.
- Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

## PULSED BAITING

- Replace eaten bait only after 3 days and then at maximum 7 day intervals. Collect any spilled bait and dead rodents.
- [When available] Follow the specific instructions provided by the applicable code of good practice at national level.

### **4.2.2. Use-specific risk mitigation measures**

### **4.2.3. Use-specific risk mitigation measures**

---

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

#### SEWER USE

- [If national policy or legislation requires it] Place baits only in sewer systems which are connected to the sewage treatment plant.

#### DIRECT APPLICATION IN BURROWS

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign [in accordance with the applicable code of good practice, if any]

#### **4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

##### OUTDOOR

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### **4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging**

N/A

#### **4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

N/A

---

## 5. GENERAL DIRECTIONS FOR USE<sup>1</sup>

### 5.1. Instructions for use

#### PROFESSIONALS

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened
- [If national policy or legislation require it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- [Wear protective chemical resistant gloves during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.

#### TRAINED PROFESSIONALS

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened.

---

<sup>1</sup>Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

- 
- [If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
  - Bait should be secured so that it cannot be dragged away from the bait station.
  - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
  - Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
  - Wear protective chemical resistant gloves during product handling phase.
  - When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
  - The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
  - If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
  - If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

## **5.2. Risk mitigation measures**

### **PROFESSIONALS**

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any]
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week). [Where relevant, specify if more frequent or daily inspection is required].
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:  
the product shall not be supplied to the general public (e.g. "for professionals only").  
the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").  
users shall properly label bait stations (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

### **TRAINED PROFESSIONALS**

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign [in accordance with the applicable code of good practice, if any]".
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment

- 
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
  - Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
  - Dispose dead rodents in accordance with local requirements.

### **5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
  - Antidote: Vitamin K1 administered by medical/veterinary personnel only.
  - In case of:
    - Dermal exposure, wash skin with water and then with water and soap.
    - Eye exposure, rinse eyes with eye-rinse liquid or water, keep eyes lids open at least 10 minutes.
    - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person.
- Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [*insert national phone number*]".
  - Hazardous to wildlife.

### **5.4. Instructions for safe disposal of the product and its packaging**

- At the end of the treatment, dispose of the uneaten bait and the packaging in accordance with local requirements.

### **5.5. Conditions of storage and shelf-life of the product under normal conditions of storage**

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 2 years



---

## 6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.
- Use Biocides Safely and Sustainably
- It is illegal to use this product for uses or in a manner other than that prescribed on this label.
- Poison Information: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.
- Please note IRELAND applies a minimum Professional pack size of 2.5 Kg instead of 3.0 kg.