

Annex XV Report

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE AS A CMR CAT 1 OR 2, PBT, VPVB OR A SUBSTANCE OF AN EQUIVALENT LEVEL OF CONCERN

Substance Name: 1,3,5-Trichlorobenzene

EC Number: 203-608-6

CAS Number: 108-70-3

Submitted by: Germany

Version: August 2010

CONTENTS

PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE AS A CMR CAT 1 OR 2, PBT, VPVB OR A SUBSTANCE OF AN EQUIVALENT LEVEL OF CONCERN	4
PART I.....	5
JUSTIFICATION	5
1 IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES	5
1.1 Name and other identifiers of the substance	5
1.2 Composition of the substance	5
1.3 Physico-chemical properties.....	5
2 HARMONISED CLASSIFICATION AND LABELLING.....	8
3 ENVIRONMENTAL FATE PROPERTIES.....	9
3.1 Degradation	9
3.1.1 Abiotic degradation	9
3.1.1.1 Hydrolysis	9
3.1.1.2 Phototransformation /Photolysis.....	9
3.1.1.2.1 Phototransformation in air.....	9
3.1.1.2.2 Phototransformation in water.....	9
3.1.2 Biodegradation	9
3.1.2.1 Screening tests.....	9
3.1.2.2 Simulation tests	10
3.1.3 Summary and discussion on degradation.....	11
3.2 Environmental distribution	11
3.2.1 Adsorption/desorption	11
3.2.2 Volatilisation	11
3.2.3 Distribution modelling	11
3.3 Bioaccumulation.....	16
3.3.1 Aquatic bioaccumulation.....	16
3.3.1.1 Bioaccumulation estimation	16
3.3.1.2 Measured bioaccumulation data	17
3.3.2 Summary and discussion of bioaccumulation	18
4 HUMAN HEALTH HAZARD ASSESSMENT.....	18
5 ENVIRONMENTAL HAZARD ASSESSMENT	19
5.1 Aquatic compartment (including sediment).....	19
5.1.1 Toxicity data.....	19
5.1.1.1 Fish.....	19
5.1.1.1.1 Long-term toxicity to fish.....	19
5.1.1.2 Aquatic invertebrates.....	20
5.1.1.2.1 Long-term toxicity to aquatic invertebrates.....	20
5.1.1.3 Algae and aquatic plants.....	20
5.1.1.4 Sediment organisms	20
5.1.1.5 Other aquatic organisms	20
6 CONCLUSIONS ON THE SVHC PROPERTIES	21

6.1	PBT, vPvB assessment	21
6.2	CMR assessment.....	21
6.3	Substances of equivalent level of concern.	21
6.3.1	Assessment of PBT/vPvB properties - comparison with the criteria of Annex XIII	21
6.3.1.1	Persistence	21
6.3.1.2	Bioaccumulation.....	21
6.3.1.3	Toxicity	22
6.3.1.4	Long-range transport potential	22
6.3.2	Summary and overall conclusions on the PBT/vPvB properties	22
PART II		23
INFORMATION ON USE, EXPOSURE, ALTERNATIVES AND RISKS		23
REFERENCES		25

TABLES

Table 1: Summary of physico- chemical properties	5
Table 2: Half-lives of TCB in the sewage sludge-amended soil and standard spiked soil in microcosm experiments (Wang and Jones, 1994)	10
Table 3: Distribution in WWTP (Simple Treat 3.0)	11
Table 4: Input parameters for both models	12
Table 5: Characteristic Travel Distances [km]	13
Table 6: Spatial ranges.....	14
Table 7: Available bioconcentration factors (BCF)	17
Table 8: Long-term toxicity to fish.....	19
Table 9: Toxicity to algae and aquatic plants	20
Table 10: Ecotoxicity to other aquatic organisms.....	20

FIGURES

Figure 1: Characteristic Travel Distance (CTD) and overall persistence (P_{ov}) of 1,3,5-Trichlorobenzene	13
Figure 2: Characteristic Travel Distances compared to PCB 28, PCB 153 and PCB 101	14
Figure 3: Relative steady-state concentration distribution of 1,3,5-Trichlorobenzene	15
Figure 4: Relative steady-state concentration distribution of PCB 28.....	16

**PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE AS A
CMR CAT 1 OR 2, PBT, VPVB OR A SUBSTANCE OF AN
EQUIVALENT LEVEL OF CONCERN**

Substance Name: 1,3,5-Trichlorobenzene

EC Number: 203-608-6

CAS number: 108-70-3

- *It is proposed to identify the substance as substance of equivalent concern according to Article 57 (f).*

Summary of how the substance meets the CMR (Cat 1 or 2), PBT or vPvB criteria, or is considered to be a substance of an equivalent level of concern

Taking into account all available information on persistent, bioaccumulative and toxic properties of 1,3,5-trichlorobenzene, it is concluded that 1,3,5-trichlorobenzene is a PBT-substance. The substance is considered to fulfil the P criterion, and also the vP-criterion. In addition to that, also the B criterion is fulfilled. In case that lipid normalization is performed, 1,3,5-trichlorobenzene still meets the B criterion of Annex XIII. When focusing on data with higher reliability, 1,3,5-trichlorobenzene would not clearly fulfil the B criterion. However, if doing so the maximum BCF will still be close to the B criterion of Annex XIII. Although 1,3,5-trichlorobenzene is highly toxic for aquatic organisms tested on the long-term, it does not formally meet the T-criterion. The overall conclusion is however drawn in a weight of evidence approach, taking supporting evidence into account relating to the very high long-range environmental transport potential of this substance. It raises an equivalent concern according to Article 57 (f).

Registration number(s) of the substance or of substances containing the substance:

PART I

JUSTIFICATION

1 IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES

1.1 Name and other identifiers of the substance

Chemical Name: Benzene, 1,3,5-trichloro-

EC Name: 1,3,5-trichlorobenzene

EC Number 203-608-6

CAS Number: 108-70-3

IUPAC Name: 1,3,5-trichlorobenzene

1.2 Composition of the substance

The detailed composition of the substance is confidential and provided in the technical dossier.

Chemical Name: Benzene, 1,3,5-trichloro-


EC Number: 203-608-6

CAS Number: 108-70-3

IUPAC Name: 1,3,5-trichlorobenzene

Molecular Formula: $C_6H_3Cl_3$

Structural Formula:



Molecular Weight: 181.45 g/mol

Typical concentration (% w/w): Min. 99 % (w/w)

Concentration range (% w/w): -

1.3 Physico-chemical properties

Table 1: Summary of physico- chemical properties

REACH ref Annex, §	Property	IUCRID section	Value	[enter comment/reference or delete column]
VII, 7.1	Physical state at 20°C and 101.3 kPa	4.1	Solid, white crystals or needles	HSDB, 2010
VII, 7.2	Melting/freezing point	4.2	62.8 °C	HSDB, 2010
VII, 7.3	Boiling point	4.3	208 °C	HSDB, 2010
VII, 7.5	Vapour pressure	4.6	0.24 mm Hg at 25 °C	HSDB, 2010
VII, 7.7	Water solubility	4.8	6.01 mg/L at 25 °C	HSDB, 2010
VII, 7.8	Partition coefficient n-octanol/water (log value)	4.7 partition coefficient	log K _{OW} = 4.19	HSDB, 2010
XI, 7.16	Dissociation constant	4.21	---	
VII, 7.4	Density	4.4	1.456 g/m ³ at 20 °C	HSDB, 2010
IX, 7.17	Viscosity	4.22	0.841 mPa x s at 70 °C	HSDB, 2010
	Henry's Law Constant	5.4.2	966.03 Pa x m ³ / mol at 25 °C	Calculation
VII, 7.9	Flash point	4.11	107 °C,(close cup)	PTB-AG 3.43, CHEMSAFE, 2009
VII, 7.10	Flammability	4.13	<p>Pyrophoric properties: The classification procedure needs not to be applied because the organic substance is known to be stable into contact with air at room temperature for prolonged periods of time (days).</p> <p>Flammability in contact with water: The classification procedure needs not to be applied because the organic substance does not contain metals or metalloids.</p>	<p>BAM, II.2 (2010)</p> <p>BAM, II.2 (2010):</p>
VII, 7.11	Explosive properties	4.14	The classification procedure needs not to be applied because there are no chemical groups present in the molecule which are associated with explosive properties.	BAM, II.2 (2010)
VII, 7.12	Relative Self-ignition temperature for solids	4.12	The classification procedure needs not to	BAM, II.2 (2010)

ANNEX XV REPORT – IDENTIFICATION OF SVHC

			be applied because the organic substance has a melting point ≤ 160 °C,	
VII, 7.13	Oxidizing properties	4.15	The classification procedure needs not to be applied because contained chlorine atoms are chemically bonded only to carbon.	BAM, II.2 (2010)

2 HARMONISED CLASSIFICATION AND LABELLING

No entry in Table 3.1/3.2 of Annex VI of Regulation (EC) 1272/2008 as amended by the 1st ATP.

3 ENVIRONMENTAL FATE PROPERTIES

3.1 Degradation

3.1.1 Abiotic degradation

3.1.1.1 Hydrolysis

Due to its chemical structure 1,3,5-trichlorobenzene (1,3,5-TCB) is not expected to hydrolyse under normal environmental conditions. This assumption is supported by the experimental data (Korte and Freitag, 1986) on hydrolysis of 1,2,4-trichlorobenzene (1,2,4-TCB) at 50°C, which has the same chemical composition and nearly the same chemical structure.

3.1.1.2 Phototransformation /Photolysis.

3.1.1.2.1 Phototransformation in air.

Indirect photochemical degradation in the troposphere is considered to be very slow based on an estimated OH radical reaction rate of $0.679 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1} \text{ molecules}^{-1}$, which corresponds to an atmospheric half-life of about 24 days for the reaction with OH-radicals (AOPWIN v1.91; 24 h day^{-1} ; OH radical concentration of $5 \cdot 10^5 \text{ OH} \cdot \text{cm}^{-3}$). A QSAR Model Reporting Format and a QSAR Prediction Reporting Format are annexed to this document in order to document the model prediction.

3.1.1.2.2 Phototransformation in water.

The environmental exposure occurs in the whole water column. Because of the substance's behaviour it will predominantly volatilize from water or adsorb at suspended organic matter and sediment. Photodegradation of 1,3,5-TCB is expected to be a relevant degradation process only in very shallow clear waters and in the first few centimetres layer of the water column. Therefore aquatic photodegradation is not considered to have relevant impact on the overall persistency of 1,3,5-TCB in the environment.

Based on the available data, 1,2,3-TCB has a low degradation rate in the atmosphere and is not expected to hydrolyse under environmental conditions.

3.1.2 Biodegradation

3.1.2.1 Screening tests

The ready biodegradability was studied with a method corresponding to the OECD TG 301C, Modified MITI (I) test (MITI-List, 2002). The test concentration was 100 mg/l and activated sludge concentration 30 mg/l. In the aerobic study, the degradation measured as Biochemical Oxygen Demand (BOD) was 0% after 14 days.

3.1.2.2 Simulation tests

Ramanand et al. studied the metabolism of individual chlorinated benzenes in mineral salts medium (Ramanand et al., 1993). 1,2,4-TCB were dechlorinated to 1,4 dichlorobenzene and 1,3 dichlorobenzene and 1,2,3 TCB were dechlorinated to 1,2 dichlorobenzene and 1,3dichlorobenzene. 1,3,5-TCB was not transformed after an incubation of 5 months. A reason for no transformation could be that a chlorine substituent, which is located between other chlorine substituents is lost more readily (Fathepure et al., 1988). Also Morita observed that the reaction of hydroxylation depends on the configuration of the chlorine substituents in the benzene ring (Morita, 1977).

All three isomers of TCB were reductively dechlorinated by enrichment cultures in the presence of lactate, glucose, ethanol or isopropanol as electron donator (Holliger et al., 1992). The inoculum was material from percolation columns filled with river sediment in which dechlorination of TCB occurred. With first-generation enrichment and lactate as electron donator, the lag phase for combined TCB isomers was 100 days. With second-generation enrichment and lactate, the lag phase for 1,2,3-TCB was 27 days, >138 days for 1,2,4-TCB and >365 days for 1,3,5-TCB. For dechlorination a temperature optimum between 25 and 30 °C and a pH-optimum of about 7.2 was found. The dechlorinating activity was lost if no electron donator was added.

The behaviour and fate of chlorobenzenes in spiked and sewage sludge-amended soil was tested by Wang and Jones (Wang and Jones, 1994). For testing the fate, four different experimental conditions were investigated: (1) normal; (2) sterilized; (3) sterilized and shaded; (4) sterilized, shaded and sealed. Under condition (4) no decrease of concentration was observed. Whereas under condition (1)-(3) the concentration of TCB decreased. This indicated volatilization as main loss process. The loss process in soil occurs in two steps (two step first order kinetic model). In the first step the TCB volatilize to the atmosphere over short periods. The second step was much slower and was presumably controlled by the rate of compound desorption from soil. A proportion of the TCB will stay in the soil for much longer periods. The concentration in spiked soil decreased faster than in sludge amended soil.

Table 2: Half-lives of TCB in the sewage sludge-amended soil and standard spiked soil in microcosm experiments (Wang and Jones, 1994)

Compound	First step				Second step			
	Sewage sludge-amended soil		Standard spiked soil		Sewage sludge-amended soil		Standard spiked soil	
	Loss (%)	DisDT ₅₀ (d)	Loss (%)	DisDT ₅₀ (d)	Loss (%)	DisDT ₅₀ (d)	Loss (%)	DisDT ₅₀ (d)
1,2,4-TCB	83.2	22.5	64.5	12.5	2.26	49500	19.8	194
1,2,3-TCB	88.6	22.2	73.0	15.0	4.45	1370	12.8	222
1,3,5-TCB	85.3	23.7	71.2	10.5	6.22	239	14.8	238

Additionally, a read-across approach was carried out, comparing 1,3,5-TCB with its structural analogues 1,2,4-TCB and 1,2,3-TCB. A Reporting format is annexed to this document which contains a more detailed discussion of the read-across approach (Reporting format for the analogue approach).

Biodegradability was assessed based on structural considerations, evaluation of the available data and evaluation of possible biodegradation pathways.

It is expected that environmental degradation of all three trichlorobenzene isomers proceeds via the same biodegradation pathways. The observed biodegradation is generally slow, with 1,3,5-trichlorobenzene being slightly more persistent than the other isomers. The most experimental results are available for 1,2,4-trichlorobenzene, which is persistent in soil and sediment. 1,3,5-

trichlorobenzene is expected to show very similar properties with respect to biodegradability, but is expected to degrade slower to some extent.

3.1.3 Summary and discussion on degradation.

1,3,5-TCB has a low degradation rate in the atmosphere and is not expected to hydrolyse under environmental conditions. The substance was shown to be persistent in soil. Read-across shows that the biodegradability is similar to the close structural analogue 1,2,4-TCB, which is persistent in soil and sediment. The degradation rate of 1,3,5-TCB is expected to be slightly slower than the degradation rate of 1,2,4-TCB.

In conclusion, 1,3,5-TCB is regarded to fulfil the P and the vP criteria of Annex XIII of the REACH regulation.

3.2 Environmental distribution

3.2.1 Adsorption/desorption

Based on model estimates a K_{OC} of 3118 L/kg has been estimated (ECHA, 2008). Therefore a high adsorption potential and a low mobility in soil is expected for 1,3,5-Trichlorobenzene. Due to the slow degradation in soil, the substance may contaminate groundwater by leaching through sandy soils with low organic carbon content.

3.2.2 Volatilisation

Calculation of the Henry constant using the equation $HENRY=VP*MOLW/SOL$ from the Guidance document for Chemical Safety Assessment (ECHA, 2008) and the substance properties from table 1 results in a Henry constant of $966.03 \text{ Pa}\cdot\text{m}^3\cdot\text{Mol}^{-1}$. This indicates that the substance is highly volatile from water. Due to the high adsorption potential to organic matter, volatilisation from soil and sludge is expected to be lower.

The distribution coefficient $K_{air,water}$ (Henry coefficient) calculated from the Henry constant is 0.397.

3.2.3 Distribution modelling

Distribution in Waste Water Treatment Plant

The modelling of the distribution in a municipal Waste Water Treatment Plant was done with SimpleTreat 3.0 (debugged version, 7 Feb 1997) and the result of the Screening Test on ready biodegradation (substance is not biodegradable; $k=0/h$).

Table 3: Distribution in WWTP (Simple Treat 3.0)

Summary of distribution	(Percent)
To air	72.2
To water	7.3
Via primary sludge	19.5
Via surplus sludge	0.9
Degraded	0.0
Total	100

The result of the distribution modelling in a WWTP indicates that nearly 72 percent of the 1,3,5-TCB found in the influent waste water is emitted to air.

Long-range Transport

An important indicator for the environmental persistence can be the long-range transport to remote areas. For the calculation of the long-range transport potential of 1,3,5-Trichlorobenzene two different multimedia models have been used. On the one hand the OECD Pov and LRTP Screening Tool (OECD, 2006) has been used and on the other hand the model ChemRange (Scheringer, 1996) has been used. Both models require only a few input parameters. These are the partition coefficients octanol/water, K_{ow} , the dimensionless Henry's Law Constant, K_{aw} and the environmental degradation half-lives in air, water and soil.

Unfortunately there is only one measured DisT50 available for 1,3,5-Trichlorobenzene. This is the DisT50 in soil. Although, it is well known that this DisT50 represents not only degradation this value was used in a calculation. 1,3,5-Trichlorobenzene is benchmarked as "non" biodegradable. According to the TGD (ECHA 2008) the half-lives in soil and water have to be considered as infinite. However the models require a discrete value for the simulation. Due to this fact the half-lives according to the TGD for inherent biodegradation in water of 150 d and 3000 d in soil have been used. Furthermore, the half-lives in water have been calculated on the one hand according to Arnot (Arnot et al., 2005) and on the other hand using the QSAR-model CATALOGIC (OASIS Catalogic v5.10.7, 2009). Finally four calculations with different half-lives for water and soil has been carried out to show the influence of these half-lives on the long-range transport potentials of 1,3,5-Trichlorobenzene.

Table 4: Input parameters for both models

Input parameter	1,3,5-TCB (TGD)	1,3,5-TCB (Arnot)	1,3,5-TCB (Catalogic)	1,3,5-TCB (measured soil)
logK _{ow}	4.19	4.19	4.19	4.19
logK _{aw}	-0.402	-0.402	-0.402	-0.402
t _{1/2} (air) [d]	23.637	23.637	23.637	23.637
t _{1/2} (water) [d]	150	91,025	82	150
t _{1/2} (soil) [d]	3000	273,075	246	238

Results using the OECD P_{ov} and LRTP Screening Tool:

Figure 1 shows that all four calculations yield nearly the same values for characteristic travel distance (CTD) and overall persistency (P_{ov}). Substances with CTD and P_{ov} values of this magnitude are considered to show a high long-range transport potential.




Figure 1: Characteristic Travel Distance (CTD) and overall persistence (P_{ov}) of 1,3,5-Trichlorobenzene

In order to classify the calculated long-range transport potential of 1,3,5-Trichlorobenzene the results have been compared with the long-range transport potential of the well known POPs PCB 28, PCB 101 and PCB 153, regulated under the Stockholm Convention.. In Table 5 the characteristic travel distances (CTD) are listed:

Table 5: Characteristic Travel Distances [km]

Substance	CTD [km]
1,3,5-Trichlorobenzene (Arnot)	11638
1,3,5-Trichlorobenzene (Catalogic)	11635
1,3,5-Trichlorobenzene (measured soil)	11648
1,3,5-Trichlorobenzene (TGD)	11649
PCB 153	12629
PCB 101	8595
PCB 28	3298

Figure 2 illustrates that the long-range transport potential of 1,3,5-Trichlorobenzene is in the same order of magnitude as the long-range transport potential of all three PCBs.




Figure 2: Characteristic Travel Distances compared to PCB 28, PCB 153 and PCB 101

Results using ChemRange:

In contrast to the OECD P_{ov} and LRTP Screening Tool, ChemRange calculates the spatial range of a chemical. The spatial range is calculated as the fraction of earth circumference which can be reached by a substance. The long-range transport is calculated for the surface compartments soil and water and air. In order to benchmark the calculated long-range transport potential of 1,3,5-Trichlorobenzene the results have been compared with the long-range transport potential of the well known POPs PCB 28, PCB 101 and PCB 153 regulated under the Stockholm Convention. Using a release scenario to air the calculated spatial ranges are listed in Table 6:

Table 6: Spatial ranges

Substance	Range soil (Fraction of earth circumference)	Range water (Fraction of earth circumference)	Range air (Fraction of earth circumference)
1,3,5-Trichlorobenzene (Arnot)	0.36	0.36	0.36
1,3,5-Trichlorobenzene (Catalogic)	0.36	0.36	0.36
1,3,5-Trichlorobenzene (measured soil)	0.36	0.37	0.36
1,3,5-Trichlorobenzene (TGD)	0.36	0.37	0.36
PCB 28	0.29	0.30	0.29
PCB 101	0.32	0.33	0.32

PCB 153	0.40	0.40	0.40
---------	------	------	------

Comparison of results

The calculated spatial ranges for 1,3,5-TCB using a release scenario to soil are very similar.

The high long-range transport potential of 1,3,5-Trichlorobenzene and PCB 28 is illustrated by Figure 3 and Figure 4: Both diagrams are similar and show a shape which is characteristic for substances with a high long-range transport potential. In case of a substance with a low long-range transport potential the shape of the plot would show just a sharp peak at the source point.




Figure 3: Relative steady-state concentration distribution of 1,3,5-Trichlorobenzene




Figure 4: Relative steady-state concentration distribution of PCB 28

Conclusions:

Due to its high volatility (Henry constant of $966.03 \text{ Pa}\cdot\text{m}^3\cdot\text{Mol}^{-1}$), 1,3,5-TCB is expected to be distributed from water to air. In the air, 1,3,5-TCB is persistent (estimated half life 24 days) and may be transported via air for several thousand kilometres. Distributed by this pathway also to remote areas, 1,3,5-TCB will be deposited there again to soil or water, where it may be even more persistent, e.g. due to lower temperatures.

Model calculations confirm the expected potential for long-range transport, as both the OECD P_{ov} and LRTP Screening Tool and ChemRange yield a long-range transport potential that is in the same order of magnitude for 1,3,5-Trichlorobenzene and for the POPs PCB 101, 153 and 28. Thus, it can be concluded that 1,3,5-Trichlorobenzene is persistent in air and has a high long-range transport potential.

3.3 Bioaccumulation

3.3.1 Aquatic bioaccumulation

3.3.1.1 Bioaccumulation estimation

The log K_{ow} of 4.19 is indicating that bioconcentration in aquatic organisms may occur.

3.3.1.2 Measured bioaccumulation data

Table 7: Available bioconcentration factors (BCF)

Organism	Exposure mg/L	Exposure days	BCF whole	Lipidcontent	Reference	Rel.
<u>Fish:</u>						
<i>Salmo gairdneri</i>	2.3 *10 ⁻⁶	119	1800 ¹ (ss)	8.3%	(Oliver and Niimi, 1983)	3
<i>Salmo gairdneri</i>	45 *10 ⁻⁶	105	4100 ¹ (ss)	8.8%		
<i>Poecilia reticulata</i>	note ²	not given	756 ²	5.4%	(Geyer et al., 1985)	4
<i>Cyprinus carpio</i>	0.0025	42	150-1700 ³	not given	(MITI-List, 2002)	2
	0.025	42	620-1620 ³			
<u>Invertebrates:</u>						
<i>Lymnaea stagnalis</i> (Pond snail)	0.022	10	2	0.5%	(Legierse et al., 1998)	2

Reliability scores: 1) reliable without restriction, 2) reliable with restriction, 3) not reliable, 4) not assignable,

¹ This study was deemed not reliable because fish were exposed simultaneously to 12 substances and the overall concentration may have elicited adverse effects on fish (conc. of 1,3,5-trichlorobenzene = 2.3 / 45 ng/L, overall conc. = 116 / 2544 ng/L); ² Comparison of literature-reported BCF values for diverse fish species and varying lipid contents; ³ A mixture of three isomeric trichlorobenzenes was administered and the individual BCF were calculated. There is no further information given on lipid normalization of BCF.

There are only few data on bioaccumulation of 1,3,5-trichlorobenzene available. In a study conducted by Oliver and Niimi (1983) trout with an initial weight of about 250 g were exposed to two different overall concentrations of a mixture containing 12 chlorinated organic compounds. The uptake period was extended to 119 days and 105 days, respectively, and the trout reached a final weight of about 400 g. Food was also analyzed for contamination with trichlorobenzenes and blank values were considered in BCF calculation. For the lower exposure concentration a BCF of 1800 was reported while for the higher exposure concentration the BCF reached 4100. However, this study was deemed not reliable because 12 substances were simultaneously administered and effects regarding mixture toxicity influencing bioaccumulation can not be excluded due to the elevated background level of additional contaminants.

Geyer et al. (1985) compared a selection of literature reported BCF values for trichlorobenzenes and the correlation with the lipid content of fish. For 1,3,5-trichlorobenzene a single BCF of 700 (determined using Guppy) is included in this overview.

The MITI database also provides BCF values for 1,3,5-trichlorobenzene determined for carp while using a standard test system (MITI-List, 2002). Though a mixture of all three isomeric trichlorobenzenes was administered in two different overall concentrations, the individual concentrations of each isomer in exposure medium as well as in fish were analyzed. The reported BCF values range between 150 and 1700 but no information on neither lipid content nor lipid normalization are given.

A further bioaccumulation test on pond snails revealed a negligible bioaccumulation (Legierse et al., 1998).

In summary, BCF values with at least reliability scoring 4 can be stated within the range from 150 to 1700. Notwithstanding the low reliability of the study by Oliver and Niimi a maximum BCF of 4100 was also observed. Therefore a BCF of >2000 can be hypothesized as a realistic worst case estimate.

In case that lipid normalization to 5% is performed the maximum BCF will reach 2330 (Oliver and Niimi, 1983). Unfortunately a normalization of the BCF with the highest reliability (MITI list, 2002) can not be performed because no original lipid contents of fish are provided. When solely considering BCF from MITI database the maximum BCF still range to 1600 – 1700 indicating a high bioaccumulation potential of 1,3,5-trichlorobenzene.

However, as the most of the few available data are not reliable it is a preferable option to assess the bioaccumulation potential through read-across using the better suited data on 1,2,4-trichlorobenzene.

read across

From a mechanistic point of view the three trichlorobenzene isomers are expected to show similarly high BCF values. The vast majority of deviations in determined BCF may be attributed to the inherent variability of experimental studies (different fish species etc.) rather than different physico-chemical properties of these constitutional isomers. For each isomer the observed BCF values are of limited reliability. Nonetheless, the results indicate that all three isomers show a similar, high bioaccumulation potential. Furthermore, the data indicate that 1,3,5-trichlorobenzene is slightly more bioaccumulative than 1,2,4- and 1,2,3-trichlorobenzene. Most results are available for 1,2,4-trichlorobenzene. These data have already been evaluated in the EU Risk Assessment and the PBT Working Group. A BCF of >2000 is derived as a realistic worst case for this substance. In analogy to this approach, a BCF of >2000 is also expected for 1,3,5-trichlorobenzene. These estimates are considered reliable within the given variance of data.

3.3.2 Summary and discussion of bioaccumulation

The log K_{ow} of 4.19 is indicating that bioconcentration in aquatic organisms may occur. The few observed BCF values are of limited reliability, with most BCF values ranging from 150 up to 1700. Although of low reliability, in one study (Oliver and Niimi, 1983) a maximum BCF of 4100 was also reported, so a BCF of > 2000 can be derived as a realistic worst case estimate. This is also confirmed by the read-across approach, which indicates that all three trichlorobenzene isomers show a similar, high bioaccumulation potential, with 1,3,5-trichlorobenzene being potentially slightly more bioaccumulative than its structural analogues due to its higher log K_{ow} . Hence, the variability of BCF can rather be attributed to the inherent variability of experimental studies (different fish species etc.) than to different inherent properties of these constitutional isomers. According to the read-across approach, 1,3,5-trichlorobenzene is expected to be slightly more bioaccumulative than 1,2,4-trichlorobenzene, for which a BCF of >2000 is assumed.

In case that lipid normalization to 5% is applied a maximum BCF of 2330 will result. Unfortunately for the second highest BCF values (MITI List, 2002) having reliability scoring 2 no lipid normalization can be performed due to lacking original lipid content of fish.

4 HUMAN HEALTH HAZARD ASSESSMENT

Not relevant for this dossier.

5 ENVIRONMENTAL HAZARD ASSESSMENT

5.1 Aquatic compartment (including sediment)

5.1.1 Toxicity data

There is very few and only less reliable data on the ecotoxicity of 1,3,5-TCB available. The assessment of the ecotoxicity of 1,3,5-TCB is difficult because the effect concentrations were in the range or in excess of the solubility of the test substance.

However, a read-across approach was carried out, comparing 1,3,5-TCB with its close structural analogues 1,2,4-TCB and 1,2,3-TCB. A Reporting format is provided as Annex 1 to this report, which contains a more detailed discussion of the read-across approach (Reporting format for the analogue approach).

Based on the chemical structure, all three trichlorobenzene isomers are expected to exhibit baseline (nonpolar narcosis) toxicity. Toxicity of nonpolar narcotics is largely determined by lipophilicity and thus by the $\log K_{ow}$. As all trichlorobenzenes are isomers they show very similar $\log K_{ow}$ values and thus, based on lipophilicity, their ecotoxicity is expected to be very similar.

In this section only test results of long-term studies are reported as those are relevant for comparison with the aquatic T-criterion of Annex XIII of the Reach Regulation.

5.1.1.1 Fish

5.1.1.1.1 Long-term toxicity to fish

For the investigation of long-term toxicity of 1,3,5-TCB to fish there is only one test available. This test is not a real long-term toxicity test and derives only an EC_{50} and not a NOEC. Thus, a read-across approach to 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene is applied.

With respect to long-term toxicity to fish, comparable values for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene are available. Both substances show a high long-term toxicity to *Danio rerio*, with 1,2,4-trichlorobenzene having a slightly smaller NOEC (28d-NOEC = 0.077 mg/l for 1,2,3-TCB and 21d-NOEC = 0.04 mg/l for 1,2,3-TCB; see more details in the Reporting format for the analogue approach). Based on structural considerations and the available data, the corresponding NOEC for 1,3,5-trichlorobenzene is expected to be in the same range, but potentially slightly higher. Thus, for the assessment, a NOEC of about 0.077 mg/l is used.

Table 8: Long-term toxicity to fish

Species	Duration	EC_{50} (mg l ⁻¹)	NOEC (mg l ⁻¹)	Method, conditions	Reference
<i>Poecilia reticulata</i>	14 d	3.3		2-3 month old fish; 25 mg CaCO ₃ /L; covered with glass; acetone or propanol-2; semi-static test; 8 fish per concentration; 22°C	(Könemann, 1981)

5.1.1.2 Aquatic invertebrates

For aquatic invertebrates there is no test available with 1,3,5-TCB. Data on toxicity to *Daphnia magna* are available for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene.

5.1.1.2.1 Long-term toxicity to aquatic invertebrates

With respect to long-term toxicity, 1,2,3 -trichlorobenzene and 1,2,4-trichlorobenzene show also similar toxicity (21d-NOEC = 0.03 mg/l and 21d-NOEC = 0.06 mg/l). For 1,3,5-trichlorobenzene, toxicity is expected to be in a similar range, but potentially slightly lower. Thus, a NOEC of 0.06 mg/L (read across from 1,2,4-TCB) is used for 1,3,5-trichlorobenzene. Further details are documented in the Reporting format for the analogue approach.

5.1.1.3 Algae and aquatic plants

There are two tests with algae available. Both are shorter than recommended by OECD test methods. The test performed by Chen et al., 2009 is valid despite the short duration of 48 hours. A closed test system is used. The result is an EC₅₀ of 1.68 mg/L and an EC₁₀ of 0.398 mg/L. The NOEC could not be estimated but is below 0.730 mg/L. This is in good agreement with the NOEC (96h) of 0.37 mg/L from 1,2,4-TCB with the same species.

Table 9: Toxicity to algae and aquatic plants

Species	Duration	EC ₅₀ (mg l ⁻¹)	NOEC (mg l ⁻¹)	Method, conditions	Reference
<i>Ankistrodesmus falcatus</i>	4 h	9.07		5000 lux; 4-7 10 ⁵ cells/mL; Acetone; flasks capped with rubber stoppers wrapped in aluminium foil	(Wong et al., 1984)
<i>Pseudokirchmeriella subcapitata</i>	48 h	1.68	< 0.730 EC ₁₀ =0.398	Triplicate; closed-system with no headspace and low bicarbonate buffer content, growth rate, biomass	(Chen et al., 2009)

5.1.1.4 Sediment organisms

5.1.1.5 Other aquatic organisms

Table 10: Ecotoxicity to other aquatic organisms

Species	Duration	EC ₅₀ (mg l ⁻¹)	Method, conditions	Reference
<i>Tetrahymena pyriformis</i> (Protozoa)	24 h	30	30 °C; DMSO; microscope and Coulter Counter;	(Yoshioka et al., 1985)

6 CONCLUSIONS ON THE SVHC PROPERTIES

6.1 PBT, vPvB assessment

Not relevant for this dossier. See section 6.3.

6.2 CMR assessment

Not relevant for this dossier.

6.3 Substances of equivalent level of concern.

6.3.1 Assessment of PBT/vPvB properties - comparison with the criteria of Annex XIII

6.3.1.1 Persistence

1,3,5-trichlorobenzene (1,3,5-TCB) is not subject to hydrolysis. The screening biodegradation tests indicate that 1,3,5-TCB is not readily biodegradable. Based on data from soil degradation tests half-lives of more than 200 days have been determined.

Due to its high volatility (Henry constant of $966.03 \text{ Pa}\cdot\text{m}^3\cdot\text{Mol}^{-1}$), 1,3,5-TCB is expected to be distributed from water to air. In the air, 1,3,5-TCB is persistent (estimated half life 24 days) and may be transported via air for several thousand kilometres. Distributed by this pathway also to remote areas, 1,3,5-TCB will be deposited again to soil or water being even more persistent in regions with lower temperatures.

Thus, considering all information available, 1,3,5-TCB is concluded to fulfil the P-criterion and the vP criterion.

6.3.1.2 Bioaccumulation

The log Kow of 4.19 is indicating that bioconcentration in aquatic organisms may occur. Available BCF values range from 150 up to 1800 and 4100, though the latter value is of limited reliability. From these data a BCF of > 2000 can be derived as a realistic worst case estimate, but it is also possible to derive additional supporting BCF values via read-across from 1,2,4-trichlorobenzene.

From a mechanistic point of view the three trichlorobenzene isomers are expected to show similarly high BCF values. Hence, the variability of BCF can rather be attributed to the inherent variability of experimental studies (different fish species etc.) than to different inherent properties of these constitutional isomers. Most experimental data are available for 1,2,4-trichlorobenzene. A BCF of >2000 was derived as a realistic worst case for this substance, leading to the conclusion that the substance is bioaccumulative. In analogy to this approach, a BCF of > 2000 is also expected for 1,3,5-trichlorobenzene. Thus, based on a weight of evidence approach considering both the available data for 1,3,5-trichlorobenzene and data for its structural analogues, it is concluded that 1,3,5-trichlorobenzene fulfils the B-criterion of Annex XIII.

In case that lipid normalization to 5% is applied, the maximum BCF of 1,3,5-trichlorobenzene will reach 2330. When focusing on data with higher reliability the maximum BCF are still at about 1600 – 1700, although no lipid normalization can be done due to lacking lipid contents of fish. This high BCF still supports an equivalent concern according to Article 57 (f) of REACH regulation.

6.3.1.3 Toxicity

Only few data of limited reliability are available for the ecotoxicity of 1,3,5-TCB. Thus, a read-across approach was carried out, comparing 1,3,5-TCB with its close structural analogues 1,2,4-TCB and 1,2,3-TCB. All three trichlorobenzene isomers are expected to exhibit baseline (nonpolar narcosis) toxicity which is largely determined by lipophilicity and thus by the $\log K_{ow}$. As all trichlorobenzenes are isomers they show very similar $\log K_{ow}$ values. Hence, based on lipophilicity, the ecotoxicity is expected to be very similar for all three TCB. 1,2,4 and 1,2,3-TCB show a high long-term toxicity. Based on structural considerations and the available data, the corresponding NOEC for 1,3,5-trichlorobenzene is expected to be in the same range. Thus, for the assessment, a long-term NOEC of about 0.06 mg/l for zebra fish is derived from the read-across approach.. Although the available information from read-across indicates a high toxicity for aquatic organisms on the long-term, the T-criterion is formally not fulfilled.

6.3.1.4 Long-range transport potential

Due to its volatility and persistence to atmospheric degradation, 1,3,5-TCB has a very high potential for long-range atmospheric transport. Based on model calculations, estimated travelling distances are several thousand kilometres.

6.3.2 Summary and overall conclusions on the PBT/vPvB properties

Taking into account all available information on persistent, bioaccumulative and toxic properties of 1,3,5-trichlorobenzene, it is concluded that 1,3,5-trichlorobenzene is a PBT- substance. The substance is considered to fulfil the P criterion, and also the vP-criterion. In addition to that, also the B criterion is fulfilled. In case that lipid normalization is performed, 1,3,5-trichlorobenzene still meets the B criterion of Annex XIII. When focusing on data with higher reliability, 1,3,5-trichlorobenzene would not clearly fulfil the B criterion. However, if doing so the maximum BCF will still be close to the B criterion of Annex XIII. Although 1,3,5-trichlorobenzene is highly toxic for aquatic organisms tested on the long-term, it does not formally meet the T-criterion. The overall conclusion is however drawn in a weight of evidence approach, taking supporting evidence into account relating to the very high long-range environmental transport potential of this substance. It raises an equivalent concern according to Article 57 (f).

PART II

INFORMATION ON USE, EXPOSURE, ALTERNATIVES AND RISKS

INFORMATION ON MANUFACTURE, IMPORT/EXPORT AND USES – CONCLUSIONS ON EXPOSURE

1,2,4-TCB is mainly used as intermediate and as a process solvent in closed systems. In addition, the RAR identified the following minor uses that are by now restricted in the EU (Directive 2005/59/EC): solvent, dye carrier, corrosion inhibitor and lubricant. In the past, 1,2,4-TCB was used as an additive in dielectric fluids. 1,2,3-TCB and 1,3,5-TCB have a lower production volume, but are assigned to the same uses. As trichlorobenzenes are priority substances under the Water Framework Directive, their occurrence in rivers is controlled by monitoring programmes.

All substances are preregistered with an envisaged registration deadline of 2010. Though the preregistration data have to be interpreted cautiously, this indicates a potential production volume of more than 100 tons for 1,2,4-TCB (classified as R50-53 according to directive 67/548/EEC) and more than 1000 tons for 1,2,3-TCB and 1,3,5-TCB.

The main uses as an intermediate and as a process solvent in closed applications are not considered to result in a release of 1,2,4-TCB into the environment. All other uses are restricted.

Despite the extensive restriction on the marketing and use of 1,2,4-TCB in the EU, the substance may enter the EU market via imported articles. For example, textiles may contain 1,2,4-TCB from use as a dye carrier. However, there is no information on the amount of imported articles containing 1,2,4-TCB.

Furthermore, 1,2,3-TCB and 1,3,5-TCB are not restricted and may be used as substitutes for 1,2,4-TCB. Again, updated information would be helpful to clarify this concern.

With regard to the known uses of the three Trichlorobenzenes, current exposure is expected to be low, but trichlorobenzenes should be considered for further regulatory steps due to their SVHC properties: Both 1,2,4-TCB and 1,2,3-TCB have been identified as PBTs by the TCNES PBT subgroup. 1,3,5-TCB shows similar properties and should thus also be regarded as PBT. These substances pose a risk to human health and the environment that is not predictable in the long-term. Once released into the environment, adverse effects caused by PBTs cannot be remediated. Consequently, strong precautionary measures are necessary to prevent the release of these substances, though it is expected that only small amounts of trichlorobenzenes are emitted in the EU. As no safe concentration may be derived for PBTs, even low emissions give rise to concern and need to be limited further. Nonetheless, the production volume has to be considered when setting priorities for regulatory measures for concurrent SVHCs. In addition to that, the long-range transport potential of the trichlorobenzenes needs to be considered in a weight of evidence assessment of the PBT-properties including all available information.

CURRENT KNOWLEDGE ON ALTERNATIVES

RISK-RELATED INFORMATION

REFERENCES

- Arnot J, Gouin T, Mackay D. 2005. Practical method for estimating environmental biodegradation rates.
- BUA. 1988. 1,3,5-Trichlorbenzene. GDCh-Advisory Committee on Existing Chemicals of Environmental Relevance (Beratergremium für Umweltrelevante Altstoffe).
- Chen CY, Wang YJ, Yang CF. 2009. Estimating low-toxic-effect concentrations in closed-system algal toxicity tests. *Ecotoxicology and Environmental Safety* 72(5): 1514-1522.
- ECHA. 2008. Guidance on information requirements and chemical safety assessment: Chapter R.16: Environmental Exposure Estimation.
- Fathepure BZ, Tiedje JM, Boyd SA. 1988. Reductive dechlorination of hexachlorobenzene to tri- and dichlorobenzenes in anaerobic sewage sludge. *Appl Environ Microbiol* 54(2): 327-330.
- Furay VJ, Smith S. 1995. Toxicity and QSAR of Chlorobenzenes in Two Species of Benthic Flatfish, Flounder (*Plachthys flesus* L.) and Sole (*Solea solea* L.). *Bull Environ Contam Toxicol* 54(1): 36-42.
- Gaffney PE. 1976. Carpet and rug industry case study II. biological effects. *J Water Pollut Control Fed* 48(12): 2731-2737.
- Geyer H, Scheunert I, Korte F. 1985. Relationship between the lipid content of fish and their bioconcentration potential of 1,2,4-trichlorobenzene. *Chemosphere* 14(5): 545-555.
- Holliger C, Schraa G, Stams AJM, Zehnder AJB. 1992. Enrichment and properties of an anaerobic mixed culture reductively dechlorinating 1,2,3-trichlorobenzene to 1,3-dichlorobenzene. *Appl Environ Microbiol* 58(5): 1636-1644.
- HSDB - Hazardous Substances Data Bank, USA, 2010.
- Hulzebos EM, Adema DMM, Dirven-van Breemen EM, Henzen L, van Dis WA, Herbold HA, Hoekstra JA, Baerselman R, van Gestel CAM. 1993. Phytotoxicity studies with *Lactuca sativa* in soil and nutrient solution. *Environ Toxicol Chem* 12: 1079-1094.
- Könemann H. 1981. Quantitative structure-activity relationships in fish toxicity studies Part 1: Relationship for 50 industrial pollutants. *Toxicology* 19(3): 209-221.
- Korte F, Freitag D. 1986. Kriterien zur Auswahl umweltgefährlicher alter Stoffe: Mobilität einschließlich Abbaubarkeit und Akkumulation. *Umweltforschungsplan des Bundesministerium des Inneren, Chemikalien*.
- Legierse KCHM, Sijm DTHM, Van Leeuwen CJ, Seinen W, Hermens JLM. 1998. Bioconcentration kinetics of chlorobenzenes and the organophosphorus pesticide chlorthion in the pond snail *Lymnaea stagnalis* - A comparison with the guppy *Poecilia reticulata*. *Aquatic Toxicology* 41(4): 301-323.
- MITI-List. 2002. Biodegradation and Bioaccumulation of Existing Chemical Substances under the Chemical Substance Control Law. National Institute of Technology and Evaluation, Japan.

- Morita M. 1977. Chlorinated benzenes in the environment. *Ecotoxicology and Environmental Safety* 1(1): 1-6.
- OASIS Catalogic v5.10.7. 2009.
- OECD. 2006. The OECD Pov and LRTP Screening Tool 2.0. Software and Manual. OECD, Paris.
- Oliver BG, Niimi AJ. 1983. Bioconcentration of chlorobenzenes from water by rainbow trout: correlations with partition coefficients and environmental residues. *Environ Sci Technol* 17(5): 287-291.
- Ramanand K, Balba MT, Duffy J. 1993. Reductive dehalogenation of chlorinated benzenes and toluenes under methanogenic conditions. *Appl Environ Microbiol* 59(10): 3266-3272.
- Ribo JM, Kaiser KLE. 1983. Effects of selected chemicals to photoluminescent bacteria and their correlations with acute and sublethal effects on other organisms *Chemosphere* 12(11-12): 1421-1442.
- Scheringer M. 1996. Persistence and spatial range as endpoints of an exposure-based assessment of organic chemicals. *Environmental Science and Technology* 30(5): 1652-1659.
- Wong PTS, Chau YK, Rhamey JS, Docker M. 1984. Relationship between water solubility of chlorobenzenes and their effects on a freshwater green alga. *Chemosphere* 13(9): 991-996.
- Yoshioka Y, Ose Y, Sato T. 1985. Testing for the toxicity of chemicals with *Tetrahymena pyriformis*. *The Science of The Total Environment* 43(1-2): 149-157.

ANNEX 1

Reporting Format for the analogue approach for 1,2,3-Trichlorobenzene and 1,3,5-Trichlorobenzene

CHEMICALS: 1,2,3-Trichlorobenzene and 1,3,5-Trichlorobenzene

EC No: 201-757-1 and 203-608-6

CAS No: 87-61-6 and 108-70-3

HYPOTHESIS OF THE ANALOGUE APPROACH

The analogue approach is mainly based on the structural isomer 1,2,4-trichlorobenzene, which was already evaluated in the EU Risk Assessment and the PBT Working Group. For 1,2,3-trichlorobenzene, read-across to 1,2,4-trichlorobenzene was already carried in the PBT Working Group. To support the approach, data on further chlorobenzenes were taken into account for some endpoints.

Mechanistic reasoning

Trichlorobenzenes have comparable structural features as they only differ in the substitution pattern of the three chlorine atoms and show very similar physico-chemical properties (see data matrix). For Trichlorobenzenes no specific metabolism or mode of action is expected for bioconcentration and aquatic ecotoxicity, respectively. Thus, these properties are assumed to depend mainly on physico-chemical properties like lipophilicity, which allows to read-across from 1,2,4-trichlorobenzene to its structural isomers.

The position of substituents may influence the chemical reactivity of aromatic compounds. For chlorine, the negative inductive effect (electron-withdrawal due to higher electronegativity) outweighs the mesomeric effect (electron-donation due to resonance effects), resulting in a weak electron-withdrawing effect, which decreases reactivity towards electrophiles. Furthermore, chlorine atoms are ortho- and para-directing substituents, i.e. substituents that direct electrophiles to attack preferably at the ortho- and para position. For 1,3,5-trichlorobenzene, each of the three unsubstituted carbon atoms is in ortho- or para-position to all three substituted carbon atoms. Contrary, for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene, each of the three unsubstituted carbon atoms is in meta-position to either one or two substituted carbon atoms. Thus, the overall reactivity of chlorobenzenes depends primarily on the number, but also on the position of substituents, with the deactivating effect with respect to reaction with electrophiles being weaker for 1,3,5-trichlorobenzene as compared to the other trichlorobenzene isomers. Consequently, 1,3,5-trichlorobenzene is less susceptible to nucleophilic attacks as compared to the other two isomers.

Endpoint	Hypothesis
k_{OH}	All trichlorobenzene isomers show the same mechanism with respect to OH radical reaction, which is OH radical addition to the aromatic ring. Chlorine substituents are expected to reduce reactivity due to the electron-withdrawing-effect. Thus, reaction rate constants are expected to be smaller for the highly chlorinated benzenes. The varying position of the chlorine substituents is expected to cause only minor differences with

	respect to reaction rates. The reaction rate constant for 1,3,5-trichlorobenzene is expected to be slightly higher than the respective rate constants of the other two trichlorobenzene isomers. This assumption is supported by QSAR results.
DT50	Biodegradation of trichlorobenzenes proceeds via reductive dechlorination or hydroxylation; a slightly larger half-life for 1,3,5-trichlorobenzene is supposed due to its reduced susceptibility to both degradation pathways.
BCF	Increasing BCF, correlated with increasing $\log K_{ow}$ within a $\log K_{ow}$ range of 1- 6. All trichlorobenzene isomers show very similar $\log K_{ow}$ values. Experimentally determined BCF values of substances with a $\log K_{ow}$ of about 6 and beyond repeatedly showed a levelling-off. This phenomenon, referred to as a “hydrophobicity cut-off”, has been attributed to diminished gill permeation ability with increasing molecular dimensions (size exclusion). However, a proceeding linear correlation beyond $\log K_{ow}$ of 6 (to at least 7 or 8) has also been hypothesized. The trichlorobenzene isomers are expected to show only slight differences with respect to these factors. Furthermore, the impact of molecular size and metabolism on the bioaccumulation potential of trichlorobenzenes is expected to be minor as compared with lipophilicity. Thus, the three trichlorobenzene isomers are supposed to have very similar BCF values.
Aquatic toxicity	The trichlorobenzene isomers are expected to show unpolar baseline (narcosis) toxicity due to the lack of specific molecular interaction potential. Due to their similarity with respect to lipophilicity they are expected to be similarly toxic.










Endpoints for which read-across approach is applied

Read across is used in conjunction with QSAR results to support the validity of experimental data and QSAR results for the following endpoints:

- Persistence: biodegradation (DT50) and OH radical reaction (k_{OH})
- Bioconcentration: BCF
- Ecotoxicity (read across for 1,3,5 trichlorobenzene only)
 - Long-term toxicity to fish
 - Long-term toxicity to invertebrates

SOURCE CHEMICAL(S)

The following chemicals were used for the read-across approach:

1,2,3-Trichlorobenzene 	1,3,5-Trichlorobenzene 	1,2,4-Trichlorobenzene 
CAS 87-61-6	CAS 108-70-3	CAS 120-82-1
1,2-Dichlorobenzene 	1,3-Dichlorobenzene 	1,4-Dichlorobenzene 
CAS 95-50-1	CAS 541-73-1	CAS 106-46-7
Benzene 	Monochlorobenzene 	Hexachlorobenzene 
CAS 71-43-2	CAS 108-90-7	CAS 118-74-1

PURITY /IMPURITIES

Trichlorobenzenes are produced mainly by chlorination of benzene or dichlorobenzene. The mixture of trichlorobenzenes can be separated and purified by crystallization and distillation. Purity of the respective isomers is greater than or equal 99 %. The impurities expected for the individual trichlorobenzenes are tetrachlorobenzenes, respective other trichlorobenzene isomers, dichlorobenzenes, dichlorotoluenes and bromo-chlorobenzenes. Impurities are not expected to significantly influence results.

ANALOGUE APPROACH JUSTIFICATION

Due to mechanistic differences and available experimental data, approaches and substances used for read across differed between endpoints. Approaches and justifications are described below.

Reaction with OH radicals

All trichlorobenzene isomers show the same mechanism with respect to OH radical reaction, which is OH radical addition to the aromatic ring. Chlorine substituents are expected to reduce reactivity due to the electron-withdrawing-effect. Thus, reaction rate constants are expected to be smaller for the highly chlorinated benzenes, which is confirmed by both the QSAR results and the available experimental data on other chlorobenzenes and benzene. The QSAR predictions for 1,2,3- and 1,3,5-trichlorobenzene are in accordance with this trend. The varying position of the chlorine substituents is expected to cause only minor differences with respect to reaction rates. This assumption is supported by both experimental data on dichlorobenzenes and QSAR results. However, as mentioned before, 1,3,5-trichlorobenzene is more susceptible to electrophilic attacks as compared to the other two trichlorobenzene isomers, which is also reflected by the QSAR results. A similar effect is observed for 1,3-dichlorobenzene which is more reactive towards OH radicals than 1,2- and 1,4-dichlorobenzene. Accordingly, all trichlorobenzene isomers are expected to show similar reactivity, with 1,3,5-trichlorobenzene being slightly more reactive. It is proposed to use the values predicted by EPISUITE's AOPWIN for 1,2,3- and 1,3,5-trichlorobenzene. A QSAR Model Reporting Format (QMRF) and a QSAR Prediction Reporting Format (QPRF) are given to document the reliability of the results (Annexes 2 and 3).

Biodegradation

The available data indicate that environmental degradation of all three trichlorobenzene isomers proceeds via the same biodegradation pathways. The observed biodegradation is generally slow, with 1,3,5-trichlorobenzene being slightly more persistent. Most results are available for 1,2,4-trichlorobenzene. These data were already evaluated in the EU Risk Assessment and the PBT Working Group. The substance was found to be persistent in soil and sediment.

In several studies, reductive dechlorination of trichlorobenzenes was observed. This biodegradation pathway is assumed to proceed via a nucleophilic aromatic substitution mechanism (Bosma et al., 1988; Marinucci and Bartha, 1979). Thus, biodegradation of 1,3,5-trichlorobenzene is expected to be more slowly as compared to the other two trichlorobenzene isomers, which are assumed to show similar degradation kinetics. This assumption is supported by the available data (Bosma et al., 1988; Marinucci and Bartha, 1979; Fathepure et al., 1988).

Another observed biodegradation pathway is hydroxylation through oxygenase enzymes (Ballschmiter and Scholz, 1980). Again, 1,3,5-trichlorobenzene is expected to be degraded more slowly than 1,2,4- and 1,2,3-trichlorobenzene, as the 1,3,5-substitution blocks the reaction site more effectively (Morita, 1977). 1,2,4- and 1,2,3-trichlorobenzene are expected to show very similar biodegradability.

Bioaccumulation

Only few data of limited reliability are available on the bioconcentration of trichlorobenzenes. For *Oncorhynchus mykiss*, *Poecilia reticulata* and *Cyprinus carpio*, data are available for two or three trichlorobenzene isomers. However, values for *Poecilia reticulata* differ strongly with respect to exposure time which complicates interpretation of the results. The values observed for *Oncorhynchus mykiss* and *Cyprinus carpio* are in the same range, with all three isomers showing a very similar, high bioaccumulation potential. Furthermore, the data indicate that 1,3,5-trichlorobenzene is slightly more bioaccumulative than 1,2,4- and 1,2,3-trichlorobenzene. Most results are available for 1,2,4-trichlorobenzene. These data were already evaluated in the EU Risk Assessment and the PBT Working Group. A BCF of > 2000 is derived as a realistic worst case for this substance.

Trichlorobenzenes are unpolar, lipophilic substances. Bioconcentration of these substances is expected to be mainly determined by their lipophilicity, which is reflected by the $\log K_{ow}$. Within a $\log K_{ow}$ range of 1- 6, bioconcentration of unpolar compounds is expected to show a linear increase with increasing $\log K_{ow}$. All trichlorobenzene isomers show very similar $\log K_{ow}$ values and thus, based on lipophilicity, their bioconcentration factors are expected to be in the same range. All trichlorobenzene isomers have a $\log K_{ow}$ about 4 and are thus potentially bioaccumulating based on lipophilicity.

Further factors like molecular size or metabolism are proposed to influence bioconcentration, though they are expected to be of minor importance as compared to lipophilicity. A large molecular size is proposed to decrease BCF due to reduced bioavailability. However, the molecular weight is clearly smaller than the cut-off values recommended in the technical guidance (MW>700 indicating not vB, MW>1100 indicating not B). Furthermore, the trichlorobenzene isomers are expected to show only marginal differences with respect to molecular size.

The metabolism of a substance may reduce or limit its bioaccumulation potential. A possible metabolism pathway for trichlorobenzenes is hydroxylation by oxygenase enzymes, following a similar pattern as the abovementioned biodegradation pathway. Analogously, 1,3,5-trichlorobenzene is probably metabolized slightly slower than 1,2,3-trichlorobenzene and 1,2,4-trichlorobenzene.

In summary it can be stated that with respect to the factors relevant for bioconcentration, there are no significant differences between 1,2,3-trichlorobenzene and 1,2,4-trichlorobenzene. For 1,3,5-trichlorobenzene, slight deviations with respect to metabolism are assumed. The impact of this factor on the BCF is expected to be minor or negligible as compared to lipophilicity. Thus, from a mechanistic point of view, the three trichlorobenzene isomers are expected to show very similar BCF values.

Thus, a BCF of >2000 is assumed for 1,2,4-trichlorobenzene and is also expected for 1,2,3- and 1,3,5-trichlorobenzene, with 1,3,5-trichlorobenzene potentially showing a slightly higher BCF.

Ecotoxicity

Toxic Mode of Action. Based on the chemical structure, all three trichlorobenzene isomers are expected to exhibit baseline (nonpolar narcosis) toxicity. This hypothesis is supported by various expert systems (so called profiler) implemented in the OECD QSAR Application Toolbox (beta Version 2.0, Mai 2010).

Toxicity of nonpolar narcotics is largely determined by lipophilicity and thus by the $\log K_{ow}$. As all trichlorobenzenes are isomers they show very similar $\log K_{ow}$ values and thus, based on lipophilicity, their ecotoxicity is expected to be very similar. Reactivity might influence ecotoxicity of these isomers. While the OECD QSAR Application toolbox gives some hints, that reactivity might be higher for 1,2,4 trichlorobenzene (expected mode of protein binding nucleophilic substitution) available data do not support this hypothesis (see data below).

Fish. Comparable experimental data on acute toxicity of 1,2,3 and 1,2,4 trichlorobenzene for *Danio rerio* and *Oncorhynchus mykiss*. show that toxicity of both compounds is in the same concentration range. Available data for 1,3,5, trichlorobenzene with *O.mykiss* are in the same range of toxicity. Small differences in toxicity (max. factor 8 between 1,2,3 and 1,3,5 trichlorobenzene) are expected to be due to interlaboratory variance (factor 3 between) resulting in 1,2,3 trichlorobenzene being slightly more toxic than 1,2,4 trichlorobenzene (factor 3 and between) , however data might indicate, that 1,3,5 trichlorobenzene is slightly less toxic than expected due to the hypothesis of similar toxicity (factor 3).

With respect to long-term toxicity, data for 1,2,4 and 1,2,3 trichlorobenzene again support hypothesis of comparable toxicity. No data for 1,3,5 trichlorobenzene are available, thus read-across needs to be applied.

Daphnia. Data on acute toxicity to *Daphnia magna* for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene support similarity of toxicity of both chemicals.

With respect to long-term effects, available data for an exposure time of 14 days also indicate similar toxicity for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene. However, considering data for an exposure time of 21 days, 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene differ to some extent (factor 13).

Algae. Comparable data on *Pseudokirchneriella subcapitata* are available for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene. For an exposure time of 96 h, both isomers show very similar NOEC values with 1,2,3-trichlorobenzene being slightly more toxic. For 1,3,5-trichlorobenzene, an EC₁₀ is given from a 48 h test, from which a maximum NOEC is derived. Comparing this result with a NOEC derived from a 48 h test on 1,2,3-trichlorobenzene, both 1,3,5-trichlorobenzene and 1,2,3-trichlorobenzene are in the same range of toxicity.

Summary on Ecotoxicity Available data for 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene show that toxicity of both isomers is very similar, with the most sensitive NOEC values being 0.04 mg/l and 0.03 mg/l. Results support the hypothesis of baseline toxicity for all three isomers and all organisms groups tested. Neither one of the isomers nor one of the organisms groups showed unexpected high toxicity.

For 1,3,5-trichlorobenzene, only few data of limited reliability are available. However, experimental results for 1,2,3 and 1,2,4 trichlorobenzene and similarity of the expected toxic mode of action for 1,3,5 Trichlorobenzene provide confidence, that a read-across for 1,3,5 trichlorobenzene based on data for 1,2,3 and 1,2,4 trichlorobenzene is reasonable. This is supported by experimental data for 1,3,5 trichlorobenzene for acute fish toxicity and algae toxicity showing toxicity comparable to those of 1,2,3 and 1,2,4 trichlorobenzene .

DATA MATRIX

	Target 1	Target 2	Source 1	Source 2	Source 3	Source 4	Source 5	Source 6	Source 7
Chemical Name	1,2,3-Tri-chlorobenzene	1,3,5-Tri-chlorobenzene	1,2,4-Tri-chlorobenzene	1,2-Dichloro-benzene	1,4-Dichloro-benzene	1,3-Dichloro-benzene	Benzene	Monochloro-benzene	Hexachloro-benzene
CAS RN	000087-61-6	000108-70-3	000120-82-1	000095-50-1	000106-46-7	000541-73-1	000071-43-2	000108-90-7	000118-74-1
Physico-chemical data									
Molecular Weight	181,45	181,45	181,45	147.01	147.01	147.01	78.11	112.56	284.78
Partitioning Coefficient logKow	4.05	4.19	3.9-4.3 ¹ 4.7 ^{RAR}						
Fate									
k_{OH} -Exp. Value in $10^{-12} \text{ cm}^3 \text{ s}^{-1} \text{ molecules}^{-1}$			0.532 ^{RAR} 0.5 ^{RAR} 0.6 ^{RAR} 0.55 ^{PysProp}	0.42 ^{PhysProp}	0.32 ^{RAR,PhysProp} 0.48 ^{RAR}	0.72 ^{PhysProp}	1.2 ^{RAR} 1.23 ^{PhysProp}	0.77 ^{PhysProp}	0.027 ^{PhysProp}
-Calc. value in $10^{-12} \text{ cm}^3 \text{ s}^{-1} \text{ molecules}^{-1}$, AOPWIN	0.282	0.679	0.282	0.4	0.4	0.965-	1.95	1.372	0.0169
ready biodegradability screening test	not readily biodegradable (MITI)	not readily biodegradable (MITI)	not readily biodegradable (MITI)						
DT50									
-Exp. Value soil	DisT50 = 222 d (259 d study, no significant)	DisT50 = 238 d (259 d study, no significant)	DisT50 = 194 d (259 d study, no significant)						

¹ The IUCLID file contains several experimental values, most of which are in the abovementioned range from 3.9 to 4.4. The overall range of values is 2.33 to 4.8.

RAR Taken from the respective EU Risk Assessment Report, recommended values, in some cases further data available

PysProp Taken from the PhysProp Database (PhysProp, 2001, US EPA, 2008)

	Target 1	Target 2	Source 1	Source 2	Source 3	Source 4	Source 5	Source 6	Source 7
Chemical Name	1,2,3-Tri-chlorobenzene	1,3,5-Tri-chlorobenzene	1,2,4-Tri-chlorobenzene	1,2-Dichloro-benzene	1,4-Dichloro-benzene	1,3-Dichloro-benzene	Benzene	Monochloro-benzene	Hexachloro-benzene
	bio-degradation)	bio-degradation)	bio-degradation)						
-Exp. Value sediment	DT50 = 202 d DT50 = 212 d (two sediment samples, ^{RAR})								
BCF									
-Exp. value <i>Oncorhynchus mykiss</i> (<i>Salmo gairdneri</i>)	1200 (119 d) 2600 (105 d)	1800 (119 d) 4100 (105 d)	1300 (119 d) 3200 (105 d)						
-Exp. value <i>Poecilia reticulata</i>	1860 (4 d) 702 ²	756 ²	1139 (17 d)						
-Exp. value (<i>Brachy)danio rerio</i>			574-1412 (28 d)						
-Exp. value <i>Jordanella floridae</i>			2026 (28 d)						
-Exp. value <i>Cyprinus carpio</i>	130-1200 (42 d)	150-1700 (42 d)	120-1320 (42 d) 805-830 ³						
<u>Environmental Toxicity</u>									
Mode of action									
-Verhaare scheme	Class 1 narcosis	Class 1 narcosis	Class 1 narcosis						
-ECOSAR Classification	Neutral organics	Neutral organics	Neutral organics						
-OASIS acute toxicity MOA	Basesurface narcotics	Basesurface narcotics	Basesurface narcotics						

² test duration not given, original literature not reviewed³ test duration not given

	Target 1	Target 2	Source 1	Source 2	Source 3	Source 4	Source 5	Source 6	Source 7
Chemical Name	1,2,3-Tri-chlorobenzene	1,3,5-Tri-chlorobenzene	1,2,4-Tri-chlorobenzene	1,2-Dichloro-benzene	1,4-Dichloro-benzene	1,3-Dichloro-benzene	Benzene	Monochloro-benzene	Hexachloro-benzene
-OASIS Protein binding	No binding	No binding	Nucleophilic substitution on aromatic halogens						
Acute Toxicity to Fish, LC ₅₀ , [mg/L]									
-Exp. value <i>Poecilia reticulata</i>	0.35 (96 h)								
-Exp. value <i>Danio rerio</i>	3.1 (24/48 h)		6.3 (24/48 h)						
-Exp. value <i>Oncorhynchus mykiss (Salmo gairdneri)</i>	0.71 (24/48 h)	5.48 (96 h)	1.95 (24/48 h)						
			1.32 (96h)						
-Exp. value <i>Leuciscus idus</i>		7.05 (96h) ⁴	0.7 (48 h)						
-Exp. value <i>Solea solea</i>		1.96 (96 h)							
Long-term Toxicity to Fish NOEC, [mg/L]									
-Exp. value, <i>Danio rerio</i>	0.077 (28 d)		0.04 (21 d)						
	0.25 (28 d)								
-Exp. value, <i>Poecilia reticulata</i>		3.3 (14 d, EC ₅₀)	0.11 (14 d)						
- Exp. Value <i>Gambusia affinis</i>	0.022 (42d, EC ₁₀)								
Acute Toxicity to Aquatic									

⁴ Test not validated because data not available

	Target 1	Target 2	Source 1	Source 2	Source 3	Source 4	Source 5	Source 6	Source 7
Chemical Name	1,2,3-Tri-chlorobenzene	1,3,5-Tri-chlorobenzene	1,2,4-Tri-chlorobenzene	1,2-Dichloro-benzene	1,4-Dichloro-benzene	1,3-Dichloro-benzene	Benzene	Monochloro-benzene	Hexachloro-benzene
Invertebrates, L(E)C₅₀, [mg/L]									
Exp. value <i>Daphnia magna</i>	1.66 (24-48h, arithmetic mean of 5 tests)		2.1 (48 h) geom. mean used in RAR						
Exp. value <i>Mysidopsis bahia</i>			0.45 (96 h)						
Long-term Toxicity to Aquatic Invertebrates. NOEC, [mg/L]									
Exp. value <i>Daphnia magna</i>	0.08 (14 d, NOEC=EC ₁₆) 0.03 (21 d)		0.06 (14 d, NOEC=EC ₁₆) 0.4 (21 d)						
Exp. value <i>Mysidopsis bahia</i>			≤ 0.064						
Toxicity to aquatic Plants. NOEC, [mg/l]									
Exp. value <i>Pseudokirchneriella subcapitata</i> (<i>Selenastrum capricornutum</i>)	0.22 (96 h) 0.2 (48 h)	< 0.73 (48 h, EC ₁₀ = 0.398)	0.37 (96 h)						
Toxicity to sediment organisms, [mg/l]									
Exp. value <i>Chironimus riparius</i>	1.7 (48 h, LC ₅₀) 0.018 (96 h, NOEC)								

Unless otherwise mentioned, all data are taken from the Annex V-Dossiers for 1,2,3-, 1,2,4- and 1,3,5-trichlorobenzenes. Choice of endpoints used for read-across: For the trichlorobenzenes, the key studies /most sensitive tests were reported respectively. Further tests were chosen if results are reported for more than one trichlorobenzene isomer and allow comparison between the substances. The other chlorinated benzenes and benzene were only used to support the estimation of $\log k_{OH}$ and thus, only calculated and experimental $\log k_{OH}$ values are considered for these substances.

CONCLUSIONS PER ENDPOINT

Reaction with OH radicals

All trichlorobenzene isomers are expected to show similar reactivity, with 1,3,5-trichlorobenzene being slightly more reactive. The values predicted by AOPWIN for 1,2,3- and 1,3,5-trichlorobenzene can be regarded as valid within the variation of the model for aromatic compounds. Results are additionally supported by experimental data from structural analogues. A QSAR Model Reporting Format (QMRF) and a QSAR Prediction Reporting Format (QPRF) are given to document the reliability of the results.

Biodegradation

The observed biodegradation is generally slow for all three trichlorobenzene isomers, with 1,3,5-trichlorobenzene being slightly more persistent. This assumption is supported by the evaluation of possible biodegradation pathways.

Most results are available for 1,2,4-trichlorobenzene. These data were already evaluated in the EU Risk Assessment and the PBT Working Group. The substance was found to be persistent in soil and sediment. As data are not standardized, default half-lives for non biodegradable substances are applied in the assessment. 1,2,3-trichlorobenzene is assumed to be as persistent as 1,2,4-trichlorobenzene, whereas 1,3,5-trichlorobenzene is probably even more persistent. Thus, default half-lives for non biodegradable substances can also be applied for 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene.

Bioaccumulation

From a mechanistic point of view, the three trichlorobenzene isomers are expected to show high BCF values, with differences between the isomers being marginal. The vast majority of deviations in determined BCF may be attributed to the inherent variability of experimental studies (different fish species etc.) rather than different physico-chemical properties of these constitutional isomers. For each isomer the observed BCF values are of limited reliability. Nonetheless, the results indicate that all three isomers show a similar, high bioaccumulation potential. Furthermore, the data indicate that 1,3,5-trichlorobenzene is slightly more bioaccumulative than 1,2,4- and 1,2,3-trichlorobenzene. Most results are available for 1,2,4-trichlorobenzene. These data were already evaluated in the EU Risk Assessment and the PBT Working Group and a BCF of 2000 was used as a realistic worst case for this substance. In analogy to this approach, a BCF of 2000 is also expected for 1,2,3- and 1,3,5-trichlorobenzene, with 1,3,5-trichlorobenzene potentially showing a slightly higher BCF. These estimates are considered as reliable within variation of the data.

Aquatic Toxicity

Long-term toxicity to fish

Taking into account data from all structural isomers, one of the most sensitive species is expected to be *Danio rerio*, for which long-term data on 1,2,4-trichlorobenzene and 1,2,3-trichlorobenzene are available. Based on structural considerations and the available data, the corresponding NOEC for 1,3,5-trichlorobenzene is expected to be in the same range, but potentially slightly higher. Thus, for the assessment, a NOEC of 0.077 mg/l is estimated.

Long-term toxicity to invertebrates

According to data available for 1,2,3 and 1,2,4 trichlorobenzene long-term toxicity of 1,3,5 trichlorobenzene to *daphnia magna* is expected to be in the range of 0.03 to 0.06 mg/L.

Based on data for 1,2,4 trichlorobenzene *Mysidopsis bahia* is expected to be more sensitive or in the same range as *Daphnia magna*.

As it is expected according to structural considerations and the available data that 1,3,5-TCB is potentially slightly less toxic, a NOEC of 0.06 mg/L (read across from 1,2,4-TCB) is used for 1,3,5-trichlorobenzene.

REFERENCES

Bosma TNP, van der Meer JR, Schraa G, Tros ME, Zehnder AJB. 1988. Reductive dechlorination of all trichloro- and dichlorobenzene isomers. *FEMS Microbiol Lett* 53(3-4):223-229.

Marinucci AC, Bartha R. 1979 Nov. Biodegradation of 1,2,3- and 1,2,4-trichlorobenzene in soil and in liquid enrichment culture. *Appl Environ Microbiol* 38(5):811-817.

Fathepure BZ, Tiedje JM, Boyd SA. 1988. Reductive dechlorination of hexachlorobenzene to tri- and dichlorobenzenes in anaerobic sewage sludge. *Appl Environ Microbiol* 54(2):327-330.

Ballschmiter K, Scholz C. 1980. Mikrobieller Abbau von chlorierten Aromaten: VI. Bildung von Dichlorphenolen und Dichlorbrenzkatechinen aus Dichlorbenzolen in mikromolarer Lösung durch *Pseudomonas* Sp. *Chemosphere* 9(7-8):457-467.

Morita M. 1977. Chlorinated benzenes in the environment. *Ecotoxicology and Environmental Safety* 1(1):1-6.



Beauman, J. A.; Howard, P. H. 2001. Physprop database. Syracuse Research Corporation: Syracuse (NY, USA).

US EPA. 2008. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.00. United States Environmental Protection Agency, Washington, DC, USA .
<http://www.epa.gov/oppt/exposure/pubs/episuitedl.htm>

ANNEX 2

QSAR Prediction Reporting Format (QPRF) (version 1.1, May 2008)

1. Substances**1.1 CAS number:** 87-61-6 and 108-70-3**1.2 EC number:****1.3 Chemical name:** 1,2,3-Trichlorobenzene and 1,3,5-Trichlorobenzene**1.4 Structural formula:**

1,2,3-Trichlorobenzene	1,3,5-Trichlorobenzene
	
CAS 87-61-6	CAS 108-70-3

1.5 Structure codes:**a. SMILES:**

1,2,3-Trichlorobenzene	1,3,5-Trichlorobenzene
<chem>c(c(c(cc1Cl)Cl)Cl)(c1)Cl</chem>	<chem>c(cc(cc1Cl)Cl)(c1)Cl</chem>

*(used for the model prediction)***b. InChI:** *not available/ not used for the model prediction***c. Other structural representation:** *not available/ not used for the model prediction***d. Stereochemical features:** *the substance is no stereo-isomer***2. General information****2.1 Date of QPRF:** 7 May 2010**2.2 QPRF author and contact details:**

Anna Bönhardt
 Section IV 2.3 Chemicals
 Federal Environment Agency/ Umweltbundesamt
 Wörlitzer Platz 1
 06844 Dessau-Roßlau
 Germany
 Phone: +49-340-2103-3029
 Email: Anna.Boehnhardt@uba.de

3. Prediction

3.1 Endpoint (OECD Principle 1)

- a. **Endpoint:** OH-radical reaction rate constant (k_{OH}) of organic compounds
- b. **Dependent variable:** $\log k_{OH}$ (AOPWIN equation for addition to aromatic rings)

3.2 Algorithm (OECD Principle 2)

- a. **Model or submodel name:** AOPWIN⁵ for k_{OH}
(references see corresponding QMRF)
- b. **Model version:** AOPWIN v1.92
- c. **Reference to QMRF:** The corresponding QMRF named ‘AOPWIN for k_{OH} ’ has been newly compiled.
- d. **Predicted value (model result):**

1,2,3-Trichlorobenzene	1,3,5-Trichlorobenzene
$0.282 \cdot 10^{-12}$	$0.679 \cdot 10^{-12}$

- e. **Predicted value (comments):** k_{OH} given in $\text{cm}^3 \text{s}^{-1} \text{molecules}^{-1}$
- f. **Input for prediction:** SMILES
- g. **Descriptor values:** electrophilic substituent constants σ^+ for chlorine, referring to attack at the meta (σ_m^+) position and at the ortho or para position (σ_p^+)

σ_m^+	σ_p^+
0.399	0.114

1,2,3-Trichlorobenzene	1,3,5-Trichlorobenzene
$\sum \sigma^+ = 2 \cdot \sigma_p^+ + 1 \cdot \sigma_m^+$	$\sum \sigma^+ = 3 \cdot \sigma_p^+$

AOPWIN equation for addition to aromatic rings:

$$\log k_{OH} = -11.71 - 1.34 \cdot \sum \sigma^+$$

3.3 Applicability domain (OECD principle 3)

a. **Domains:** 1,2,3-Trichlorobenzene and 1,3,5-Trichlorobenzene are considered to be in the applicability domain as described in the corresponding QMRF and the original Literature (Atkinson 1991).

- i. *descriptor domain.* The molecular weight (MW) of both isomers is 181.45. This is in the range of the training set that was used to derive for the model equation for monocyclic aromatic compounds and biphenyls (Atkinson 1991).

Training set: $78.12 \leq \text{MW} \leq 204.01$, Average MW: 125.68

- ii. *structural fragment domain.* The fragments of the chemical structure are three chlorine atoms, that are substituted to different positions of the benzene ring. These fragments were also included in the training set, which contains

⁵ US EPA. 2008. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.00. United States Environmental Protection Agency, Washington, DC, USA . <http://www.epa.gov/oppt/exposure/pubs/episuitd.htm>

1,2,4-trichlorobenzene, 1,2-dichlorobenzene, 1,4-dichlorobenzene, 1,3-dichlorobenzene and chlorobenzenes. Furthermore, their occurrence per molecule is in the range of the training set.

iii. *mechanism domain.* The mechanistic domain of the model can be defined clearly by the different reaction mechanisms that are accounted for by respective model equations: H-atom abstraction from C-H and O-H bonds; OH radical addition to double and triple bonds; OH radical addition to aromatic rings; and OH radical interaction with N-,S-, and P- atoms. Accordingly, Therefore, 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene can be considered to be in the mechanistic domain of the model.

iv. *metabolic domain.* Not relevant.

b. Structural analogues: Experimental data on structural analogues are given in tables 1 and 2.

Table 1: Calculated and experimental k_{OH} for structural analogues (experimental data taken from the PhysProp Database⁶, k_{OH} given in $10^{-12} \cdot \text{cm}^3 \text{s}^{-1} \text{molecules}^{-1}$).

CAS No.	Chemical Name	SMILES	Exp k_{OH}	Est k_{OH}	Training Set
000120-82-1	1,2,4-Trichlorobenzene	c(ccc(c1Cl)Cl)(c1)Cl	0.550	0.282	yes
000095-50-1	1,2-Dichlorobenzene	c(c(ccc1)Cl)(c1)Cl	0.420	0.400	yes
000106-46-7	1,4-Dichlorobenzene	c(ccc(c1)Cl)(c1)Cl	0.320	0.400	yes
000541-73-1	1,3-Dichlorobenzene	c(cccc1Cl)(c1)Cl	0.720	0.965	yes
000071-43-2	Benzene	c(cccc1)c1	1.230	1.950	yes
000108-90-7	Chlorobenzene	c(cccc1)(c1)Cl	0.770	1.372	yes
000118-74-1	Hexachlorobenzene	c(c(c(c(c1Cl)Cl)Cl)Cl)(c1Cl)Cl	0.027	0.017	no

Table 2: Calculated and experimental $\log k_{OH}$ for structural analogues.


CAS No.	Chemical Name	SMILES	Exp $\log k_{OH}$	Est $\log k_{OH}$	Training Set
000120-82-1	1,2,4-Trichlorobenzene	c(ccc(c1Cl)Cl)(c1)Cl	-12.26	-12.55	yes
000095-50-1	1,2-Dichlorobenzene	c(c(ccc1)Cl)(c1)Cl	-12.38	-12.40	yes
000106-46-7	1,4-Dichlorobenzene	c(ccc(c1)Cl)(c1)Cl	-12.49	-12.40	yes
000541-73-1	1,3-Dichlorobenzene	c(cccc1Cl)(c1)Cl	-12.14	-12.02	yes
000071-43-2	Benzene	c(cccc1)c1	-11.91	-11.71	yes
000108-90-7	Chlorobenzene	c(cccc1)(c1)Cl	-12.11	-11.86	yes
000118-74-1	Hexachlorobenzene	c(c(c(c(c1Cl)Cl)Cl)Cl)(c1Cl)Cl	-13.57	-13.77	no

⁶ Beauman, J. A.; Howard, P. H. Physprop database. Syracuse Research Corporation: Syracuse (NY, USA), 2001.


US EPA. 2008. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.00. United States Environmental Protection Agency, Washington, DC, USA . <http://www.epa.gov/oppt/exposure/pubs/episuitd.htm>

c. Considerations on structural analogues: Figure 1 shows calculated vs. experimental k_{OH} values for structural analogues. The correlation between calculated and experimental values is good ($r^2 \approx 0.89$). The largest deviation is observed for hexachlorobenzene, which was not used in the training set. Hexachlorobenzene is outside the molecular weight range of the training set and has more chlorine substituents per molecule than the substances in the training set. However both deviations from the training set range are not considered to affect model performance significantly. Note further that the prediction error for this substance is still small. Further, due to the rather small errors, the variation of experimental data also contributes to the observed deviations. As k_{OH} values of the training set cover several orders of magnitude, the model predicts k_{OH} values in logarithmic units and commonly, QSARs for predicting a wide range of k_{OH} values are evaluated in logarithmic units.

The predicted order of k_{OH} values is:



Analysis of experimental data yields:



Thus, the overall trends for calculated and experimental k_{OH} values are in good agreement. The only deviation is 1,2,4-trichlorobenzene, that shows a slightly higher experimental k_{OH} than expected from an analysis of the other experimental data. However, differences in reactivity are determined solely by the number of chlorine substituents and the substitution pattern, and thus, this deviation from the expected trend is assumed to result from variation of the experimental data.

Figure 1: Calculated vs. experimental k_{OH} for structural analogues. The dashed line indicates conformity between calculated and experimental results.




Figure 2 shows the resulting plot of calculated vs. experimental $\log k_{OH}$ values, giving a very good correlation between calculated and experimental values ($r^2 \approx 0.94$). With respect to $\log k_{OH}$, all calculation errors are within a range of 0.3 and the mean absolute

error is 0.15. This is in accordance with the very good performance for aromatics that is documented in the QSAR Model Reporting Format. As can be seen from figure 3, AOPWIN tends to overestimate k_{OH} slightly for lowly chlorinated structural analogues, whereas the highly chlorinated hexachlorobenzene is slightly underestimated. Given the small calculation errors and the fact that trichlorobenzenes can be considered to have a medium number of chlorine substituents, this finding is not considered to impact reliability for 1,2,3- and 1,3,5-trichlorobenzenes.

Figure 2: Calculated vs. experimental $\log k_{OH}$ for structural analogues. The dashed line indicates conformity between calculated and experimental results.





Figure 3: Calculation error vs. experimental $\log k_{OH}$ for structural analogues



3.4 The uncertainty of the prediction (OECD principle 4)


Given the calculation errors and the variability observed for the structural analogues, it is expected that the QSAR prediction is correct within a range of 0.3 (see 3.3c).

3.5 The chemical and biological mechanisms according to the model underpinning the predicted result (OECD principle 5).

The reaction of chlorobenzenes with OH radicals proceeds via addition of the OH radical to the aromatic ring. Due to their electron-withdrawing effect, chlorine substituents are generally expected to reduce reactivity towards OH radicals, resulting in decreasing $\log k_{\text{OH}}$ values with increasing chlorine content. Furthermore, the position of the substituents has an impact on susceptibility to electrophilic attacks. For example, 1,3-dichlorobenzene and 1,3,5-trichlorobenzene are expected to be more reactive than their respective isomers. This may lead to slight deviations from the aforementioned trend. Taking into account the experimental data from structural analogues, the predicted $\log k_{\text{OH}}$ values for 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene are in accordance with this trend.

This is illustrated in figure 4, which shows the experimental $\log k_{\text{OH}}$ values of the structural analogues and the predicted $\log k_{\text{OH}}$ values for 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene, plotted vs. the number of chlorine atoms.

Figure 4: Experimental $\log k_{\text{OH}}$ for structural analogues plotted vs. number of chlorine atoms. The red dots represent predicted $\log k_{\text{OH}}$ values for 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene.



4. Adequacy (Optional)

- 4.1 Regulatory purpose:** REACH Annex XV Report. Proposal for Identification of a substance as a PBT.
- 4.2 Approach for regulatory interpretation of the model result:** The model result is a key factor determining atmospheric residence time and thus an important input when modelling and assessing the atmospheric long-range transport potential of the substances.

- 4.3 Outcome:** The k_{OH} values of 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene are $0.282 \cdot 10^{-12}$ and $0.679 \cdot 10^{-12} \text{ cm}^3\text{s}^{-1}\text{molecules}^{-1}$. They are considered to be correct within a range of 0.3 logarithmic units.
- 4.4 Conclusion:** The k_{OH} values of 1,2,3-trichlorobenzene and 1,3,5-trichlorobenzene are $0.282 \cdot 10^{-12}$ and $0.679 \cdot 10^{-12} \text{ cm}^3\text{s}^{-1}\text{molecules}^{-1}$. These k_{OH} values can be used when assessing the atmospheric long-range transport potential of the substances. The resulting atmospheric half-lives depend on the underlying assumptions on the OH radical concentration.

ANNEX 3

QSAR Model Reporting Format (QMRF) for AOPWIN

Text version; original QMRF was generated in xml-Format, using the QMRF Editor v1.2, which is available free of charge <http://ecb.jrc.ec.europa.eu/qsar/>

1.QSAR identifier

1.1.QSAR identifier (title):

AOPWIN for k_{OH}

1.2.Other related models:

AOPWIN method for estimating k_{O_3} (the rate constant for the gas-phase reaction between ozone and olefinic/acetylenic compounds)

1.3.Software coding the model:

EPISUITE Freely available software from the US EPA and the Syracuse Research Corporation

<http://www.epa.gov/oppt/exposure/pubs/contact.htm>

<http://www.epa.gov/oppt/exposure/pubs/episuitedl.htm>

2.General information

2.1.Date of QMRF:

23.04.2010

2.2.QMRF author(s) and contact details:

Anna Böhnhardt

German Federal Environment Agency (UBA)

Wörlitzer Platz 1

06844 Dessau, Germany,

Phone: + 4934021033029

Anna.Boehnhardt@uba.de

2.3.Date of QMRF update(s):

2.4.QMRF update(s):

2.5.Model developer(s) and contact details:

Roger Atkinson

Air Pollution Research Center

Department of Environmental Sciences

University of California

Riverside, CA 92521

<http://www.envisci.ucr.edu/faculty/atkinson.html>

2.6.Date of model development and/or publication:

Original model published in the years 1986 and 1987, update published in 1995. Software is regularly updated.

2.7.Reference(s) to main scientific papers and/or software package:

- [1] Atkinson R 1986. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. Chemical Reviews 86 (1): 69-201. <http://dx.doi.org/10.1021/cr00071a004>
- [2] Atkinson R 1987. A structure-activity relationship for the estimation of rate constants for the gas-phase reactions of OH radicals with organic compounds. International Journal of Chemical Kinetics 19 (9): 799-828. <http://dx.doi.org/10.1002/kin.550190903>
- [3] Kwok ESC and Atkinson R 1995. Estimation of hydroxyl radical reaction rate constants for gas-phase organic compounds using a structure-reactivity relationship: An update. Atmospheric Environment 29 (14): 1685-1695. [http://dx.doi.org/10.1016/1352-2310\(95\)00069-B](http://dx.doi.org/10.1016/1352-2310(95)00069-B)
- [4] US EPA 2008. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.00. United States Environmental Protection Agency, Washington, DC, USA. <http://www.epa.gov/oppt/exposure/pubs/episutedl.htm>
- [5] US EPA. 2000. On-Line AOPWIN™ User's Guide (accessed on April 22nd, 2010). In: US EPA. 2008. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.00. United States Environmental Protection Agency, Washington, DC, USA. <http://www.epa.gov/oppt/exposure/pubs/episutedl.htm>
- [6] Atkinson R 1991. Atmospheric lifetimes of dibenzo-p-dioxins and dibenzofurans. The Science of the Total Environment 104: 17-33. [http://dx.doi.org/10.1016/0048-9697\(91\)90005-Y](http://dx.doi.org/10.1016/0048-9697(91)90005-Y)

2.8.Availability of information about the model:

The model equations and increments are documented in the original papers (Atkinson1986, Atkinson1987, Kwok&Atkinson1995) and in the On-Line AOPWIN User's Guide that is included in the EPISUITE software package. The software and the user's guide are available free of charge (<http://www.epa.gov/oppt/exposure/pubs/episutedl.htm>). Most information given in this QMRF is taken from the original papers and the On-Line AOPWIN User's Guide. According to the AOPWIN User's Guide, the complete training set is not available. However, the training set may be partly reconstructed by a careful analysis of the original papers and the literature cited therein. For example, the substances used to derive the regression equation for aromatic reactivity are given in a study on atmospheric lifetimes of dibenzo-p-dioxins and dibenzofurans (Atkinson 1991).

2.9.Availability of another QMRF for exactly the same model:

no other QMRF known

3.Defining the endpoint - OECD Principle 13.1.Species:3.2.Endpoint:

- 2.Environmental fate parameters
 2.Persistence: Abiotic degradation in air (Phototransformation)
 2.2.b.Indirect photolysis (OH-radical reaction rate constant k_{OH})

3.3.Comment on endpoint:

k_{OH} values cover several orders of magnitude and are thus evaluated as $\log k_{OH}$ in the section on validation

3.4.Endpoint units:

k_{OH} in $\text{cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$

3.5.Dependent variable:

$\log k_{OH}$ for aromatic reactivity; k_{OH} or $\log k_{OH}$ for other reaction mechanisms (probably k_{OH})

3.6.Experimental protocol:

Reaction rate constants can be determined directly by absolute rate techniques (pulsed photolysis, flow discharge techniques) or by relative rate methods.

3.7.Endpoint data quality and variability:

Data mainly taken from the PhysProp Database (SRC PhysProp Database.Syracuse Research Corporation) and the monographs of Atkinson (Atkinson 1989, Atkinson 1994). These sources contain collected experimental data from the scientific literature and have been carefully evaluated by atmospheric chemists.

4.Defining the algorithm - OECD Principle 24.1.Type of model:

QSAR estimating the OH-radical reaction rate (k_{OH}) of organic compounds

4.2.Explicit algorithm:

Determination of the overall reaction rate constant k_{OH}

Equation 1: Determination of the overall reaction rate constant (k_{OH}) by summation over the partial rate constants of all reaction sites r ($k_{\text{mech},r}$), accounting for the respective reaction mechanisms.

$$k_{OH} = \sum k_{\text{mech}} = \sum \sum k_{\text{mech},r} \quad (1)$$

Estimation of local reactivity I

Equation 2: Estimation of local reactivity for hydrogen abstraction & addition to double and triple bonds ($k_{\text{mech},r}$) through group rate constants ($k_{\text{mech,group}}$) that reflect the basic reactivity towards OH. The influence of substituents (X) is accounted for by substituent factors ($F(X)$) that are specific for the respective mechanism. Both group rate constants and substituent values are incremental values and have been published in the original literature.

$$k_{\text{mech},r} = k_{\text{mech,group}} \cdot \prod_X F(X) \quad (2)$$

Estimation of local reactivity II

Equation 3: Estimation of local reactivity for addition to aromatic rings ($k_{\text{aro},r}$) from sum of electrophilic substituent constants $\Sigma\sigma^+$

$$\log k_{\text{aro},r} = -11.71 - 1.34 \sum \sigma^+ \quad (3)$$

4.3.Descriptors in the model:

- group rate constant ($k_{\text{mech,group}}$)
increment; values given in the original literature and in the user's guide

- substituent factor (F(X))
increment; values given in the original literature and in the user's guide
- electrophilic substituent constants σ^+
increment; values given in user's guide

4.4.Descriptor selection:

Electrophilic substituent constants describe aromatic reactivity (Brown and Okamoto 1958) and were thus selected as descriptors for aromatic rings.

4.5.Algorithm and descriptor generation:

Values for group rate constants and substituent factors were derived partly from individual molecules and partly from a nonlinear least-squares fit. The model equation for aromatic reactivity was derived from a least-squares analysis on a set of monocyclic aromatics and biphenyls.

4.6.Software name and version for descriptor generation:

4.7.Chemicals/Descriptors ratio:

5.Defining the applicability domain - OECD Principle 3

5.1.Description of the applicability domain of the model:

The compound to be predicted should have similar structural features as the compounds from the training set (chemical domain). The chemical domain can be estimated from the group rate constants and substituent factors applied in the model. If it is not possible to describe all features of the chemical structure by group rate constants and/or substituent factors, the compound is outside the applicability domain. However, this approach only allows for a rather coarse estimation of the chemical domain. For a more precise description, knowledge of the training set is required. Furthermore, the compound to be predicted should be in the so-called mechanistic domain. The latter can be defined clearly by the different reaction mechanisms that are accounted for by respective model equations: H-atom abstraction from C-H and O-H bonds; OH radical addition to double and triple bonds; OH radical addition to aromatic rings; and OH radical interaction with N-, S-, and P- atoms. Finally, the compound to be predicted should be shown to be in the same parameter range (parameter domain).

As the training set is not available, a precise description of the chemical and the parameter domain is not possible. However, subsets of the training set can be reconstructed by a careful analysis of the original papers and the literature cited therein. For example, the substances used to derive the regression equation for aromatic reactivity are given in a study on atmospheric lifetimes of dibenzodioxins and dibenzofurans (Atkinson 1991). This way, a "subdomain" may be derived from a set of substances that are verified to belong to the training set. For example, the regression equation for aromatic reactivity was derived from a set of monocyclic aromatic compounds and biphenyls. Thus, for applying this equation, the compound to be predicted should be chemically similar to the respective training set of monocyclic aromatic compounds and biphenyls. The reaction rate constants of these compounds are available in the PhysProp Database and range from $1.4 \cdot 10^{-13} \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ (Nitrobenzene) to $1.13 \cdot 10^{-10} \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ (3,5-Dimethylphenol). According to the user's guide, further substituent constants and increments, e.g. for condensed aromatics have been derived from experimental data to extend the applicability of the model.

5.2.Method used to assess the applicability domain:

As the training set is not available, the applicability domain has to be assessed by the user in a case-to-case decision, e.g. by studying the original literature to find out whether substances similar to the compound under study were included in the training set.

Subsets of the training set can be reconstructed by a careful analysis of the original papers and the literature cited therein. Thus, it would probably be possible to reconstruct large parts of the training set and thus derive a more comprehensive description of the applicability domain.

5.3.Software name and version for applicability domain assessment:

5.4.Limits of applicability:

If it is not possible to describe all features of the chemical structure by respective increments (group rate constants and/or substituent factors), the compound is considered outside the applicability domain. Generally, the method can be considered to describe organic compounds that may contain the following atoms: carbon, hydrogen, oxygen, nitrogen, sulfur, phosphorus. A substance containing other atoms should be regarded as being outside the application domain. An exception are lead and mercury: In the user's guide, it is mentioned that additional substituent factors F(Pb) and F(Hg) were added. However, it should be checked carefully whether the chemical structure can be adequately described by the respective increments.

6.Internal validation - OECD Principle 4

6.1.Availability of the training set:

No

6.2.Available information for the training set:

CAS RN: No

Chemical Name: No

Smiles: No

Formula:No

INChI:No

MOL file:No

6.3.Data for each descriptor variable for the training set:

No

6.4.Data for the dependent variable for the training set:

No

6.5.Other information about the training set:

6.6.Pre-processing of data before modelling:

6.Internal validation - OECD Principle 4

6.7.Statistics for goodness-of-fit:

6.8.Robustness - Statistics obtained by leave-one-out cross-validation:

6.9. Robustness - Statistics obtained by leave-many-out cross-validation:

6.10. Robustness - Statistics obtained by Y-scrambling:

6.11. Robustness - Statistics obtained by bootstrap:

6.12. Robustness - Statistics obtained by other methods:

7. External validation - OECD Principle 4

7.1. Availability of the external validation set:

No

7.2. Available information for the external validation set:

CAS RN: No

Chemical Name: Yes

Smiles: No

Formula: No

INChI: No

MOL file: No

7.3. Data for each descriptor variable for the external validation set:

No

7.4. Data for the dependent variable for the external validation set:

All

7.5. Other information about the external validation set:

partly published (Meylan and Howard 1993); refers to the original version of the model (before 1995 update)

7.6. Experimental design of test set:

7.7. Predictivity - Statistics obtained by external validation:

Overall dataset:

$n = 77$; $r^2 = 0.89$; $stdv = 0.24$; $me = 0.17$, $mne = -2.56$; $mpe = 2.2$ (referring to $\log k_{OH}$)

n = number of compounds; $stdv$ = standard deviation; me = mean error; mne = maximum negative error (largest underestimation); mpe = maximum positive error (largest overestimation)

7.8. Predictivity - Assessment of the external validation set:

The published part of the test set contains 35 substances (5 aliphatic compounds, 19 oxygenated organic compounds, 8 aromatics, 1 alkyne, 2 substances containing N or S functional groups but no C-C multiple bonds).

The reaction mechanism represented best in this subset is hydrogen abstraction from C-H and O-H bonds (24 compounds), followed by aromatics. OH radical addition to double and triple bonds is only represented by one alkyne with no olefinic compounds reported. The unpublished part of the test set cannot be assessed.

7.9. Comments on the external validation of the model:

Model performance was also validated using a larger data set of 805 compounds (Böhnhardt et al 2008), referring to the updated AOPWIN model. However, it should be noted that this data set contains external data as well as data that were used for model development. The test set contains data from the PhysProp Database as well as further experimental reaction rates collected from literature. It covers the following reaction mechanisms: H-atom abstraction from C-H and O-H bonds; OH radical addition to double and triple bonds; OH radical addition to aromatic rings. The test set does only contain N, S and P containing functional groups that are substituted to aromatic rings and thus, model performance for other N, S and P containing compounds is not tested.

The following statistics are reported for the overall dataset (referring to $\log k_{OH}$):
 $n = 805$; $r^2 = 0.93$; $q^2 = 0.92$; $rms = 0.35$; $bias = 0$; $me = 0.2$, $mne = -2.56$; $mpe = 2.2$

The test set was divided into subsets according to chemical compound classes and thus, compound-class specific statistics are also available in the original study (Böhnhardt et al 2008). The model performance was found to depend on the compound-class under consideration. The best performance is observed for the subset of alkynes ($r^2 = 0.99$), however, this data set contains only 7 substances (most of which were probably used to derive the model). This QMRF gives statistics for the compound classes with the weakest and the second best performance, respectively:

Alkenes:

$n = 124$; $r^2 = 0.62$; $q^2 = 0.51$; $rms = 0.48$; $bias = -0.05$; $me = 0.23$, $mne = -2.56$; $mpe = 1.29$

Aromatics:

$n = 134$; $r^2 = 0.94$; $q^2 = 0.94$; $rms = 0.21$; $bias = -0.02$; $me = 0.14$, $mne = -0.65$; $mpe = 0.83$

n = number of compounds; q^2 = predictive squared correlation coefficient; rms = root-mean-square error; $bias$ = systematic error; me = mean error; mne = maximum negative error (largest underestimation); mpe = maximum positive error (largest overestimation)

8. Providing a mechanistic interpretation - OECD Principle 5

8.1. Mechanistic basis of the model:

The model is developed based on profound knowledge of the different reaction pathways and assigns incremental values that are specific for the mechanism under consideration. The overall reaction rate is assumed to be the sum of four reaction constants corresponding to the following reaction mechanisms: H-atom abstraction from C-H and O-H bonds; OH radical addition to double and triple bonds; OH radical addition to aromatic rings; and OH radical interaction with N-, S-, and P atoms. These mechanism-specific rate constants are assumed to be the sum of partial rate constants that can be assigned to the individual reaction sites and that are described by increment values.

8.2. A priori or a posteriori mechanistic interpretation:

a priori

8.3. Other information about the mechanistic interpretation:

9. Miscellaneous information

9.1. Comments:

9.2. Bibliography:

- [1] Meylan WM and Howard PH 1993. Computer estimation of the atmospheric gas-phase reaction rate of organic compounds with hydroxyl radicals and ozone. Chemosphere 26: 2293-99. [http://dx.doi.org/10.1016/0045-6535\(93\)90355-9](http://dx.doi.org/10.1016/0045-6535(93)90355-9)
- [2] Böhnhardt A, Kühne R, Ebert R-U, Schüürmann G 2008. Indirect photolysis of organic compounds – prediction of OH reaction rate constants through molecular orbital calculations. J. Phys. Chem. A. 112: 11391-11399. <http://dx.doi.org/10.1021/jp8052218>
- [3] Brown HC and Okamoto Y 1958. Electrophilic Substituent Constants. Journal of the American Chemical Society 80 (18):4979-4987. <http://dx.doi.org/10.1021/ja01551a055>

9.3. Supporting information:

Training set(s) Test set(s) Supporting information

10. Summary (JRC Inventory)

10.1. QMRF number:

To be entered by JRC

10.2. Publication date:

To be entered by JRC

10.3. Keywords:

To be entered by JRC

10.4. Comments:

To be entered by JRC