

# CLH report

## Proposal for Harmonised Classification and Labelling

Based on Regulation (EC) No 1272/2008 (CLP Regulation),  
Annex VI, Part 2

### International Chemical Identification:

**(1,3,4,5,6,7-hexahydro-1,3-dioxo-2*H*-isoindol-2-yl)methyl (1*R*-*trans*)-  
2,2-dimethyl-3-(2-methylprop-1-enyl)cyclopropanecarboxylate; d-  
*trans*-tetramethrin**

**EC Number:** 214-619-0

**CAS Number:** 1166-46-7

**Index Number:** -

#### Contact details for dossier submitter:

**BAuA**  
Federal Institute for Occupational Safety and Health  
Federal Office for Chemicals  
Friedrich-Henkel-Weg 1-25  
44149 Dortmund  
e-mail: chemg@baua.bund.de

**Version number:** 2 (post Accordance Check)

**Date:** September 2015

# CONTENTS

<b>1. IDENTITY OF THE SUBSTANCE .....</b>	<b>1</b>
1.1 NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE .....	1
1.2 COMPOSITION OF THE SUBSTANCE.....	2
<b>2. PROPOSED HARMONISED CLASSIFICATION AND LABELLING.....</b>	<b>3</b>
2.1 PROPOSED HARMONISED CLASSIFICATION AND LABELLING ACCORDING TO THE CLP CRITERIA.....	3
<b>3. HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING.....</b>	<b>5</b>
<b>4. JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL .....</b>	<b>5</b>
<b>5. IDENTIFIED USES .....</b>	<b>5</b>
<b>6. DATA SOURCES .....</b>	<b>5</b>
<b>7. PHYSICOCHEMICAL PROPERTIES .....</b>	<b>5</b>
<b>8. EVALUATION OF PHYSICAL HAZARDS .....</b>	<b>8</b>
8.1 EXPLOSIVES.....	8
8.1.1 <i>Short summary and overall relevance of the provided information on explosive properties</i> .....	8
8.1.2 <i>Comparison with the CLP criteria</i> .....	8
8.1.3 <i>Conclusion on classification and labelling for explosive properties</i> .....	8
8.2 FLAMMABLE GASES (INCLUDING CHEMICALLY UNSTABLE GASES) .....	8
8.3 OXIDISING GASES .....	8
8.4 GASES UNDER PRESSURE .....	8
8.5 FLAMMABLE LIQUIDS .....	9
8.5.1 <i>Short summary and overall relevance of the provided information on flammable liquids</i> .....	9
8.5.2 <i>Comparison with the CLP criteria</i> .....	9
8.5.3 <i>Conclusion on classification and labelling for flammable liquids</i> .....	9
8.6 FLAMMABLE SOLIDS.....	9
8.7 SELF-REACTIVE SUBSTANCES.....	9
8.7.1 <i>Short summary and overall relevance of the provided information on self-reactive substances</i> .....	9
8.7.2 <i>Comparison with the CLP criteria</i> .....	9
8.7.3 <i>Conclusion on classification and labelling for self-reactive substances</i> .....	10
8.8 PYROPHORIC LIQUIDS .....	10
8.8.1 <i>Short summary and overall relevance of the provided information on pyrophoric liquids</i> .....	10
8.8.2 <i>Comparison with the CLP criteria</i> .....	10
8.8.3 <i>Conclusion on classification and labelling for pyrophoric liquids</i> .....	10
8.9 PYROPHORIC SOLIDS .....	10
8.10 SELF-HEATING SUBSTANCES.....	10
8.11 SUBSTANCES WHICH IN CONTACT WITH WATER EMIT FLAMMABLE GASES.....	11
8.11.1 <i>Short summary and overall relevance of the provided information on substances which in contact with water emit flammable gases</i> .....	11
8.11.2 <i>Comparison with the CLP criteria</i> .....	11
8.11.3 <i>Conclusion on classification and labelling for substances which in contact with water emit flammable gases</i> .....	11
8.12 OXIDISING LIQUIDS .....	11
8.12.1 <i>Short summary and overall relevance of the provided information on oxidising liquids</i> .....	11
8.12.2 <i>Comparison with the CLP criteria</i> .....	11
8.12.3 <i>Conclusion on classification and labelling for oxidising liquids</i> .....	12
8.13 OXIDISING SOLIDS .....	12
8.14 ORGANIC PEROXIDES .....	12
8.15 CORROSIVE TO METALS.....	12
<b>9. TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION) .....</b>	<b>13</b>

9.1	SHORT SUMMARY AND OVERALL RELEVANCE OF THE PROVIDED TOXICOKINETIC INFORMATION ON THE PROPOSED CLASSIFICATION(S).....	16
<b>10.</b>	<b>EVALUATION OF HEALTH HAZARDS.....</b>	<b>19</b>
10.1	ACUTE TOXICITY - ORAL ROUTE .....	21
10.1.1	Short summary and overall relevance of the provided information on acute oral toxicity.....	22
10.1.2	Comparison with the CLP criteria.....	23
10.1.3	Conclusion on classification and labelling for acute oral toxicity.....	23
10.2	ACUTE TOXICITY - DERMAL ROUTE.....	23
10.2.1	Short summary and overall relevance of the provided information on acute dermal toxicity .....	24
10.2.2	Comparison with the CLP criteria.....	24
10.2.3	Conclusion on classification and labelling for acute dermal toxicity .....	24
10.3	ACUTE TOXICITY - INHALATION ROUTE.....	25
10.3.1	Short summary and overall relevance of the provided information on acute inhalation toxicity.....	26
10.3.2	Comparison with the CLP criteria.....	27
10.3.3	Conclusion on classification and labelling for acute inhalation toxicity .....	28
10.4	SKIN CORROSION/IRRITATION .....	28
10.4.1	Short summary and overall relevance of the provided information on skin corrosion/irritation.....	28
10.4.2	Comparison with the CLP criteria.....	29
10.4.3	Conclusion on classification and labelling for skin corrosion/irritation.....	29
10.5	SERIOUS EYE DAMAGE/EYE IRRITATION .....	29
10.5.1	Short summary and overall relevance of the provided information on serious eye damage/eye irritation .....	30
10.5.2	Comparison with the CLP criteria.....	30
10.5.3	Conclusion on classification and labelling for serious eye damage/eye irritation.....	30
10.6	RESPIRATORY SENSITISATION.....	31
10.6.1	Short summary and overall relevance of the provided information on respiratory sensitisation.....	32
10.6.2	Comparison with the CLP criteria.....	32
10.6.3	Conclusion on classification and labelling for respiratory sensitisation .....	32
10.7	SKIN SENSITISATION .....	32
10.7.1	Short summary and overall relevance of the provided information on skin sensitisation .....	33
10.7.2	Comparison with the CLP criteria.....	34
10.7.3	Conclusion on classification and labelling for skin sensitisation .....	34
10.8	GERM CELL MUTAGENICITY .....	35
10.8.1	Short summary and overall relevance of the provided information on germ cell mutagenicity.....	38
10.8.2	Comparison with the CLP criteria.....	39
10.8.3	Conclusion on classification and labelling for germ cell mutagenicity.....	40
10.9	CARCINOGENICITY .....	40
10.9.1	Short summary and overall relevance of the provided information on carcinogenicity.....	42
10.9.2	Comparison with the CLP criteria.....	45
10.9.3	Conclusion on classification and labelling for carcinogenicity.....	47
10.10	REPRODUCTIVE TOXICITY .....	48
10.10.1	Adverse effects on sexual function and fertility.....	48
10.10.2	Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility.....	51
10.10.3	Comparison with the CLP criteria.....	52
10.10.4	Adverse effects on development .....	52
10.10.5	Short summary and overall relevance of the provided information on adverse effects on development .....	57
10.10.6	Comparison with the CLP criteria.....	59
10.10.7	Adverse effects on or via lactation.....	59
10.10.8	Short summary and overall relevance of the provided information on effects on or via lactation.....	60
10.10.9	Comparison with the CLP criteria.....	60
10.10.10	Conclusion on classification and labelling for reproductive toxicity.....	60
10.11	SPECIFIC TARGET ORGAN TOXICITY-SINGLE EXPOSURE.....	61
10.11.1	Short summary and overall relevance of the provided information on specific target organ toxicity – single exposure .....	63
10.11.2	Comparison with the CLP criteria.....	64
10.11.3	Conclusion on classification and labelling for STOT SE .....	65
10.12	SPECIFIC TARGET ORGAN TOXICITY-REPEATED EXPOSURE .....	66

# CLH REPORT FOR D-TRANS-TETRAMETHRIN


---

10.13	SHORT SUMMARY AND OVERALL RELEVANCE OF THE PROVIDED INFORMATION ON SPECIFIC TARGET ORGAN TOXICITY – REPEATED EXPOSURE .....	69
10.13.1	Comparison with the CLP criteria.....	74
10.13.2	Conclusion on classification and labelling for STOT RE.....	75
10.14	ASPIRATION HAZARD .....	75
<b>11.</b>	<b>EVALUATION OF ENVIRONMENTAL HAZARDS .....</b>	<b>76</b>
11.1	ACUTE AQUATIC HAZARD .....	76
11.1.1	Acute (short-term) toxicity to fish .....	77
11.1.2	Acute (short-term) toxicity to algae or other aquatic plants.....	78
11.2	LONG-TERM AQUATIC HAZARD .....	78
11.3	BIOACCUMULATION.....	79
11.3.1	Estimated bioaccumulation .....	79
11.3.2	Measured partition coefficient and bioaccumulation test data.....	80
	<b>NAME OF COMPOUND USED IN REPORTS.....</b>	<b>84</b>
	<b>CONCENTRATION AT END OF UPTAKE PHASE (DAY 28)<sup>1</sup>.....</b>	<b>84</b>
	<b>CONCENTRATION AT END OF DEPURATION PHASE.....</b>	<b>84</b>
11.4	RAPID DEGRADABILITY OF ORGANIC SUBSTANCES .....	85
11.4.1	Ready biodegradability .....	87
11.4.2	BOD <sub>5</sub> /COD.....	88
11.4.3	Other convincing scientific evidence.....	88
11.4.3.1	Field investigations and monitoring data (if relevant for C&L) .....	88
11.4.3.2	Inherent and Enhanced Ready Biodegradability tests .....	88
11.4.3.3	Soil and sediment degradation data .....	88
11.4.3.4	Hydrolysis .....	89
11.4.3.5	Photochemical degradation .....	90
11.5	ENVIRONMENTAL FATE AND OTHER RELEVANT INFORMATION.....	91
11.5.1	Adsorption/desorption.....	91
11.6	COMPARISON WITH THE CLP CRITERIA .....	92
11.6.1	Acute aquatic hazard .....	92
11.6.2	Long-term aquatic hazard (including bioaccumulation potential and degradation) .....	92
11.7	CONCLUSION ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS .....	93
<b>12.</b>	<b>EVALUATION OF ADDITIONAL HAZARDS .....</b>	<b>93</b>
12.1	HAZARDOUS TO THE OZONE LAYER .....	93
<b>13.</b>	<b>DETAILED STUDY SUMMARIES.....</b>	<b>93</b>
<b>14.</b>	<b>REFERENCES.....</b>	<b>93</b>
<b>15.</b>	<b>ANNEXES.....</b>	<b>94</b>

## 1. IDENTITY OF THE SUBSTANCE

### 1.1 NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE

Table 1: Substance identity and information related to molecular and structural formula of the substance

<b>Name(s) in the IUPAC nomenclature</b>	(1,3-dioxo-1,3,4,5,6,7-hexahydro-2 <i>H</i> -isoindol-2-yl)methyl (1 <i>R</i> ,3 <i>R</i> )-2,2-dimethyl-3-(2-methylprop-1-en-1-yl)cyclopropanecarboxylate
<b>Other names (usual name, trade name, abbreviation)</b>	<i>d-trans</i> -Tetramethrin; CAS name: Cyclopropanecarboxylic acid, 2, 2-dimethyl-3-(2-methyl-1-propen-1-yl)-, (1, 3, 4, 5, 6, 7- hexahydro-1, 3- dioxo-2 <i>H</i> -isoindol-2-yl) methyl ester, (1 <i>R</i> , 3 <i>R</i> )-
<b>EC number (if available and appropriate)</b>	214-619-0
<b>EC name (if available and appropriate)</b>	(1,3,4,5,6,7-hexahydro-1,3-dioxo-2 <i>H</i> -isoindol-2-yl)methyl (1 <i>R-trans</i> )-2,2-dimethyl-3-(2-methylprop-1-enyl)cyclopropanecarboxylate
<b>CAS number (if available)</b>	1166-46-7
<b>Molecular formula</b>	C <sub>19</sub> H <sub>25</sub> NO <sub>4</sub>
<b>Structural formula</b>	
<b>Molecular weight or molecular weight range</b>	331.41 g/mol

## 1.2 COMPOSITION OF THE SUBSTANCE

Table 2: Constituents (non-confidential information)

Constituent (Name and numerical identifier)	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self-classification and labelling (CLP)
(1,3-dioxo-1,3,4,5,6,7-hexahydro-2 <i>H</i> -isoindol-2-yl)methyl (1 <i>R</i> ,3 <i>R</i> )-2,2-dimethyl-3-(2-methylprop-1-en-1-yl)cyclopropanecarboxylate; EC number: 214-619-0			

For further information: Please refer to the confidential annex.

Table 3: Impurities (non-confidential information) if relevant for the classification of the substance

Impurity (Name and numerical identifier)	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self-classification and labelling (CLP)	The impurity contributes to the classification and labelling
For further information: Please refer to the confidential annex.				

Table 4: Additives (non-confidential information) if relevant for the classification of the substance

Additive (Name and numerical identifier)	Function	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self-classification and labelling (CLP)	The additive contributes to the classification and labelling
For further information: Please refer to the confidential annex.					

Table 5: Test substances (non-confidential information)

Identification of test substance	Purity	Impurities and additives (identity, %, classification if available)	Other information

## 2. PROPOSED HARMONISED CLASSIFICATION AND LABELLING

### 2.1 PROPOSED HARMONISED CLASSIFICATION AND LABELLING ACCORDING TO THE CLP CRITERIA

Table 6: Proposed harmonised classification and labelling according to the CLP criteria

	Index No	International Chemical Identification	EC No	CAS No	Classification		Labelling			Specific Conc. Limits, M-factors	Notes
					Hazard Class and Category Code(s)	Hazard statement Code(s)	Pictogram, Signal Word Code(s)	Hazard statement Code(s)	Suppl. Hazard statement Code(s)		
Current Annex VI entry	-										
Dossier submitters proposal	607-RST-00-Y	(1,3,4,5,6,7-hexahydro-1,3-dioxo-2H-isoindol-2-yl)methyl (1R-trans)-2,2-dimethyl-3-(2-methylprop-1-enyl)cyclopropane carboxylate; d- <i>trans</i> -tetramethrin	214-619-0	1166-46-7	Acute Tox. 4 Carc. 2 STOT SE 2 Aquatic Acute 1 Aquatic Chronic 1	H332 H351 H371 H400 H410	GHS07 GHS08 GHS09 Warning	H332 H351 H371 H410		M = 100 (acute) M = 100 (chronic)	
Resulting Annex VI entry if agreed by RAC and COM	607-RST-00-Y	(1,3,4,5,6,7-hexahydro-1,3-dioxo-2H-isoindol-2-yl)methyl (1R-trans)-2,2-dimethyl-3-(2-methylprop-1-enyl)cyclopropane carboxylate; d- <i>trans</i> -tetramethrin	214-619-0	1166-46-7	Acute Tox. 4 Carc. 2 STOT SE 2 Aquatic Acute 1 Aquatic Chronic 1	H332 H351 H371 H400 H410	GHS07 GHS08 GHS09 Warning	H332 H351 H371 H410		M = 100 (acute) M = 100 (chronic)	

Table 7: Reason for not proposing harmonised classification and status under public consultation

<b>Hazard class</b>	<b>Reason for no classification</b>	<b>Within the scope of public consultation</b>
<b>Explosives</b>	data conclusive but not sufficient for classification	Yes
<b>Flammable gases (including chemically unstable gases)</b>	hazard class not applicable	
<b>Oxidising gases</b>	hazard class not applicable	
<b>Gases under pressure</b>	hazard class not applicable	
<b>Flammable liquids</b>	data conclusive but not sufficient for classification	Yes
<b>Flammable solids</b>	hazard class not applicable	
<b>Self-reactive substances</b>	data conclusive but not sufficient for classification	Yes
<b>Pyrophoric liquids</b>	data conclusive but not sufficient for classification	Yes
<b>Pyrophoric solids</b>	hazard class not applicable	
<b>Self-heating substances</b>	hazard class not applicable	
<b>Substances which in contact with water emit flammable gases</b>	data conclusive but not sufficient for classification	Yes
<b>Oxidising liquids</b>	data conclusive but not sufficient for classification	Yes
<b>Oxidising solids</b>	hazard class not applicable	
<b>Organic peroxides</b>	hazard class not applicable	
<b>Corrosive to metals</b>	hazard class not assessed in this dossier	Yes
<b>Acute toxicity via oral route</b>	data conclusive but not sufficient for classification	Yes
<b>Acute toxicity via dermal route</b>	data conclusive but not sufficient for classification	Yes
<b>Acute toxicity via inhalation route</b>	harmonised classification proposed	Yes
<b>Skin corrosion/irritation</b>	data conclusive but not sufficient for classification	Yes
<b>Serious eye damage/eye irritation</b>	data conclusive but not sufficient for classification	Yes
<b>Respiratory sensitisation</b>	data lacking	Yes
<b>Skin sensitisation</b>	data conclusive but not sufficient for classification	Yes
<b>Germ cell mutagenicity</b>	data conclusive but not sufficient for classification	Yes
<b>Carcinogenicity</b>	harmonised classification proposed	Yes
<b>Reproductive toxicity</b>	data conclusive but not sufficient for classification	Yes
<b>Specific target organ toxicity-single exposure</b>	harmonised classification proposed	Yes
<b>Specific target organ toxicity-repeated exposure</b>	data conclusive but not sufficient for classification	Yes
<b>Aspiration hazard</b>	data lacking	Yes
<b>Hazardous to the aquatic environment</b>	harmonised classification proposed	Yes
<b>Hazardous to the ozone layer</b>	hazard class not applicable	



**3. HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING****4. JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL**

The substance is an active substance in the meaning of Directive 91/414/EEC (repealed by the Regulation EC 1107/2009) or Directive 98/8/EC (will be repealed by Regulation (EU) No 528/2012 on 1 September 2013) shall normally be subject to harmonised classification and labelling, and justification is not required. (Article 36 CLP Regulation)

There is no requirement for justification that action is needed at Community level.

**5. IDENTIFIED USES****6. DATA SOURCES****7. PHYSICOCHEMICAL PROPERTIES**

Table 8: Summary of physicochemical properties

Property	Value	Reference	Comment (e.g. measured or estimated)
<b>Physical state at 20°C and 101,3 kPa</b>	extremely viscous liquid (T = 25 °C) (88.7% <i>1R-trans-Isomer</i> )	O'Donnell, R. T. et al Sumitomo Report No. SVP-0008, 1999	measured (visual assessment)
<b>Melting/freezing point</b>	19 °C ± 3 °C (pour point) (89.5% <i>1R-trans-Isomer</i> )	Evans A.J. et al, SPL Project No: 1430/011, 2002	measured (92/69/EEC, A.1; pour point)
<b>Boiling point</b>	263 °C (pressure: 988 hPa) (100% <i>1R-trans-Isomer</i> )	Malinkski, M. F. Sumitomo Report No. SVP-0003, 1999	measured (92/69/EEC, A.2 ; DSC)
<b>Relative density</b>	1.11 g/cm <sup>3</sup> (92.1% <i>1R-trans-Isomer</i> )	Lentz NR (2008), Sumitomo Report No SVP-0020	measured (92/69/EEC, A.3; Pycnometer method)
<b>Vapour pressure</b>	4.03*10 <sup>-6</sup> Pa (T = 25 °C), 1.56 * 10 <sup>-6</sup> Pa (T = 20 °C) (100% <i>1R-trans-Isomer</i> )	Schetter, J. E. Sumitomo Report No. SVP-0006, 1999	measured (92/69/EEC, A.4; gas saturation method)
<b>Surface tension</b>	55.9 mN/m (mean) (T = 20.4 °C, c = 1.58 mg/l) (92.1 % <i>1R-trans-Isomer</i> )	Lentz NR (2008), Study No. 13048.6593	measured (92/69/EEC, A.5; ring method)
<b>Water solubility</b>	1.60 ± 0.21 mg/l (mean) at T = 20 °C, pH = 6 (unbuffered) (92.1 % <i>1R-trans-Isomer</i> )	Lentz NR (2008), Study No. 13048.6571	measured (92/69/EEC, A.6; column elution method)
<b>Partition coefficient <i>n</i>-octanol/water</b>	log Pow = 4.3 (T = 25 °C) (100% <i>1R-trans-Isomer</i> )	Dudones, L. P. Sumitomo Report No. SVP-0004 (1999)	measured (92/69/EEC, A.8; HPLC method)

CLH REPORT FOR D-TRANS-TETRAMETHRIN

<b>Flash point</b>	148 °C (± 2 °C) (Purity: 96.0 %)	White D. F., Mullee D. M., SPL Project No: 0555/0076, 2006	measured (92/69/EEC, A.9: ISO 3679:1983 Setaflash closed-cup apparatus)
<b>Flammability</b>	<p>Since the flash point for d-tetramethrin was found to be 148 ± 2 °C, a flammability study is scientifically unjustified.</p> <p>Flammability in contact with water: The classification procedure needs not to be applied because the organic substance does not contain metals or metalloids.</p> <p>Pyrophoric properties: The classification procedure needs not to be applied because the organic substance is known to be stable into contact with air at room temperature for prolonged periods of time (days).</p>	<p>Sumitomo Chemical (2006)</p> <p>BAM 2.2 (2014)</p>	<p>Justification for non-submission of data - Method 92/69/EEC, A.10, A.11</p> <p>Justification for non-submission of data - Method 92/69/EEC, A.12, A.13</p>
<b>Explosive properties</b>	d-Tetramethrin does not have any functional groups such as diazo, azide, polynitro or peroxide, which are found in chemically explosive compounds. Therefore, chemical explosion reactions are considered not to occur under ordinary circumstances. A study to determine explosive properties would therefore be scientifically unjustified.	Sumitomo Chemical (2006)	Justification for non-submission of data - Method 92/69/EEC, A.14
<b>Self-ignition temperature</b>	366 °C (± 5°C) (atmospheric pressure: 101.25 to 102.73 kPa) (Purity: 96.0 %)	White D.F., , Mullee D. M., SPL Project No: 0555/0076, 2006	measured (92/69/EEC, A.15)
<b>Oxidising properties</b>	d-Tetramethrin does not have any functional groups associated with oxidising action. It does not undergo substitution, addition or elimination reactions. Therefore, oxidising action is considered not to occur under ordinary circumstances. A study to determine oxidising properties would therefore be scientifically unjustified.	Sumitomo Chemical (2006)	Justification for non-submission of data - Method 92/69/EEC, A.21
<b>Granulometry</b>	Not applicable. <i>(The substance is a liquid under ambient conditions.)</i>	-	-

CLH REPORT FOR D-TRANS-TETRAMETHRIN

<p><b>Stability in organic solvents and identity of relevant degradation products</b></p>	<p>The test substance is stable at T = 40 °C for 3 month in the following solvents: Xylene, Deobase, Methyl isobutyl Ketone, Ethyl acetate, Acetonitrile, Chloroform, Propionic Acid, Dimethylformamide, Isopropyl alcohol, Kerosene, Methyl Chlorofom</p> <p>The test substance is not stable at T = 60 °C for 3 month in the following solvents: Methanol, Ethylcellosorb, Cyclohexanone, Chloroform, Propionic acid, Dimethylformamide, Corn oil (74.9 % 1R-trans-Isomer, 17.5 % 1R-cis-Isomer )</p>	<p>Sumitomo Chemical Co., Sumitomo Report No. IP-10-0054, 1986</p>	<p>measured (<i>Gaschromatography</i>)</p>
<p><b>Dissociation constant</b></p>	<p><math>pK_a = -2.55 \pm 0.20</math> (<i>calculation</i>)</p>	<p>Roth H., Safepharm Laboratories, 2006</p>	<p><i>calculated</i></p>
<p><b>Viscosity</b></p>	<p><math>8.89 \times 10^4</math> mPa.s temperature: 20.0°C (89.7 % 1R-trans-Isomer)</p>	<p>White D.F., SPL Project No: 0555/0076, 2006</p>	<p>measured (<i>OECD 114; rotational viscometer</i>)</p>

## 8. EVALUATION OF PHYSICAL HAZARDS

### 8.1 EXPLOSIVES

Table 9: Summary table of studies on explosives

Method	Results	Remarks	Reference
Justification for data waiving	d-Tetramethrin does not have any functional groups such as diazo, azide, polynitro or peroxide, which are found in chemically explosive compounds. Therefore, chemical explosion reactions are considered not to occur under ordinary circumstances. A study to determine explosive properties would therefore be scientifically unjustified.		Sumitomo Chemical (2006)

#### 8.1.1 Short summary and overall relevance of the provided information on explosive properties

Justification is acceptable because there are no chemical groups present in the molecule which are associated with explosive properties.

#### 8.1.2 Comparison with the CLP criteria

The classification procedure needs not to be applied due to justification for data waiving.

#### 8.1.3 Conclusion on classification and labelling for explosive properties

Should not be classified as explosives according to Annex I part 2 of the CLP regulation.

### 8.2 FLAMMABLE GASES (INCLUDING CHEMICALLY UNSTABLE GASES)

Hazard class not applicable.

### 8.3 OXIDISING GASES

Hazard class not applicable.

### 8.4 GASES UNDER PRESSURE

Hazard class not applicable.

## 8.5 FLAMMABLE LIQUIDS

Table 10: Summary table of studies on flammable liquids

Method	Results	Remarks	Reference
92/69/EEC, A.9: ISO 3679:1983 Setaflash closed-cup apparatus	148 °C ( $\pm 2$ °C)		White D. F., Mullee D. M.,SPL Project No: 0555/0076, 2006

### 8.5.1 Short summary and overall relevance of the provided information on flammable liquids

A single study is available. The flash point for d-tetramethrin was found to be  $148 \pm 2$  °C.

### 8.5.2 Comparison with the CLP criteria

A test was conducted according to EU Method A.9. Based on the experimental data it is concluded that d-tetramethrin with a flash point of more than 60 °C does not meet the criteria for classification as flammable liquid according to Annex I part 2 of the CLP regulation.

### 8.5.3 Conclusion on classification and labelling for flammable liquids

Should not be classified as flammable liquid according to Annex I part 2 of the CLP regulation.

## 8.6 FLAMMABLE SOLIDS

Hazard class not applicable.

## 8.7 SELF-REACTIVE SUBSTANCES

Table 11: Summary table of studies on self-reactivity

Method	Results	Remarks	Reference
Justification for data waiving	The classification procedure needs not to be applied because there are no chemical groups present in the molecule which are associated with explosive or self-reactive properties.		BAM 2.2 (2014)

### 8.7.1 Short summary and overall relevance of the provided information on self-reactive substances

Justification is acceptable.

### 8.7.2 Comparison with the CLP criteria

The classification procedure needs not to be applied due to justification for data waiving.

**8.7.3 Conclusion on classification and labelling for self-reactive substances**

Should not be classified as self-reactive substance according to Annex I part 2 of the CLP regulation.

**8.8 PYROPHORIC LIQUIDS**

Table 12: Summary table of studies on pyrophoric liquids

Method	Results	Remarks	Reference
Justification for data waiving	The classification procedure needs not to be applied because the organic substance is known to be stable into contact with air at room temperature for prolonged periods of time (days).		BAM 2.2 (2014)

**8.8.1 Short summary and overall relevance of the provided information on pyrophoric liquids**

Justification is acceptable.

**8.8.2 Comparison with the CLP criteria**

The classification procedure needs not to be applied due to justification for data waiving.

**8.8.3 Conclusion on classification and labelling for pyrophoric liquids**

Should not be classified as pyrophoric liquid according to Annex I part 2 of the CLP regulation.

**8.9 PYROPHORIC SOLIDS**

Hazard class not applicable.

**8.10 SELF-HEATING SUBSTANCES**

Hazard class not applicable.

## 8.11 SUBSTANCES WHICH IN CONTACT WITH WATER EMIT FLAMMABLE GASES

Table 13: Summary table of studies on substances which in contact with water emit flammable gases

Method	Results	Remarks	Reference
Justification for data waiving	The classification procedure needs not to be applied because the organic substance does not contain metals or metalloids.		BAM 2.2 (2014)

### 8.11.1 Short summary and overall relevance of the provided information on substances which in contact with water emit flammable gases

Justification is acceptable.

### 8.11.2 Comparison with the CLP criteria

The classification procedure needs not to be applied due to justification for data waiving.

### 8.11.3 Conclusion on classification and labelling for substances which in contact with water emit flammable gases

Should not be classified as substance which in contact with water emits flammable gases according to Annex I part 2 of the CLP regulation.

## 8.12 OXIDISING LIQUIDS

Table 14: Summary table of studies on oxidising liquids

Method	Results	Remarks	Reference
Justification for data waiving	d-Tetramethrin does not have any functional groups associated with oxidising action. It does not undergo substitution, addition or elimination reactions. Therefore, oxidising action is considered not to occur under ordinary circumstances. A study to determine oxidising properties would therefore be scientifically unjustified.		Sumitomo Chemical (2006)

### 8.12.1 Short summary and overall relevance of the provided information on oxidising liquids

Justification is acceptable because there are no chemical groups present in the molecule which are associated with oxidising properties.

### 8.12.2 Comparison with the CLP criteria

The classification procedure needs not to be applied due to justification for data waiving.

### **8.12.3 Conclusion on classification and labelling for oxidising liquids**

Should not be classified as oxidising liquid according to Annex I part 2 of the CLP regulation.

### **8.13 OXIDISING SOLIDS**

Hazard class not applicable.

### **8.14 ORGANIC PEROXIDES**

Hazard class not applicable.

### **8.15 CORROSIVE TO METALS**

Hazard class not assessed in this dossier.



## 9. TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)

Table 15: Summary table of toxicokinetic studies

Method, guideline	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Results	Remarks	Reference
Similar to OECD 417, oral, gavage	Rat, Sprague-Dawley, Low, high and repeated dose groups: 5 M + 5 F  Controls: 3 M + 3 F	[1RS, trans]-tetramethrin in in corn oil, C-14-labelled and unlabelled	<u>Low single dose group with labelled material:</u> 2 mg/kg bw  <u>High single dose group with labelled material:</u> 250 mg/kg bw  <u>Repeated dose group:</u> 14 days pre-treatment with unlabelled material at 2 mg/kg bw/day, followed by single dose of labelled material (2 mg/kg bw)  <u>Control group:</u> 14 days pre-treatment with corn oil at 5 ml/kg/day, followed by single dose of labelled material (2 mg/kg bw)	Radiocarbon absorption 42-71 % (based on C-14 excretion in urine),  94-100 % and 95-101 % excretion of radioactivity within 2 and 7 days, respectively (urine: 42.3-71.4 %, faeces: 29-57.9 %, air: < 0.1 %),  0.2-0.4 % C-14 tissue residues after 7 days, widely distributed, highest residues in blood cells  Excreted at >5 % in urine plus faeces within 2 days after dosing of radioactive compound in any group: 1-sulfo-cyclohexane-dicarboximide, 3-hydroxy-cyclohexane-dicarboximide  Excreted unmetabolised parent in urine plus faeces: 5-23 % in single dose groups, < 3 % in repeated dose groups	C-14 label at phthalimide moiety only (alcohol label)  48 % of radioactivity in unknown or non-extractable metabolites	Shiba K, 1992, Sumitomo Report No. IM-20-0015  Parts of study also published in: Tomigahara et al., 1994, Xenobiotica 24(12): 1205-1214
Sim. to OECD 417, Oral, gavage	Rat, Sprague-Dawley, Low, high and repeated dose groups: 5 M + 5 F	[1RS, cis]-tetramethrin in in corn oil, C-14-labelled and unlabelled	<u>Low single dose group with labelled material:</u> 2 mg/kg bw  <u>High single dose group with labelled material:</u>	Radiocarbon absorption 9-32 % (based on C-14 excretion in urine),  95-101 % and 96-102 % excretion of radioactivity within 2 and 7 days, respectively (urine: 8.5-32.4, faeces: 65.9-91.3, air: < 0.1 %),	C-14 label at phthalimide moiety only (alcohol label)  42-68 % unknown, non-extractable or other radioactive	Shiba K, 1992, Sumitomo Report No. IM-20-0016  Parts of study also published in: Tomigahara

Method, guideline	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Results	Remarks	Reference
	Controls: 3 M + 3 F		250 mg/kg bw  <u>Repeated dose group:</u> 14 days pre-treatment with unlabelled material at 2 mg/kg bw/day, followed by single dose of labelled material (2 mg/kg bw)  <u>Control group:</u> 14 days pre-treatment with corn oil at 5 ml/kg/day, followed by single dose of labelled material (2 mg/kg bw)	0.2-0.4 % C-14 tissue residues after 7 days, widely distributed, highest residues in blood cells  Excreted at > 5 % in urine plus faeces within 2 days after dosing of radioactive compound in any group: 3-hydroxy-cyclohexane-dicarboximide, 1-sulfo-cyclohexane-dicarboximide, N-(hydroxymethyl)-3-hydroxy-1-sulfo-cyclohexane-dicaboximide, unknown compounds: 30, 34 and 39  Excreted unmetabolised parent in urine plus faeces: 13-34 % in single dose groups, < 4 % in repeated dose groups	metabolites	et al., 1994, Xenobiotica 24(12): 1205-1214
Supplementary information  Non guideline, non-GLP  Oral and subcutaneous	Rat, Sprague-Dawley, 3-4 M + 3-4 F	C-14 [1R, trans]- and [1R, cis]-tetramethrin, respectively, each preparation suspended in 10% Tween 80	<u>Single dose:</u> 3.2 to 5.3 mg/kg bw	27-46 and 42-67 % radiocarbon absorption after oral administration of [1R, cis]- and [1R, trans]-isomer (urinary C-14 excretion);  > 90 % C-14 excretion within 7 d of oral or subcutaneous administration (cis/trans in urine: 27-49/42-74 %, faeces: 45-68/21-55 %, air: < 1 - 3 %);  2-3-fold slower initial metabolism and C-14 elimination after s.c. application of the cis-isomer as compared to the trans-isomer,  ~5-fold lower C-14 residues (day 7) for	C-14 label at phthalimide (alcohol) or chrysanthemic acid moiety	Sumitomo Report No: IM-10-0006  Summary of study published: Kaneko H, Ohkawa H, Miyamoto J, 1981, J. Pesticide Sci. 6 (4): 425-435

## CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method, guideline	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Results	Remarks	Reference
				acid than alcohol labelled tetramethrin isomers		
Supplementary information  Non guideline, non-GLP  <i>In vitro</i> (rat liver microsomes)	Rat, Sprague-Dawley, male, (no. not specified, for microsome preparation only)	[1RS, trans]-tetramethrin	1 mM (331 ppm),  C-14 [RS, trans]-tetramethrin,  1 hour incubation (37 °C)	68.0/58.5 % of acid/alcohol labelled substance degraded within 1 h by 7 mg protein eq. of rat liver microsomes in the presence of NADPH,  ~50 % NADPH-dependent oxidation,  ~50 % paraoxon-sensitive ester hydrolysis,  non-enzymatic decomposition of N-(hydroxymethyl)-tetrahydrophthalimide to tetrahydrophthalimide	C-14 [1RS, trans]-tetramethrin, labelled at alcohol or acid moiety:  Alcohol labelled: <sup>14</sup> C at the carbonyl group in the tetrahydrophthalimide moiety  Acid labelled: <sup>14</sup> C at the carboxyl position in the chrysanthemumic acid moiety	Suzuki T, Miyamoto J, 1974, Pesticide Biochem. Physiol. 4(1): 86-97
Supplementary information  Non guideline, non-GLP, oral and i.v.	Rat, Wistar, Male, (no. not specified)	C-14 [1RS, trans]-tetramethrin emulsion,	<u>Single dose:</u> 1.5 (i.v.) or 500 (oral) mg/kg bw	Oral radiocarbon absorption approx. 50 % (based on metabolites in urine);  ~90 and > 95 % radiocarbon excretion within 2 and 5 d after oral admin., respectively (urine: 42.3-71.4, faeces: 29-57.9, air: < 0.1 %);  sustained oral C-14 absorption with t <sub>1/2</sub> of ~ 2 h, rapid systemic degradation of parent compound with t <sub>1/2</sub> of ~ 10 min (i.v.), limiting the percentage of parent compound to < 1 % of C-14 from 1 h after oral administration	C-14 label at phthalimide moiety (alcohol label)	Miyamoto J et al., 1968, Agr. Biol. Chem. 32(5): 628-640 Sumitomo Report No: IM-80-0003
Supplementary information  Non guideline, non-GLP	Mice, male albino,  Rat, male albino (no. not	C-14 tetramethrin	Single dose:  30µl/20 g mouse  150 µl/200 g rat	Tetramethrin underwent Michael addition with thiols,  tetramethrin-gluthathione conjugate is formed in the	C-14 label at acid or alcohol moiety	Smith, I.H., Wood, E.J., & Casida, J.E., 1982, J. agri. food Chem. 30: 598-600 (not

Method, guideline	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Results	Remarks	Reference
i. p.	specified)			presence of mouse liver homogenate,  mercapturic acid or tetramethrin-gluthathion conjugates were not found in bile or urine of rats or mice,  biliary excretion of C-14 tetramethrin: 6% in 2 h and 51% in 24 h  50-66 % urinary excretion of C-14 tetramethrin within 24 h		submitted by applicant)
Dermal absorption study <i>in vitro</i> OECD 428	Heat separated human epidermis, <i>in vitro</i>	Tetramethrin, dissolved in ethanol	20 µL/cm <sup>2</sup> (200 µg/cm <sup>2</sup> in ethanol), 24 h	22.3 µg/cm <sup>2</sup> (11.2 % of dose) absorbed		Hadfield N, 2006, Central Toxicology Laboratory Report No. JV1900; Sumitomo
Dermal absorption study <i>in vitro</i> OECD 428	Heat separated human epidermis, <i>in vitro</i>	d-Tetramethrin, dissolved in ethanol	20 µL/cm <sup>2</sup> (200 µg/cm <sup>2</sup> in ethanol), 24 h	12.4 µg/cm <sup>2</sup> (6.2 % of dose) absorbed		Hadfield N, 2006, Central Toxicology Laboratory Report No. JV1901; Sumitomo

### 9.1 SHORT SUMMARY AND OVERALL RELEVANCE OF THE PROVIDED TOXICOKINETIC INFORMATION ON THE PROPOSED CLASSIFICATION(S)

A number of studies have been performed in the rat to address the toxicokinetics of tetramethrins using [1RS, trans]-, [1RS, cis]-, [1R, trans] and [1R, cis]-isomers labelled at their alcohol (phthalimide) or acid (chrysanthemetic acid) moiety. From the absence of detectable differences between the 1RS- and the 1R-isomers, it may be concluded that the general toxicokinetic behaviour of the 1R- and the 1S-isomers is similar. Minor differences relating to the relative amount of radiocarbon metabolites excreted via urine and the initial elimination velocity were observed for the trans- and cis-isomers (Kaneko et al., 1981; Sumitomo). Considering the low content of cis-isomers of 20 % and 3 % in technical products of tetramethrin and d-tetramethrin, respectively, these minor differences appear not to be of significant practical relevance.

Radiocarbon absorption after administration of acid- and alcohol-labelled C-14 tetramethrin may be estimated from the relative amount excreted in urine and amounted to 42-74 and 9-46 % of trans- and cis-isomers, respectively, depending on dose and pre-treatment. Considering the prevalence of the trans-isomers in tetramethrin technical products (see Table 3-1), the oral absorption is estimated to be 50 % of C-14 tetramethrin.

Radioactivity in blood and tissues suggests sustained absorption of orally administered tetramethrin isomers with an estimated absorption half-life of approx. 2 h. In the body tetramethrin is degraded very rapidly as indicated by a  $t_{\max}$  of 1 h for the parent compound compared to 8 h for radiocarbon. Analysis of blood levels of the parent compound and metabolites following i.v. administration support rapid degradation with a  $t_{1/2}$  of approx. 10 min. When administered orally, the percentage of undegraded a.s. was less than 1 % of total radiocarbon in blood as early as 1 hour following administration, although it remained unclear whether the responsible metabolic reactions occurred systemically or pre-systemically (Miyamoto et al., 1968).

The bioavailability of orally administered (d-)tetramethrin for the systemic circulation as undegraded substance presents a very rough estimate of 0.5 % to 50 %. Consequently, internal reference doses are extrapolated from inhalation rather than from oral studies.

Total excretion of orally or subcutaneously administered radiocarbon was generally  $\geq 90$  % after 2 days and  $\geq 95$  % after 1 week. Detectable amounts in exhaled air were reported for acid-labelled tetramethrin only: up to 3 % of the orally (but not subcutaneously) administered dose of cis- as well as trans-isomers were exhaled as  $\text{CO}_2$  (Kaneko et al., 1981; Sumitomo). Remaining radioactive tissue residues 5-7 days after dosing were generally low with 0.2-0.4 % and widely distributed. Highest residue concentrations were found in blood cells. Residues after administration of acid-labelled tetramethrins were approx. 5 times lower than for alcohol-labelled isomers (Kaneko et al., 1981).

Initial metabolism and elimination of the cis-isomers after subcutaneous administration was 2-3 times slower than for the trans-isomers (Kaneko et al., 1981; Sumitomo).

Main metabolic reactions include ester hydrolysis yielding chrysanthemic acid (or its corresponding oxidation products, see below) and N-(hydroxymethyl)-3,4,5,6-tetrahydrophthalimide (MTI). MTI decomposed, presumably non-enzymatically (Suzuki and Miyamoto, 1974), into 3,4,5,6-tetrahydrophthalimide (TPI) which was subject to further extensive metabolic reactions, including reduction of the 1,2-double bond, hydroxylation at position 2 and 3, and full or partial hydrolysis of the carboxydimide moiety. Chrysanthemic acid was metabolised primarily by oxidation of the isobutenyl group. Subsequent conjugation reactions were reported for various phase I metabolites. *In vitro* studies using rat liver microsomes further demonstrated that oxidation reactions were NADPH-dependent, whereas a paraoxon-sensitive enzyme apparently mediated ester hydrolysis.

As a result of the extensive metabolism of tetramethrin, a large percentage of the resulting products (~50 %) remained unknown, unidentified or not extracted. Major labelled species identified at  $> 5$  % in urine or faeces include the parent compound (5-30 % in faeces only), 3-hydroxy-cyclohexane-dicarboximide, 1-sulfo-cyclohexane-dicarboximide, N-(hydroxymethyl)-3-hydroxy-1-sulfo-cyclohexane-dicarboximide and three unknown metabolites. The sulfonic acid products were reported almost exclusively in faeces and result chemically from addition of sulfite to the 1,2-unsaturated bond of the tetrahydrophthalimide moiety.




Figure 1: Metabolic pathways of tetramethrin in mammals (adopted from Environmental Health Criteria 98, World Health Organisation, Geneva, 1990).

In summary, the studies indicate sustained absorption of tetramethrin isomers from the G.I. tract which is estimated to be 50 % at doses in the range of the relevant oral NOAELs. Moreover, a very rapid systemic and/or pre-systemic metabolic degradation by oxidation and ester hydrolysis has been shown which is slower for cis- than for trans-isomers. Hence, the resulting levels of undegraded a.s. in blood and nervous system remain very low after oral administration (below 1 % of total C-14 at 1 h). However, bioavailability and systemic levels of parental compound may be different with other routes of application (inhalative, dermal). Excretion occurs via faeces and urine. Sulfonated metabolites in faeces are suggested to be a product of (intra)intestinal metabolism. Radiocarbon excretion is almost completed after 2-3 days. After 1 week C-14 tissue residues decreased to 0.4 % or below. There is no information that would justify usage of other assessment factors for inter- and intraspecies variation in toxicokinetics than the default values of 4 and 3.2, respectively. Mechanistically, there is potential for interference of organophosphates (paraoxon) with tetramethrin toxicokinetics.

### Percutaneous absorption

Sumitomo provided a study on percutaneous absorption of technical products containing d-tetramethrin or tetramethrin. Substances were tested with 1 % solutions in ethanol on human heat-separated epidermis over 24 h, using a static diffusion cell set-up. Absorption into the receptor fluid (40 % acetone) was continuous over 24 h for d-tetramethrin but increased from 0.1 (0-8 h) to 0.3 (20-24 h)  $\mu\text{g}/\text{cm}^2/\text{h}$  for tetramethrin. As post-exposure separation of the upper stratum corneum from the remaining epidermis was not reliable due to deficiencies of the methodology (tissue samples disintegrated during tape stripping), both fractions were included in the calculation of overall skin absorption. For d-tetramethrin, a total of  $12.4 \pm 9.8 \mu\text{g}/\text{cm}^2$ , corresponding to  $6.2 \pm 4.9$  % of the applied dose, was absorbed into and through the skin.  $1.1 \pm 1.3$  and  $3.5 \pm 5.3$  % of the doses were found in tissue layers attributed to stratum corneum and remaining epidermis, respectively. Totals of absorbed and absorbable dose were higher for tetramethrin, with  $22.3 \pm 5.8 \mu\text{g}/\text{cm}^2$ , corresponding to  $11.2 \pm 2.9$  % of the applied dose, after 24 h. Apparent residues in the stratum corneum were higher than for d-tetramethrin with  $13.8 \mu\text{g}/\text{cm}^2$  (6.9 %), but lower in the remaining epidermis with  $3.5 \mu\text{g}/\text{cm}^2$  (1.8 %).

Applying the read-across concept established for d-tetramethrin and tetramethrin, and considering the limitations of the *in vitro* test system, an overall dermal absorption of 10 % is derived for tetramethrin and the technical product. Taking an estimated oral absorption of 50 % into account, the proposed value for dermal absorption (10 %) would also be supported by comparison of oral vs. dermal acute neurotoxicity of d-tetramethrin in the mouse (with adverse effects observed at 200-385 vs. 2500 mg/kg bw, see Evaluation of acute toxicity, Tables 25a and 26a).

### **Inhalative absorption**

No experimental data are available for the derivation of internal doses achieved by absorption of inhaled tetramethrin or d-tetramethrin aerosol from the lung. Therefore, physiologically-based default assumptions must be used. As outlined in the corresponding Technical Guidance Document, respirable fractions of the test aerosol were calculated from measured aerodynamic particle sizes and, given the molecular weight of 331 g/mol and a logP of 4.3-4.6, 100 % of the respirable fraction was assumed to be the absorbable dose (European Commission, EUR 20418 EN/1, 2003). These assumptions are supported by experimental data obtained for other substances with similar physicochemical properties, e.g. prochlorperazine (MW 374 g/mol, LogP 4.6) with > 80 % inhalative absorption from a fine aerosol (Avram et al., 2007).

## **10. EVALUATION OF HEALTH HAZARDS**

For assessment of human health hazards, read-across was performed between tetramethrin and d-tetramethrin.

### ***Justification for read-across between tetramethrin and d-tetramethrin***

Tetramethrins are esters of chrysanthemic acid with 3, 4, 5, 6-tetrahydrophthalimidomethyl alcohol and are classified as type I pyrethroids, which lack a cyano group within the alcohol moiety. Tetramethrin and d-tetramethrin are isomeric mixtures of [1R, cis], [1S, cis], [1R, trans] and [1S, trans]-tetramethrin, differing in the ratios of individual stereoisomers. The average isomeric composition of 5 batches of the technical products tetramethrin (Neo-Pynamin) and d-tetramethrin (Neo-Pynamin forte; both Sumitomo) as well as the range of isomeric composition derived from four batches of tetramethrin (Duracide A; Endura) are shown in the table below. [1R, trans] and



[1S, trans]-isomers prevailed in an analysis of the average batch composition of the corresponding technical product tetramethrin, while [1R, trans] is the main component (> 90 %) in d-tetramethrin.

Table 16: Average isomeric composition of technical products containing d-tetramethrin and tetramethrin (n=5; according to Fujita, Sumitomo Report No. SUP-0014, 2006; Endura: range derived from 4 batches)

Technical product	1R, cis (%)	1S, cis (%)	1R, trans (%)	1S, trans (%)
d-tetramethrin; Sumitomo	2.8	0.2	93.1	3.9
tetramethrin; Sumitomo	10.1	10.0	39.7	40.2
tetramethrin; Endura	9.1-10.3	7.6-10.3	40.0-46.9	35.5-40.4

The [1R, trans] isomer has been reported to be the isomer displaying the highest potential for activation of invertebrate axonal sodium channels (Lund and Narahashi, *NeuroTox* (1982) 3: 11-24).

All toxicological studies in mammals relate to isomeric mixtures. Thus, in the absence of comparative toxicological studies on the four isolated isomers, the toxicological potential of individual isomers in mammals has not been defined.

A review comparing the physico-chemical and biological properties of d-tetramethrin and tetramethrin was provided by the applicant Sumitomo (H. Roth, 2007/03/05, Safepharm Laboratories Ltd., Shardlow, UK). It was concluded that the two substances are essentially similar and “data on each of the compounds may be used to predict the toxicology and fate of the other”.

Indeed, a comparison of the submitted toxicological studies on d-tetramethrin and tetramethrin showed that, in cases in which both substances were investigated, the observed effects as well as NOAEL/LOAEL values were similar.

In rats, acute poisoning syndrome associated with type I pyrethroids typically is characterised by neurological effects such as aggressive sparring, whole body tremor and prostration (Verschoyle and Aldridge, 1980). Symptoms of neurotoxicity were observed for both tetramethrin and d-tetramethrin as a result of acute exposure and as acute effects during repeated inhalative exposure, including irregular respiration, bradypnoe and decreased spontaneous activity, whereas symptoms such as tremor, muscular fibrillation, urinary incontinence, hyperexcitability, ataxia, and limb paralysis were only observed in the acute inhalative toxicity study with d-tetramethrin.

After single oral administration, both tetramethrin and d-tetramethrin displayed only low acute toxicity in rats, with similar LOAELs for neurotoxicity (>2000 (Endura) 2500 mg/kg bw and 5000 mg/kg bw, resp.; both Sumitomo).

Regarding inhalative toxicity in rats, the LOAECs (converted to inhaled doses) for acute neurotoxic effects observed during exposure were within overlapping ranges for tetramethrin (13 mg/kg bw/d, 7 days, neonate mouse; 29.8 mg/kg bw/d, 90 d, rat) and d-tetramethrin (16.8 mg/kg bw, acute, rat; 11.2 mg/kg bw/d, 28 d, rat).

In repeated-dose rat studies, relevant adverse effects involved neurotoxic effects (proprioceptive alterations), liver changes (weight increase, enlargement) supported by alterations of parameters in haematology and clinical chemistry (e.g. cholesterol, liver enzymes), although acute neurotoxic effects were also observed during daily exposure in inhalation toxicity studies. Only few studies with d-tetramethrin are available on subchronic/chronic toxicity (one 3/6 month rat oral toxicity study and a 2-generation reproduction toxicity study). Notably, no carcinogenicity study has been provided for d-tetramethrin. However, on the basis of overlapping NOAEL-LOAEL intervals for



tetramethrin and d-tetramethrin in medium-term (subacute/subchronic) and long-term studies, it can be concluded that tetramethrin and d-tetramethrin are of similar toxicological potency regarding liver changes and haematological findings.

For both tetramethrin and d-tetramethrin, no evidence for toxic effects on foetuses below doses causing maternal toxicity was provided in developmental toxicity studies.

Finally, neither tetramethrin nor d-tetramethrin met the criteria for classification as skin or eye irritating or as genotoxic. Both substances did not lead to sensitisation in guinea pig (Buehler) tests.

In summary, the comparison of the available data appears to scientifically justify risk assessment involving i) read-across of acute toxicity data, ii) unidirectional read-across from tetramethrin to d-tetramethrin for chronic toxicity, and iii) derivation of mutual NOAEL/LOAEL for all other types of toxicity.

## Acute toxicity

### 10.1 ACUTE TOXICITY - ORAL ROUTE

Table 17: Summary table of animal studies on acute oral toxicity

Method, guideline, deviation(s) if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Value LD <sub>50</sub>	Reference
OECD 423 Oral, gavage	Rat, Wistar  3 F+3 M	Tetramethrin Vehicle: corn oil	2000 mg/kg bw	LD <sub>50</sub> : > 2000 mg/kg bw  No deaths and no toxic signs observed	Venugopala R K, 2002. Rallis Ltd, Endura Study No. 3335/01
Sim. to OECD 420 Oral, gavage	Rat, Sprague-Dawley, 5 M + 5 F	Tetramethrin Vehicle: corn oil	0-2500-5000 mg/kg bw	LD <sub>50</sub> : > 5000 mg/kg bw  No deaths  ≥ 2500 mg/kg bw: Decrease in spontaneous activity, urinary incontinence, excretion of oily substance	Kawasaki H, 1990, Sumitomo Report No. IT- 00-0224
Sim. to OECD 420 Oral, gavage	Rat, Crj:CD (SD), 5 M + 5 F	d-Tetramethrin Vehicle: corn oil	0-2500-5000 mg/kg bw	LD <sub>50</sub> : > 5000 mg/kg bw;  5000 mg/kg bw: tremor, urinary incontinence, ataxic gait; 1 single death in the high dose group	Misaki Y, 1999, Sumitomo Report No. SVT- 0008
Pre-guideline, sim. to OECD	Mouse, ddY,	d-Tetramethrin	0-100-150-200- 285-385-500-	LD <sub>50</sub> : 1060/1040 mg/kg bw (M/F);	Kohda H, Misaki Y, Suzuki T,

Method, guideline, deviation(s) if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Value LD <sub>50</sub>	Reference
401, non-GLP Oral, gavage	10 M + 10 F	Vehicle: corn oil	650-845-1000- 1300-1700 mg/kg bw	200 + 285 mg/kg bw/d: Slight decrease in spontaneous activity; ≥ 385 mg/kg bw: Hyperexcitaion, muscular fibrillation, ataxic gait, irregular respiration; ≥ 845 mg/kg bw: Whole body ataxia, weak respiration, salivation, mortalities	1980, Sumitomo Report No. IT- 00-0086

Table 18: Summary table of human data on acute oral toxicity

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data		No cases of poisoning of humans with d-tetramethrin or tetramethrin have been reported		

### 10.1.1 Short summary and overall relevance of the provided information on acute oral toxicity

In rats, tetramethrin displayed very low acute toxicity after oral administration, with no deaths at the limit dose of 2000 mg/kg bw (LD<sub>50</sub> > 2000 mg/kg bw, Venugopala, 2002; Endura). This study was performed in accordance with OECD guideline 423 and confirmed the results of earlier studies in which d-tetramethrin and tetramethrin were tested up to oral doses of 5000 mg/kg bw and no mortality was observed (LD<sub>50</sub> > 5000 mg/kg bw, Misaki, 1999; Kawasaki, 1990; both Sumitomo). Toxic effects were reversible within 3 days and comprised neurological symptoms (tremor, decrease in spontaneous activity, urinary incontinence) and excretion of oily substance.

In mice, the LD<sub>50</sub> value for d-tetramethrin was 1040 mg/kg bw after oral administration and signs of toxicity were seen at 200 mg/kg bw and above. No cases of poisoning of humans with d-tetramethrin or tetramethrin have been reported.

### 10.1.2 Comparison with the CLP criteria

Table 19: Results of acute oral toxicity studies in comparison to the CLP criteria

Toxicological results	CLP criteria
Oral LD <sub>50</sub> rat: > 2000 mg/kg bw (tetramethrin, d-tetramethrin)	Cat 4 (H302): 300 < LD <sub>50</sub> ≤ 2000 mg/kg bw (acute oral)
Oral LD <sub>50</sub> mouse: 1040 mg/kg bw (F) (d-tetramethrin)	Cat 3 (H301): 50 < LD <sub>50</sub> ≤ 300 mg/kg bw (acute oral)
	Cat 2 (H300): 5 < LD <sub>50</sub> ≤ 50 mg/kg bw (acute oral)
	Cat. 1 (H300): LD <sub>50</sub> ≤ 5 mg/kg bw (acute oral)

### 10.1.3 Conclusion on classification and labelling for acute oral toxicity

Based on the available acute oral rat studies, tetramethrin / d-tetramethrin do not meet the criteria according to the CLP regulation for classification for acute oral toxicity. Although the oral LD<sub>50</sub> value for mice was in the range for classification as "Acute Tox. 4", no classification for acute oral toxicity is proposed, taking into account that the single mouse study is a pre-guideline study and that results in mice are usually considered less relevant for classification. However, the final decision is within the remit of RAC/ECHA.

## 10.2 ACUTE TOXICITY - DERMAL ROUTE

Table 20: Summary table of animal studies on acute dermal toxicity

Method, guideline, deviation(s) if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Value LD <sub>50</sub>	Reference
OECD 402, dermal	Rat, Wistar  5 M + 5 F	Tetramethrin  Vehicle: corn oil	2000 mg/kg bw;  24 h	LD <sub>50</sub> : > 2000 mg/kg bw  No toxic signs observed	Venugopala RK, 2002, Rallis Ltd., Endura Study No.° 3336/01, 14
Pre-guideline, sim. to OECD 402, non-GLP, dermal	Rat, Sprague-Dawley, 10 M + 10 F	d-Tetramethrin  Vehicle: corn oil	0-2500-5000 mg/kg bw;  24 h	LD <sub>50</sub> : > 5000 mg/kg bw  No deaths and no toxic signs observed	Kohda H, Misaki Y, Suzuki T, 1980, Sumitomo Report No. IT-00-0086
Pre-guideline, sim. to OECD 402, non-GLP, dermal, occlusive	Rabbit, New Zealand White, 5 M + 5 F	Tetramethrin  Vehicle: corn oil	0-2000 mg/kg bw;  24 h	LD <sub>50</sub> : > 2000 mg/kg bw  No deaths and no toxic signs observed	Suzuki T et al., 1987, Sumitomo Report No. IT-70-0207

Pre-guideline, sim. to OECD 402, non-GLP, dermal	Mouse, ddY, 10 M + 10 F	d-Tetramethrin Vehicle: corn oil	0-1000-2500-5000 mg/kg bw; 24 h	LD <sub>50</sub> : > 5000 mg/kg bw; ≥ 2500 mg/kg bw: muscular fibrillation, irregular respiration	Kohda H, Misaki Y, Suzuki T, 1980, Sumitomo Report No. IT-00-0086
--	-------------------------	-------------------------------------	------------------------------------	--	---

Table 21: Summary table of human data on acute dermal toxicity

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data		No cases of poisoning of humans with d-tetramethrin or tetramethrin have been reported		

### 10.2.1 Short summary and overall relevance of the provided information on acute dermal toxicity

Dermal exposure of rats or rabbits to tetramethrin yielded no toxic effects at the limit dose of 2000 mg/kg bw (LD<sub>50</sub> > 2000 mg/kg bw, Venugopala, 2002; Suzuki et al. 1987). Dermal exposure of rats to d-tetramethrin yielded no relevant toxic effects up to 5000 mg/kg bw in rats (Kohda et al., 1980). By contrast, mice displayed toxicity after dermal exposure to 2500 mg/kg bw d-tetramethrin (muscular fibrillation, irregular respiration), but the LD<sub>50</sub> was estimated to be above 5000 mg/kg bw (Kohda et al. 1980).

### 10.2.2 Comparison with the CLP criteria

Table 22: Results of acute dermal toxicity studies in comparison to the CLP criteria

Toxicological results	CLP criteria
Dermal LD <sub>50</sub> rat: > 2000 mg/kg bw (tetramethrin)	Cat 4 (H312): 1000 < LD <sub>50</sub> ≤ 2000 mg/kg bw (acute dermal)
Dermal LD <sub>50</sub> rat: > 5000 mg/kg bw (d-tetramethrin)	Cat 3 (H311): 200 < LD <sub>50</sub> ≤ 1000 mg/kg bw (acute dermal)
Dermal LD <sub>50</sub> rabbit: > 2000 mg/kg bw (tetramethrin)	Cat 2 (H310): 50 < LD <sub>50</sub> ≤ 200 mg/kg bw (acute dermal)
Dermal LD <sub>50</sub> mouse: > 5000 mg/kg bw (d-tetramethrin)	Cat. 1 (H310): LD <sub>50</sub> ≤ 50 mg/kg bw (acute dermal)

### 10.2.3 Conclusion on classification and labelling for acute dermal toxicity

Based on the submitted acute dermal toxicity studies, tetramethrin and d-tetramethrin do not meet the criteria for classification for acute dermal toxicity.

## 10.3 ACUTE TOXICITY - INHALATION ROUTE

Table 23: Summary table of animal studies on acute inhalation toxicity

Method, guideline, deviation(s) if any	Species, strain, sex, no/group	Test substance, reference to table 5, form and particle size (MMAD)	Dose levels, duration of exposure	Value LC <sub>50</sub>	Reference
<p>OECD 403</p> <p>Acute inhalation toxicity study with Tetramethrin in Wistar rats.</p> <p>Head and nose exposure</p> <p>Technical deficiencies:</p> <p>Volume of air chamber 500 L, suggesting long time to achieve steady-state of concentration; no indication of whether concentration measurements performed in breathing zone; dense aerosol accumulation in chamber of G2 groups</p>	<p>Rat, Wistar</p> <p>Pre-study (G1): 2 M + 2 F</p> <p>Main study (G2): 5 M + 5 F</p>	<p>Tetramethrin aerosol in cyclohexanone (50% w/v);</p> <p>mean aerosol particle size:</p> <p>G1: 0.67 ± 0.26 µm;</p> <p>G2: 0.68 ± 0.26 µm</p>	<p>0-5.63 ± 0.86 mg/L;</p> <p>4 h</p>	<p>LC<sub>50</sub>: &gt; 5.63 mg/L</p> <p>Slight lacrimation and nasal discharge on day 1 (all rats in G2), normal from day 2 onwards), otherwise no toxic signs observed</p> <p>Absence of signs of acute neurotoxicity in comparison to other studies suggests that concentration of 5.63 mg/L may not have been achieved in the breathing zone, study not appropriate for classification purposes</p>	<p>Venugopala RK, 2006, Toxicology Department, Advinus Therapeutics Private Limited., Endura Study N° 4414/05</p>
<p>Pre-guideline, non-GLP.</p> <p>Acute inhalation, whole body</p> <p>3 h exposure instead of 4 h;</p> <p>Dose-finding study for a subacute (28 day) study</p>	<p>Rat, Sprague-Dawley, 10 M + 10 F</p>	<p>d-Tetramethrin dissolved in: deodorised kerosene</p> <p>mist, particle size 1.23-1.51 µm</p>	<p>0-0.026-0.131-0.243-0.595-1.18 mg/L</p> <p>(approx. 0-3.3-16.8-32.3-76.3-151 mg/kg bw);</p> <p>3 h</p>	<p>LC<sub>50</sub>: &gt; 1.18 mg/L (151 mg/kg bw);</p> <p>≥ 0.131 mg/L: Muscular fibrillation, urinary incontinence, limb paralysis, bradypnoe, irregular respiration respiration (no. of affected animals and severity of findings not reported; toxic signs began to appear 15 to 30</p>	<p>Suzuki T, Kohda H, Misaki Y, Okuno Y, Koyama Y, Miyamoto J, 1981, Sumitomo Report No. IT-10-0144</p>

Method, guideline, deviation(s) if any	Species, strain, sex, no/group	Test substance, reference to table 5, form and particle size (MMAD)	Dose levels, duration of exposure	Value LC <sub>50</sub>	Reference
				min, after initiation of exposure and disappeared 1 to 2 hours after exposure)  at 1.18 mg/L: 1/10 females died	
28 day inhalation, whole body  3 h / day exposure	Rat, Sprague-Dawley  10 M + 10 F	d-Tetra-methrin  dissolved in: deodorised kerosene  mist, particle size 1.23-1.51 µm	0-0.026-0.049-0.087 mg/L  3 h / day exposure for 7 days aweek	≥ 0.087 mg/L:  Slight bradypnea, irregular respiration, salivation directly after exposure (no. of affected animals and severity of findings not reported), no cumulative effect  Increase in leucocyte and decrease in eosinophils count at 0.087 mg/l  NOAEL: 0.049 mg/L	Suzuki T, Kohda H, Misaki Y, Okuno Y, Koyama Y, Miyamoto J, 1981, Sumitomo Report No. IT-10-0144

Table 24: Summary table of human data on acute inhalation toxicity

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data		No cases of poisoning of humans with d-tetramethrin or tetramethrin have been reported.		

### 10.3.1 Short summary and overall relevance of the provided information on acute inhalation toxicity

An acute rat inhalation toxicity study with d-tetramethrin and exposure time of 3 hours revealed moderate toxicity, with systemic effects occurring in the groups at 0.131 mg/L (corresponding to approx. 16.8 mg/kg bw) and above: decreases of spontaneous activity, salivation, hyperexcitability, hyperpnoea, irregular respiration, urinary incontinence, muscular fibrillation, ataxia, limb paralysis and other toxic signs

were observed (Suzuki et al. 1981) (no. of affected animals and severity of findings not reported). The NOAEC of 0.026 mg/L was estimated to correspond to 3.3 mg/kg bw, the conversion of inhaled concentrations into inhaled doses being based on default assumptions regarding body weight, inhalation volume and 100 % availability. The LC50 could only be estimated as being > 1.18 mg/L (> 151 mg/kg bw), with 1/20 mortality at this concentration. The RMS notes that the duration of exposure in the study was shorter than recommended according to OECD 403 (only 3 hours instead of 4) and that more severe effects might be anticipated for the standard exposure of 4 hours. Higher concentrations for refining the LC50 were not tested in this study by Suzuki et al. 1981, but appear technically feasible, since an almost linear relationship was recognised between substance concentration injection and aerial concentration. By contrast, the results of the study by Suzuki et al. 1981 were not confirmed in an acute rat inhalation toxicity study using tetramethrin and an exposure time of 4 hours in accordance with OECD guideline 403. No mortality was observed up to the highest concentration tested. Hence, the LC50 was estimated to be > 5.63 mg/L. Slight lacrimation and nasal discharge occurred on day 1, which was not observed from day 2 post exposure onwards. No other clinical signs were observed (Venugopala 2006; Endura). Due to dense aerosol accumulation inside the chamber, observation of the animals during exposure was not possible. Thus, potential clinical signs that may occur only during but not after exposure could not have been recorded. In addition, it is not clear which aerosol concentrations were actually achieved in the breathing zone. In contrast to other inhalation studies performed with d-tetramethrin (the acute rat study by Suzuki et al. 1981 and the 28-day study by Suzuki et al. 1981) or tetramethrin (90-day study performed by Kawaguchi 1991, see Table 38d), no clinical signs of acute neurotoxicity were reported in the study by Venugopala. In conclusion, the acute rat inhalation study by Venugopala is regarded by the RMS as being less relevant to considerations concerning classification and labelling, due to deficiencies in conduction of the study and the total absence of expected acute neurotoxicity.

### 10.3.2 Comparison with the CLP criteria

Table 25: Results of acute inhalation toxicity studies in comparison to the CLP criteria

Toxicological results	CLP criteria
Inhalative LD <sub>50</sub> rat: > 1.18 mg/L, (d-tetramethrin mist; 1/20 mortality at 1.18 mg/L, exposure for only 3 h)	Cat 4 (H332): 10 < LD <sub>50</sub> ≤ 20 mg/L (acute inhalation, vapours) 1 < LD <sub>50</sub> ≤ 5 mg/L (dusts and mists)  Cat 3 (H331): 2.0 < LD <sub>50</sub> ≤ 10 mg/L (vapours) 0.5 < LD <sub>50</sub> ≤ 1 mg/L (dusts and mists)  Cat 2 (H330): 0.5 < LD <sub>50</sub> ≤ 2 mg/L (vapours) 0.05 < LD <sub>50</sub> ≤ 0.5 mg/L (dusts and mists)  Cat. 1 (H330): LD <sub>50</sub> ≤ 0.5 mg/L (vapours) LD <sub>50</sub> ≤ 0.05 mg/L (dusts and mists)  Exposure generally for 4 h

### 10.3.3 Conclusion on classification and labelling for acute inhalation toxicity

From the data presented in the study by Suzuki et al. 1981 for d-tetramethrin, indicating an LD50 above 1.18 mg/L (observed mortality at 1.18 mg/L), and considering that rats were only exposed for 3 h, it cannot be ruled out that the LC50 is  $\leq 5$  mg/L. Thus, classification of d-tetramethrin and tetramethrin as “Acute Tox. 4; H332” is proposed.

### 10.4 SKIN CORROSION/IRRITATION

Table 26: Summary table of animal studies on skin corrosion/irritation

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results and time point of onset -Observations -Mean scores/animal -Reversibility	Reference
OECD 404	Rabbit, New Zealand White, 3 M	Tetramethrin, as paste with corn oil	500 mg, 4 h	1, 24, 48, 72 h: No erythema, oedema or any other effects on skin observed; Reversibility: N/A Not irritating	Mohan Kumar, 2002, Rallis Ltd., Endura Study N° 3337/01
OECD 404	Rabbit, New Zealand White, 3 M + 3 F	Tetramethrin, Moistened with corn oil	500 mg, 4h	0.5, 24, 48, 72 h: No erythema, oedema or any other effects on skin observed; Reversibility: N/A Not irritating	Nakanishi T, 1990, Sumitomo Report No. IT-00-0217
Pre-guideline, sim. to OECD 404, non-GLP	Rabbit, Albino, 6 M	d-Tetramethrin	0.5 ml of formulation 24 h, Intact and abraded skin, occlusive	24, 48, 72 h and 1 week: No erythema, oedema or any other effects on skin observed; Reversibility: N/A Not irritating	Hara S, Suzuki T, 1980, Sumitomo Report No. IT-00-0073

Table 27: Summary table of human data on skin corrosion/irritation

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

#### 10.4.1 Short summary and overall relevance of the provided information on skin corrosion/irritation

Studies on skin irritation revealed that neither d-tetramethrin nor tetramethrin is irritating to the skin of rabbits.



### 10.4.2 Comparison with the CLP criteria

Table 28: Results of skin irritation studies in comparison to the CLP criteria

Toxicological results	CLP criteria
No erythema, oedema or any other skin effects observed	Irritating to skin (Category 2, H315): at least in 2/3 tested animal a positive response of: Mean value of $\geq 2.3 - \leq 4.0$ for erythema/eschar or for oedema

### 10.4.3 Conclusion on classification and labelling for skin corrosion/irritation

No classification/labelling for skin irritation is proposed for tetramethrin or d-tetramethrin

## 10.5 SERIOUS EYE DAMAGE/EYE IRRITATION

Table 29: Summary table of animal studies on serious eye damage/eye irritation

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results and time point of onset -Observations -Mean scores/animal -Reversibility	Reference
OECD 405	Rabbit, New Zealand white, 3 M	Tetramethrin, mixed in corn oil	53 mg (equivalent to 0.1 ml, test substance mixed in corn oil)  24 h	1, 24, 48, 72 h: No ocular lesions observed; Reversibility: N/A Not irritating	Mohan Kumar SB, 2002, Rallis Ltd., Endura Study N° 3338/01
OECD 405	Rabbit, New Zealand White, 3 M + 3 F	Tetramethrin	100 mg  Exposure period: N/A	Observation times (times after application): 1, 24, 48, 72 h; <u>Corneal opacity</u> : 2/6 animals with grade 1 at 24 h; <u>Iris</u> : No signs of irritation observed; <u>Conjunctival redness</u> : 6/6 animals with grade 1 at 1 h; 3/6 animals with grade 1 at 24 h; <u>Conjunctival chemosis</u> : 5/6 animals, grade 1 at 1 h; Reversibility of all effects: Yes, within 48 h of application Not irritating (below classification threshold)	Nakanishi T, 1990, Sumitomo Report No. IT-00-0217
Pre-guideline, non GLP	Rabbit, Albino, 6 M (eyes not washed after applicatio	d-Tetramethrin	0.1 ml instilled  Exposure period: Eyes of 6 animals	Observation times at 1, 24, 48 h and at 1 week <u>Cornea</u> : No signs of irritation observed <u>Iris</u> : No signs of irritation observed <u>Conjunctiva</u> : slight hyperaemia and/or	Hara S, Suzuki T, 1980, Sumitomo Report No. IT-00-0073

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results -Observations and time point of onset -Mean scores/animal -Reversibility	Reference
	n)  3M (eyes washed application)		unwashed after application;  Eyes of 3 animals washed 30 seconds after application	chemosis 1 h (grade 1) after application (unwashed group); slight hyperaemia (grade 1) 1 h after application in the washed group  Reversibility: Yes, within 48 h  Not irritating (below classification threshold)	

Table 30: Summary table of human data on serious eye damage/eye irritation

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

### 10.5.1 Short summary and overall relevance of the provided information on serious eye damage/eye irritation

When applied to rabbit eyes, tetramethrin displayed no eye irritation (Mohan Kumar, 2002) or minimal eye irritating potential (Naganishi, 1990). Similarly, in a pre-guideline study, d-tetramethrin elicited only minimal signs of irritation (Hara and Suzuki, 1980). Any slight effects observed in the rabbit studies (on cornea and conjunctiva for tetramethrin, or on conjunctiva for d-tetramethrin) were reversible by 48 hours after application.

### 10.5.2 Comparison with the CLP criteria

Table 31: Results of eye irritation studies in comparison to the CLP criteria

Toxicological results	CLP criteria
Tetramethrin: Study 1: Not irritating (Mohan Kumar, 2002); Study 2: 2/6 animals positive for corneal opacity, grade 1, at 24 h and 6/6 animals positive for conjunctival redness and/or chemosis, grade 1, at 1 h (Nakanishi, 1990)	Irritating to eyes (Category 2, H319):  at least in 2/3 tested animal a positive response of: corneal opacity: $\geq 1$ and/or iritis: $\geq 1$ and/or conjunctival redness: $\geq 2$ and/or conjunctival oedema (chemosis): $\geq 2$
d-Tetramethrin: Slight hyperaemia and/or chemosis, grade 1, 1 h after application	

### 10.5.3 Conclusion on classification and labelling for serious eye damage/eye irritation

Based on the available data, classification of tetramethrin and d-tetramethrin for serious eye damage/eye irritation is not proposed.

### 10.6 RESPIRATORY SENSITISATION

Table 32: Summary table of animal studies on respiratory sensitisation

Method, guideline, deviations if any	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels, duration of exposure	Results	Reference
No studies available					

Table 33: Summary table of human data on respiratory sensitisation

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
Factory workers' surveillance data	Tetramethrin	Regular medical examination, blood, hepatic, renal and urine analysis, spirometry, biological monitoring, audiometry, ergovision  65 workers	No findings attributable to exposure	Dr. Savron L, 2006, Endura S.p.A. - Ravenna Plant Medical Data. Safety, Environment and quality department, 11 April 2006
Factory worker examination review		Regular medical check-up (bw, visual and auditory acuity, chest x-ray, blood pressure, urinalysis, serum biochemistry), 7 workers exposed to pyrethroids incl. tetramethrin dermally and by inhalation during packaging	No findings attributable to exposure	Shono F, 2005, Sumitomo Report No. SVT-0009
Case report	Tetramethrin	Single case of professional (M) developing asthma after 6 years of work as exterminator  <u>Inhalation challenges:</u>  1 <sup>st</sup> challenge: formulation containing tetramethrin + organophosphate;  2 <sup>nd</sup> challenge (5 mo. after first):  Tetramethrin powder diluted 1/10 in lactose powder	Skin prick testing to tetramethrin negative  Challenge-provoked reactions: Reduced respiratory function (reduced forced expiratory volume in 1 second; asthma)  Patient treated with beta-agonist when required	Vandenplas O, Delwiche J P, Auverdin J, Caroyer U M, Canghai F B, 2000, Allergy 55(4): 417-418

### 10.6.1 Short summary and overall relevance of the provided information on respiratory sensitisation

Medical surveillance of workers exposed to pyrethroids, including d-tetramethrin and tetramethrin, did not provide evidence for significant adverse effects. A single case study (Vandenplas et al., 2000) indicates that individual asthmatic reactions against tetramethrin may be possible. However, with regard to tetramethrin, epidemiological studies on exposed populations are not available. No animal studies involving respiratory sensitisation are available.

### 10.6.2 Comparison with the CLP criteria

Table 34: Information on respiratory sensitisation in comparison to the CLP criteria

Toxicological results	CLP criteria
One case study on single individual, showing reaction after challenge to tetramethrin-containing formulation or tetramethrin powder (reduced respiratory function, asthma)	Cat 1: Based on evidence in humans that the substance can lead to specific respiratory hypersensitivity and/or Positive results from appropriate animal test

### 10.6.3 Conclusion on classification and labelling for respiratory sensitisation

Due to paucity of data, no classification of tetramethrin or d-tetramethrin for respiratory sensitisation is proposed.

## 10.7 SKIN SENSITISATION

Table 35: Summary table of animal studies on skin sensitisation

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
OECD 406, Buehler	Guinea pig, Albino, NIH (Dunkin Hartley), Vehicle: 5 M+5 F Pos.Co: 5 M+5 F Treatment group: 10 M+10 F	Tetramethrin,	0.5 g, as paste in deionised water  6 h	No animals sensitised to tetramethrin (0/20); 2-MBT control: 08/10  Not sensitising	Prakash P.J., (2006). Toxicology Department, Advinus Therapeutics Private Limited, Endura Study N° 4415/05
Buehler, pre-guideline, non-GLP	Guinea pig, Hartley, 10 M	d-Tetramethrin	50% (0.5 ml of test material diluted in acetone); Time of removal of test substance not specified	No animals sensitised to d-tetramethrin (0/10); DNCB control: 10/10  M+K test not possible due to strong irritation reaction after intradermal application	Hara S, Suzuki T, 1980, Sumitomo Report No. IT-00-0082

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
				Not sensitising	
Buehler, sim. to OECD 406 (deficiencies)	Guinea pig, Hartley, 10 M positive control: 3 M	Tetramethrin	500 mg, applied undiluted, time of removal of test substance not specified	No animals sensitised to tetramethrin (0/10); positive control DNCB 3/3	Nakanishi T, 1990, Sumitomo Report No. IT-00-0218
Pre-guideline, non-GLP  Severe deviations from M + K protocol	Guinea pig, Hartley, 7 M  positive control: 5 M	Tetramethrin	1 % solution in corn oil, 10 intracutaneous injections over 23 d, challenge 14 d later	No animals sensitised to tetramethrin (0/7); Positive control DNCB 5/5  Severe deviations from M + K protocol  Not sensitising	Okuno Y et al., 1976, Sumitomo Report No. IT-60-0013  Reviewed in WHO, IPCS, 1990, Environmental Health Criteria 98, Tetramethrin

Table 36: Summary table of human data on skin sensitisation

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
Human patch test, modified Schwartz Peck method, pre-guideline, non-GLP	Tetramethrin	Human 23 M + 177 F, Semi-occlusive  Composition of test formulation and concentration of tetramethrin not clear	No cases of sensitisation observed  Not reliable, due to ethic and scientific deficiencies  No conclusion possible	Osborn R, 1966, Sumitomo Report No. IT-61-0008

### 10.7.1 Short summary and overall relevance of the provided information on skin sensitisation

No evidence for skin sensitisation potential of tetramethrin or d-tetramethrin was observed in a total of three Buehler tests (Prakash 2006; Hara and Suzuki, 1980; Nakanishi 1990). The test substances were applied diluted in acetone (d-tetramethrin, 0/10 animals sensitised), undiluted, apparently as solid substance (tetramethrin, 0/10 animals sensitised), and as paste in deionised water (tetramethrin, 0/20 animals sensitised). DNCB (2,4-dinitrochlorobenzene; 10/10 resp. 3/3 animals sensitised) and 2-MBT (2-mercaptobenzothiazole; 8/10 sensitised) were used as positive controls, respectively, and showed the expected skin reactions.

The application of (solid) tetramethrin to the skin in its undiluted form may have hampered skin penetration as a prerequisite for sensitisation. Nevertheless, lack of sensitising potential for tetramethrin is in line with the outcome of the test performed with d-tetramethrin. An additional study involving intracutaneous injections of tetramethrin diluted in corn oil yielded no indication for skin sensitisation (Okuno et al., 1976). However, low reliability is assigned to this study, since it shows severe deviations from the Magnusson-Kligman protocol in OECD 406 (i. e. no use of adjuvant, no justification for selection of concentration).

A study with 200 humans (Osborn 1966) was judged as not reliable. The substance tested was a tetramethrin formulation of unknown specification which was heated as well as autoclaved without analysing the actual tetramethrin content prior to application. Furthermore, use of the study is ethically not justifiable as no statement of informed consent of the study subjects (some of them pregnant women) was provided.

### 10.7.2 Comparison with the CLP criteria

Table 37: Results of skin sensitisation studies in comparison to the CLP criteria

Toxicological results	CLP criteria
No animals sensitised to tetramethrin (result of pre-guideline study with deviations from M+K protocol)	Guinea pig maximisation test Category 1A (H317): ≥ 30 % responding at ≤ 0.1 % intradermal induction dose or ≥ 60 % responding at > 0.1 % to ≤ 1 % intradermal induction dose  Category 1B (H317): ≥ 30 % to < 60 % responding at > 0,1 % to ≤ 1 % intradermal induction dose or ≥ 30 % responding at > 1 % intradermal induction dose
No animals sensitised to tetramethrin or d-tetramethrin (results of 3 Buehler tests)	Buehler assay Category 1A (H317): ≥ 15 % responding at ≤ 0.2 % topical induction dose or ≥ 60 % responding at > 0.2 % to ≤ 20 % topical induction dose  Category 1B (H317): ≥ 15 % to < 60 % responding at > 0.2 % to ≤ 20 % topical induction dose or ≥ 15 % responding at > 20 % topical induction dose

### 10.7.3 Conclusion on classification and labelling for skin sensitisation

No classification/labelling for skin sensitisation is proposed for tetramethrin or d-tetramethrin.

## 10.8 GERM CELL MUTAGENICITY

Table 38: Summary table of mutagenicity/genotoxicity tests in vitro

Method, guideline, deviation s if any	Test system (Organism, strain)	Test substance, reference to table 5	Concentrations tested (range)	Results		Remarks (information on cytotoxicity)	Reference
				+ S9	- S9		
Bacterial reverse mutation test  OECD 471	<i>Salmonella typhimurium</i> : TA98 TA100 TA102 TA1535 TA1537	Tetramethrin, dissolved in DMSO	Pre- test: 50-5000 µg/plate  Main assay: 313-5000 µg/plate	+/- TA100 Neg. for all other strains	-	TA100: two-fold increase in the number of revertant colonies (+S9), no dose-response relationship  No cytotoxicity up to the highest concentration tested  Outcome inconclusive	Scarcella O, 2004, Research Toxicology Centre S.p.A., Endura Study No. 9324
Ames test, pre-guideline, sim. to OECD 471, non-GLP	<i>S. typhimurium</i> : TA98 TA100 TA1535 TA1537 TA1538 <i>E. coli</i> : WP2 <u>uvr A</u>	d-Tetramethrin, dissolved in DMSO	10-5000 µg/plate	-	-	No cytotoxicity up to the highest dose tested  Negative for genotoxicity	Kishida F, Suzuki H, 1980, Sumitomo report No. IT-00-0101
<i>Bacillus subtilis</i> rec assay, pre-guideline, sim. to OECD 471, non-GLP	<i>B. subtilis</i> : M45 rec <sup>-</sup> H17 (wild type)	d-Tetramethrin, dissolved in DMSO	1-500 mg/ml  (10-5000 µg/plate)	-	-	No cytotoxicity up to the highest dose tested  Negative for genotoxicity	Kishida F, Suzuki H, 1980, Sumitomo report No. IT-00-0101
Bacterial reverse mutation test  Non-GLP, pre-guideline, sim. to OECD 471	<i>S. typhimurium</i> : TA1535, TA1537, TA97, TA98, TA100 <i>E. coli</i> : WP2 <u>uvrA</u>	Tetramethrin, dissolved in DMSO	100-5000 µg/plate	-	-	Precipitations of test substance on plates at ≥ 2000 µg/plate (without S9) and at 5000 µg/plate (with S9)  <u>Cytotoxicity</u> : TA97 (-S9): (> 500 µg/plate)  Negative for genotoxicity	Kogiso S, Yoshitake A, 1987, Sumitomo Report No. IT-70-0205

CLH REPORT FOR D-TRANS-TETRAMETHRIN

DNA-repair test Non-guideline, Non-GLP	<i>E. coli</i> : W3110/polA <sup>+</sup> and p3478/polA <sup>-</sup>	Tetramethrin, dissolved in DMSO	100- 33333 µg/ plate	-	-	No cytotoxicity up to the highest concentration tested  Negative for genotoxicity	McGregor DB, 1984, Inveresk Research International, Endura Report No. 2988
Chromosomal aberration test  OECD 473	Chinese hamster Ovary (CHO) cells	Tetramethrin, dissolved in DMSO	Experiment I: 0.781- 200 µg/ml, treatment time 3 h  Experiment II: 1.56-200 µg/ml, treatment time 21 h	+/-	+/-	<u>Genotoxicity:</u> Increase in CA at: <u>+S9:</u> ≥ 50 µg/mL  <u>-S9:</u> ≥100 µg/mL (Exp. I), ≥50 µg/mL (Exp. II)  <u>Cytotoxicity:</u> <u>+S9:</u> ≥ 100 µg/ml, <u>-S9</u> ≥ 50 µg/ml  Result: equivocal	Cilutti P, 2004, Research Toxicology Centre S.p.A., Endura Study No. 9325
Chromosomal aberration study, sim. to OECD 473	Chinese hamster ovary (CHO) cells	Tetramethrin, dissolved in ethanol	<u>-S9:</u> 15.1- 80.3 µg/ml  <u>+ S9:</u> 50.2- 151 µg/ml	+	-	<u>Cytotoxicity:</u> ≥ 20.1 µg/ml ( <u>-S9</u> , 20 h, ≥ 75.3 µg/ml ( <u>+S9</u> , 20 h) ≥ 101 µg/ml ( <u>+S9</u> , 30h)  Test considered positive under metabolic activation conditions	Murli H, 1989, Sumitomo Report No. IT- 91-0216
Gene mutation assay  OECD 476	Mouse lymphoma L5178Y TK <sup>±</sup> cells	Tetramethrin, dissolved in DMSO	Cytotoxicity assay: 1.56-400 µg/ml  Experiment I: 1.56-40 µg/ml, treatment for 3 h  Experiment II: 3.13-75 µg/ml, treatment for 3 or 24 h	+/-	-	<u>Genotoxicity:</u> <u>+ S9:</u> ≥ 50µg/ml (high concentrations only tested in Exp. II, 3 h); <u>-S9:</u> ≥ 50 µg/ml (high concentrations only tested in Exp. II, 24 h)  <u>Cytotoxicity:</u> <u>+S9:</u> 75 µg/ml (Exp. II, 3 h):  <u>-S9:</u> ≥ 50µg/ml (Exp. II, 24 h  Outcome: equivocal	Cinelli S, 2004, Research Toxicology Centre S.p.A., Endura Study No. 9326



CLH REPORT FOR D-TRANS-TETRAMETHRIN

Mutation test, sim. OECD 476, non-GLP	Chinese hamster V79 cells	Tetramethrin, dissolved in DMSO	- S9: 3.75-30 µg/ml + S9: 25-200 µg/ml	-	-	<u>Cytotoxicity:</u> ≥ 3.75 µg/ml (- S9), ≥ 100 µg/ml (+ S9)  Test result: negative	Kogiso S, 1989, Sumitomo Report No. IT-90-0214
UDS assay, sim. to OECD 482 non-GLP	Primary rat hepatocytes	Tetramethrin, dissolved in DMSO	0.2-100 µg/ml	N/A	-	<u>Cytotoxicity:</u> 30-33 % viability at 100 µg/ml  Test result: negative	Kogiso S, 1988, Sumitomo Report No. IT-80-0213

Table 39: Summary table of mutagenicity/genotoxicity tests in mammalian somatic or germ cells *in vivo*

Method, guideline, deviation s if any	Species, strain, sex, No/group	Test substance, reference to table 5	Route and frequency of application	Dose levels and sampling times	Results	Reference
Micro-nucleus test  OECD 474	Mice, Swiss albino NsdOla: MF1  5 M+5 F	Tetramethrin  200 mg/ml in vehicle (0.5% aqueous carboxymethyl cellulose with Tween 80, 1 mL/L)	Oral gavage,  twice at interval of 24 h	2000 mg/kg bw (Limit dose)  Sampling at 24 h after 2 <sup>nd</sup> treatment	Negative,  At all sampling times	Badarinath J.C., (2006). Toxicology Department, Advinus Therapeutics Private Limited, Endura Study No. 4416/05
Chromosomal aberration test, sim. to OECD 475, non-GLP	Mouse, ICR, 4-6 M	d-Tetramethrin, vehicle: corn oil	Single intraperitoneal dose	Sampling times: 6, 12, 24, 48 h  Doses: 150, 300 mg/kg bw (24 h), 600 mg/kg bw (all sampling times)	Negative,  at all doses and sampling times	Hara M, Suzuki H, 1981, Sumitomo Report No. IT-10-0102
Chromosomal aberration test, sim. to OECD 475	Mouse, IRC, 5 M + 5 F	Tetramethrin, Vehicle: corn oil	Single intraperitoneal dose	0-500-1000-2000 mg/kg bw  Sampling 6, 18, 30 h after injection	Negative,  at all doses and sampling times	Murli H, 1992, Sumitomo Report No. IT-21-0254

Table 40: Summary table of human data relevant for germ cell mutagenicity

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

### 10.8.1 Short summary and overall relevance of the provided information on germ cell mutagenicity

#### *Summary of results obtained in vitro:*

In general, negative results were obtained with d-tetramethrin and tetramethrin in bacterial mutation assays, irrespective of metabolic activation (Kishida and Suzuki 1980, Kogiso and Yoshitake 1987, both Sumitomo; McGregor 1984, Scarcella 2004; both Endura). Only one exception was reported in the study by Scarcella 2004, Endura, in that inconclusive results for the tester strain TA 100 were obtained with metabolic activation.

Regarding *in vitro* testing with mammalian cell systems, negative results for tetramethrin were supported by one *in vitro* test using Chinese hamster ovary (CHO) cells (Kagiso 1989; Sumitomo). None of the concentrations tested exhibited chromosomal aberrations, irrespective of metabolic activation. In addition, an UDS assay in primary rat hepatocytes (Kogiso 1988; Sumitomo) showed negative results without S9-mix (not tested with S9 mix).

By contrast, test results for tetramethrin were considered positive in a chromosomal aberrations assay in CHO cells after metabolic activation (Murli 1989; Sumitomo). Equivocal results were obtained in a further chromosomal aberration assay using CHO cells (Cilutti 2004; Endura) as well as in one mouse lymphoma assay using L5178Y TK<sup>±</sup> cells (Cinelli 2004; Endura). In the study performed by Cilutti, 2004, increases of chromosomal aberrations were observed at the highest concentrations tested with and without metabolic activation, and were accompanied by increasing cytotoxicity.

In the gene mutation assay by Cinelli, 2004, a statistically significant increase in mutant frequency was observed in the presence of S9-mix at the highest tetramethrin concentrations tested (50 and 75 µg/mL). At 50 µg/mL a 1.7-fold increase in mutant frequency was observed with a calculated survival rate of 81 %. At 75 µg/mL, mutant frequency increased to 1.9-fold of the control values but cytotoxicity was also high (57 % survival rate). As the results were observed with increasing cytotoxicity and the experiment was not repeated with the two highest concentrations, the results were regarded as equivocal. Increases of mutant frequency in the assay without metabolic activation at ≥ 50 µg/mL are not considered positive, as they were only observed at cytotoxic levels (relative survival of 25 and 0 % at 50 resp. 75 µg/mL).

In summary, there is some indication for a slight mutagenic potential of tetramethrin in *in vitro* test systems involving mammalian cells. However, as effects were observed at increasing cytotoxic dose levels, the results are regarded as equivocal.

#### *Summary of results obtained in vivo:*

Tetramethrin and d-tetramethrin showed negative results in all *in vivo* genotoxicity assays performed. Tetramethrin displayed negative results in the Mammalian Erythrocyte Micronucleus test following oral administration of 2x 2000 mg/kg bw (limit dose) at an interval of 24 hours (Badarinath 2006; Endura). The results were supported by two earlier chromosomal aberration studies with tetramethrin (Murli 1992; Sumitomo) and d-tetramethrin (Hara and Suzuki 1981; Sumitomo). In both assays no chromosomal aberrations were observed in mammalian bone marrow after single intraperitoneal application of dose levels between 150 and 600 mg/kg bw (d-tetramethrin) and between 500 and 2000 mg/kg bw (tetramethrin).

However, the negative results for tetramethrin in the micronucleus assay have to be interpreted with care, as no data for the oral bioavailability of tetramethrin were provided in the study. Moreover,

vehicle effects on oral absorption cannot be excluded. Carboxymethyl cellulose was used as the dosing vehicle which has been found to decrease the absorption rate of pyrethroids (e.g. deltamethrin) and may thereby reduce the toxic potential of pyrethroid compounds (Crofton et al. 1995; USEPA, 2007). Indeed, the limit dose used in the micronucleus assay (2x 2000 mg/kg) is above the oral LD<sub>50</sub> of d-tetramethrin for mice (LD<sub>50</sub> ~ 1000 mg/kg; Kohda et al. 1980; Sumitomo) and no significant toxic effects were observed (Badarinath 2006; Endura). By contrast, toxic signs were observed in all dosage groups of mice (500, 1000, 2000 mg/kg bw) after single i.p. application of tetramethrin in corn oil (Murli 1992, Sumitomo). The maximal tolerated dose after single i.p. application of d-tetramethrin was 600 mg/kg (7/10 deaths at 1200 mg/kg bw; Hara and Suzuki, 1981; Sumitomo). Consequently, the validity of the two *in vivo* studies carried out with i.p. application is regarded as being higher with respect to assessment of the genotoxic potential (negative) of tetramethrin, although effects of corn oil as vehicle for intraperitoneal application cannot be ruled out completely.

Overall, negative *in vivo* findings for chromosomal aberrations and micronucleus formations outweigh the positive test results of the above mentioned inconclusive or positive *in vitro* tests, although results of *in vivo* testing after oral application of tetramethrin can only be considered reliable if evidence is provided in the assay for the systemic bioavailability of tetramethrin after oral application.

### 10.8.2 Comparison with the CLP criteria

Table 41: Results of genotoxicity studies in comparison to the CLP criteria

Toxicological results	CLP criteria
<p><u>Testing <i>in vitro</i>:</u></p> <p>Bacterial mutation assays: Generally negative for tetramethrin and d-tetramethrin</p> <p>Tests involving mammalian cells: - Negative (mutation test with CHO cells, UDS test with rat hepatocytes, tetramethrin)</p> <p>- Positive with S9-mix (chromosomal aberration test, tetramethrin)</p> <p>- Equivocal (chromosomal aberration test, mouse lymphoma cell gene mutation test, tetramethrin)</p> <p><u>Testing <i>in vivo</i> (experiments in mammals):</u></p> <p>Negative (micronucleus test and chromosomal aberration test, tetramethrin; chromosomal aberration test, d-tetramethrin)</p>	<p>The classification in Category 1A is based on positive evidence from human epidemiological studies. Substances to be regarded as if they induce heritable mutations in the germ cells of humans.</p> <p>The classification in Category 1B is based on: — positive result(s) from <i>in vivo</i> heritable germ cell mutagenicity tests in mammals; or — positive result(s) from <i>in vivo</i> somatic cell mutagenicity tests in mammals, in combination with some evidence that the substance has potential to cause mutations to germ cells. It is possible to derive this supporting evidence from mutagenicity/genotoxicity tests in germ cells <i>in vivo</i>, or by demonstrating the ability of the substance or its metabolite(s) to interact with the genetic material of germ cells; or — positive results from tests showing mutagenic effects in the germ cells of humans, without demonstration of transmission to progeny; for example, an increase in the frequency of aneuploidy in sperm cells of exposed people.</p> <p>The classification in Category 2 is based on: — positive evidence obtained from experiments in mammals and/or in some cases from <i>in vitro</i> experiments, obtained from: — somatic cell mutagenicity tests <i>in vivo</i>, in mammals; or — other <i>in vivo</i> somatic cell genotoxicity tests which are supported by positive results from <i>in vitro</i> mutagenicity assays.</p> <p>Note: Substances which are positive in <i>in vitro</i> mammalian mutagenicity assays, and which also show chemical structure activity relationship to known germ cell mutagens, shall be considered for classification as Category 2 mutagens.</p>

### 10.8.3 Conclusion on classification and labelling for germ cell mutagenicity

No human data are available for tetramethrin or d-tetramethrin, hence a classification in category 1A is not possible.

On the basis of the negative results from *in vivo* animal studies (especially after i.p. administration), no further classification or labelling of tetramethrin and d-tetramethrin for genotoxicity is proposed.

## 10.9 CARCINOGENICITY

Table 42: Summary table of animal studies on carcinogenicity

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
Sim. to OECD 453, Oral, dietary, 104 weeks	Mouse, B6C3F1, 50 M + 50 F, satellite group: 40 M + 40 F	Tetramethrin	0-12-60-300-1500 ppm  (0-2.4/3.5-12/17-61/85-300/430 mg/kg bw/d (M/F)),  104 weeks	<u>No non-neoplastic effects</u>  <u>No neoplastic effects</u> , no increased tumour rate	Cox R, 1986, Sumitomo Report No. IT-61-0193
Sim. to OECD 453, non-GLP, Oral, dietary, 104 weeks	Rat, CRL:SD:C OBS, 50 M + 50 F, controls: 60 M + 60 F  F1A litters from a one-generation study (animals exposed also throughout gestation and weaning)	Tetramethrin	0-1000-3000-5000 ppm  (0-42/55-125/165-230/300 mg/kg bw/d (M/F))  104 weeks post weaning	<u>Non-neoplastic effects:</u> ≥ 3000 ppm: Cytoplasmic vacuolation of midzonal hepatocytes (M), testes enlarged (M), reduced bw and food consumption  <u>Neoplastic effects:</u> ≥ 3000 ppm: Increased incidence of interstitial adenomas of the testis  5000 ppm: Decrease in mammary tumour incidence (F) (see Table 33a-1)  Tetramethrin was administered with the diet to rats at levels of 0, 1000, 3000, and 5000 ppm for 104 weeks post-weaning. These treated rats were obtained as F <sub>1a</sub> weanlings from parental animals which had been treated with the compound at levels of 0, 1000, 3000, and 6000 ppm until sexual maturity prior to mating and which continued to receive the test compound during mating and throughout the gestation and nursing period	Rutter H A, 1974, Sumitomo Report No. IT-41-0024
Sim. to OECD 453,	Rat, CR CD and Long Evans	Tetramethrin	0-200-1000-5000 ppm	<u>Non-neoplastic effects:</u> 5000 ppm: Reduced bw gain, slight increase in incidence of testicular degeneration with	Pence, D H, 1981, Sumitomo Re-

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method, guideline, deviations if any	Species, strain, sex, no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
non-GLP, Oral, dietary, 104 weeks	Hooded, 50 M per strain/dose group		(CR CD: 0-7.5-35-180 mg/kg bw/d, Long Evans Hooded: 0-8-40-205 mg/kg bw/d)  Parental: 0-1000-3000-6000 ppm	associated hypospermatogenesis or aspermatogenesis; increased weight of testis with epididymis (LE), increased absolute and relative liver weight  <u>Neoplastic effects:</u> 5000 ppm: Statistically significant increased incidence of interstitial tumours of the testis (see Table 33a-2)  Rats were exposed to the test substance maternally from conception to weaning and via diet for 104 weeks thereafter  Follow-up on the results of the study by Rutter 1974	port No. IT-11-0097

Table 43: Supplementary Information: Incidences of testicular and mammary gland tumours (Rutter, 1974; Report No. IT-41-0024)

Parameter	Control		1000 ppm		3000 ppm		5000 ppm	
	M	F	M	F	M	F	M	F
Number of animals examined (after 2 years / interim sacrifice)	50/10	50/10	40/10	40/10	40/10	40/10	40/10	40/10
Mortality after 2 years	17/50	14/50	23/40	14/40	11/40	9/40	18/40	17/40
No. of animals with neoplasms (including interim sacrifice)	30/60	44/60	19/50	31/50	19/50	28/50	22/50	31/50
No. of animals with only benign tumours	26	41	17	26	17	23	19	24
No. of animals with malignant tumours	4	3	2	5	2	5	3	7
Interstitial cell adenoma in the testis	2/50	-	3/40	-	9/40	-	14/40	
Tumours in the mammary gland	1/50	31/50	0/40	26/40	0/40	21/40	1/40	12/40

No statistical analysis performed.

Table 44: Supplementary Information: Incidences of interstitial cell tumors in the testis (Pence, 1981, Report No. IT-11-0097)

Parameter	Control		200 ppm		1000 ppm		5000 ppm	
	CRCD	LE	CRCD	LE	CRCD	LE	CRCD	LE
Survival data at week 104	30/50	37/50	26/50	37/50	26/50	34/50	30/50	34/50
Interstitial cell tumour in the testis, unilateral	3	4	5	3	1	2	5	10
Interstitial cell tumour in the testis, bilateral	4	0	2	0	2	2	11	12

Total interstitial tumours of testes	7	4	7	3	3	4	16	22
--------------------------------------	---	---	---	---	---	---	----	----

No statistical significance reported

CRCD: Sprague-Dawley derived male rats

LE: Male rats of Long Evans hooded strain

Table 45: Summary table of human data on carcinogenicity

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

### 10.9.1 Short summary and overall relevance of the provided information on carcinogenicity

In a chronic (2 years) oral toxicity study on tetramethrin in mice, no adverse effects (neither neoplastic nor non-neoplastic) were observed up to the highest dose of 1500 ppm (300/430 mg/kg bw/d; Cox 1986; Sumitomo).

Two pre-guideline chronic toxicity studies were performed on tetramethrin with F<sub>1A</sub> rat weanlings. In the first study by Rutter (1974, Sumitomo), F<sub>1A</sub> litters from a one-generation study were further exposed to tetramethrin via the diet for 104 weeks. Since parental animals had already been orally exposed to tetramethrin, it is assumed that the F<sub>1A</sub> animals were exposed from conception onward (*in utero*, via milk), although the extent of exposure prior to weaning is not known. In the second study by Pence (1981, Sumitomo), which was conducted as a follow-up on the results of the Rutter study, only male animals of two rat strains (CR:CD and Long Evans Hooded) were exposed to tetramethrin from conception to two years after birth. The exposure estimates in both of the chronic studies refer to the two year dietary period.

The results of both chronic rat studies are essentially consistent: For the study by Rutter (1974), the LOAEL was set to 3000 ppm (125/165 mg/kg bw/d) for non-neoplastic effects (lower bw gain and histological abnormalities in liver) and for neoplastic effects in males. Testicular tumours were observed already at the lowest dose (1000 ppm), but achieved statistical significance at  $\geq 3000$  ppm ( $\geq 125$  mg/kg bw/d). Apart from the dose-dependent incidence, interstitial cell adenoma occurred bilaterally only at the medium and high doses. In addition, female rats displayed a significantly decreased incidence in benign mammary tumours (i. e. fibroadenomas) at the highest tetramethrin dose (5000 ppm).

In the study by Pence, a statistically significant increase in testicular interstitial cell tumours was confirmed in both rat strains for groups receiving the high dose (5000 ppm). Furthermore, a slight increase in the incidence of testicular degeneration with hypospermatogenesis and aspermatogenesis as compared to control rats as well as an increase in testis and liver weight were observed (group-wise incidences not reported in the original report). The NOAELs of 35 and 40 mg/kg bw/d derived for non-neoplastic and neoplastic effects in male rats are in agreement with the results of Rutter (1974). In contrast to the chronic toxicity studies performed with tetramethrin, no striking tumour-related effects were seen in a two-generation reproduction toxicity study conducted with d-tetramethrin in rats (see Pence, 1986). However, indirect parameters for hyperplasia or neoplastic effects such as testis enlargement were not specified in the two-generation study. Furthermore, oral

exposure of animals to the test substance did not exceed 38 weeks (F1 generation). In the chronic (2 year) toxicity study by Rutter (1974), the time at which testicular interstitial cell tumours were suspected (from external signs) did not pre-date week 83 of dietary exposure. The inability of the two-generation toxicity study by Pence (1986) to detect tumourigenic effects may therefore also be due to an insufficient duration in individual animal exposure.

Rats appear to be particularly susceptible towards induction of Leydig cell adenomas, and exhibit a relatively high spontaneous incidence as compared to humans (Clegg et al., 1997). However, the mode of action by which tetramethrin leads to Leydig cell adenoma has not been specified. From the presented data for tetramethrin and d-tetramethrin concerning genotoxicity testing, it appears unlikely that tetramethrin acts tumourigenic via a genotoxic mechanism. A common basis for several groups of other non-genotoxic Leydig cell adenoma-inducing agents comprises disturbance of endocrine homeostasis, e. g. perturbation of the hypothalamic-pituitary-gonadal axis. Mechanisms leading to disruption of the hypothalamic-pituitary-gonadal axis in rodents may include a decrease in circulating sex (steroid) hormones, anti-estrogenic activity or enhancement of central dopamine receptor-dependent signalling (Clegg et al., 1997; ECBI/61/03 (“thought starter” for developing an agreed position on the relevance of Leydig cell tumours in rats to humans)).

Limited evidence exists that one or more of these listed mechanisms may be involved in the mode of action by which tetramethrin enhances testicular interstitial cell tumour incidence in rats. Although the effect of tetramethrin on hepatic cytochrome P-450 expression has not been investigated, pyrethrins and individual synthetic pyrethroids have been identified as inducers in rat liver enzymes (Price et al., 2007; [Krechniak](#) and [Wrzesniowska](#), 1991). Possible enzyme induction by tetramethrin would be consistent with the observation of liver weight increase in repeated-dose studies, including the study by Pence (1981). Hepatic enzyme induction would be expected to lead to enhanced androgen catabolism, resulting in a decrease in circulating androgen levels. As a consequence, androgen-dependent negative feed-back at the hypothalamic/anterior pituitary level would be diminished, and thus enhanced secretion of GnRH and LH would occur, ultimately leading to stimulation of Leydig cell proliferation.

Substances exhibiting anti-estrogenic or selective estrogen receptor modulating activity such as tamoxifen have been associated with Leydig cell tumours in mice (Clegg et al., 1997). A disruption of the negative feed-back exerted by estrogens on the hypothalamic-pituitary system, resulting in elevation of circulating LH, may provide an explanation for anti-estrogen-dependent increase in Leydig cell tumours. A possible anti-estrogenic activity of tetramethrin is indicated by the finding within the study by Rutter (1974), that occurrence of mammary tumours was reduced in the tetramethrin high dose group. In this case, tetramethrin would be mimicking effects of the selective estrogen receptor modulator tamoxifen. Furthermore, while estradiol showed a uterotrophic action in immature female rats, tetramethrin suppressed uterine weight as compared to untreated controls, also suggesting an anti-estrogenic activity (Kim et al., 2005), although an earlier study indicated that tetramethrin did not directly compete with estradiol concerning estrogen receptor binding (Kim et al., 2004).

Furthermore, for the rat, dopamine receptor agonists have been associated with Leydig cell tumours (reviewed by Clegg et al., 1997), presumably by causing a decrease in pituitary prolactin secretion. Since the number of LH receptors on rat Leydig cells is regulated by prolactin, a reduction in circulating prolactin results in a reduction in functional LH receptors and thus to a decreased stimulation of testosterone synthesis, which ultimately leads to enhanced GnRH release and stimulation of Leydig cell proliferation. This mechanism is regarded as not being relevant for the development of Leydig cell tumours in humans (Clegg et al., 1997; ECBI/61/03; ECBI/08/04 Add.4), since LH receptor density in humans appears not to depend on prolactin stimulation. Although dopaminergic system modulation by pyrethroids has been suggested (Hossain et al.,



2006), it is unknown whether tetramethrin itself leads to stimulation of dopamine receptor-dependent pathways. Therefore, it is unclear whether this mechanism is relevant in tetramethrin-dependent support of Leydig cell tumour development.

In summary, several mechanisms by which tetramethrin enhances Leydig cell tumour incidence in male rats may be considered, but additional information is required to define the precise mode(s) of action and to specify whether or not these mechanisms are of relevance to human health. As long as the mode(s) of action has/have not been specified and their relevance to humans has not been ruled out, tetramethrin should be classified as a Category 2 carcinogen based on the findings in rats. This conclusion is in accordance with recommendations along the IPCS framework for analysing the relevance of a cancer mode of action for humans (Boobis et al., 2006), with ECBI/08/04 Add. 4 (Conclusions from the Specialised Experts Meeting concerning Leydig cell tumours) and ECBI/61/03 (“thought starter” for developing an agreed position on the relevance of Leydig cell tumours in rats to humans).

Table 46: Compilation of factors to be taken into consideration in the hazard assessment

Species and strain	Tumour type and background incidence	Multi-site responses	Progression of lesions to malignancy	Reduced tumour latency	Responses in single or both sexes	Confounding effect by excessive toxicity?	Route of exposure	MoA and relevance to humans
Rat, CRL:SD:COBS	Leydig cell tumours	Increase in Leydig cell tumours in males;  Decrease in female mammary tumours	Based on microscopic evaluation at week 104, no apparent capsular invasion or metastasis of adenomas observed		Male  (Decrease in female mammary tumours)	No	oral	Not specified
Rat, CR and Long Evans Hooded	Leydig cell tumours		No differentiation between adenomas and carcinomas in the report		Male	No	Oral	Not specified



**10.9.2 Comparison with the CLP criteria**

Table 47: Results of carcinogenicity studies in comparison to the CLP criteria

Toxicological results	CLP criteria
<p>No data on carcinogenicity of tetramethrin or d-tetramethrin in humans, e. g. in form of epidemiological studies, are available.</p> <p>Two independent rat studies show evidence for a tumourigenic effect of tetramethrin in animals (rat males, 3 rat strains, statistically significant increase in interstitial tumours of the testes).</p> <p>Despite the statistically significant increase in testicular interstitial cell tumours in two independent rat studies, evidence is not considered sufficient to place tetramethrin in Category 1B.</p> <p>As the mode of action leading to Leydig cell tumours in rats has not been specified, relevance to humans is unclear. However, in the absence of information demonstrating non-relevance to humans, relevance is assumed by default.</p> <p>Based on the available animal studies, there is some uncertainty as to which extent promotion of adenomas may have occurred to malignancy (no differentiation between adenoma and carcinomas in the study by Pence).</p>	<p>A substance is classified in Category 1 (known or presumed human carcinogens) for carcinogenicity on the basis of epidemiological and/or animal data. A substance may be further distinguished as:</p> <p>Category 1A, known to have carcinogenic potential for humans, classification is largely based on human evidence, or</p> <p>Category 1B, presumed to have carcinogenic potential for humans, classification is largely based on animal evidence.</p> <p>The classification in Category 1A and 1B is based on strength of evidence together with additional considerations (see section 3.6.2.2). Such evidence may be derived from:                      — human studies that establish a causal relationship between human exposure to a substance and the development of cancer (known human carcinogen); or                      — animal experiments for which there is sufficient (1) evidence to demonstrate animal carcinogenicity (presumed human carcinogen).</p> <p>In addition, on a case-by-case basis, scientific judgement may warrant a decision of presumed human carcinogenicity derived from studies showing limited evidence of carcinogenicity in humans together with limited evidence of carcinogenicity in experimental animals.</p> <p>The placing of a substance in Category 2 (suspected human carcinogens) is done on the basis of evidence obtained from human and/or animal studies, but which is not sufficiently convincing to place the substance in Category 1A or 1B, based on strength of evidence together with additional considerations (see section 3.6.2.2). Such evidence may be derived either from limited (1) evidence of carcinogenicity in human studies or from limited evidence of carcinogenicity in animal studies.                      [...]</p>
	<p>3.6.2.2.3. Strength of evidence involves the enumeration of tumours in human and animal studies and determination of their level of statistical significance. Sufficient human evidence demonstrates causality between human exposure and the development of cancer, whereas sufficient evidence in animals shows a causal relationship between the substance and an increased incidence of tumours. Limited evidence in humans is demonstrated by a positive association between exposure and cancer, but a causal relationship cannot be stated. Limited evidence in animals is provided when data suggest a carcinogenic effect, but are less than sufficient. The terms ‘sufficient’ and</p>

Toxicological results	CLP criteria
	<p>'limited' have been used here as they have been defined by the International Agency for Research on Cancer (IARC) and read as follows:</p> <p>(a) Carcinogenicity in humans The evidence relevant to carcinogenicity from studies in humans is classified into one of the following categories:</p> <ul style="list-style-type: none"> <li>— sufficient evidence of carcinogenicity: a causal relationship has been established between exposure to the agent and human cancer. That is, a positive relationship has been observed between the exposure and cancer in studies in which chance, bias and confounding could be ruled out with reasonable confidence;</li> <li>— limited evidence of carcinogenicity: a positive association has been observed between exposure to the agent and cancer for which a causal interpretation is considered to be credible, but chance, bias or confounding could not be ruled out with reasonable confidence.</li> </ul> <p>(b) Carcinogenicity in experimental animals Carcinogenicity in experimental animals can be evaluated using conventional bioassays, bioassays that employ genetically modified animals, and other in-vivo bioassays that focus on one or more of the critical stages of carcinogenesis. In the absence of data from conventional long-term bioassays or from assays with neoplasia as the end-point, consistently positive results in several models that address several stages in the multistage process of carcinogenesis should be considered in evaluating the degree of evidence of carcinogenicity in experimental animals. The evidence relevant to carcinogenicity in experimental animals is classified into one of the following categories:</p> <ul style="list-style-type: none"> <li>— sufficient evidence of carcinogenicity: a causal relationship has been established between the agent and an increased incidence of malignant neoplasms or of an appropriate combination of benign and malignant neoplasms in (a) two or more species of animals or (b) two or more independent studies in one species carried out at different times or in different laboratories or under different protocols. An increased incidence of tumours in both sexes of a single species in a well-conducted study, ideally conducted under Good Laboratory Practices, can also provide sufficient evidence. A single study in one species and sex might be considered to provide sufficient evidence of carcinogenicity when malignant neoplasms occur to an unusual degree with regard to incidence, site, type of tumour or age at onset, or when there are strong findings of tumours at multiple sites;</li> </ul>
	<ul style="list-style-type: none"> <li>— limited evidence of carcinogenicity: the data suggest a carcinogenic effect but are limited for making a definitive evaluation because, e.g. (a) the evidence of carcinogenicity is restricted to a single experiment; (b) there are unresolved questions regarding the adequacy of the design, conduct or interpretation of the studies; (c) the agent increases the incidence only of benign neoplasms or lesions of uncertain neoplastic potential; or (d) the evidence of carcinogenicity is restricted to studies that demonstrate only promoting activity in a narrow range of tissues or organs.</li> </ul> <p>3.6.2.2.4. Additional considerations (as part of the weight of evidence approach (see 1.1.1)). Beyond the determination of the</p>

Toxicological results	CLP criteria
	<p>strength of evidence for carcinogenicity, a number of other factors need to be considered that influence the overall likelihood that a substance poses a carcinogenic hazard in humans. The full list of factors that influence this determination would be very lengthy, but some of the more important ones are considered here.</p> <p>3.6.2.2.5. The factors can be viewed as either increasing or decreasing the level of concern for human carcinogenicity. The relative emphasis accorded to each factor depends upon the amount and coherence of evidence bearing on each. Generally there is a requirement for more complete information to decrease than to increase the level of concern. Additional considerations should be used in evaluating the tumour findings and the other factors in a case-by-case manner.</p> <p>3.6.2.2.6. Some important factors which may be taken into consideration, when assessing the overall level of concern are:</p> <ul style="list-style-type: none"> <li>(a) tumour type and background incidence;</li> <li>(b) multi-site responses;</li> <li>(c) progression of lesions to malignancy;</li> <li>(d) reduced tumour latency;</li> <li>(e) whether responses are in single or both sexes;</li> <li>(f) whether responses are in a single species or several species;</li> <li>(g) structural similarity to a substance(s) for which there is good evidence of carcinogenicity;</li> <li>(h) routes of exposure;</li> <li>(i) comparison of absorption, distribution, metabolism and excretion between test animals and humans;</li> <li>(j) the possibility of a confounding effect of excessive toxicity at test doses;</li> <li>(k) mode of action and its relevance for humans, such as cytotoxicity with growth stimulation, mitogenesis, immunosuppression, mutagenicity.</li> </ul> <p>Mutagenicity: it is recognised that genetic events are central in the overall process of cancer development. Therefore evidence of mutagenic activity <i>in vivo</i> may indicate that a substance has a potential for carcinogenic effects.</p>

### 10.9.3 Conclusion on classification and labelling for carcinogenicity

Classification/labelling of tetramethrin and d-tetramethrin for carcinogenicity, “Carc. 2”, is proposed, based on two independent rat studies demonstrating a statistically significant increase in incidence in Leydig cell tumours in male rats, without the knowledge whether the underlying mode of action may be of non-relevance to humans.

**10.10 REPRODUCTIVE TOXICITY**

**10.10.1 Adverse effects on sexual function and fertility**

Table 48: Summary table of animal studies on adverse effects on sexual function and fertility

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
Pre-guideline Sim. to OECD 415 non-GLP One-generation. Reproduction test of Neo-Pynamin, Part 1: Fertility study in rats,	The test substance was not administered to maternal animals beyond implantation (day 7 of pregnancy), and was not administered throughout pregnancy and nursing periods. Dosing of males started 9 weeks prior to the mating period. Dosing of females commenced 2 weeks prior to the mating period.	Rat; Slc:SD 20 male and 20 female per dose group	Tetramethrin (Sumitomo) in 0.5% sodium carboxymethylcellulose (vehicle)	0, 100, 300 or 1000 mg/kg bw/d gavage Dosing started at 6 weeks of age (males) or 11 weeks (females). Continued to confirmation of mating success (males) or day 7 of pregnancy (females).	<b>Parental:</b> <u>1000 mg/kg bw/d:</u> Increase in liver weight (M), lower body weight gain after tetramethrin withdrawal (F) <b>Reproductive:</b> <u>1000 mg/kg bw/d:</u> Delayed mating, lower numbers of corpora lutea and implantation sites <b>Offspring (F1)</b> 1000 mg/kg bw/d: Number of surviving fetuses significantly reduced; number of corpora lutea and number of implantations reduced, lower pup weight and body length, delayed ossification <b>Parental and offspring NOAEL</b> 300 mg/kg b w/d	Sato T, Tagewa G, Narama K, (1980), Sumitomo Report No. IT-01-0075
Preguideline		Rat Slc: SD	Tetramethrin in 0.5%	100, 300 and 1000 mg/kg bw/d	300 and 1000 mg/kg bw:	Sato T, Tagawa G and Narama K,

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
Sim. to OECD 415  Non-GLP		(SPF) Male (for mating) and 11-13 (pretest)/20 (main study) females	carboxymethylcellulose	gavage day 7 of pregnancy to day 21 of lactation	<p><u>Dams:</u> slight liver swelling and statistically sign. increased absolute liver weight at autopsy at weaning.</p> <p><u>Offspring:</u> : Sign. increase of stillborn pups, post-implantation loss in the high-dose group in pre-study</p> <p>No embryotoxic effects in main study</p> <p>Maternal NOAEL: 300 mg/kg bw/d</p> <p>Offspring NOAEL: 100 mg/kg bw/d</p>	(1980), Reproduction test of Neo-Pynamin, Part 4: Perinatal and postnatal study in rats, Sumitomo Report No. IT-01-0078

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
Two-generation reproduction toxicity  Sim. to OECD 416, non-GLP	Sperm parameters not determined, longer pre-mating dose period	Rat, Sprague-Dawley,  13 male + 26 female (F <sub>0</sub> )  15 male + 26 female (F <sub>1</sub> )	d-Tetramethrin	0-100-500-3000 ppm 0, 7, 35, 210 mg/kg bw/d (M)  0, 9, 45, 270 mg/kg bw/d (F)  Duration of exposure not mentioned	<u>3000 ppm:</u> Parental: Reduced bw gain (F: F0/F1) during all stages  (F1): Bile duct hyperplasia in maternal females  Offspring (F1/F2): Reduced body weight gain during lactation (M/F)  <b>Parental and offspring</b> NOAEL: 500 ppm <b>NOAEL reproductive</b> : 3000 ppm	Pence D (1986) Two generation reproduction study in rats, Sumitomo Report No. IT-61-0201
Pre-guideline, non-GLP  One-generation reproduction toxicity	Suppl. study with deficiencies, e.g. parental weight development after pre-mating, pathology/histopathology not reported	Rat, Sprague-Dawley,  15 male + 30 female per dose group  Controls: 20 males + 40 females	Tetramethrin	0, 1000, 3000, 6000 ppm  0, 65, 185, 390 mg/kg bw/d (M)  0, 75, 227,482 mg/kg bw/d (F)  Dietary administration  Duration of exposure not mentioned.	Parental: None  Offspring (F1): <u>≥ 3000 ppm:</u> Pup weight at weaning reduced  <u>6000 ppm:</u> Lower lactation index  <b>NOAEL reproductive</b> : <u>1000 ppm</u>	Rutter H A (1974) One generation reproduction study – Rats, Sumitomo Report No. IT-41-0042

Table 49: Summary table of human data on adverse effects on sexual function and fertility

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

**10.10.2 Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility**

Of the submitted rat reproduction toxicity studies, one represents a two-generation study on d-tetramethrin (Pence, 1986; Sumitomo). No effects on reproductive performance were observed within this study up to the highest dose tested (3000 ppm, 210/270 mg/kg bw/d). The parental (maternal) NOAEL of 500 ppm (35/45 mg/kg bw/d) was based on reduced body weight gain in maternal animals and on bile duct hyperplasia in maternal F1 females. The offspring toxicity NOAEL of 45 mg/kg bw/d was based on reduced body weight gain in F1 and F2 generations during lactation and after feeding of 3000 ppm to maternal animals.

The other rat studies represented one-generation studies conducted with tetramethrin, administered orally either by gavage (Sato et al., 1980; Report No. 01-0075 and 01-0078, Sumitomo) or with the diet (Rutter, 1974; Sumitomo). Lower numbers of *corpora lutea* and of resulting implantations/live foetuses were reported in females at the highest dose of 1000 mg/kg bw/d in the one-generation study (01-0075), which may point to a tendency for inhibition of ovulation. However, this effect was slight (about 10 % difference) and occurred only at the highest dose applied. A reduction in litter size is not confirmed by read-across from the two-generation study for d-tetramethrin (Pence, 1986; Sumitomo). Delayed ossification in offspring in this study is attributed to reduced growth and appears to be secondary in nature, since tetramethrin was not administered to maternal animals beyond day 7 of gestation. Beside slight liver swelling and increases of liver weight as maternal toxicity at 300 and 1000 mg/kg bw/, no embryotoxic effects were seen in the second 1-generation study (01-0078).

The study conducted by Rutter (1974) is considered supplemental only. Due of lack of detail to the study report, it is unknown whether parental effects might have occurred. For example, weight development of parental animals was not recorded after the pre-mating phase and pathology/histopathology was not performed. Due to lack of detail to study, the assignment of a parental NO(A)EL is impeded. Anyhow, male and female pup weights at weaning was significantly lower in the 3000 and 6000 ppm groups. At 6000 ppm lactation index (number of pups weaned / number of pups left to nurse) was reduced.

In summary, the available data does not provide evidence for toxic effects of tetramethrin on fertility below doses causing maternal toxicity.

### Comparison with the CLP criteria

Table 50: Results of studies on sexual function and fertility in comparison to the CLP criteria

Toxicological results	CLP criteria
	<p>Category 1A: Known human reproductive toxicant</p> <p>Category 1B: Presumed human reproductive toxicant largely based on data from animal studies</p> <ul style="list-style-type: none"> <li>- clear evidence of an adverse effect on sexual function and fertility in the absence of other toxic effects, or</li> <li>- the adverse effect on reproduction is considered not to be a secondary non-specific consequence of other toxic effects</li> </ul> <p>Category 2: Suspected human reproductive toxicant</p> <ul style="list-style-type: none"> <li>- some evidence from humans or experimental animals, possibly supplemented with other information, of an adverse effect on sexual function and fertility</li> <li>- where the evidence is not sufficiently convincing to place the substance in Category 1 (deficiencies in the study).</li> <li>- the adverse effect on reproduction is considered not to be a secondary non-specific consequence of the other toxic effects</li> </ul>

### 10.10.3 Adverse effects on development

Table 51: Summary table of animal studies on adverse effects on development

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
Teratogenicity study in the rabbit  OECD 414 GLP	Loss of a data book concerned with logging the quantities of test article used. This deviation does not affect the interpretation of this study.	Rabbit, New Zealand white, 20/21 females per dose group	Tetramethrin in 0.5% Carboxymethylcellulose	0, 30, 100, 300, 500 mg/kg bw/d  gavage  day 7-19 of gestation	In a range finding study (500, 1000, 1500 mg/kg bw/d) lack of bw gain ( $\geq$ 500 mg/kg bw/d), deaths (1, 4, 1 from 500 mg/kg bw/d onwards and abortions (2, 4, 5 from 500 mg/kg bw/d onwards) in females  NOAEL maternal: 300 mg/kg bw/d NOAEL developmenta	Robinson K, Washer G and Noveroske J W, (1991) An oral teratology study of Neo-Pynamin in the Rabbit, Sumitomo Report No. IT-11-0234



## CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
					l: 500 mg/kg bw/d	
Teratogenicity study in the rat  OECD 414 GLP	None	Rat, CrI:COBS VAF CD(SD)BR  25 females per dose group	Tetramethrin in 0.5% Carboxymethylcellulose	0, 150, 500, 1000 mg/kg bw gavage day 6-15 of gestation	No treatment-related effects  NOAEL maternal/developmental: 1000 mg/kg bw/d	Robinson K, Washer G and Noveroske J W, (1991), An oral teratology study of Neo-Pynamin in the Rat, Sumitomo Report No. IT-11-0241
Preguideline Sim. to OECD 414 Non-GLP		Rat Slc: SD (SPF) rats 11-13 females (preliminary test), 20 females (main study)	Tetramethrin in 0.5% Carboxymethylcellulose	0, 100, 300, 1000 mg/kg bw gavage day 7 of pregnancy to day 21 of lactation	Slightly lower food consumption and higher liver weights at 1000 mg/kg bw/d  NOAEL maternal 300 mg/kg bw/d  NOAEL developmental: 1000 mg/kg bw/d	Sato T, Tagawa G and Narama K (1980), Reproduction test of Neo-Pynamin, Part 4: Perinatal and postnatal study in rats, Sumitomo Report No. IT-01-0078
Preguideline Sim. to OECD 414 GLP	Subcutaneous administration of the test compound, number of animals insufficient, reporting lacks details on visceral examinations of fetuses	Rabbit New Zealand white  11 females per dose group	d-Tetramethrin in corn oil	0, 30, 100, 300 mg/kg bw/d s.c. injection once daily day 6-18 of gestation	In a range finding study (3 pregnant dams), daily dosing on days 6-18 resulted in all dams dying at 1000 and 2000 mg/kg bw/d. At 500 mg/kg bw/d lower maternal bw, lesser food consumption, and lower number of live fetuses. No effects at 250 mg/kg bw/d.	Satoh R., Kashima M., Takahashi M. and Satoh H.(1982), Teratogenicity study of Neo-Pynamin Forte in rabbits, Sumitomo Report No IT-11-0142

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
					Main study: No effects in dams and offspring.  NOAEL maternal/developmental: 300 mg/kg bw/d	
Preguideline Sim. to OECD 414  GLP	Subcutaneous administration of d-Tetramethrin.  23 dams for examination at term (day 21) and 14 dams for examination of offspring at weaning. Additional groups in which post natal development was examined	Rat Sprague Dawley  37 females per dose group	d-Tetramethrin in corn oil	0, 100, 300, 1000 mg/kg bw/d  s.c. injection once daily  day 7-17 of gestation	1000 mg/kg bw/d: clonic convulsions, lower food consumption and decreases in bw, increases in liver and kidney weights of dams.  300 mg/kg bw/d:  Lower food consumption  No treatment-related effects in fetuses.  NOAEL maternal: 300 mg/kg bw/d  NOAEL developmental: 1000 mg/kg bw/d	Satoh R., Kashima M., Takahashi M. and Satoh H.(1982), Teratogenicity study of Neopynamin Forte in rats, Sumitomo Report No IT-11-0141
Preguideline  Sim. to OECD 414,  non-GLP		Rat, SD (SPF)  30 females per dose group  (20 females: Sacrifice at Caesarean section, 10 females sacrifice at weaning)	Tetramethrin in 0.5% carboxymethyl-cellulose	0, 100, 300, 1000 mg/kg bw/d  gavage  day 7-17 of gestation	1000 mg/kg bw/d: Lower bw gain; liver: swelling, weight increase of liver and kidney.  No treatment-related effects in fetuses.	Sato T and Narama K, (1980b), Reproduction Test of Neopynamin Part 2: Teratology study in rats, Sumitomo Report No. IT-01-0076

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
					NOAEL maternal: 300 mg/kg bw/d NOAEL developmental: 1000 mg/kg bw/d	
Preguideline, Sim. to OECD 414, non-GLP		Rabbit, Japanese White,  10 females per dose group  <u>Range-finding study</u> : 6 females per dose group	Tetramethrin in 0.5% carboxymethylcellulose	0, 50, 150, 500 mg/kg bw/d (main study)  (range finding study: 0, 150, 500, 1500 mg/kg bw/d)  gavage,  day 6-18 of gestation	Range-finding study at 1500 mg/kg bw/d: In dams, decreased bw gain  500 and 1500 mg/kg bw/d: Liver weight increase  Main study at 500 mg/kg bw/d: In dams, decreased bw gain  Main study at 500 mg/kg bw/d: Lower bw of fetuses  Skeletal anomalies (stat. not significant: 3 litter (1 foetus each) with different anomalies at 500 mg/kg bw/d)  NOAEL maternal / developmental: 150 mg/kg bw/d	Sato T and Narama K, (1980c), Reproduction Test of Neopynamin Part 3: Teratology study in rabbits, Sumitomo Report No. IT-01-0077
Preguideline, non-GLP		Rabbit, New	Tetramethrin in corn oil	0-30-90 mg/kg bw/d (in add. 90 mg/kg bw/d)	No treatment-related critical	Dudeck T (1978), Reproduction

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
		Zealand white, 9 females per dose group	Dosing lower than for expected effects	Pyrethrin) Oral, capsule, day 8-16 of gestation	effects  NOAEL maternal/developmental: 90 mg/kg bw/d	Study Neo-Pynamin and Pyrethrin – Rabbits, Sumitomo Report No. IT-61-0009

Table 52: Summary table of human data on adverse effects on development

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

Table 53: Summary table of other studies relevant for developmental toxicity

Type of study/data	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
OECD rodent uterotrophic and Hershberger assays  Non-GLP	Tetramethrin in corn oil	Uterotrophic assay administration was subcutaneous injection  Hershberger assay administration was oral gavage  Rats Sprague-Dawley, Number of animals not specified.  5 to 800 mg/kg in the uterotrophic assay  10, 50 or 100 mg/kg in the Hershberger assay	Tetramethrin may exert endocrine-disrupting effects on female rats through anti-estrogenic action.  No indications for androgenic or antiandrogenic effects	Kim, S. S. et al. (2005) Assessment of estrogenic and androgenic activities of tetramethrin in vitro and in vivo assays. Journal of Toxicology and Environmental Health, 68: 2277-2289.
Soto's E-screen assay (estrogen receptor affinity), Non-guideline Non-GLP	Tetramethrin	In vitro MCF-7 BUS human breast cancer cells	No evidence for estrogenic activity or estrogen receptor affinity	Kim Y. et al. (2004), Assessing Estrogenic Activity of Pyrethroid Insecticides Using In Vitro Combination assays. Journal of Reproduction and development, 50 (2): 245-255

#### 10.10.4 Short summary and overall relevance of the provided information on adverse effects on development

From pre-guideline developmental toxicity studies in rabbits subcutaneously exposed to d-tetramethrin (Sato et al., 1982b and respective range-finding study 1982c), a maternal and fetal NOAEL of 300 mg/kg bw/d was defined, based on mortality of all pregnant dams at 1000 and 2000 mg/kg bw/d daily subcutaneous dosing. At 500 mg/kg bw/d lower maternal body weight, lesser food consumption, and lower number of live foetuses were noted. Subcutaneous dosing of 250 mg/kg bw/d d-tetramethrin resulted in no effects. In the main study daily subcutaneous injections of 0, 30, 100, 300 mg/kg bw/d d-tetramethrin during days 6-18 of gestation no toxicological effects were seen in dams and in the offspring.

Abnormal kidney position was observed in a foetus in the 30 mg/kg group and in 3 foetuses in the 300 mg/kg group but the frequencies were not of statistical significance and therefore considered not *substance-related*. Therefore, the NOAEL (maternal / developmental) was set at 300 mg/kg bw/d.

Two further studies involving tetramethrin administration to rabbits via the oral (gavage) route support a LOAEL of 500 mg/kg bw/d for maternal toxicity and a NOAEL of 300 mg/kg bw/d. In the study by Robinson et al., (1991 a), maternal animals exhibited weight loss at this dose level, one death and two abortions occurred at 500 mg/kg bw/d in the corresponding dose range-finding study. In the study by Sato and Narama (1980c), decreased body weight gain and an increase in liver weight were observed for dams at 500 mg/kg bw/d.

In the studies by Robinson et al. (1991a, 1991, Sumitomo), oral administration of tetramethrin revealed no developmental toxicity up to 500 mg/kg bw/d and 1000 mg/kg bw/d. Likewise, administration of 1000 mg/kg/day resulted in no maternal toxicity.

A statistically non-significant increase in the incidence of rabbit foetal skeletal anomalies was noted at 500 mg/kg bw/d tetramethrin per os (Sato and Namara, 1980c) which were not observed in the control group. As the study was not performed according to GLP and OECD guidelines (e.g. 10 animals per group only) and due to insufficient reporting and recording the study cannot be regarded as a key study. However, the results are regarded as supportive and the NOAEL of 150 mg/kg bw/d is used to establish the lowest NOAEL for developmental toxicity since it cannot be excluded that the rabbit strain used (Japanes White) is the most sensitive. In addition, the study supports the fetal LOAEL demonstrated for subcutaneously administered d-tetramethrin. Thus, from the developmental toxicity studies performed in rabbits, an overall NOAEL of 300 mg/kg bw/d was established for maternal toxicity whereas the lowest NOAEL for developmental toxicity was established at 150 mg/kg bw/d (Sato and Namara, 1980c) of tetramethrin/d-tetramethrin in the rabbit.

Four rat developmental toxicity studies were compared, one involving subcutaneous administration of d-tetramethrin and three dealing with orally administered tetramethrin. In the pre-guideline study with subcutaneous d-tetramethrin (Sato et al., 1982a), clonic convulsions, indications for liver and kidney toxicity, as well as reduced body weight gain were reported for the dams at 1000 mg/kg bw/d, yielding a maternal NOAEL of 300 mg/kg bw/d.

Although no substance-related maternal toxicity was observed up to 1000 mg/kg bw/d in two of the studies with oral tetramethrin (Robinson et al., 1991b; Sato et al., 1980), lower body weight gain and liver/kidney weight increase at this dose in the study by Sato and Narama (1980b) are in line with the LOAEL of 1000 mg/kg bw/d set for subcutaneous d-tetramethrin. Thus, an overall maternal NOAEL of 300 mg/kg bw/d for d-tetramethrin/tetramethrin was established in rats, which is in accordance with the rabbit NOAEL for maternal toxicity.

In all of the four rat developmental toxicity studies, no treatment-related embryotoxic or teratogenic effects were noted up to the highest dose of 1000 mg/kg bw/d. Thus, the developmental NOAEL was set at 1000 mg/kg bw/d for rats, which is higher than the developmental toxicity NOAEL established for rabbits (300 mg/kg bw/d), indicating that susceptibility towards fetal toxicity may differ between species.

Evidence for an antiestrogenic activity of tetramethrin at doses of 5 mg/kg bw/d and above in female Sprague Dawley rats was provided by Kim, S.S. et al. (2005). However, antagonism to oestrogenic actions may not be mediated by binding of tetramethrin to oestrogen receptors, but rather be functional or indirect in nature as tetramethrin appeared not to compete with estradiol for estrogen receptor binding (Kim, I. Y. et al., 2004).

### 10.10.5 Comparison with the CLP criteria

Table 54: Results of developmental toxicity studies in comparison to the CLP criteria

Toxicological results	CLP criteria
<p>No human data are available for tetramethrin or d-tetramethrin, hence a classification in category 1A is not possible.</p> <p>On the basis of the absence of embryotoxic effects and effects on sexual function and fertility especially below maternal toxicity, no classification or labelling of tetramethrin and d-tetramethrin for reproductive toxicity is proposed.</p>	<p>Category 1A: Known human reproductive toxicant</p> <p>Category 1B: Presumed human reproductive toxicant largely based on data from animal studies - clear evidence of an adverse effect on development in the absence of other toxic effects, or - the adverse effect on reproduction is considered not to be a secondary non-specific consequence of other toxic effects</p> <p>Category 2: Suspected human reproductive toxicant - some evidence from humans or experimental animals, possibly supplemented with other information, of an adverse effect on development and - the evidence is not sufficiently convincing to place the substance in Category 1 (deficiencies in the study). - the adverse effect on reproduction is considered not to be a secondary non-specific consequence of the other toxic effects</p>

### 10.10.6 Adverse effects on or via lactation

Table 55: Summary table of animal studies on effects on or via lactation

Method Guideline	Deviation(s) from the guideline (if any)	Species Strain Sex no/group	Test substance, reference to table 5	Dose levels duration of exposure	Results	Reference
Refer to reproductive studies with oral application						

Table 56: Summary table of human data on effects on or via lactation

Type of data/report	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
No data				

**10.10.7 Short summary and overall relevance of the provided information on effects on or via lactation**

Refer to reproductive studies with oral (dietary or gavage) application

**10.10.8 Comparison with the CLP criteria**

No data

**10.10.9 Conclusion on classification and labelling for reproductive toxicity**

No human data are available for tetramethrin or d-tetramethrin, hence a classification in category 1A is not possible.

On the basis of the absence of embryotoxic effects and effects on sexual function and fertility especially below maternal toxicity, no further classification or labelling of tetramethrin and d-tetramethrin for reproductive toxicity is proposed.



**10.11 SPECIFIC TARGET ORGAN TOXICITY-SINGLE EXPOSURE**

Table 57: Summary table of animal studies on STOT SE

Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
OECD 403 Acute inhalation toxicity study with Tetramethrin in Wistar rats. GLP Technical deficiencies: Volume of air chamber 500 L, suggesting long time to achieve steady-state of concentration; no indication of whether concentration measurements performed in breathing zone; dense aerosol accumulation in chamber of G2 groups	Tetra-methrin aerosol in cyclo-hexanone (50% w/v); mean aerosol particle size: G1: 0.67 ± 0.26 µm; G2: 0.68 ± 0.26 µm	Rat, Wistar Pre-study (G1): 2 M + 2 F Main study (G2): 5 M + 5 F	Head and nose exposure	0-5.63 ± 0.86 mg/L 4 h	LC <sub>50</sub> : > 5.63 mg/L  Slight lacrimation and nasal discharge on day 1 (all rats in G2), normal from day 2 onwards, otherwise no toxic signs observed.  Absence of signs of acute toxicity in comparison to other studies suggests that concentration of 5.63 mg/L may not have been achieved in the breathing zone, study not appropriate for classification purposes	Venugopala RK, 2006, Toxicology Department, Advinus Therapeutics Private Limited., Endura Study N° 4414/05
Pre-guideline OECD 403, non-GLP. Acute study, inhalative, whole body 3 h exposure instead of 4 h in TG 403, 1981; Dose-finding study for a	d-Tetra-methrin dissolved in: deodorised kerosene mist, particle size 1.23-1.51 µm	Rat, Sprague-Dawley 10 M + 10 F	Inhalative, whole body	0-0.026-0.131-0.243-0.595-1.18 mg/L  (approx. 0-3.3-16.8-32.3-76.3-151 mg/kg bw); 3 h	LC <sub>50</sub> : > 1.18 mg/L (151 mg/kg bw); ≥ 0.131 mg/L: Muscular fibrillation, urinary incontinence, limb paralysis, bradypnoe, irregular respiration (no. of affected animals and severity of findings not reported; toxic signs began to appear 15 to 30 min, after initiation of exposure	Suzuki T, Kohda H, Misaki Y, Okuno Y, Koyama Y, Miyamoto J, 1981, Sumitomo Report No. IT-10-0144 (Doc III 6.1.3)

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
subacute (28 day) study					and disappeared 1 to 2 hours after exposure)  at 1.18 mg/L: 1/10 females died	
Pre-guideline OECD 412 Non-GLP 28 day inhalation 3 h / day exposure	d-Tetramethrin  dissolved in: deodorised kerosene  mist, particle size 1.23-1.51 µm	Rat, Sprague-Dawley  10 M + 10 F	Inhalative, whole body	0-0.026-0.049-0.087 mg/L  3 h / day exposure for 7 days a week	≥ 0.087 mg/L:  Slight bradypnea, irregular respiration, salivation directly after exposure (no. of affected animals and severity of findings not reported), no cumulative effect.  Increase in leucocyte and decrease in eosinophils count at 0.087 mg/l  NOAEL: 0.049 mg/L	Suzuki T, Kohda H, Misaki Y, Okuno Y, Koyama Y, Miyamoto J, 1981, Sumitomo Report No. IT-10-0144 (Doc III 6.1.3)
90 day inhalation OECD 413 GLP	Tetramethrin in corn oil  Mist particles size 0.65 – 0.95 µm	Rat, Crj:CD(SD)  10 M + 10 F	Inhalative, whole body	0-0.020-0.134-0.824 mg/L  (~0-4.5-29.8-183 mg/kg bw/d)  6 h/day for 5 days a week over 13-weeks	≥ 0.020 mg/L: increased liver and kidney weights (M/F)  In addition at ≥ 0.134 mg/L (29.8 mg/kg bw/d): irregular respiration, bradypnea, decrease bw, changes of haematological (increased fibrinogen, prolongation of blood clotting activities) and blood chemistry parameters (increase in cholesterol and phospholipid levels (M) increased GGT and leucine aminotransferase levels (M)), dark-red discoloration of liver, soft and large liver, hepatocellular hypertrophy, hyaline droplets in renal tubules (M)  In addition at 0.824 mg/L: decrease spontaneous activity. Nasal discharge, salivation, red tears, urinary incontinence.  Increased AIP, AST (M), and gamma-GT (M + F) levels, increased cholesterol levels (M + F), focal necroses in liver, hyaline	Kawaguchi (1991) IT-10-0239 ((Doc III 6.4.3.(1)) Sumitomo

Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
					casts in renal tubules (M) NOAEC: 0.02 mg/L (4.5 mg/kg bw/d) (for incidences and severity see Table 38d)	
90 day inhalation  Determination of the NOEL  OECD 413  GLP	Tetra-methrin  in corn oil  Mist particles size 0.65 – 0.95 µm	Rat, Crj:CD(SD)  10 M + 10 F	Inhalative, whole body	0-0.002-0.004-0.02 mg/L  (0-0.42-0.98-4.4 mg/kg bw/d)  6 h/day for 5 days a week over 13-weeks	Beside increase of liver weights in F F (absolute and relative), no adverse effects observed  NOAEC: 0.02 mg/L (4.5 mg/kg bw/d)	Kawaguchi (1991 b)  IT-10-0238 (Doc III 6.4.3.(2))  Sumitomo

Table 58: Summary table of human data on STOT SE

Type of data/report	Test substance, reference to table 5	Route of exposure	Relevant information about the study (as applicable)	Observations	Reference
No data					

### 10.11.1 Short summary and overall relevance of the provided information on specific target organ toxicity – single exposure

Clinical signs of neurotoxicity were observed in the acute inhalation study performed by Suzuki et al. (1981; Sumitomo) with d-tetramethrin and have to be considered for classification as some of them were also observed in the 28-day study by Suzuki et al. (1981; Sumitomo) with d-tetramethrin as well as in the 90-day study performed by Kawaguchi (1991; Sumitomo) with tetramethrin. Significant neurotoxic effects (e.g. muscular fibrillation, urinary incontinence, limb paralysis, bradypnoe, irregular respiration) were observed in the acute and 90-day inhalative toxicity study with d-tetramethrin at concentrations of 0.131 mg/L or higher.

Slight bradypnea, irregular respiration and salivation directly after exposure were noted in the 28-day inhalative study (Suzuki et al., 1981) at  $\geq 0.087$  mg/L. A cumulative effect was not noted.

The observed effects can be presumed to have the potential to produce significant toxicity in humans on the basis of evidence from studies in experimental animals at a moderate concentration (in accordance with Guidance to Regulation (EC) No 1272/2008 on Classification, Labelling and Packaging of substances and mixtures; November 2013). Hence the proposed classification is “STOT SE 2; H371” for inhalative exposure.

According to guidance values in the above cited guidance document to Regulation (EC) No 1272/2008, classification as “STOT SE 1; H370” (inhalation rat: dust/mist/fume cat. 1:  $C \leq 1.0$  mg/L; Cat. 2:  $5.0 \geq C > 1.0$ ) might be considered. However, since the severity of the effects was not clearly documented in the acute inhalation study (study was performed to estimate the  $LC_{50}$ ), classification as “STOT SE 2; H371” is proposed. Although neurotoxic effects may have been responsible for the observed mortality resulting in the  $LC_{50} > 1.18$  mg/L additional classification as “STOT SE 2; H371” is justified as the neurotoxic effects were already observed at approximately 10-fold lower concentrations. The severity of the effects cannot be estimated from the provided data.

### 10.11.2 Comparison with the CLP criteria

Table 59: Results of toxicity studies relevant for STOT SE in comparison to the CLP criteria

Toxicological results	CLP criteria	
<p>Data on significant toxicity in humans are lacking.</p> <p>Since the severity of neurotoxic effects was not clearly documented in the acute inhalation study (study was performed to estimate the <math>LC_{50}</math>), classification as “STOT SE 1; H370” is not proposed.</p>	<p>Category 1 (H370)</p> <p>Oral (rat): <math>C \leq 300</math> mg/kg bw</p> <p>Dermal (rat or rabbit): <math>C \leq 1000</math> mg/kg bw</p> <p>Inhalative (rat, dust/mist/fume): <math>\leq 1</math> mg/L/4 h</p>	<p>Substances that have produced significant toxicity in humans</p> <p>or that, on the basis of evidence from studies in experimental animals, can be presumed to have the potential to produce significant toxicity in humans following single exposure</p> <ul style="list-style-type: none"> <li>- reliable and good quality evidence from human cases or epidemiological studies; or</li> <li>- observations from appropriate studies in experimental animals in which significant and/or severe toxic effects of relevance to human health were produced at generally low exposure concentrations.</li> </ul>
<p>On the basis of evidence from studies in experimental animals at a moderate concentration (especially noted under specific target organ toxicity – repeated exposure), the observed neurotoxic effects were essentially acute effects and can be presumed to have the potential to produce significant toxicity in humans (in accordance with Guidance to Regulation (EC) No 1272/2008 on Classification, Labelling and Packaging of substances and mixtures; November 2013). Hence the proposed classification is “STOT SE 2; H371” for inhalative exposure.</p>	<p>Category 2 (H371)</p> <p>Oral (rat): <math>2000 \geq C &gt; 300</math> mg/kg bw</p> <p>Dermal (rat or rabbit): <math>2000 \geq C &gt; 1000</math> mg/kg bw</p> <p>Inhalative (rat, dust/mist/fume): <math>5 \geq C &gt; 1</math> mg/L/4 h</p>	<p>Substances that, on the basis of evidence from studies in experimental animals can be presumed to have the potential to be harmful to human health following single exposure</p> <ul style="list-style-type: none"> <li>- observations from appropriate studies in experimental animals in which significant toxic effects, of relevance to human health, were produced at generally moderate exposure concentrations.</li> </ul>
	<p>Category 3 (H335/H336)</p> <p>Guidance values do not apply (mainly based on</p>	<p>Transient target organ effects</p> <p>This category only includes narcotic effects and respiratory tract irritation. These are target</p>

Toxicological results	CLP criteria	
	human data)	organ effects for which a substance does not meet the criteria to be classified in Categories 1 or 2 indicated above. These are effects which adversely alter human function for a short duration after exposure and from which humans may recover in a reasonable period without leaving significant alteration of structure or function.

**10.11.3 Conclusion on classification and labelling for STOT SE**

STOT SE 2; H371 (May cause damage to the central nervous system if inhaled).

## 10.12 SPECIFIC TARGET ORGAN TOXICITY-REPEATED EXPOSURE

Table 60: Summary table of animal studies on STOT RE

Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
Sim. to OECD 407, non-GLP	d-Tetra-methrin	Rat, CRJ:CD (SD), 10 M + 10 F / dose group	Oral, dietary,	0-300-3000-10000 ppm (0-30-290/295-965/940 mg/kg bw/d (M/F))  28 days	From 3000 ppm onwards: Increased cholesterol, albumin, total protein, and glucose levels , increased liver weights (regarded as adaptive)  10000 ppm: increased ALT and AST levels; Liver: Focal necrosis, (M), enlargement (M + F), peripheral lobular hypertrophy of parenchymal cells Haematology: Increased thrombocyte levels (M), decreased haematocrit and haemoglobin (M) Other: Increased urinary protein levels (M), decreased bw gain  NOAEL: 3000 ppm (290/295 mg/kg bw/d (M/F))	Hosokawa, S. (1985) Sumitomo report No. IT-20-0188 (Su: Doc III-A6.3.1)
OECD 408 GLP	Tetra-methrin in food	Rat, Wistar 10 M+10 F /group	Oral, dietary, Including 4-week recovery period	0-500-1000-2000 ppm (0-38-76-151 mg/kg bw/d), Recovery group at 0 + 2000 ppm  90 days	1000 + 2000 ppm: Increased liver weights (rel./abs; M) and cholesterol (M/F), hypertrophy of hepatocytes (M)  2000 ppm: Neurological effects: significant increase in landing foot splay (M) increased liver weight (F)  NOAEL: 1000 ppm (76 mg/kg bw/d)	Malleshappa, H.N. (2002). Endura Study N°3343/01, 24 (En: Doc III-A6.4.1.1)
Sim. to OECD 408, non-GLP	d-Tetra-methrin in corn oil	Rat, Sprague-Dawley, 3 mo: 12 M + 12 F/dose group (satellite	Oral, dietary	0-100-300-1000-3000 ppm (0-5.8/7-17/21-58/71-178/214 mg/kg bw/d	1000 ppm onwards: Slightly increased urinary protein and cholesterol levels (M/F), increased liver (M/F) and kidney weights (M) In addition at 3000 ppm: Increased kidney weights	Hosokawa S, Hiromori T, Seki T, Okuno Y, Miyamoto J, 1980, Sumitomo Report No.

## CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
		groups) 6 mo: 20 M + 20 F / dose group (main groups)		(M/F) 3 months + 6 months	(MF), significant reduced bw (M/F) , swelling and 'luster surface' of the liver (M), eosinophilic bodies in renal tubular epithelium (M)  NOAEL: 1000 ppm (58/71 mg/kg bw/d (M+F))	IT-00-0139 (Su: Doc III-A6.4.1)
Sim. to OECD 408, non-GLP	Tetra-methrin	Rat, Sprague-Dawley,  16 M + 16 F/dose group	Oral, dietary	0-500-1500-5000 ppm  (0-30-95-325 mg/kg bw/d)  6 months	<u>5000 ppm:</u> Increased cholesterol levels (M/F), increase in absolute (M) and relative (M/F) liver weight, decreased haemoglobin levels (M), decreased AIP and ALAT (M), decreased body weight gain (M+F)  NOAEL 1500 ppm (95 mg/kg bw/d)	Suzuki, T, Okuno Y., 1977, Sumitomo Report No. IT-60-0015 (Su: Doc III-A6.4.1(2))
Sim. to OECD 409, non-GLP	Tetra-methrin	Dog, Beagle, 6 M + 6 F / dose group	Oral, dietary	0-1250-2500-5000 ppm  (0-45-90-180 mg/kg bw/d)  6 months	<u>2500 ppm onwards:</u> Decrease in albumin/globulin ratio (M/F), increase in cholesterol (F) increased liver weight (M), increased nervousness, tremors (M+F)  <u>5000 ppm:</u> Decrease in total protein (M/F), albumin (M/F); increase in cholesterol (M/F); decrease blood urea nitrogen (M), decreased haematocrit and erythrocyte counts (M); relative liver weight increase (M/F); decrease in absolute + relative ovary weight; Ovary, mammary gland: absence of histopathological changes associated with oestrus (i.e. corpora lutea; endometrial hypertrophy; mammary gland hyperplasia, secretory activity, and stromal and ductal proliferation)  NOAEL: 2500 ppm (90	Pence D, 1981, Sumitomo Report No. IT-11-0098 (Su: Doc III-A6.4.1(1))

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
					mg/kg bw/d)	
Sim. to OECD 452 GLP	Tetra-methrin	Dog, Beagle,  4 M + 4 F/dose group	Oral, dietary	0-300-1200-5000-10000 ppm  (0-8/9-36/36-147/157-286/325 mg/kg bw/d (M/F))  1 year	1200 ppm: Slightly increased cholesterol and phospholipids (M), increased liver weight (M)  <u>≥ 5000 ppm:</u> Lower bw (F), increased cholesterol, phospholipid and AIP levels (M+F), decreased albumin levels (F), decreased erythrocyte count, haemoglobin and haematocrit levels (F), increased liver weight (M)  NOAEL: 1200 ppm (36 mg/kg bw/d)  Slight changes in liver physiology and clinical chemistry were already observed at 1200 ppm in males. However, the deduced NO(A)EL is based on aggravation of liver effects (increase in liver weight also in females, increase in AIP) and on occurrence of haematological effects (increased platelet count).	Walker M D, 1996, Sumitomo Report No. IT-0276 (Doc. III-A6.5.4)
90 day inhalation  OECD 413  GLP	Tetra-methrin  in corn oil  Mist particles size 0.65 – 0.95 µm	Rat, Crj:CD(SD)  10 M + 10 F	Inhalative, whole body	0-0.020-0.134-0.824 mg/L  (~0-4.5-29.8-183 mg/kg bw/d)  6 h/day for 5 days a week over 13-weeks	≥ 0.020 mg/L: increased liver and kidney weights (M/F)  In addition at ≥ 0.134 mg/L (29.8 mg/kg bw/d): irregular respiration, bradypnea, decrease bw, changes of haematological (increased bilirubin and urobilinogen) and blood chemistry parameters (increased cholesterol and phospholipid levels (M) increased GGT and leucine aminotransferase levels (M)), dark-red discoloration of liver, soft and large liver, hepatocellular hypertrophy, hyaline droplets in renal tubules (M)  In addition at 0.824 mg/L: decrease spontaneous	Kawaguchi (1991 a)  IT-10-0239 (Doc. III 6.4.3.1)  Sumitomo



Method Guideline, Deviation(s) from the guideline (if any)	Test substance, reference to table 5	Species Strain Sex no/group	Route of exposure	Dose levels duration of exposure	Results	Reference
					activity. Nasal discharge, salivation, red tears, urinary incontinence.  Increased ALP, AST (M), and gamma-GT (M + F) levels, increased cholesterol levels (M + F), focal necroses in liver, hyaline casts in renal tubules (M)  NOAEC: 0.02 mg/L (4.5 mg/kg bw/d)	

Table 61: Summary table of human data on STOT RE

Type of data/report	Test substance, reference to table 5	Route of exposure	Relevant information about the study (as applicable)	Observations	Reference
No Data					

**10.13 SHORT SUMMARY AND OVERALL RELEVANCE OF THE PROVIDED INFORMATION ON SPECIFIC TARGET ORGAN TOXICITY – REPEATED EXPOSURE**

Toxicity of orally administered substance was tested in rats over 4 weeks in a study with d-tetramethrin (Hosokawa, 1985; Sumitomo), in a 90-day study with tetramethrin (Mallehappa, 2002; Endura) as well as in one 3-6-month study in rats with d-tetramethrin (Hosokawa, 1980; Sumitomo) and in a study over 6 months with tetramethrin (Suzuki et al., 1977; Sumitomo). Common effects observed in these studies included a decrease in body weight gain, haematological alterations, changes in clinical chemistry, increased liver weight and dose-dependent liver toxicity. The LOAELs were predominantly based on liver toxicity (gross and histopathology and clinical chemistry) and haematological findings. The derived subacute and subchronic NOAELs were 290 mg/kg bw/d for exposure over 4 weeks (d-tetramethrin; Hosokawa 1985, Sumitomo), 90 mg/kg bw/d (2500 ppm) and 36 mg/kg bw/d (1200 ppm) for exposure of Beagle dogs to tetramethrin over one year (Walker 1986; Sumitomo) and 6 months (Pence, 1981, Sumitomo).

Changes in blood chemistry (decreased albumin/globulin ratio, decrease in total protein and albumin, increased cholesterol levels) and haematological parameters (decreased blood urea nitrogen decreased haematocrit and erythrocyte counts), increased liver weight as well as increased nervousness and tremors were noted at 36 mg/kg bw/d (Walker, 1996) and 90 mg/kg bw/d (Pence 1981). In the study conducted by Pence (1981) female dogs showed a decrease in absolute + relative ovary weight as well as absence of histopathological changes associated with oestrus (i.e. corpora lutea; endometrial hypertrophy; mammary gland hyperplasia, secretory activity, and stromal

and ductal proliferation) at the highest dose of 5000 ppm (180 mg/kg bw/d). This finding was not reproducible in the study performed by Walker (1996).

Increased ALP levels, increased liver weights, and decreased erythrocyte count, haemoglobin and hematocrit levels, were observed from dose level of 5000 ppm (147/157 mg/kg bw/d; M/F) onwards.

These values were supported by NOAELs of 76 mg/kg bw (1000 ppm; Malleshappa 2002; Endura) and 95 mg/kg bw (1500 ppm; Suzuki and Okuno 1977; Sumitomo) in subchronic toxicity studies between 3 and 6 months in rats.

Furthermore, two overlapping studies were performed to test subchronic (3 months) inhalative toxicity of tetramethrin in Sprague Dawley Crj:CD rats (Kawaguchi, 1991a and b; Sumitomo), the study IT-10-238 (Kawaguchi 1991 b) displaying a refined spacing of the low concentration range as compared to the study IT-10-239 (Kawaguchi 1991 a). In the study No. -238 no adverse effects were observed up to a concentration of 0.02 mg tetramethrin /L air (4.4 mg/kg bw/d). In the study -0239 (Kawaguchi 1991 a), adverse effects relevant for the LOAEC of 0.134 mg/L comprised lower body weight gain (males, mid and high dose; females high dose without statistical significance), changes in clinical chemistry (higher total protein, lower albumin, higher alpha2-globulin resulting in lower A/G ratio in both sexes of the high dose group, higher cholesterol, phospholipid and  $\Gamma$ -Glutamyl transpeptidase in males of the mid and high dose and in females of the high dose), urinalysis (significantly increased bilirubin and urobilinogen in both sexes, mid and high dose), haematology (prolonged prothrombin time, higher activated partial thromboplastin time (APTT) and fibrinogen in both sexes, high dose) organ toxicity (increased absolute and relative liver weights in both sexes from the low dose onwards), macro- (dark red liver, soft and large liver in males of the mid and high dose, with lesser extent also in females of these dose groups), and histopathological organ abnormalities (In the liver: focal necrosis and bile duct hyperplasia in males of the high dose group, massive necrosis in 1/10 female of the high dose group, hepatocellular hypertrophy in both sexes from the mid and high dose group. In the kidney: Basophilic tubules, eosinophilic bodies, hyaline casts and hyaline droplets in tubules with higher incidence and/or severity grading in males of the mid and high dose). Acute clinical signs occurred during the daily exposure period (irregular respiration, bradypnoe). Both 3-month studies pointed to a NOAEC of 0.02 mg/L (approximate daily dose of 4.5 mg/kg bw) and a LOAEC of 0.134 mg/L (29.8 mg/kg bw/d). Conversion of inhaled concentrations into inhaled daily doses was based on default assumptions regarding body weight, inhalation volume and 100 % availability and suggests a significantly higher sensitivity to d-tetramethrin and tetramethrin by the inhalative than the oral route of administration in the rat (NOAEL for tetramethrin of 4.5 mg/kg bw/d, 3 months inhalative versus 95 mg/kg bw/d, in subchronic oral study in rats (6 months; Suzuki et al. 1977; Sumitomo).

The effects in the 3-month inhalation toxicity study (Kawaguchi 1991; see table 38d) were more pronounced in male rats of the mid and high dose groups than in female rats of these dose groups and were considered adverse. Anyhow, in comparison with the criteria for specific target organ toxicity after repeated exposure, on the basis of the weight of evidence and by the use of expert judgement the observed changes in liver physiology and clinical chemistry were considered not adequate for a classification of tetramethrin as specific target organ toxicants following repeated exposure (STOT RE).

The observed neurotoxic effects seen in the 28-day and 90-day studies were essentially acute effects and they were considered relevant for classification as specific target organ toxicant – single exposure (STOT SE) (for neurotoxicity). In the 90-d rat study with inhalative exposure (Kawaguchi et al. 1991a) concentration dependent incidences of irregular respiration were observed in mid- and high concentration groups (LOAEC:

CLH REPORT FOR D-TRANS-TETRAMETHRIN

0.134 mg/L) over the entire period of exposure. Bradypnoea and decrease of spontaneous activity occurred in a concentration dependent manner and were considered adverse in the highest exposure group (animals affected per exposure day, days with affected animals). Bradypnoea was observed up to day 12 (males) resp. day 45 (females), decrease of spontaneous activity up to day 45 of exposure (males and females). The results are summarised in Table 38d. The severity of effects was not reported.

Table 62: Summary table of selected adverse effects in the 3-month inhalative toxicity study in rats (Kawaguchi; 1991a)

Males Effect	Dose groups			
	0 mg/L (Vehicle control)	0.02 mg/L (4.5 mg/kg bw/D)	0.134 mg/L 30 mg/kg bw/d)	0.824 mg/L 183 mg/kg bw/d)
Mean body weight on day 89 ± SD (g)	552 ± 52.5	550 ± 62.4	494 * ± 31.8	454 ** ± 43.4
Final body weight (g)	524 ± 51.2	521 ± 61.9	470* ± 28.4	422** ± 39.9
Prothrombin time (sec)	17.7 ± 1.64	17.3 ± 1.95	18.9 ± 2.94	21.3** ± 3.35
Activated partial thromboplastin time (sec)	24.0 ± 1.62	23.3 ± 2.93	24.9 ± 2.37	27.3** ± 2.79
Fibrinogen (mg/dl)	233.9 ± 14.85	234.9 ± 15.48	235.2 ± 16.77	261.2** ± 22.75
Total protein (g/dl)	6.0 ± 0.25	6.2* ± 0.28	6.2* ± 0.24	6.5** ± 0.25
Albumin (%)	54.0 ± 1.54	51.5* ± 0.23	52.0 ± 1.93	50.0** ± 3.46
A2- Globulin (%)	4.8 ± 0.43	5.2 ± 0.47	5.4* ± 0.58	5.9** ± 0.96
A/G ratio	1.18 ± 0.075	1.07 ± 0.117	1.09 ± 0.085	1.05* ± 0.150
Cholesterol (mg/dl)	70 ± 18.8	72 ± 14.9	92* ± 20.3	114** ± 29.9
Phospholipid (mg/dl)	105 ± 21.8	110 ± 21.0	135* ± 29.5	172** ± 44.6
Γ-Glutamyl transpeptidase (U/l)	1 ± 1.2	1 ± 1.2	4** ± 2.1	13** ± 3.6
Liver weight (abs., g)	14.20 ± 1.945	15.58 ± 2.532	16.00* ± 1.480	17.68** ± 2.212
Liver weight (rel., g%)	2.71 ± 0.156	2.98* ± 0.217	3.42** ± 0.194	4.19** ± 0.369
<b>Gross pathological changes</b>				
Liver: Dark red	0/10	0/10	4/10	9/10
Liver: Soft	0/10	0/10	3/10	3/10
Liver: Large	0/10	0/10	4/10	6/10
<b>Histopathological examination</b>				
Liver: Focal necrosis	1/10	0/10	0/10	3/10
Liver: Hepatocellular hypertrophy	0/10	0/10	5/10 (slight)	5/10 (slight) 4/10 (mild)
Liver: Bile duct hyperplasia	0/10	0/10	0/10	4/10 (slight)
Kidney: hyaline droplets in tubules	2/10 (slight)	4/10 (slight)	3/10 (slight) 4/10(mild) 1/10 (severe)	2/10 (slight) 3/10(mild) 5/10 (severe)

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Clinical signs in first 2 weeks (10 days of exposure)				
Bradypnoea <b>days</b> with $\geq 1$ animal affected/days with exposure (mean <b>animals</b> affected per day) Exposure: days 1 - 14 <sup>§</sup>	1/10 (0)	0/10 (0)	0/10 (0)	10/10 (6)
Irregular respiration: # days with $\geq 1$ animal affected/days with exposure (mean animals affected per day) Exposure: days 1 - 14 <sup>§§</sup>	7/10 (2)	9/10 (2)	10/10 (7)	10/10 (10)
Decrease of spontaneous activity # days with $\geq 1$ animal affected/days with exposure (mean animals affected per day) Exposure: days 1 - 14 <sup>§§§</sup>	1/10 (0)	1/10 (0)	8/10 (2)	10/10 (9)

§: days 6, 7, 13, 14: non-exposure, bradypnoea occurred only incidental after day 12 of exposure

§§: days 6,7, 13, 14: non-exposure, irregular respiration: all animals in the two highest exposure groups affected on every exposure day during entire treatment period, groups vehicle control and lowest concentration: incidental following approx. 2 wks of exposure

§§§: days 6, 7, 13, 14: non-exposure, decrease in spontaneous activity, decreases in spontaneous activity observed at highest concentration up to day 45 of exposure, afterwards no more incidences reported

## CLH REPORT FOR D-TRANS-TETRAMETHRIN

Females Effect	Dose groups			
	0 mg/L	0.02 mg/L (4.5 mg/kg bw/D)	0.134 mg/L 30 mg/kg bw/d)	0.824 mg/L 183 mg/kg bw/d)
Mean body weight on day 89 ± SD (g)	310 ± 30.2	305 ± 27.7	296 ± 24.3	288 ± 22.8
Final body weight (g)	291 ± 27.9	287 ± 27.1	274 ± 22.7	259 ± 21.6
Prothrombin time (sec)	14.0 ± 0.25	13.9 ± 0.23	14.1 ± 0.25	14.5** ± 0.76
Activated partial thromboplastin time (sec)	20.7 ± 1.46	20.2 ± 1.43	20.6 ± 1.50	22.4** ± 1.59
Fibrinogen (mg/dl)	173.7 ± 17.32	161.1 ± 18.44	163.8 ± 13.49	190.8* ± 16.86
Total protein (g/dl)	6.5 ± 0.35	6.5 ± 0.45	6.6 ± 0.28	7.1** ± 0.20
Albumin (%)	59.3 ± 3.49	50.4 ± 2.26	58.3 ± 3.07	53.0** ± 2.39
A2- Globulin (%)	5.0 ± 0.77	5.5 ± 0.48	5.3 ± 0.41	6.1 ± 0.78
A/G ratio	1.47 ± 0.204	1.47 ± 0.139	1.41 ± 0.183	1.16** ± 0.109
Cholesterol (mg/dl)	80 ± 10.1	83 ± 14.1	91 ± 17.9	117** ± 17.8
Phospholipid (mg/dl)	148 ± 21.5	158 ± 35.1	162 ± 23.4	196 ± 25.4
Γ-Glutamyl transpeptidase (U/l)	1 ± 1.2	1 ± 1.2	2 ± 1.1	9** ± 7.5
Liver weight (abs., g)	7.48 ± 0.885	7.96 ± 1.341	8.78** ± 0.811	10.24** ± 0.639
Liver weight (rel., g%)	2.57 ± 0.135	2.77* ± 0.288	3.21** ± 0.071	3.97** ± 0.273
<b>Gross pathological changes</b>				
Liver: Dark red	0/10	0/10	1/10	7/10
Liver: Soft	0/10	0/10	0/10	1/10
Liver: Large	0/10	0/10	2/10	7/10
<b>Histopathological examination</b>				
Liver: Massive necrosis	0/10	0/10	0/10	1/10
Liver: Hepatocellular hypertrophy	0/10	0/10	2/10 (slight)	9/10 (slight)
Liver: Bile duct hyperplasia	0/10	0/10	0/10	1/10 (slight)
<b>Clinical signs in first 2 weeks (10 days of exposure)</b>				
Bradypnoea # days with ≥ 1 animal affected/days with exposure (mean animals affected per day) Exposure: days 1 - 14 <sup>§</sup>	0/10 (0)	0/10 (0)	1/10 (2)	10/10 (5)
Irregular respiration: # days with ≥ 1 animal affected/days with exposure (mean animals affected per day) Exposure: days 1 - 14 <sup>§§</sup>	7/10 (1)	6/10 (1)	10/10 (3)	10/10 (10)
Decrease of spontaneous activity # days with ≥ 1 animal affected/days with exposure (mean	0/10 (0)	0/10 (0)	3/10 (0)	10/10 (6)

animals affected per day) Exposure: days 1 - 14 <sup>§§§</sup>				
---	--	--	--	--

§: days 6, 7, 13, 14: non-exposure, bradypnoea occurred only incidental after day 12 of exposure  
 §§: days 6,7, 13, 14: non-exposure, irregular respiration: all animals in the two highest exposure groups affected on every exposure day during entire treatment period, groups vehicle control and lowest concentration: incidental following approx. 2 wks of exposure  
 §§§: days 6, 7, 13, 14: non-exposure, decrease in spontaneous activity, decreases in spontaneous activity observed at highest concentration up to day 45 of exposure, afterwards no more incidences reported  
 \*\* Significantly different from vehicle control (P<0.01)  
 \* Significantly different from vehicle control (P<0.05)

Table 63: Correction of the effective dose

Study reference	Effective dose (mg/kg/d)	Length of exposure	Correction of the effective dose when extrapolated to 90-day exposure	Classification supported by the study
Not necessary				

### 10.13.1 Comparison with the CLP criteria

Table 64: Results of toxicity studies relevant for STOT RE in comparison to the CLP criteria

Toxicological results	CLP criteria
Data on significant toxicity in humans are lacking and guidance values are not applicable. <sup>74</sup>	<p>Category 1 (H372):                      Substances that have produced significant toxicity in humans or that, on the basis of evidence from studies in experimental animals, can be presumed to have the potential to produce significant toxicity in humans following repeated exposure.                      Substances are classified in Category 1 for target organ toxicity (repeat exposure) on the basis of:                      reliable and good quality evidence from human cases or epidemiological studies; or observations from appropriate studies in experimental animals in which significant and/or severe toxic effects, of relevance to human health, were produced at generally low exposure concentrations.                      Equivalent guidance values for 28-day and 90-day studies:                      Inhalation dust/mist/fume, rat:                      28-day: ≤ 0.06 mg/L/6 h/d                      90-day: ≤ 0.02 mg/L/6 h/d</p>
On the basis of evidence from studies with repeated exposure in experimental animals at a moderate concentration, the observed effects (liver) can be presumed to represent adaptative effects only and not to have the potential to produce significant toxicity in humans (in accordance with Guidance to Regulation (EC) No 1272/2008 on Classification, Labelling and Packaging of substances and mixtures; November 2013). Hence, no classification for “STOT-RE” for oral exposure is proposed.	<p>Category 2 (H373):                      Substances that, on the basis of evidence from studies in experimental animals can be presumed to have the potential to be harmful to human health following repeated exposure.                      Substances are classified in category 2 for target organ toxicity (repeat exposure) on the basis of observations from appropriate studies in experimental animals in which significant toxic effects, of relevance to human health, were produced at generally moderate exposure concentrations.                      Guidance dose/concentration values are provided below (see 3.9.2.9) in order to help in classification.                      In exceptional cases human evidence can also be used to place a substance in Category 2.                      Equivalent guidance values for 28-day and 90-day studies:                      Inhalation dust/mist/fume, rat:                      28-day: ≤ 0.6 mg/L/6 h/d                      90-day: ≤ 0.2 mg/L/6 h/d</p>

**10.13.2 Conclusion on classification and labelling for STOT RE**

Classification and labelling for STOT RE is not proposed.

**10.14 ASPIRATION HAZARD**

Table 65: Summary table of evidence for aspiration hazard

Type of study/data	Test substance, reference to table 5	Relevant information about the study (as applicable)	Observations	Reference
Not applicable				

## 11. EVALUATION OF ENVIRONMENTAL HAZARDS

### 11.1 ACUTE AQUATIC HAZARD

The applicant provided ecotoxicological tests performed with tetramethrin for the effect assessment of d-trans-tetramethrin. Both substances consist of the same 4 isomers (1R trans, 1R cis, 1S trans, 1S cis). However, there is a difference in the ratio of the isomers. Tetramethrin contains the isomers 1R trans, 1R cis, 1S trans, 1S cis in the ratio 4:1:1:4. D-trans-tetramethrin contains mainly the isomer 1R trans with > 90 %. In general, different isomeric mixtures may result in different ecotoxicological effect values dependant on which isomer is the most active and on the composition of the mixture. Compilation of the available data gives the following picture:

Table 66:

	<b>Fish (LC<sub>50</sub>)</b>	<b>Daphnia (EC<sub>50</sub>)</b>	<b>Algae (E<sub>r</sub>C<sub>50</sub>)</b>
tetramethrin	3.7 µg/L	0.11 mg/l	> 0.25 mg/l
d-trans-tetramethrin	5.9 µg/L	-	> 1.25 mg/L

For fish (*O. mykiss*) and green algae effect values for both compounds are available. The LC<sub>50</sub>/EC<sub>50</sub> values are quite similar. Although the data basis is quite scarce, it is concluded that both compounds are ecotoxicologically equivalent and the effect values for tetramethrin are used for the effects assessment of d-trans-tetramethrin. The table below summarises the available key studies.



Table 67: Summary of relevant information on acute aquatic toxicity

Method	Species	Test (endpoint, design, duration)	Results <sup>1</sup>	Key or Supportive study	Remarks	Reference
OECD 203; FIFRA 72-1; OPPTS 850.1075	<i>Oncorhynchus mykiss</i>	mortality flow-through 96 h	LC <sub>50</sub> = 5.9 µg/l (m)	key	Solvent DMF used	York D., 2008 d-trans-Tetramethrin_14.4.1_03_short-term toxicity fish
EPA OPP 72-2	<i>Daphnia magna</i>	Immobility flow-through 48 h	EC <sub>50</sub> = 0.11 mg/l (m)	key	Solvent DMF used; abnormal behaviour observed: surfacing, daphnids on bottom of the vessel, trailing extraneous material, moving slower; test substance <b>Tetramethrin</b> .	Blasberg, 1993 Tetramethrin_14.4.2_02_short-term toxicity invertebrates_Sumitomo
OECD 201	<i>Pseudokirchneriella subcapitata</i>	growth inhibition static 72 h	ErC <sub>50</sub> > 1.25 mg/l (MMC) NOErC = 0.25 mg/l (m)	key	Solvent DMF used	Hoberg 2002; d-trans-Tetramethrin_14.4.3_growth inhibition algae

<sup>1</sup> Indicate if the results are based on the measured (m) or on the nominal (n) concentration

### 11.1.1 Acute (short-term) toxicity to fish

An acute fish test using *Oncorhynchus mykiss* as test species was performed with d-trans-tetramethrin as test substance (York, 2008). 10 fish per replicate (2 replicates per concentration) were exposed in a flow-through system to 5 concentrations of the test substance (2.0, 4.0, 8.0, 16 and 32 µg/L (nominal), a control and a solvent control. Dimethylformamide (DMF) was used as a solvent. The solvent control chamber received an aliquot of 0.05 mL of DMF which was equivalent to that received by the highest test concentration. Mean measured concentrations ranged between 1.6 and 31 µg/L. All effect values are based on mean measured concentrations. Mortality first occurred at a concentration of 7.8 µg/L (mean measured). The 96h-LC<sub>50</sub> was determined to be 5.9 µg/L. The applicant provided two further acute fish tests with d-trans tetramethrin (cf. study summaries “d-trans-Tetramethrin\_14.4.1\_01\_short-term toxicity fish” and “d-trans-Tetramethrin\_14.4.1\_02\_short-term toxicity fish”, see confidential annex). However, both studies by Sousa & LeBlanc (1981) were regarded as not valid by the eCA as both tests were performed in a static test system without analytical monitoring of the test substance concentration. For the ecotoxicological equivalent compound tetramethrin, a 96h-LC<sub>50</sub> of 3.7 µg/L was obtained for *Oncorhynchus mykiss* (cf. study summary “Tetramethrin\_14.4.1\_03\_short-term toxicity fish\_Sumitomo”, see confidential annex).

The lowest available aquatic endpoint for tetramethrin is the  $LC_{50} = 5.9 \mu\text{g/l}$  for *Oncorhynchus mykiss*. However, the classification of d-trans-tetramethrin is based on the 96h- $LC_{50}$  of  $3.7 \mu\text{g/L}$  for tetramethrin, as the ecotoxicological equivalence of both compounds was accepted.

#### Acute (short-term) toxicity to aquatic invertebrates

No acute aquatic toxicity study with invertebrates has been provided by the applicant for d-trans-tetramethrin. Instead, an acute toxicity study with *Daphnia magna* according to EPA guideline was performed for tetramethrin, using a flow-through system (Blasberg, 1993). The test substance concentration was measured at test start and end. 20 daphnids divided in 2 replicates were exposed to 5 test substance concentrations (0.06 – 1 mg/l nominal), a control and a solvent control. DMF was used as solvent. The solvent control chamber received an aliquot of 0.1 ml of DMF which was approximately equivalent to that received by the highest test concentration. The result is based on mean measured concentration as the measured concentrations significantly deviate from the nominal concentrations. The mean measured concentration was 31 ( $\pm 5$ ) % of the nominal concentration. The 24h- $EC_{50}$  was  $> 0.38 \text{ mg/l}$ , the highest concentration tested. Although the study was performed with tetramethrin instead of d-trans-tetramethrin, it can be used for the effect assessment of d-trans-tetramethrin, as the available studies with fish and algae allow the conclusion, that the toxicity of both compounds is comparable.

#### 11.1.2 Acute (short-term) toxicity to algae or other aquatic plants

An algae growth inhibition test according to OECD 201 is available for d-trans-tetramethrin (Hoberg, 2002). Green algae *Pseudokirchneriella subcapitata* were exposed to 5 test concentrations (0.38, 0.75, 1.5, 3.0 and 6 mg/l nominal), a control and a solvent control (3 replicates each) over 72 h. DMF was used as a solvent. The solvent control chamber received an aliquot of 0.1 ml of DMF which was approximately equivalent to that present in the test solutions. Test substance concentration was measured at test start and end. At test start measured concentrations were in the range of 40 – 84 % of nominal. After 72 hours for all test concentrations the concentration was below the limit of detection. Due to the strong deviation between measured and nominal concentrations, the effect values have to be based on mean measured concentration. According to the OECD Guidance Document on Difficult Substances the detection limit should be used, if the concentration at test end is not detectable. Mean values of cell concentration ( $\times 10^4$  cells/ml) were measured at test start and after 24, 48 and 72 hours. The  $NOE_{RC}$  is 0.25 mg/l and the  $ER_{C50} > 1.25 \text{ mg/l}$ .

### 11.2 LONG-TERM AQUATIC HAZARD

No long term aquatic data are available for both d-trans-tetramethrin and tetramethrin.

### 11.3 BIOACCUMULATION

Table 41: Summary of relevant information on bioaccumulation

Method	Species	Results	Key or Supportive study	Remarks	Reference
OECD 305	Bluegill Sunfish ( <i>Lepomis macrochirus</i> )	827 L*kg <sub>wet fish</sub> <sup>-1</sup>	Key	Information is available for (1RS)-trans-tetramethrin as isomeric mixture, not for the single isomers. But, read across between the both trans-isomers is possible due to information on the optical trans-isomer ratios in the water and fish samples.	Saito S, Miyamoto M, Tagawa Y and Hagino S (1994); Report No. IM-40-0019
QSAR (according to Guidance on the Biocidal Products Regulation, Volume IV )	-	902 L*kg <sub>wet fish</sub> <sup>-1</sup>	Supportive	Log K <sub>ow</sub> : 4.3 (25°C)	Roth, H. (2006)

#### 11.3.1 Estimated bioaccumulation

Table 68: Estimations on aquatic bioconcentration

Basis for estimation	log P <sub>OW</sub> (measured)	Estimated BCF for fish (freshwater)	Reference
Standard equation (74), TGD on Risk Assessment (2003), Part II, chapter 3.8.3.2	4.3	902 L*kg <sub>wet fish</sub> <sup>-1</sup>	Roth, H. (2006)

Based on the physicochemical properties an approximate estimation of the bioconcentration factor BCF can be calculated according to Guidance on the Biocidal Products Regulation (2015, volume IV, chapter 3.8.3.2, equation 74, p. 144). Taking into account the log K<sub>OW</sub> value of 4.3 given by the company Sumitomo the BCF<sub>fish</sub> amounts to:

$$\text{Log BCF}_{\text{fish}} = 0.85 * \text{log K}_{\text{OW}} - 0.70$$

$$\text{Log BCF}_{\text{fish}} = 0.85 \cdot (4.3) - 0.70$$

$$\text{Log BCF}_{\text{fish}} = 2.955$$

$$\text{BCF}_{\text{fish}} = 902 \text{ L*kg}_{\text{wet fish}}^{-1}$$

K<sub>OW</sub> Octanol-water partition coefficient

BCF<sub>fish</sub> Bioconcentration factor for fish on wet weight basis

Summarised the calculated  $BCF_{fish}$  for D-trans-tetramethrin amounts to  $902 L \cdot kg_{wet\ fish}^{-1}$ , indicating the active substance as potentially bioaccumulative.

### 11.3.2 Measured partition coefficient and bioaccumulation test data

Table 69: Measurements of aquatic bioconcentration/depuration

Guideline/ Test method	Exposure [days]	Initial concentr. (nominal)	Measured Steady state $BCF_L^1$	Calculated Kinetics $BCF_L^1$	$T_{1/2}$ for clearance [days]	Identified Metabolites	Reference	
US EPA Subdivision N, § 165-4, Laboratory Studies of Pesticide Accumulation in Fish (1982)	28	1 µg/L	[Alc- <sup>14</sup> C]-(1RS)-trans-tetramethrin <sup>2</sup>				Whole fish and edible portion: THPI, HPI Non-edible portion: 1 Unknown	SUMI-TOMO: Saito, S., Miyamoto, M., Tagawa, Y. and Hagino, S. (1994); DocNo. IM-40-0019; Doc III A 7.4.3.3.1
			827	642	6.06			
			[Acid- <sup>14</sup> C]-(1RS)-trans-tetramethrin <sup>2</sup> :				Whole fish: CRA, COOH- CRA conj.-1 & conj.-2 Edible portion: CRA, CH <sub>2</sub> OH- CRA Non-edible portion: CRA, CRA conj., COOH-CRA, COOH-CRA conj.-1 & conj.-2	
			722	695	1.37			

<sup>1</sup> BCF values related to total radioactive residues, whole fish and a lipid content of 5%.

<sup>2</sup> 1R-trans-tetramethrin: This isomer is contained with 89 % in the active substance D-trans-tetramethrin.

In consequence of the log  $K_{OW}$  above 3 an experimental study with fish is required. A study on bioaccumulation in aquatic organisms is missing for the active substance d-trans-tetramethrin (isomeric ratio: 1R-trans-tetramethrin 89%, 1S-trans-tetramethrin 4%, 1R-cis-tetramethrin 2%, 1S-cis-tetramethrin 0.1%). Instead an aquatic bioaccumulation study with (1RS)-trans-tetramethrin (1:1 mixture of the both trans-isomers) was delivered by the company Sumitomo. An individual  $BCF_{fish}$  value for 1R-trans-tetramethrin is lacking. However, taking into account information on the optical trans-isomer ratios in the samples of the available study the BCF value derived for (1RS)-trans-tetramethrin can be used for the environmental risk and hazard assessment of d-trans-tetramethrin (explanation see below).

A dynamic 42-day study was conducted to evaluate the accumulation and the elimination of (1RS)-trans-tetramethrin as well as its metabolism in bluegill sunfish (*Lepomis macrochirus*). All test fish were from the same year class (less than 1 year of age) and held 8 months prior to testing. Their initial mean body weight was  $1.9 \pm 0.2$  g, they had an initial mean standard body length of  $4.1 \pm 0.1$

cm, and a mean fat content of 3.7%. The fish sampled during the study had a body weight and a standard body length of  $2.49 \pm 0.47$  g and  $4.4 \pm 0.2$  cm. Further information on weight, length and fat content are not available. Due to this a growth correction of the resulting BCF values is not possible (see Annex V of OECD 305). The lipid normalisation to 5% was related to the initial mean fat content of 3.7% as further information on fat content is not available. In the uptake phase 85 fish were exposed each to either [Alc- $^{14}\text{C}$ ]- or [Acid- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin at the concentration of 1  $\mu\text{g/L}$  under a flow-through test condition for 28 days at 25 °C. The measured concentrations for  $^{14}\text{C}$  and (1RS)-trans-tetramethrin in water were 0.899 – 1.10  $\mu\text{g/L}$  and 0.549 – 0.766  $\mu\text{g/L}$  for [Alc- $^{14}\text{C}$ ] preparation, and 0.922 – 1.10  $\mu\text{g/L}$  and 0.536 – 0.792  $\mu\text{g/L}$  for [Acid- $^{14}\text{C}$ ] preparation, respectively. On day 28 the fish were transferred to fresh running water to determine elimination rates. For analysis of the  $^{14}\text{C}$  compounds in fish, 3 fish for whole fish analysis and 3 additional fish for edible portion (body, muscle, skin and skeleton) and non-edible portion (fin, head and internal organs) analyses were randomly picked up from the treated aquaria on days 3, 7, 14, 21 and 28 of exposure. On days 1, 3, 7, 10 and 14 of the recovery, three fish for whole fish analysis and additional 3 fish for edible portion and non-edible portion analyses were randomly sampled from the treated and control aquaria.

For whole fish and total radioactivity the uptake rate and depuration rate constants were estimated with BIOFAC at  $54.2 \text{ L kg}^{-1} \text{ day}^{-1}$  in fish and  $0.114 \text{ day}^{-1}$  for [Alc- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin and  $260 \text{ L kg}^{-1} \text{ day}^{-1}$  and  $0.506 \text{ day}^{-1}$  for [Acid- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin, respectively.

The values of  $\text{BCF}_{\text{SSL}}$  and the kinetic  $\text{BCF}_L$  related to total radioactive residues, whole fish and a lipid content of 5% for [Alc- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin amounts to 827 and  $642 \text{ L*kg}_{\text{wet fish}}^{-1}$ , respectively.

The values of  $\text{BCF}_{\text{SSL}}$  and the kinetic  $\text{BCF}_L$  related to total radioactive residues, whole fish and a lipid content of 5% derived for [Acid- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin amounts to 722 and  $695 \text{ L*kg}_{\text{wet fish}}^{-1}$ , respectively.

A stable steady state concentration was reached in the study conducted with [Alc- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin, but not in the study conducted with [Acid- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin. As additionally the BCF values derived from the study conducted with [Alc- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin are higher than the BCF values derived for [Acid- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin the environmental risk and hazard assessment for d-trans-tetramethrin will be based on the BCF values determined for [Alc- $^{14}\text{C}$ ]- (1RS)-trans-tetramethrin as a worst-case approach.

The  $\text{BCF}_{\text{SSL}}$  results with  $827 \text{ L*kg}_{\text{wet fish}}^{-1}$  in a higher value than the  $\text{BCF}_{\text{KL}}$  with  $642 \text{ L*kg}_{\text{wet fish}}^{-1}$ . Due to this the  $\text{BCF}_{\text{SSL}}$  of  $827 \text{ L*kg}_{\text{wet fish}}^{-1}$  related to total radioactive residues, whole fish and lipid content of 5% derived for (1RS)-trans-tetramethrin will be used as a worst case in the environmental risk and hazard assessment.

For edible portion and total radioactivity the uptake and depuration rate constants were estimated at  $34.0 \text{ L kg}^{-1}$  and  $0.09 \text{ day}^{-1}$  for [Alc- $^{14}\text{C}$ ] preparation as well as  $46.5 \text{ L kg}^{-1}$  and  $0.270 \text{ day}^{-1}$  for [Acid- $^{14}\text{C}$ ] preparation, respectively. For BCF in edible portion a steady state was only reached in the approach with [Alc- $^{14}\text{C}$ ] preparation. Moreover, the derived BCF values for the [Alc- $^{14}\text{C}$ ] preparation are higher than that derived for the [Acid- $^{14}\text{C}$ ] preparation, so the results from the [Alc- $^{14}\text{C}$ ] preparation are used for BCF derivation related to edible portion. The  $\text{BCF}_{\text{KL}}$  is with  $511 \text{ L*kg}_{\text{wet fish}}^{-1}$  higher than the  $\text{BCF}_{\text{SSL}}$  with  $453 \text{ L*kg}_{\text{wet fish}}^{-1}$  and can be used as the worst-case value for edible portion. Both BCF values are normalised to a fat content of 5%.

*For non-edible portion and total radioactivity the uptake and depuration rate constants were estimated at  $99.1 \text{ L kg}^{-1}$  and  $0.144 \text{ day}^{-1}$  for [Alc- $^{14}\text{C}$ ] preparation as well as  $424 \text{ L kg}^{-1}$  and  $0.380 \text{ day}^{-1}$  for [Acid- $^{14}\text{C}$ ] preparation, respectively. Regarding BCF in non-edible portion steady state*

was only reached in the approach with [Acid-<sup>14</sup>C] preparation. As additionally the resulting BCF values for the [Acid-<sup>14</sup>C] preparation are higher than that derived for the [Alc-<sup>14</sup>C] preparation, the BCF derivation for non-edible portion is based on the results derived for [Acid-<sup>14</sup>C] preparation. The  $BCF_{KL}$  is with  $1508 \text{ L} \cdot \text{kg}_{\text{wet fish}}^{-1}$  higher than the  $BCF_{SSL}$  with  $1338 \text{ L} \cdot \text{kg}_{\text{wet fish}}^{-1}$  and represents the worst-case value for non-edible portion. Both BCF values are normalised to a fat content of 5%.

Despite the study on bioaccumulation was conducted with (1RS)-trans-tetramethrin (1:1 mixture of the both trans-isomers) the derived BCF values can be used for the active substance d-trans-tetramethrin containing mainly the 1R-trans-isomer (isomeric ratio: 1R-trans-tetramethrin 89%, 1S-trans-tetramethrin 4%, 1R-cis-tetramethrin 2%, 1S-cis-tetramethrin 0.1%), if taking the available information on optical trans-isomer ratios in the samples into account for read-across between the both trans-isomers. Measurement of the ratios of the both isomers R-trans-tetramethrin and S-trans-tetramethrin verifies that they remain at ca. 1/1 ratio in water and fish during the study (see table 4-14). This means that the BCF value of  $827 \text{ L} \cdot \text{kg}_{\text{wet fish}}^{-1}$  derived for (1RS)-trans-tetramethrin is also valid for the single trans-isomers.

Table 70: Optical isomer ratio in the samples

Label position	Sample	R/S ratio
[Acid- <sup>14</sup> C]	Water (exposure 0 days)	53 / 47
	Water (exposure 28 days)	48 / 52
	Fish, non edible (exposure 21 days)	54 / 46
	Fish, edible (exposure 21 days)	51 / 49
	Fish, whole (exposure 14 days)	40 / 60
	Fish, whole (exposure 28 days)	46 / 54
[Alc- <sup>14</sup> C]	Fish, non-edible (exposure 21 days)	53 / 47
	Fish, whole (exposure 28 days)	52 / 48

For the reasons described above, the BCF values derived for (1RS)-trans-tetramethrin (1:1 mixture of the both trans-isomers) are also valid for the active substance d-trans-tetramethrin.









The half-life for clearance related to total radioactive residues and whole fish amounts to 6.06 days and 1.37 days for [Alc-<sup>14</sup>C]- (1RS)-trans-tetramethrin and [Acid-<sup>14</sup>C]- (1RS)-trans-tetramethrin, respectively. The time to reach 90% of steady-state related to total radioactivity was determined to be 20.1 days for [Alc-<sup>14</sup>C]- (1RS)-trans-tetramethrin and 4.55 days for [Acid-<sup>14</sup>C]- (1RS)-trans-tetramethrin.

For edible portion half-life for clearance related to total radioactive residues was determined to be 7.70 days and 2.57 days for [Alc-<sup>14</sup>C] and [Acid-<sup>14</sup>C] preparation, respectively. The time to reach 90% of steady-state related to total radioactivity was determined to be 25.6 days for [Alc-<sup>14</sup>C]- (1RS)-trans-tetramethrin and 8.52 days for [Acid-<sup>14</sup>C]- (1RS)-trans-tetramethrin.

*For non-edible portion half-life for clearance related to total radioactive residues was determined to be 4.81 days and 1.82 days for [Alc-<sup>14</sup>C] and [Acid-<sup>14</sup>C] preparation, respectively. The time to reach 90% of steady-state related to total radioactivity was determined to be 16.0 days for [Alc-<sup>14</sup>C]- (IRS)-trans-tetramethrin and 6.06 days for [Acid-<sup>14</sup>C]- (IRS)-trans-tetramethrin.*

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Table 71: List of metabolites found in bioaccumulation study with (1R)-trans-tetramethrin in whole fish, edible and non-edible portion:

Name of compound used in reports	Structural formula	Maximum concentration <sup>1</sup>	Concentration at end of uptake phase (day 28) <sup>1</sup>	Concentration at end of depuration phase (day 7 or day 10) <sup>1</sup>	Related to:
<b>THPI</b> (3,4,5,6-tetrahydrophthalimide)		13.1 % (day 3)	2.3%	0.3%	<b>Whole fish</b>
		22.0% (day 3)	3.3%	0.2%	<b>Edible portion</b>
		-	-	-	<b>Non-edible portion</b>
<b>HPI</b> (cyclohexane-1,2-dicarboxyimide)		13.0 % (day 3)	2.4%	n.d.	<b>Whole fish</b>
		20.8% (day 3)	3.6%	n.d.	<b>Edible portion</b>
		-	-	-	<b>Non-edible portion</b>
<b>CRA</b> (1R, 3R)-2,2-dimethyl-3-(2-methylprop-1-enyl)cyclopropane-carboxylic acid)		20.1% (day 3)	8.0%	n.d.	<b>Whole fish</b>
		50.4% (day 3)	11.0%	n.d.	<b>Edible portion</b>
		16.1% (day 3)	5.9%	n.d.	<b>Non-edible portion</b>
<b>CRA conj.</b>		-	-	-	<b>Whole fish</b>
		-	-	-	<b>Edible portion</b>
		11.2% (day 7)	4.5%	n.d.	<b>Non-edible portion</b>
<b>COOH-CRA</b> (1R, 3R)-2,2-dimethyl-3-(E-2-carboxyprop-1-enyl)cyclopropane-carboxylic acid)		-	-	-	<b>Whole fish</b>
		-	-	-	<b>Edible portion</b>
		10.6% (day 7)	5.3%	3.4%	<b>Non-edible portion</b>
<b>COOH-CRA conj.-1</b>		15.6% (day 7)	5.2%	n.d.	<b>Whole fish</b>
		-	-	-	<b>Edible portion</b>
		21.1% (day 7)	8.1%	n.d.	<b>Non-edible portion</b>
<b>COOH-CRA conj-2</b>		10.8% (day 7)	4.9%	n.d.	<b>Whole fish</b>
		-	-	-	<b>Edible portion</b>
		11.3% (day 3)	7.8%	n.d.	<b>Non-edible portion</b>
<b>CH<sub>2</sub>OH-CRA</b> (1R, 3R)-2,2-dimethyl-3-(E-2-hydroxymethylprop-1-enyl)cyclopropanecarboxylic acid)		-	-	-	<b>Whole fish</b>
		14.6% (day 7, inclusive unknown metabolite A)	9.4%	n.d.	<b>Edible portion</b>
		-	-	-	<b>Non-edible portion</b>
<b>Unknown</b> (14 unidentified spots pooled)	-	-	-	-	<b>Whole fish</b>
		-	-	-	<b>Edible portion</b>
		10.1% (day 3 of depuration phase, max. of single 14 unidentified spots)	Not applicable	3.3%	<b>Non-edible portion</b>



The metabolites in whole fish, edible and non-edible portions were analysed by TLC for the fish samples on days 3, 7, 14, 21, and 28 of exposure (see table 4-15). Metabolites detected for both  $^{14}\text{C}$  preparations in whole fish throughout the uptake phase with  $\geq 10\%$  of total radioactivity are THPI (max. 13.1%, day 3), HPI (max. 13.0%, day 3), CRA (max. 20.1%, day 3) and COOH-CRA conj.-1 (max. 15.6%, day 7) and conj.-2 (max. 10.8%, day 7). All metabolites detected with  $\geq 10\%$  in whole fish decreased to levels between 2.3 and 8.0 at the end of uptake phase and 0.3% and not detectable between day 7 and day 10 of depuration phase. In edible portion 4 metabolites  $\geq 10\%$  were detected for both  $^{14}\text{C}$  preparations during the uptake phase, THPI (max. 22.0%, day 3), HPI (max. 20.8%, day 3), CRA (max. 50.4%, day 3) and  $\text{CH}_2\text{OH-CRA}$  with unknown metabolite A (max. 14.6%, day 7). At the end of the uptake phase all metabolites were observed with 3.3 – 11.0%. After 7 – 10 days of depuration levels of these metabolites decreased to 5.3% and not detectable in edible portion. In non-edible portion detected metabolites  $\geq 10\%$  during the uptake phase are Unknown (maximum value of the single 14 unidentified spots pooled in Unknown is 10.1%, day 3 of depuration phase), CRA (max. 16.1%, day 3), CRA conj. (max. 11.2%, day 7), COOH-CRA (10.6%, day 7), COOH-CRA conj.-1 (21.1%, day 7) and COOH-CRA conj.-2 (11.3%, day 3) for both  $^{14}\text{C}$  preparations. All these metabolites decrease to levels between 4.5 – 8.1% at the end of uptake phase and 3.4% and not detectable after 7 – 10 days of depuration. Based on these results further investigation on the bioaccumulation behavior of the observed metabolites seems not to be necessary.

Information about metabolites detected in the bioaccumulation study conducted with (1RS)-trans-tetramethrin may also be used for the assessment of the active substance d-trans-tetramethrin as it is not expected that the metabolisation process will result in building of different metabolites for the single isomers. However, it cannot be excluded that the built amounts will be different for d-trans-tetramethrin.

Despite the study on bioaccumulation was conducted with (1RS)-trans-tetramethrin (1:1 mixture of the both trans-isomers) the derived BCF values can be used also to assess the bioaccumulation behaviour of the cis-isomers contained in the active substance d-trans-tetramethrin, if taking the available information on toxicokinetics of the cis- and trans-isomers in rat into account for read-across between cis- and trans-isomers. Toxicokinetics of the cis- and trans-isomers of tetramethrin in rat were almost identical. For details see chapter 3.2 “Toxicokinetics, Metabolism and Distribution”. Minor differences relating to the relative amount of radiocarbon metabolites excreted via urine and the initial elimination velocity were observed for the trans- and cis-isomers. Very rapid systemic and/or pre-systemic metabolic degradation by oxidation and ester hydrolysis has been shown which is slower for cis- than for trans-isomers. However, for cis- as well as trans-isomers a rapid elimination from the body could be observed with excretion of  $>94\%$  of the dosed  $^{14}\text{C}$  into faeces and urine within 2-7 days. As only observed difference trans-isomers are more excreted via urine (trans: 42 – 74%, cis: 9 – 49%) and cis-isomers more via faeces (trans: 21 – 58%, cis: 45 – 91%). For cis- and trans-isomers the remaining  $^{14}\text{C}$  tissue residues were widely distributed with highest concentrations in blood cells and generally low, with 0.2 – 0.4% after 7 days.

The major metabolic pathway of [ $\text{Alc-}^{14}\text{C}$ ]-trans-tetramethrin was confirmed to be common between fish and rat. Following cleavage of the ester linkage, the liberated alcohol moiety was further transformed by N-dealkylation, followed by reduction of the 1,2-double bond and/or cleavage of the imide linkage. Taking the information on similarity of rapid elimination from the body and low remaining  $^{14}\text{C}$  tissue residues of cis- and trans-isomers in rat as well as the common major metabolic pathway shown for the trans-isomer in rat and fish into account, read-across between cis- and trans-isomers related to bioaccumulation behaviour in fish seems reasonable.

Additionally, the similar  $K_{OC}$  values of the cis- (2045 mL/g) and trans-isomers (2754 mL/g) may indicate, that BCF values of the trans- and cis-isomers might also be similar. QSAR calculation related to single isomers for support of this conclusion is not possible, as SMILES codes are unable to indicate isomerism. More complex structure activity investigations were not possible as well, due to non availability of suitable tools.

In a publication of Corcellas et al. (Environment International 75 (2015) 110–116) bioaccumulation of 12 pyrethroids including tetramethrin were investigated in wild river fish collected in 4 different Iberian rivers. Pyrethroids were detected in all analysed samples; detection frequency of tetramethrin was 83%. Evaluation of the enantiomeric contribution of tetramethrin was not possible, but in general an accumulation preference of the cis-isomers could be observed according to this publication. Levels of pyrethroids was compared with those of other pollutants like flame retardants, personal care products, hormones and pharmaceuticals. Pyrethroids were detected more frequently in the samples than most other compounds and were also found in higher concentrations. The results of this publication should be kept in mind in future in case of renewal of the authorization procedure. Until then maybe further information might be available on monitoring data regarding tetramethrin and its isomers that should then be taken into account for the environmental risk and hazard assessment.

Summarised measured data on bioaccumulation behaviour in fish are only available for (1RS)-trans-tetramethrin. Taking the available information on optical trans-isomer ratios in the samples into account read-across between the both trans-isomers related to bioaccumulation behaviour in fish seems reasonable. Considering the information on similarity of cis- and trans-isomers regarding  $K_{OC}$  values, rapid elimination from the body and low remaining  $^{14}C$  tissue residues in rat as well as the common major metabolic pathway shown for the trans-isomer in rat and fish into account, read-across between cis- and trans-isomers related to bioaccumulation behaviour in fish seems also to be acceptable. Therefore, a  $BCF_{SS}$  of  $827 \text{ L} \cdot \text{kg}_{\text{wet fish}}^{-1}$  related to total radioactive residues, whole fish and lipid content of 5% will be used as a worst case in the environmental risk and hazard assessment for d-trans-tetramethrin. Based on this result d-trans-tetramethrin has to be considered as bioaccumulative.

A test with an appropriate invertebrate species (Doc III-A 7.4.3.3.2) was not required because no direct release to marine/brackish waters occurs. If once the application scenario is changing and release to marine/brackish water occurs an accumulation study with invertebrates might be necessary.

## 11.4 RAPID DEGRADABILITY OF ORGANIC SUBSTANCES

Table 72: Summary of relevant information on rapid degradability

Method	Results	Key or Supportive study	Remarks	Reference
OECD 301 F	Not readily biodegradable (27% degradation within 28 days)	Key	None	Grützner, I. (2002); DocNo SVM-0001

### 11.4.1 Ready biodegradability

Table 73: Ready biodegradability tests

Method /Guide-line	Test type <sup>1</sup>	Test parameter	Inoculum			Additional substrate	Test substance conc.	Degradation		Reference
			Type	Conc. <sup>3</sup>	Adap-tation			Incuba-tion period	Degree [%]	
OECD 301 F	ready	BOD <sup>2</sup>	Acti-vated sludge - pre-dominantly do-mestic	30 mg/L dw	No	No	104 & 100 mg/L	28 days	27	SUMITOMO: Grützner, I (2002); DocNo SVM-0001; Doc IIIA 7.1.1.2.1

<sup>1</sup> Test on inherent or ready biodegradability according to OECD criteria

<sup>2</sup> Biochemical Oxygen Demand

<sup>3</sup> Suspended solid concentration

A test on ready biodegradability of d-trans-tetramethrin was submitted from the company Sumitomo conducted according to OECD testguideline 301 F investigating ready biodegradability by measurement of biochemical oxygen demand. In this test a biodegradation of 27 % within 28 days was observed. The beginning of relevant biodegradation processes could be observed after 15 days. This delayed degradation may be a sign of adaption procedures.

Based on this result d-trans-tetramethrin can be considered to be not readily biodegradable.

### 11.4.2 BOD<sub>5</sub>/COD

BOD<sub>5</sub>/COD tests are not available.

### 11.4.3 Other convincing scientific evidence

Aquatic simulation tests are not available.

#### 11.4.3.1 Field investigations and monitoring data (if relevant for C&L)

Field investigations and monitoring data are not available.

#### 11.4.3.2 Inherent and Enhanced Ready Biodegradability tests

Inherent and enhanced biodegradability test data are not available.

#### 11.4.3.3 Soil and sediment degradation data

In a laboratory study (OECD 307) the route and rate of aerobic degradation of [<sup>14</sup>C]-tetramethrin in three soils was investigated using a radiolabel at the alcohol moiety of the molecule. As also the degradation of the single isomers (1R-trans-tetramethrin: contained with 89 % in d-trans-tetramethrin) was determined the results of this study can be used for the assessment of the active substance d-trans-tetramethrin. The first-order DT<sub>50</sub> values (12°C) were calculated to be 4.0 – 8.0 days for the isomeric mixture tetramethrin, 4.4 – 7.6 days for the isomer 1R-trans-tetramethrin (contained with 89 % in the active substance d-trans-tetramethrin), 2.5 – 4.9 days for 1S-trans-tetramethrin, 15.2 – 36.4 days for 1R-cis-tetramethrin and 13.3 – 21.6 days for the isomer 1S-cis-tetramethrin. Mineralisation of the active substance tetramethrin accounted for a maximum of 57.6 % AR (applied radioactivity) for the silt loam soil (day 30), 33.3 % AR for the loam soil (day 62) and 40.7 % AR for the sandy clay loam soil (day 30). These mineralisation values can also be used for assessment of the active substance d-trans-tetramethrin as the degradation kinetics are very similar between the isomer 1R-trans-tetramethrin (contained with 89 % in d-trans-tetramethrin) and the isomeric mixture tetramethrin containing additionally the slower degradable cis-isomers. Therefore, a similar carbon dioxide evolution is expectable.

In a second laboratory study (US-EPA 162-1) the route and rate of aerobic degradation of [<sup>14</sup>C]- (1RS)-trans-tetramethrin (1:1 mixture of the both trans-isomers) labeled at the alcohol as well as on the acid moiety of the molecule was investigated. As the DT<sub>50</sub> values determined for both trans-isomers of tetramethrin in the first soil simulation study are very close together (4.4 – 7.6 days and 2.5 – 4.9 days) the results of the study conducted with (1RS)-trans-tetramethrin can be used for the assessment of the active substance d-trans-tetramethrin.

In the sandy loam soil (1RS)-trans-tetramethrin undergoes degradation under aerobic dark conditions with first order DT<sub>50</sub> values of 35.6 – 50.6 days (12°C). Carbon dioxide evolution amounts to 49.6% and 49.8% after 122 days of incubation and to 69.8% and 65.0% after 365 days of incubation for [THP-<sup>14</sup>C]- (1RS)-trans-tetramethrin and [Cyclopropyl-<sup>14</sup>C] – (1RS)-trans – tetramethrin, respectively. A similar mineralisation rate is expectable for the active substance d-trans-tetramethrin compared to (1RS)-trans-tetramethrin due to the similar degradation kinetics.

Although the results of the simulation studies demonstrate a rapid to moderate primary degradation, the ultimate degradation has to be considered as rather low due to mineralisation rates below 60% (33.3 – 57.6%) after 30 to 122 days of incubation and below 70% (65.0 – 69.8%) after 365 days of

incubation. This supports the classification met by the screening test that d-trans-tetramethrin must be considered as not readily biodegradable.

#### 11.4.3.4 Hydrolysis

Table 74:

Method /Guideline	pH	Temperature [°C]	Initial concentration, C <sub>0</sub> , [µg/L]	Reaction rate constant, K <sub>h</sub> [days <sup>-1</sup> ]	Half-life, DT <sub>50</sub>	Coefficient of correlation, r <sup>2</sup>	Reference	Key or Supportive study
Trans-Neo-Pynamin [3,4,5,6-tetrahydrophthalimidomethyl (1RS)-trans-chrysanthemate or (1RS)-trans-tetramethrin] US EPA Pesticide Assessment Guidelines Subdivision N 161-1	5	25	300	$3.514 \times 10^{-2} - 4.364 \times 10^{-2}$	15.9-19.7 days	n.s.	Katagi T. et al, Sumitomo Report No: IM-10-0012, 1991	Key
	7	25	300	0.653 – 0.778	21.4-25.5 hrs	n.s.		
	9	25	300	44.2 – 75.6	13.2-21 min	n.s.		

A definitive hydrolysis study on degradation products and kinetics was conducted with (1RS)-trans-tetramethrin at pH 5, 7 and 9 at 25 °C according to US EPA N 161-1. The temperature dependence of hydrolysis has not been determined in this study. Information is only available for (1RS)-trans-tetramethrin, information on the single (1R)-trans-isomer is lacking. The eCA does not expect enantiomerism to affect significantly the hydrolysis of the parent compound under environmental conditions and, hence, considers the data as sufficient to characterise the route and kinetic of degradation and the mayor degradation products of the (1R)-trans-isomer.

The mayor degradation products of (1RS)-trans-tetramethrin are (1RS)-trans-crysamtemic acid (trans-CRA) and 3,4,5,6-tetrahydrophthalamic acid (THAM), that is finally hydrolysed to 3,4,5,6-tetrahydrophthalic acid (THPA). Trans-CRA increased to 68.85% of the applied radioactivity at pH 5, 98.42% at pH 7 and 100% at pH 9 at day 30. THMA peaked after one day to 15.43% at pH7 and to 80.6% at pH9. THPA increased to 66.07% at pH5, 95.78% at pH 7, and 68.07 % at pH9. The degradation products trans-CRA and THPA are assumed to be hydrolytic stable.

The hydrolysis half-lives of (1RS)-trans-tetramethrin were recalculated to reflect an average EU outdoor temperature of 12°C for fresh water (based on EU TGD (2003), chapter 2.3.6.1). The half-lives amount to 45.0 – 55.7 days at pH 5, 66.5 – 72.1 hours at pH 7, and 37.3 – 59.4 min at pH 9.

### 11.4.3.5 Photochemical degradation

Photolysis in water

Table 75:

Method /Guideline	Initial molar TS concentration	Total recovery of test substance [% of appl. a.s.]	Photolysis rate constant ( $k_p^c$ )	Direct photolysis sunlight rate constant ( $k_{pE}$ )	Reaction quantum yield ( $\Phi_E^c$ )	Half-life $DT_{50}$ [days]	Reference	Key or Supportive study
US EPA OPPTS 835.2210	0.2 $\mu\text{g/mL}$	95.6 $\pm$ 7.0%	0.147 $\text{h}^{-1}$	n.s.	0.19	0.46 US summer days; 0.38 Global summer days	Lopez A., PTRL Report No. 1762W-1, 2003	Key

(1R)-trans-tetramethrin undergoes photodegradation in aqueous media at pH 5. A degradation rate constant of 0.147  $\text{hours}^{-1}$  and a half-life of  $DT_{50} = 0.46$  days for a US summer day was determined. Photo-induced isomerisation to the cis-isomer was minor in light exposed samples. The other main degradation products observed were not known and were assigned as D-1, D-3 and D-6. Based on LC/MS data proposed structures are given (see confidential annex *d-trans-Tetramethrin\_14.4.17\_03*). D-1 represented an average of 24.1% of the initial dose at the end of the irradiation period (304 hours). D-3 and D-6 reached after 144 hours of irradiation 32.2% and 20.1% of the applied dose, respectively.

(1R)-trans-tetramethrin degraded significantly in dark control samples. After 312 hours of incubation at 25°C, 47% of the initial dose occurred in the non-irradiated samples. The major degradation product was THPA, reaching an average of 55.7% of the applied dose at the end of the incubation period. The corresponding half-life in dark controls was 12.3 days of incubation. Since (1R)-trans-tetramethrin degraded more rapidly when exposed to light, its hydrolytic degradation without irradiation had no significant effect on the light exposed set.

Indirect photolysis in water bodies of the active substance has not been measured. However, information on indirect photolysis is not regarded to be scientifically necessary as other degradation process (hydrolysis, direct photolysis) are not regarded to be slow.

#### Phototransformation in air

Table 76:

Guideline / Test method	Time-dependent OH radical concentration [OH radicals $\text{cm}^{-3}$ ]	Overall reaction rate constant $k$ [ $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ ]	Half-life $DT_{50}$ [h]	Chemical lifetime [h]	Reference	Key or Supportive study
AOPWIN calculation v. 1.92	5x10 <sup>5</sup> 24 h average	127.3092 E <sup>-12</sup>	3.025	4.36	(CA's estimation)	Key

(1R)-trans-tetramethrin is expected to degrade in the atmosphere by reaction with photochemically-produced hydroxyl radicals. The half-life of (1R)-trans-tetramethrin in air is estimated to be 3.025 h using the generally accepted estimation program AOPWIN, version 1.92. Referring to the vapour pressure of  $4.03 \times 10^{-6}$  Pa and a Henry's Law constant of  $0.000112 \text{ Pa m}^{-3} \text{ mol}^{-1}$  (1R)-trans-tetramethrin is not expected to volatilize. Therefore, emissions to air are expected to be low. In conclusion, due to the low volatility, the fast degradation by OH radicals in air, the hydrolysis in the presence of water, accumulation and long range transport in air are not to be expected under environmentally relevant conditions.

## 11.5 ENVIRONMENTAL FATE AND OTHER RELEVANT INFORMATION

### 11.5.1 Adsorption/desorption

Table 77: Adsorption/desorption in five soils

Method /Guideline	Tested Soils	Adsorbed a.s. [%]	$K_a^1$	$K_{aOC}^2$	$K_d^3$	$K_{dOC}^4$	$K_a / K_d^5$	Degradation products	Reference	Key or Supportive study
US EPA Pesticide Assessment Guidelines Subdivision N 163-1	A13 = Sandy loam	64.4	11.47	1289	13.61	1529	0.84	none	Yoshimura J., Sumitomo Report No: IM-10-0013, 1991	Key
	B7 = Loamy sand	59.7	12.03	2933	12.99	3167	0.93			
	F6 = Sandy clay loam	59.9	19.27	1407	21.06	1537	0.92			
	Hanford = Sandy Loam	47.2	8.54	1857	8.03	1746	1.06			
	J11= Clay	59.3	10.71	1552	11.92	1728	0.90			

<sup>1)</sup>  $K_a$  = Adsorption coefficient, mean value of duplicate measure

<sup>2)</sup>  $K_{aOC}$  = Adsorption coefficient based on organic carbon content, mean value of duplicate measure

<sup>3)</sup>  $K_d$  = Desorption coefficient, mean value of duplicate measure

<sup>4)</sup>  $K_{dOC}$  = Desorption coefficient based on organic carbon content, mean value of duplicate measure

<sup>5)</sup>  $K_a / K_d$  = Adsorption / Desorption distribution coefficient, mean value of duplicate measure

A study on adsorption/ desorption of (1RS)-trans-tetramethrin in five different soils was conducted according to EPA 163-1. Information is only available for (1RS)-trans-tetramethrin, information on the single (1R)-trans-isomer is lacking. The eCA does not expect enantiomerism to affect significantly the sorption behaviour of the parent compound in soils and, hence, considers the data as sufficient to characterise the adsorption/desorption behaviour of the (1R)-trans-isomer. Nevertheless, deviations from the OECD guideline 106 were determined regarding reporting on soil sampling and use of standard methods for soil characterization. Deficiencies were further determined regarding soil classification. Moreover, deficiencies were determined regarding the range of the five different soils as the selection does not comprises acidic soil types with a pH <6.5, soil types of high OC content >1.5% and the soil types do not comprise five different soils according to the OECD guideline 106.

Radiolabelled (1RS)-trans-tetramethrin was stable in the test system during the study as more than 99 % was identified to be (1RS)-trans-tetramethrin at the end of the study. The adsorption equilibrium coefficient  $K_a$  ranged from 8.54 to 19.32 mL/g and the desorption equilibrium coefficient  $K_d$  ranged from 8.03 to 21.06 mL/g. The  $K_{aoc}$  value was calculated to range from 1289 to 2933 mL/g and the  $K_{doc}$  value was calculated to range from 1529 to 3167 mL/g.  $K_{oc}$  values did not correlate with the OC content. The arithmetic mean  $K_{oc}$  is 1807 mL/g, with a mean  $\log K_{oc}$  of 3.26 mL/g. From the average  $K_{oc}$  values for adsorption and desorption it can be concluded that (1RS)-trans-tetramethrin is considered of low mobility in four types of soil and as slight mobility in one soil type tested.

## 11.6 COMPARISON WITH THE CLP CRITERIA

### 11.6.1 Acute aquatic hazard

For d-trans-tetramethrin an acute aquatic endpoint for fish is available, for invertebrates a read-across to Tetramethrin was accepted. For algae a test according to OECD 201 was done and an  $ErC_{50}$  and a  $NOErC$  were derived. The most sensitive endpoint is a  $LC_{50} = 5.9 \mu\text{g/l}$  for *Oncorhynchus mykiss* (cf. chapter 11.1.1). However, the classification of d-trans-tetramethrin is based on the 96h- $LC_{50}$  of  $3.7 \mu\text{g/L}$  for tetramethrin, as the ecotoxicological equivalence of both compounds was accepted. A substance has to be classified as “Aquatic Acute 1; **H400**”, if the  $LC_{50}/EC_{50}$  is  $\leq 1 \text{ mg/l}$ . This criterium is fulfilled for d-trans-tetramethrin. The corresponding **M-factor is 100**.

### 11.6.2 Long-term aquatic hazard (including bioaccumulation potential and degradation)

D-trans-tetramethrin has shown a biodegradation of only 27% in 28 days in a test according to OECD guideline 301 F and has therefore to be regarded as not readily biodegradable (cf. chapter 11.4.1). With an estimated  $BCF_{fish}$ , based on the  $\log K_{ow} = 4.3$ , of  $902 \text{ L*kg}_{wet\ fish}^{-1}$  and a measured  $BCF_{fish}$  value of  $827 \text{ L*kg}_{wet\ fish}^{-1}$  (cf. chapter 11.3) d-trans-tetramethrin is considered to be bioaccumulative.

The “Guidance on the Application of the CLP Criteria” (ECHA, 2013) gives a decision tree for the decision on categories for substances long-term hazardous to the aquatic environment (Figure 4.1.1, p. 524). According to this decision tree, if adequate chronic toxicity data are only available for one or two trophic levels, the chronic classification should be done considering both the available chronic endpoints and also, as a surrogate system, considering the acute endpoints. The most stringent outcome should be taken for the classification. In the case of d-trans-tetramethrin, long-term aquatic toxicity data are only available for algae, but algae are not the most sensitive organisms in acute toxicity tests. Taking into account the lowest available acute  $LC_{50}$   $3.7 \mu\text{g/L}$  (*Oncorhynchus mykiss*) for the ecotoxicological equivalent substance tetramethrin (read-across was accepted by the Competent Authority), d-trans-tetramethrin has to be classified as “Aquatic Chronic 1; H410.

As there are no chronic effect data available, the acute M-factor of 100 has to be applied for the chronic category as well.



## 11.7 CONCLUSION ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS

D-trans-tetramethrin should be classified as “Aquatic Acute 1; H400” – “Very toxic to aquatic organisms” (M = 100) and “Aquatic Chronic 1; H410” – “Very toxic to aquatic organisms with long lasting effects” (M = 100) for the environment. This leads to a proposed labelling of H410 (Very toxic to aquatic life with long lasting effects), which triggers the pictogram GHS09 and the signal word “Warning” on the label. The following precautionary statements are indicated: P273, P391 and P501.

## 12. EVALUATION OF ADDITIONAL HAZARDS

### 12.1 HAZARDOUS TO THE OZONE LAYER

According to Regulation EC (No) 1272/2008 a substance has to be considered as hazardous to the ozone layer if it is listed in Regulation EC (No) 2037/2000. As this is not the case for d-trans-tetramethrin, no additional labelling is necessary.

## 13. DETAILED STUDY SUMMARIES

See confidential Annexes

## 14. REFERENCES

Author(s)	Year	Title. Source (where different from company) Company, Report No. GLP (where relevant) / (Un)Unpublished	Owner
Blasberg	1993	Acute flow-through toxicity of Neo-Pynamin to <i>Daphnia Magna</i> , ABC Laboratories, Inc. Sumitomo Report No. IW-31-0017	Sumitomo Chemical Co., Ltd
Evans A.J. et al	2002	Final S-1723: Determination of Melting/Freezing Temperature, Safepharm Laboratories Ltd, SPL Project No: 1430/011, GLP, (Un)	Sumitomo Chemical Co., Ltd
Grützner I	2002	Ready biodegradability of S-1723 in a manometric respirometry test, Environmental Chemistry and Pharamalytics Division, Sumitomo Report No. SVM-0001, GLP, (Un)	Sumitomo Chemical Co., Ltd
Hohberg	2002	Toxicity to the Freshwater Green Alga ( <i>Pseudokirchneriella subcapitata</i> ), Springborn Smithers Laboratories, Sumitomo Report No. SVW-0002	Sumitomo Chemical Co., Ltd
Katagi	1991	Hydrolysis of trans-Neopynamin in buffered aqueous solution, Environmental health science laboratory, Sumitomo Report No. IM-10-0012, 1991	Sumitomo Chemical Co., Ltd
Lentz NR	2008	Final Neo-Pynamin Forte PAI – Determination of the Density of a Test Substance Following OECD Guideline 109 and the Official Journal of the European Communities, L383A, Method A.3, Springborn Smithers Laboratories, Study No. 13048.6572, Sumitomo Report No. SVP-0020, GLP, (Un)	Sumitomo Chemical Co., Ltd
Lentz NR	2008	Final Neo-Pynamin Forte PAI – Determination of the Water Solubility of a Test Substance Following OECD Guideline 105 and the Official Journal of the European Communities, L383A, Method A6, Springborn Smithers Laboratories, Study No. 13048.6571, GLP, (Un)	Sumitomo Chemical Co., Ltd

CLH REPORT FOR D-TRANS-TETRAMETHRIN

Lentz NR	2008	Final Neo-Pynamin Forte PAI – Determination of Surface Tension using a Ring Tensiometer Following OECD Guideline 115, Springborn Smithers Laboratories, Study No. 13048.6593, GLP, (Un)	Sumitomo Chemical Co., Ltd
Lopez	2009	Phototransformation of [14C] Tetramethrin in Sterile Water Buffered at pH5. PTRL West, Inc., 625-B Alfred Nobel Drive, Hercules, CA 94547. PTRL Report No. 1762W-1	Sumitomo Chemical Co., Ltd
Malinski M.F.	1999	Final S-1723 - Boiling Point, Analytical Services, Sumitomo Report No. SVP-0003, GLP, (Un)	Sumitomo Chemical Co., Ltd
O'Donnell R.T. et al	1999	Final S-1723 - Colour, Physical State, Density and Solvent Solubility, Analytical Services, Sumitomo Report No. SVP-0008, GLP, (Un)	Sumitomo Chemical Co., Ltd
Poncipe	2006	Tetramethrin. Estimation of indirect photolysis in air, Safepharm laboratories	Sumitomo Chemical Co., Ltd
Roth H	2006	Safepharm Laboratories Ltd, , (Un)	Sumitomo Chemical Co., Ltd
Saito S, Miyamoto M, Tagawa Y and Hagino S	1994	Bioconcentration and metabolism of 14C-trans-Neo-Pynamin in Bluegill Sunfish ( <i>Lepomis macrochirus</i> ), Environmental Health Science Laboratory, Sumitomo Report No. IM-40-0019, GLP, (Un)	Sumitomo Chemical Co., Ltd
Schetter J.E.	1999	Final S-1723 - Vapour Pressure, Analytical Services, Sumitomo Report No. SVP-0006, GLP, (Un)	Sumitomo Chemical Co., Ltd
Sousa and LeBlanc	1981	Acute toxicity of Neo-Pynamin Forte T.G. to Bluegill sunfish ( <i>Lepomis macrochirus</i> ). EG & G Bionomics, Aquatic Toxicology Laboratory. Sumitomo Report No. IW-11-0007	Sumitomo Chemical Co., Ltd
Sousa and LeBlanc	1981	Acute toxicity of Neo-Pynamin Forte T.G. to Rainbow Trout ( <i>Salmo gairdneri</i> ). EG & G Bionomics, Aquatic Toxicology Laboratory. Sumitomo Report No. IW-11-0006	Sumitomo Chemical Co., Ltd
Sumitomo Chemical Co. Ltd	1986	Stability of Neopynamin Forte. Sumitomo Report No. IP-00-0054	Sumitomo Chemical Co., Ltd
White D.F., Mullee, D. M.	2006	D-tetramethrin: Determination of Physico-Chemical Properties, Safepharm Laboratories Ltd, Project No: 0555/0076, GLP, (Un)	Sumitomo Chemical Co., Ltd
York	2008	D-Tetramethrin/ S-1723 – Acute Toxicity To Rainbow Trout ( <i>Oncorhynchus mykiss</i> ) Under Flow-Through Conditions, Springborn Smithers Laboratories, Study No. 13048.6540.	S Sumitomo Chemical Co., Ltd
Yoshimura	1991	Adsorption and desorption of tetramethrin in water-soil suspension systems, Environmental Health Science Laboratory, Sumitomo Report No. IM-10-0013, 1991	Sumitomo Chemical Co., Ltd

15. ANNEXES

Confidential Annexes