

# Summary of product characteristics for a biocidal product

**Product name:** Rodex Whole Wheat Bait

**Product type(s):** PT14 - Rodenticides (Pest control)

**Authorisation number:** IE/BPA 70584

**R4BP 3 asset reference number:** IE-0000302-0000

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
4.1.1 Use-specific instructions for use	5
4.1.2 Use-specific risk mitigation measures	5
4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	6
4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	6
4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	6
4.2.1 Use-specific instructions for use	8
4.2.2 Use-specific risk mitigation measures	8
4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	8
4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	8
4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	9
4.3.1 Use-specific instructions for use	10
4.3.2 Use-specific risk mitigation measures	10
4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	10
4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	10
4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	10
4.4.1 Use-specific instructions for use	13
4.4.2 Use-specific risk mitigation measures	13

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	13
4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	14
4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	14
4.5.1 Use-specific instructions for use	15
4.5.2 Use-specific risk mitigation measures	16
4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	16
4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	16
4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	16
5. General directions for use	16
5.1. Instructions for use	16
5.2. Risk mitigation measures	17
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	18
5.4. Instructions for safe disposal of the product and its packaging	18
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	19
6. Other information	19

## Administrative information

### 1.1. Trade names of the product

Rodex Whole Wheat Bait
------------------------

### 1.2. Authorisation holder

<b>Name and address of the authorisation holder</b>	Name	PelGar International Limited
	Address	18 rue des Remparts d'Ainay 69002 Lyon France
<b>Authorisation number</b>	IE/BPA 70584	
<b>R4BP 3 asset reference number</b>	IE-0000302-0000	
<b>Date of the authorisation</b>	19/11/2012	
<b>Expiry date of the authorisation</b>	19/03/2023	

### 1.3. Manufacturer(s) of the biocidal products

<b>Name of the manufacturer</b>	PelGar International Limited
<b>Address of the manufacturer</b>	18 rue des Remparts d'Ainay 69002 Lyon France
<b>Location of manufacturing sites</b>	Unit 13 Newman Lane GU34 2QR Alton United Kingdom
	Doa Rat Bait Production Center, Mesokeleos street Palouriotissa, Nicosia Cyprus
	Promedivet SRL, str. Lunga nr. 46/G 545500 SOVATA Romania

#### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	12 - Bromadiolone
<b>Name of the manufacturer</b>	PelGar International Limited
<b>Address of the manufacturer</b>	Unit 13 Newman Lane GU34 2QR Alton United Kingdom
<b>Location of manufacturing sites</b>	Prazska 54 280 02 Kolin Czech Republic

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[3-(4'-Bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	Active Substance	28772-56-7	249-205-9	0.005

### 2.2. Type of formulation

RB - Bait (ready for use)

## 3. Hazard and precautionary statements

<b>Hazard statements</b>	Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction. May damage the unborn child. Causes damage to organs (blood) through prolonged or repeated exposure .
<b>Precautionary statements</b>	Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe dust.

Wear protective gloves.

Do not eat, drink or smoke when using this product.

Wash hands thoroughly after handling.

IF exposed or concerned: Get medical advice.

Get medical advice if you feel unwell.

Store locked up.

Dispose of contents in accordance with national regulations..

## 4. Authorised use(s)

### 4.1 Use description

#### Use 1 - Trained professionals – indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Mus musculus-House mouseAdults and juveniles Rattus norvegicus-Brown ratAdults and juveniles
<b>Field(s) of use</b>	Indoor Indoor
<b>Application method(s)</b>	Bait application -  - Ready-to-use bait to be used in tamper-resistant bait stations  - Covered and protected baiting points

**Application rate(s) and frequencies**

Rats: up to 200g of bait per bait point. Mice: up to 40g of bait per baiting point. - 100 -  
Rats:  
- High infestation: Up to 200 g of bait per baiting point spaced 5m apart.  
- Low infestation: Up to 200 g of bait per baiting point spaced 10m apart.  
  
Mice:  
- High infestation: 40 g of bait per baiting point spaced 2m apart.  
- Low infestation: 40 g of bait per baiting point spaced 5m apart.

**Category(ies) of users**

Trained professional

**Pack sizes and packaging material**

**1. Loose bait**

- No inner bag is to be larger than 10kg. Packs supplied either with (up to 20 kg) or without (up to 10kg) wire-tied inner bags of PET(E), HDPE or LDPE (up to 10 kg):

Polyethylene-lined polypropylene sack or multi-ply paper sack or polyethylene/polypropylene laminated sack 2.5kg to 20kg

Polyethylene lined double-walled or fibreboard carton 2.5kg to 10kg

Polyethylene/paper bag in a fibre-board carton 2.5kg to 10kg.

Polypropylene or polyethylene tub 2.5kg to 20kg

Polypropylene or polyethylene pail 2.5kg to 20kg

Polypropylene or polyethylene jerrican 2.5kg to 10kg

Polyethylene sachets or polypropylene/paper sachets or orientated polypropylene sachets or polyethylene/aluminium sachets or polyethylene/paper/aluminium sachets. Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

Polypropylene pouches or laminated polypropylene pouches or PET/polyethylene pouches or polyethylene/paper/aluminium pouches . Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

**2. Bait in sachets/place packs**

Bait in perforated or non-perforated polypropylene/paper film sachets, PET/polyethylene film sachets, polyethylene/paper film sachets or polyethylene/aluminised film sachets (10 g, 25 g, 30 g, 40 g, 50 g, 100 g or 200 g) supplied in the following outer packs:

Polypropylene woven sack or multi-ply paper sack. - 2.5kg to 25kg

Polypropylene pail - 2.5kg to 20kg

Polypropylene or polyethylene tub - 2.5kg to 10kg.

PET/polyethylene, polypropylene/polyethylene or paper/polyethylene pouch - 2.5kg to 10kg

Jerry can, with tamper-evident screw-top lid. - 2.5-2.5kg.

Fibreboard box - 2.5kg to 10 kg

### **3. Bait in trays**

((PVC or polystyrene or polypropylene or PET or PET/polyethylene or fibreboard); 25g, 30g, 40g, 50g, 100g or 200g)) with heat sealed lid (PET/polypropylene or PET/polyethylene or polypropylene) supplied in the following outer packs:

Cardboard outer - 2.5kg to 6.4kg

Single-use or multi-use tamper-proof HDPE or PP bait station, packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper. Packed in multipacks of 4, 8, 16, 24, 32, 48, 60, 72, 96, 100, 120, 144 - 2.5 kg to 20 kg.

#### **4.1.1 Use-specific instructions for use**

- Remove the remaining product at the end of treatment period.

#### **4.1.2 Use-specific risk mitigation measures**

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.  
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.  
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.  
- Do not use the product in pulsed baiting treatments.  
-Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient  
-The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

**4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

**4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See "Directions for use"

**4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

See "Directions for use"

**4.2 Use description**

**Use 2 - Trained Professionals – outdoor around buildings**

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Mus musculus-House mouseAdults and juveniles Rattus norvegicus-Brown ratAdults and juveniles
<b>Field(s) of use</b>	Outdoor Outdoors around buildings
<b>Application method(s)</b>	Bait application -  - Ready-to-use bait to be used in tamper-resistant bait stations .  - Covered and protected baiting points

**Application rate(s) and frequencies**

- Direct application of ready-to-use bait into the burrow

Rats: up to 200g of bait per bait point. Mice: up to 40g of bait per baiting point. - 100 -

**Rats:**

- High infestation: Up to 200 g of bait per baiting point spaced 5m apart.
- Low infestation: Up to 200 g of bait per baiting point spaced 10m apart.

**Mice:**

- High infestation: 40 g of bait per baiting point spaced 2m apart.
- Low infestation: 40 g of bait per baiting point spaced 5m apart.

- Burrow baiting: 20-200 g of bait per burrow.

**Category(ies) of users**

Trained professional

**Pack sizes and packaging material**

**1. Loose bait** - No inner bag is to be larger than 10kg. Packs supplied either with (up to 20 kg) or without (up to 10kg) wire-tied inner bags of PET(E), HDPE or LDPE (up to 10 kg):

Polyethylene-lined polypropylene sack or multi-ply paper sack or polyethylene/polypropylene laminated sack 2.5kg to 20kg  
Polyethylene lined double-walled or fibreboard carton 2.5kg to 10kg  
Polyethylene/paper bag in a fibre-board carton 2.5kg to 10kg.  
Polypropylene or polyethylene tub 2.5kg to 20kg  
Polypropylene or polyethylene pail 2.5kg to 20kg  
Polypropylene or polyethylene jerrican 2.5kg to 10kg  
Polyethylene sachets or polypropylene/paper sachets or orientated polypropylene sachets or polyethylene/aluminium sachets or polyethylene/paper/aluminium sachets. Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg  
Polypropylene pouches or laminated polypropylene pouches or PET/polyethylene pouches or polyethylene/paper/aluminium pouches . Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

**2. Bait in sachets/place packs**

Bait in perforated or non-perforated polypropylene/paper film sachets, PET/polyethylene film sachets, polyethylene/paper film sachets or polyethylene/aluminised film sachets (10 g, 25 g, 30 g, 40 g, 50 g, 100 g or 200 g) supplied in the following outer packs:

Polypropylene woven sack or multi-ply paper sack. - 2.5kg to 25kg  
Polypropylene pail - 2.5kg to 20kg  
Polypropylene or polyethylene tub - 2.5kg to 10kg.  
PET/polyethylene, polypropylene/polyethylene or paper/polyethylene pouch - 2.5kg to 10kg  
Jerry can, with tamper-evident screw-top lid. 2.5 - 2.5kg.  
Fibreboard box - 2.5kg to 10 kg

**3. Bait in trays** ((PVC or polystyrene or polypropylene or PET or PET/polyethylene or fibreboard); 25g, 30g, 40g, 50g, 100g or 200g)) with heat sealed lid (PET/polypropylene or PET/polyethylene or polypropylene) supplied in the following outer packs:

Cardboard outer - 2.5kg to 6.4kg  
Single-use or multi-use tamper-proof HDPE or PP bait station, packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper. Packed in multipacks of 4, 8, 16, 24, 32, 48, 60, 72, 96, 100, 120, 144 - 2.5 kg to 20 kg.

#### 4.2.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period (except when directly applied into burrows).
- Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU Code of Best Practice (or equivalent)
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting.

#### 4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product in pulsed baiting treatments.
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

#### 4.3 Use description

##### Use 3 - Trained Professionals – sewers

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Rattus norvegicus-Brown ratAdults and juveniles
<b>Field(s) of use</b>	Other  Sewers
<b>Application method(s)</b>	Bait application - - Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. - Covered and protected baiting points
<b>Application rate(s) and frequencies</b>	Up to 200 g bait per manhole - 100 - Up to 200 g bait per manhole
<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	<b>1. Loose bait</b> - No inner bag is to be larger than 10kg. Packs supplied either with (up to 20 kg) or without (up to 10kg) wire-tied inner bags of PET(E), HDPE or LDPE (up to 10 kg):  Polyethylene-lined polypropylene sack or multi-ply paper sack or polyethylene/polypropylene laminated sack 2.5kg to 20kg Polyethylene lined double-walled or fibreboard carton 2.5kg to 10kg Polyethylene/paper bag in a fibre-board carton 2.5kg to 10kg. Polypropylene or polyethylene tub 2.5kg to 20kg Polypropylene or polyethylene pail 2.5kg to 20kg Polypropylene or polyethylene jerrican 2.5kg to 10kg Polyethylene sachets or polypropylene/paper sachets or orientated polypropylene sachets or polyethylene/aluminium sachets or polyethylene/paper/aluminium sachets. Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg Polypropylene pouches or laminated polypropylene pouches or PET/polyethylene pouches or polyethylene/paper/aluminium pouches . Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

## **2. Bait in sachets/place packs**

Bait in perforated or non-perforated polypropylene/paper film sachets, PET/polyethylene film sachets, polyethylene/paper film sachets or polyethylene/aluminised film sachets (10 g, 25 g, 30 g, 40 g, 50 g, 100 g or 200 g) supplied in the following outer packs:

Polypropylene woven sack or multi-ply paper sack. - 2.5kg to 25kg

Polypropylene pail - 2.5kg to 20kg

Polypropylene or polyethylene tub - 2.5kg to 10kg.

PET/polyethylene, polypropylene/polyethylene or paper/polyethylene pouch - 2.5kg to 10kg

Jerry can, with tamper-evident screw-top lid. 2.5 - 2.5kg.

Fibreboard box - 2.5kg to 10 kg

**3. Bait in trays** ((PVC or polystyrene or polypropylene or PET or PET/polyethylene or fibreboard); 25g, 30g, 40g, 50g, 100g or 200g)) with heat sealed lid (PET/polypropylene or PET/polyethylene or polypropylene) supplied in the following outer packs:

Cardboard outer - 2.5kg to 6.4kg

Single-use or multi-use tamper-proof HDPE or PP bait station, packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper. Packed in multipacks of 4, 8, 16, 24, 32, 48, 60, 72, 96, 100, 120, 144 - 2.5 kg to 20 kg.

### **4.3.1 Use-specific instructions for use**

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.

### **4.3.2 Use-specific risk mitigation measures**

- Do not use this product in pulsed baiting treatments.

### **4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

See "Directions for use"

### **4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See "Directions for use"

### **4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

See "Directions for use"

#### 4.4 Use description

##### Use 4 - House mice and/or rats – Professional users – indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Mus musculus-House mouseAdults and juveniles Rattus norvegicus-Brown ratAdults and juveniles
<b>Field(s) of use</b>	Indoor Indoor
<b>Application method(s)</b>	Bait application -  - Ready-to-use bait to be used in tamper-resistant bait stations
<b>Application rate(s) and frequencies</b>	Rats: up to 200g of bait per bait point. Mice: up to 40g of bait per baiting point. - 100 - Rats: - High infestation: Up to 200 g of bait per baiting point spaced 5m apart. - Low infestation: Up to 200 g of bait per baiting point spaced 10m apart.  Mice: - High infestation: 40 g of bait per baiting point spaced 2m apart. - Low infestation: 40 g of bait per baiting point spaced 5m apart.
<b>Category(ies) of users</b>	Professional

## Pack sizes and packaging material

### 1. Loose bait

- No inner bag is to be larger than 10kg. Packs supplied either with (up to 20 kg) or without (up to 10kg) wire-tied inner bags of PET(E), HDPE or LDPE (up to 10 kg):

Polyethylene-lined polypropylene sack or multi-ply paper sack or polyethylene/polypropylene laminated sack 2.5kg to 20kg

Polyethylene lined double-walled or fibreboard carton 2.5kg to 10kg

Polyethylene/paper bag in a fibre-board carton 2.5kg to 10kg.

Polypropylene or polyethylene tub 2.5kg to 20kg

Polypropylene or polyethylene pail 2.5kg to 20kg

Polypropylene or polyethylene jerrican 2.5kg to 10k

Polyethylene sachets or polypropylene/paper sachets or orientated polypropylene sachets or polyethylene/aluminium sachets or polyethylene/paper/aluminium sachets. Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

Polypropylene pouches or laminated polypropylene pouches or PET/polyethylene pouches or polyethylene/paper/aluminium pouches . Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

### 2. Bait in sachets/place packs

Bait in perforated or non-perforated polypropylene/paper film sachets, PET/polyethylene film sachets, polyethylene/paper film sachets or polyethylene/aluminised film sachets (10 g, 25 g, 30 g, 40 g, 50 g, 100 g or 200 g) supplied in the following outer packs:

Polypropylene woven sack or multi-ply paper sack. - 2.5kg to 25kg

Polypropylene pail - 2.5kg to 20kg

Polypropylene or polyethylene tub - 2.5kg to 10kg.

PET/polyethylene, polypropylene/polyethylene or paper/polyethylene pouch - 2.5kg to 10kg

Jerry can, with tamper-evident screw-top lid. 2.5 - 2.5kg.

Fibreboard box - 2.5kg to 10 kg

### **3. Bait in trays**

((PVC or polystyrene or polypropylene or PET or PET/polyethylene or fibreboard); 25g, 30g, 40g, 50g, 100g or 200g) with heat sealed lid (PET/polypropylene or PET/polyethylene or polypropylene) supplied in the following outer packs:

Cardboard outer - 2.5kg to 6.4kg

Single-use or multi-use tamper-proof HDPE or PP bait station, packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper. Packed in multipacks of 4, 8, 16, 24, 32, 48, 60, 72, 96, 100, 120, 144 - 2.5 kg to 20 kg.

#### **4.4.1 Use-specific instructions for use**

- For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

#### **4.4.2 Use-specific risk mitigation measures**

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed) the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

#### **4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

**4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See "Directions for use"

**4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

See "Directions for use"

**4.5 Use description**

**Use 5 - Mice and/or rats – Professional users – outdoor around buildings**

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Mus musculus-House mouseAdults and juveniles Rattus norvegicus-Brown ratAdults and juveniles
<b>Field(s) of use</b>	Outdoor Outdoors around buildings
<b>Application method(s)</b>	Bait application -  - Ready-to-use bait to be used in tamper-resistant bait stations .
<b>Application rate(s) and frequencies</b>	Rats: up to 200g of bait per bait point. Mice: up to 40g of bait per baiting point. - 100 -  Rats: - High infestation: Up to 200 g of bait per baiting point spaced 5m apart. - Low infestation: Up to 200 g of bait per baiting point spaced 10m apart.

Mice:  
 - High infestation: 40 g of bait per baiting point spaced 2m apart.  
 - Low infestation: 40 g of bait per baiting point spaced 5m apart.

**Category(ies) of users**

Trained professional  
 Professional

**Pack sizes and packaging material**

**1. Loose bait** - No inner bag is to be larger than 10kg. Packs supplied either with (up to 20 kg) or without (up to 10kg) wire-tied inner bags of PET(E), HDPE or LDPE (up to 10 kg):

Polyethylene-lined polypropylene sack or multi-ply paper sack or polyethylene/polypropylene laminated sack 2.5kg to 20kg  
 Polyethylene lined double-walled or fibreboard carton 2.5kg to 10kg  
 Polyethylene/paper bag in a fibre-board carton 2.5kg to 10kg.  
 Polypropylene or polyethylene tub 2.5kg to 20kg  
 Polypropylene or polyethylene pail 2.5kg to 20kg  
 Polypropylene or polyethylene jerrican 2.5kg to 10kg  
 Polyethylene sachets or polypropylene/paper sachets or orientated polypropylene sachets or polyethylene/aluminium sachets or polyethylene/paper/aluminium sachets. Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg  
 Polypropylene pouches or laminated polypropylene pouches or PET/polyethylene pouches or polyethylene/paper/aluminium pouches . Supplied without outer packaging or in fibre-board carton with or without polyethylene liner or in polyethylene pack or in polypropylene pack or in blister pack 2.5kg to 10kg

**2. Bait in sachets/place packs**  
 Bait in perforated or non-perforated polypropylene/paper film sachets, PET/polyethylene film sachets, polyethylene/paper film sachets or polyethylene/aluminised film sachets (10 g, 25 g, 30 g, 40 g, 50 g, 100 g or 200 g) supplied in the following outer packs:

Polypropylene woven sack or multi-ply paper sack. - 2.5kg to 25kg  
 Polypropylene pail - 2.5kg to 20kg  
 Polypropylene or polyethylene tub - 2.5kg to 10kg.  
 PET/polyethylene, polypropylene/polyethylene or paper/polyethylene pouch - 2.5kg to 10kg  
 Jerry can, with tamper-evident screw-top lid. - 2.5kg.  
 Fibreboard box - 2.5kg to 10 kg

**3. Bait in trays** ((PVC or polystyrene or polypropylene or PET or PET/polyethylene or fibreboard); 25g, 30g, 40g, 50g, 100g or 200g)) with heat sealed lid (PET/polypropylene or PET/polyethylene or polypropylene) supplied in the following outer packs:

Cardboard outer - 2.5kg to 6.4kg  
 Single-use or multi-use tamper-proof HDPE or PP bait station, packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper. Packed in multipacks of 4, 8, 16, 24, 32, 48, 60, 72, 96, 100, 120, 144 - 2.5 kg to 20 kg.

**4.5.1 Use-specific instructions for use**

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period (except when directly applied into burrows).
- Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU Code of Best Practice (or equivalent)
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

#### **4.5.2 Use-specific risk mitigation measures**

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product in pulsed baiting treatments.
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

#### **4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### **4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

See "Directions for use"

#### **4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

See "Directions for use"

### **5. General directions for use**

#### **5.1. Instructions for use**

#### Trained professionals

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by CRRU Code of Best Practice (or equivalent).
- Sites where burrow baiting is used should be visited more frequently than those where secure bait boxes are employed
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Bait in non-emptiable sachets: Do not open the sachets containing the bait.
- Loose grains: Place the bait in the baiting point by using a dosage device. Specify the methods to minimise dust (e.g. wet wiping).

#### Professional users

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 2.3 for the information to be shown on the label).
- [If national policy or legislation require it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.
- Bait in non-emptiable sachets - Do not open the sachets containing the bait.
- Place loose bait in the baiting station by using a dosage device. Specify the methods to minimise dust (e.g. wet wiping).

## 5.2. Risk mitigation measures

#### Trained professionals

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment (unless permanent baiting)
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements

#### Professional users

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed) the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

### **5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
  - Antidote: Vitamin K1 administered by medical/veterinary personnel only.
  - In case of:
    - Dermal exposure, wash skin with water and then with water and soap.
    - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
    - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.
  - Contact a veterinary surgeon in case of ingestion by a pet.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre".
- Hazardous to wildlife.

### **5.4. Instructions for safe disposal of the product and its packaging**

Trained professionals

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended.

Professional users

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

## 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 2 years.

## 6. Other information

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.
- Use Biocides Safely and Sustainably
- It is illegal to use this product for uses or in a manner other than that prescribed on this label.
- Poison Information: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.
- Please note IRELAND applies a minimum Professional pack size of 2.5 Kg instead of 3.0 kg.