

Summary of product characteristics for a biocidal product

Product name: NYNA PELLETT 25

Product type(s): PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

Authorisation number: FR-2017-0033

R4BP 3 asset reference number: FR-0015759-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	4
2. Product composition and formulation	4
2.1. Qualitative and quantitative information on the composition of the biocidal product	4
2.2. Type of formulation	4
3. Hazard and precautionary statements	5
4. Authorised use(s)	5
5. General directions for use	25
5.1. Instructions for use	26
5.2. Risk mitigation measures	28
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	28
5.4. Instructions for safe disposal of the product and its packaging	29
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	29
6. Other information	29

Administrative information

1.1. Trade names of the product

NYNA PELLETT 25
DIFEPELLET RAT 25
DIFEPELLET RONGEUR 25
STOP DIFEPELLET 25
SUPP' DIFEPELLET 25

1.2. Authorisation holder

Name and address of the authorisation holder	Name	TRIPLAN SA
	Address	BP 258 LA POSTE FRANCAISE AD500 ANDORRA LA VELLA France
Authorisation number	FR-2017-0033	
R4BP 3 asset reference number	FR-0015759-0000	
Date of the authorisation	05/07/2017	
Expiry date of the authorisation	01/07/2024	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	NOXIMA
Address of the manufacturer	CARREFOUR JEAN MONNET, LACROIX SAINT-OUEN 60201 COMPIEGNE France
Location of manufacturing sites	CARREFOUR JEAN MONNET, LACROIX SAINT-OUEN 60201 COMPIEGNE France

Name of the manufacturer	INDUSTRIALCHIMICA SRL
Address of the manufacturer	VIA SORGAGLIA 40 I-35020 ARRE (PD) Italy
Location of manufacturing sites	VIA SORGAGLIA 40 I-35020 ARRE (PD) Italy

Name of the manufacturer	DALTA SA
Address of the manufacturer	ZONE INDUSTRIELLE – RN20-BP6 82270 MONTPEZAT DE QUERCY France
Location of manufacturing sites	ZONE INDUSTRIELLE – RN20-BP6 82270 MONTPEZAT DE QUERCY France

Name of the manufacturer	FARMAVIT OOD
Address of the manufacturer	BUL TSAR BORIS III, N° 63, OFFICE 1 1612 SOFIA Bulgaria
Location of manufacturing sites	INDULSTRIANA 2 STR ; PLEVEN DISTRICT 5960 GULIANTSI Bulgaria

Name of the manufacturer	SOFAR France
Address of the manufacturer	ZA DU DREVERS BP 02 29190 PLEYBEN France
Location of manufacturing sites	ZA DU DREVERS BP 02 29190 PLEYBEN France

Name of the manufacturer	AEDES PROTECTA
Address of the manufacturer	75 Rue d'Orgemont 95210 SAINT-GRATIEN France
Location of manufacturing sites	LIEU DIT DOUILLAC 81310 PARISOT France

Name of the manufacturer	IRIS
Address of the manufacturer	1126A, Avenue du Moulinas, Route de Saint-Privat 30340 SALINDRES France
Location of manufacturing sites	1126A, Avenue du Moulinas, Route de Saint-Privat 30340 SALINDRES France

Name of the manufacturer	HDA
Address of the manufacturer	ZA LA CHARME MENETROL 63200 RIOM France
Location of manufacturing sites	ZA LA CHARME MENETROL 63200 RIOM France

Name of the manufacturer	RATOUCY SAS
Address of the manufacturer	29 Rue de la Forêt LOOZE - BP 145 89303 JOIGNY CEDEX France
Location of manufacturing sites	29 Rue de la Forêt LOOZE - BP 145 89303 JOIGNY CEDEX France

Name of the manufacturer	EDIALUX
Address of the manufacturer	ZA MACON EST 01750 REPLONGES France
Location of manufacturing sites	Chez LARC - ZA DE KERAMPAOU 29140 MELGVEN France

Name of the manufacturer	PROFARM SA
Address of the manufacturer	Industrial Area of Thessaloniki, Entrance A, O.T. 20, Buildings 30 and 53 57022 SINDOS Greece
Location of manufacturing sites	Industrial Area of Thessaloniki, Entrance A, O.T. 20, Buildings 30 and 53 57022 SINDOS Greece

Name of the manufacturer	FARMA-CHEM SA
Address of the manufacturer	Industrial Area of Sindos, P.O. BOX 1026 Block 53 Zone C 57022 THESSALONIKI Greece
Location of manufacturing sites	Industrial Area of Sindos, P.O. BOX 1026 Block 53 Zone C 57022 THESSALONIKI Greece

Name of the manufacturer	AGROLOGY SA
Address of the manufacturer	P.O. BOX 1089, Industrial Area of Thessaloniki 57022 THESSALONIKI Greece
Location of manufacturing sites	P.O. BOX 1089, Industrial Area of Thessaloniki 57022 THESSALONIKI Greece

Name of the manufacturer	AGGRESS
Address of the manufacturer	EL. VENIZELOU 158 A 16341 ATHENES Greece
Location of manufacturing sites	ARMA THEBES 32200 VOIOTIA Greece

1.4. Manufacturer(s) of the active substance(s)

Active substance	26 - Difenacoum
Name of the manufacturer	DR TEZZA S.R.L.
Address of the manufacturer	Via Tre Ponti 22 37050 S. Maria di Zevio (VR) Italy
Location of manufacturing sites	Via Tre Ponti 22 37050 S. Maria di Zevio (VR) Italy

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Difenacoum	3-(3-biphenyl-4-yl-1,2,3,4-tetrahydro-1-naphthyl)-4-hydroxycoumarin	Active Substance	56073-07-5	259-978-4	0,0025

2.2. Type of formulation

Ready-to-use bait : pellet

3. Hazard and precautionary statements

Hazard statements

May cause damage to organs (blood) through prolonged or repeated exposure .

Precautionary statements

Do not breathe dust.

Get medical if you feel unwell.

Dispose of to .

4. Authorised use(s)

4.1 Use description

Use 1 - (Not relevant in France) House mice – professionals - indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: Mus musculus
Common name: House mice
Development stage: Juvenils and adults

Field(s) of use

Indoor

Indoor

Application method(s)

Method: Bait application
Detailed description:
Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequencies

Application Rate: 40 g of bait per bait point
Dilution (%): 0
Number and timing of application:
40 g of bait per bait point. If more than one bait point is needed, the minimum distance between bait points should be of 1 to 2 meters.

Category(ies) of users

Professional

Pack sizes and packaging material

Minimum pack size of 3kg.

The product is supplied in individual PE/PP sachets (5-100g) and in bulk.

The sachets (5-100 g) are wrapped in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Metal boxes (5-25kg)
- Bait box in PET/PP/PE/PVC

The product is supplied in bulk in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Sachet (PET/ALU/PET) (5-25kg) packed in cardboard box (5-50kg)
- Metal boxes (5-25kg)
- Pre-filled bait boxes in PET/PP/PE/PVC up to the claimed dose for mice/rats

4.1.1 Use-specific instructions for use

- The bait station should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait station are intact and to remove rodent bodies. Re-fill bait when necessary.

- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.1.2 Use-specific risk mitigation measures

-

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.2 Use description

Use 2 - (Not relevant in France) Rats - professionals - indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles and adults Scientific name: Rattus rattus Common name: Black or roof rat Development stage: Juveniles and adults
Field(s) of use	Indoor Indoor
Application method(s)	Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequencies

Application Rate: 100 g of bait per bait point
Dilution (%): 0
Number and timing of application:
100 g of bait per bait point. If more than one bait point is needed, the minimum distance between bait stations should be of 5-10 meters.

Category(ies) of users

Professional

Pack sizes and packaging material

Minimum pack size of 3kg.

The product is supplied in individual PE/PP sachets (5-100g) and in bulk.

The sachets (5-100 g) are wrapped in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Metal boxes (5-25kg)
- Bait box in PET/PP/PE/PVC

The product is supplied in bulk in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Sachet (PET/ALU/PET) (5-25kg) packed in cardboard box (5-50kg)
- Tin-plate boxes (5-25kg)
- Pre-filled bait boxes in PET/PP/PE/PVC up to the claimed dose for mice/rats

4.2.1 Use-specific instructions for use

- The bait station should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait station are intact and to remove rodent bodies. Re-fill bait when necessary.

- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.2.2 Use-specific risk mitigation measures

-

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.3 Use description

Use 3 - (Not relevant in France) House mice and rats - professionals - outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mice
Development stage: Juveniles and adults

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: Juveniles and adults

Scientific name: *Rattus rattus*
Common name: Black or roof rat
Development stage: Juveniles and adults

Field(s) of use

Outdoor

Outdoor around building

Application method(s)

Method: Bait application
Detailed description:
Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequencies

Application Rate: 40 g of bait per bait point for mice. 100 g of bait per bait point for rats
Dilution (%): 0

Number and timing of application:

- 40 g of bait per bait point for mice. If more than one bait point is needed, the minimum distance between bait points should be of 1 to 2 meters.

- 100 g of bait per bait point for rats. If more than one bait point is needed, the minimum distance between bait points should be of 5 to 10 meters.

Category(ies) of users

Professional

Pack sizes and packaging material

Minimum pack size of 3kg.

The product is supplied in individual PE/PP sachets (5-100g) and in bulk.

The sachets (5-100 g) are wrapped in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Metal boxes (5-25kg)
- Bait box in PET/PP/PE/PVC

The product is supplied in bulk in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Sachet (PET/ALU/PET) (5-25kg) packed in cardboard box (5-50kg)
- Tin-plate boxes (5-25kg)
- Pre-filled bait boxes in PET/PP/PE/PVC up to the claimed dose for mice/rats

4.3.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait station should be visited for mice at least every 2 to 3 days at the beginning of the treatment and for rats only 5 to 7 days after the beginning of the treatment and at least weekly afterwards (for mice and rats), in order to check whether the bait is accepted, the bait station are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in bait station in which bait has been damaged by water or contaminated by dirt.

4.3.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or drainage systems, ensure that bait contact with water is avoided.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.4 Use description

Use 4 - House mice and/or rats – trained professionnals - indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mice
Development stage: Juveniles and adults

Scientific name: *Rattus norvegicus*
Common name: Brown rat

Development stage: Juveniles and adults

Scientific name: *Rattus rattus*
Common name: Black or roof rat
Development stage: Juveniles and adults

Field(s) of use

Indoor

Indoor

Application method(s)

Method: Bait application
Detailed description:
Bait formulations:

- ready-to-use bait to be used in tamper-resistant bait stations

- ready-to-use bait may be used in covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait station

Application rate(s) and frequencies

Application Rate: Mice - 40 g of bait per baiting point every 1 to 2 meters. Rats - 100 g of bait per baiting point every 5 to 10 meters.

Dilution (%): 0

Number and timing of application:

Rats

- 100 g of bait per baiting point every 5 to 10 meters.

Mice

- 40 g of bait per baiting point every 1 to 2 meters.

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size of 3 kg.

(In France only : minimum pack size of 5 kg)

The product is supplied in individual PE/PP sachets (5-100g) and in bulk.

The sachets (5-100 g) are wrapped in :

- Buckets/Barrels (PE/PP) (5-30kg)

- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Metal boxes (5-25kg)
- Bait box in PET/PP/PE/PVC

The product is supplied in bulk in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Sachet (PET/ALU/PET) (5-25kg) packed in cardboard box (5-50kg)
- Tin-plate boxes (5-25kg)
- Pre-filled bait boxes in PET/PP/PE/PVC up to the claimed dose for mice/rats

4.4.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.
- Follow any additional instructions provided by the relevant code of best practice.

4.4.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

- Do not use the product in pulsed baiting treatments.

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.5 Use description

Use 5 - Mice and/or rats - trained professionals - outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mice
Development stage: Juveniles and adults

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: Juveniles and adults

Scientific name: *Rattus rattus*
Common name: Black or roof rat
Development stage: Juveniles and adults

Field(s) of use

Outdoor

Outdoor around buildings

Application method(s)

Method: Bait application

Detailed description:

Bait formulations:

- ready-to-use bait to be used in tamper-resistant bait stations

- ready-to-use bait may be used in covered and protected baiting points as long as they

provide the same level of protection for non-target species and humans as tamper-resistant bait station

Application rate(s) and frequencies

Application Rate: Mice 40 g of bait per baiting point every 1 to 2 meters. Rats 100 g of bait per baiting point every 5 to 10 meters.

Dilution (%): 0

Number and timing of application:

Rats

100 g of bait per baiting point every 5 to 10 meters.

Mice

40 g of bait per baiting point every 1 to 2 meters.

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size of 3 kg.

(in France only : Minimum pack size of 5 kg)

The product is supplied in individual PE/PP sachets (5-100g) and in bulk.

The sachets (5-100 g) are wrapped in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Metal boxes (5-25kg)
- Bait box in PET/PP/PE/PVC

The product is supplied in bulk in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)

- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Sachet (PET/ALU/PET) (5-25kg) packed in cardboard box (5-50kg)
- Tin-plate boxes (5-25kg)
- Pre-filled bait boxes in PET/PP/PE/PVC up to the claimed dose for mice/rats

4.5.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.5.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.6 Use description

Use 6 - Rats - trained professionals - Outdoor open areas

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: Brown rat Development stage: Juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: Black or roof rat Development stage: Juveniles and adults
Field(s) of use	Outdoor Outdoor open areas
Application method(s)	Method: Bait application Detailed description: Bait formulations: - ready-to-use bait to be used in tamper-resistant bait stations or in covered and protected baiting points
Application rate(s) and frequencies	Application Rate: 100 g of bait per baiting point Dilution (%): 0

Number and timing of application:
Bait products

- 100 g of bait per baiting point every 5 to 10 meters.

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size of 3 kg.

(in France only : Minimum pack size of 5 kg)

The product is supplied in individual PE/PP sachets (5-100g) and in bulk.

The sachets (5-100 g) are wrapped in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Tin -plate boxes (5-25kg)
- Bait box in PET/PP/PE/PVC

The product is supplied in bulk in :

- Buckets/Barrels (PE/PP) (5-30kg)
- Bags (paper bags with or without plastic film in PE/PP inside) (5-30kg)
- Cardboard boxes (5-50kg)
- Bags/Films PE/PP (5-25kg)
- Sachet (PET/ALU/PET) (5-25kg) packed in cardboard box (5-50kg)
- Tin-plate boxes (5-25kg)
- Pre-filled bait boxes in PET/PP/PE/PVC up to the claimed dose for mice/rats



4.6.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Follow any additional instructions provided by the relevant code of best practice.

4.6.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows

4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.7 Use description

Use 7 - House Mice – General public - Indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: House mice Development stage: Juveniles and adults
Field(s) of use	Indoor Indoor
Application method(s)	Method: Bait application Detailed description: Bait formulations: - ready-to-use bait to be used in tamper-resistant bait stations or in covered and protected baiting points
Application rate(s) and frequencies	Application Rate: 40 g of bait per bait point Dilution (%): 0 Number and timing of application: 40 g of bait per bait point for mice. If more than one bait point is needed, the minimum distance between bait points should be of 1 to 2 meters.
Category(ies) of users	General public (non-professional)
Pack sizes and packaging material	Maximum packaging 150g The product is supplied in individual PE/PP sachets (5-40g) packed in : - Buckets (PE/PP) - Bottles/Containers (PE/PP)

- Cardboard boxes
- Films (PE/PP)
- Tin-plate boxes
- Bait boxes PET/PP/PE/PVC

4.7.1 Use-specific instructions for use

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.7.2 Use-specific risk mitigation measures

-

4.7.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.7.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.7.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.8 Use description

Use 8 - Rats – General public - Indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles and adults Scientific name: Rattus rattus Common name: Black or roof rat Development stage: Juveniles and adults
Field(s) of use	Indoor Indoor
Application method(s)	Method: Bait application Detailed description: Bait formulations: - ready-to-use bait to be used in tamper-resistant bait stations or in covered and protected baiting points
Application rate(s) and frequencies	Application Rate: 100 g of bait per baiting point Dilution (%): 0 Number and timing of application: 100 g of bait per baiting point. If more than one bait point is needed, the minimum distance between bait points should be of 5 to 10 meters.
Category(ies) of users	General public (non-professional)
Pack sizes and packaging material	Maximum packaging 150g. The product is supplied in individual PE/PP sachets (5-100g) packed in : - Buckets (PE/PP) - Bottle/Containers (PE/PP) - Cardboard boxes

- Films (PE/PP)
- Tin-plate boxes
- Bait boxes PET/PP/PE/PVC

4.8.1 Use-specific instructions for use

- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.8.2 Use-specific risk mitigation measures

-

4.8.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.8.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.8.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.9 Use description

Use 9 - Rats – General public - outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: Juveniles and adults

Scientific name: *Rattus rattus*
Common name: Black or roof rat
Development stage: Juveniles and adults

Field(s) of use

Outdoor

Outdoor around buildings

Application method(s)

Method: Bait application

Detailed description:

Bait formulations:

- ready-to-use bait to be used in tamper-resistant bait stations or in covered and protected baiting points

Application rate(s) and frequencies

Application Rate: 100 g of bait per baiting point

Dilution (%): 0

Number and timing of application:

100 g of bait per baiting point.

If more than one bait point is needed, the minimum distance between bait points should be of 5 to 10 meters.

Category(ies) of users

General public (non-professional)

Pack sizes and packaging material

Maximum packaging 150g.

The product is supplied in individual PE/PP sachets (5-100g) packed in :

- Buckets (PE/PP)

- Bottles/Containers (PE/PP)

- Cardboard boxes

- Films (PE/PP)

- Tin plate boxes

- Bait boxes PET/PP/PE/PVC

4.9.1 Use-specific instructions for use

- Place the bait stations in areas not liable to flooding.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.9.2 Use-specific risk mitigation measures

-

4.9.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.9.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.9.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. General directions for use

5.1. Instructions for use

FOR PROFESSIONAL AND TRAINED PROFESSIONAL USERS

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
 - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
 - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
 - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
 - The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
 - Where possible, bait stations must be fixed to the ground or other structures.
 - Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
 - [If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
 - Bait should be secured so that it cannot be dragged away from the bait station.
 - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
 - Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

FOR TRAINED PROFESSIONAL ONLY

- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

FOR PROFESSIONALS ONLY

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
 - Remove the remaining bait or the bait stations at the end of the treatment period.
 - For non-emptiable sachets - Do not open the sachets containing the bait.
- Loose pellets-granules, grains:
- Place the bait in the baiting point by using a dosage device. Specify the methods to minimise dust (e.g. wet wiping).

- Decanting is to be avoided. In case decanting cannot be avoided, an RPE of APF 10 has to be used.

FOR NON PROFESSIONAL USERS

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Do not open the sachets containing the bait
- Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.
- Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Do not place bait stations near water drainage systems where they can come into contact with water.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

FOR PROFESSIONAL AND TRAINED PROFESSIONAL USERS

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign [in accordance with the applicable code of good practice, if any].

- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").

FOR TRAINED PROFESSIONAL ONLY

- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

FOR PROFESSIONAL ONLY

- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week). [Where relevant, specify if more frequent or daily inspection is required].
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
 - the product shall not be supplied to the general public (e.g. "for professionals only").
 - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

FOR NON PROFESSIONAL USERS

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).
- The product information (i.e. label and/or leaflet) shall clearly show that:
 - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.
- Dispose dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Ingestion: Wash out mouth with water. Contact poison treatment specialist. Seek medical advice immediately if symptoms occur and/or large quantities have been ingested.
- Skin contact: Wash contaminated skin with soap and water. Contact poison treatment specialist if symptoms occur.
- Eye contact: Immediately flush with plenty of water, occasionally lifting the upper and lower eyelids. Check for and remove any contact lenses if easy to do. Continue to rinse with warm water for at least 10 minutes. Get medical attention if irritation or vision impairment occurs.
- Antidote: vitamin K1 (phytomedione). Contact poison treatment specialist for antidote dosage and INR (or PT) monitoring.
- Keep the container or label available.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre or emergency telephone number.
- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Keep out of reach of children
- Shelf life: 2 years.

6. Other information

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.

