

## **Biocidal Products Committee (BPC)**

Opinion on the application for approval of the active substance:

**Reaction products of paraformaldehyde and 2-hydroxypropylamine  
(ratio 1:1)**

**Product type: 6**

ECHA/BPC/162/2017

Adopted

29 June 2017



## Opinion of the Biocidal Products Committee

### on the application for approval of the active substance Reaction products of paraformaldehyde and 2-hydroxy-propylamine (ratio 1:1) for product type 6

In accordance with Article 90(2) of Regulation (EU) No 528/2012 of the European Parliament and of the Council 22 May 2012 concerning the making available on the market and use of biocidal products (BPR), the Biocidal Products Committee (BPC) has adopted this opinion on the approval in product type 6 of the following active substance:

<b>Common name:</b>	<b>Formaldehyde released from the reaction products of paraformaldehyde and 2-hydroxypropylamine (ratio 1:1);</b>
	<b>RP 1:1</b>
<b>Chemical name(s):</b>	<b>Reaction products of paraformaldehyde and 2-hydroxypropylamine (ratio 1.1)</b>
<b>EC No.:</b>	<b>not applicable</b>
<b>CAS No.:</b>	<b>not applicable</b>
<b>Existing active substance</b>	

This document presents the opinion adopted by the BPC, having regard to the conclusions of the evaluating Competent Authority. The assessment report, as a supporting document to the opinion, contains the detailed grounds for the opinion.

### Process for the adoption of BPC opinions

Following the submission of an application by Task Force Lubrizol Deutschland GmbH and Schülke & Mayr GmbH. on 1 August 2007, the evaluating Competent Authority Austria submitted an assessment report and the conclusions of its evaluation to the Commission on 29 September 2016. In order to review the assessment report and the conclusions of the evaluating Competent Authority, the Agency organised consultations via the BPC (BPC-21) and its Working Groups (WG II 2017). Revisions agreed upon were presented and the assessment report and the conclusions were amended accordingly.

Information on the fulfilment of the conditions for considering the active substance as a candidate for substitution was made publicly available at <https://www.echa.europa.eu/web/guest/addressing-chemicals-of-concern/biocidal-products-regulation/potential-candidates-for-substitution-previous-consultations/-/substance-rev/5401/term%20on%209th%20February%202015>

on 4 November 2016, in accordance with the requirements of Article 10(3) of the BPR. Interested third parties were invited to submit relevant information by 3 January 2017.

## Adoption of the BPC opinion

### Rapporteur: Austria

The BPC opinion on the approval of the active substance reaction products of paraformaldehyde and 2-hydroxy-propylamine (ratio 1:1) in product type 6 was adopted on 29 June 2017.

The BPC opinion takes into account the comments of interested third parties provided in accordance with Article 10(3) of the BPR.

The BPC opinion was adopted by consensus. The opinion is published on the ECHA webpage at: <http://echa.europa.eu/regulations/biocidal-products-regulation/approval-of-active-substances/bpc-opinions-on-active-substance-approval>.

## Detailed BPC opinion and background

### 1. Overall conclusion

Since RP 1:1 fulfils the criteria set in Article 5(1) of Regulation (EU) No 528/2012, the overall conclusion of the BPC is that RP 1:1 in product type 6 should normally not be approved, unless one of the conditions for derogation in Article 5(2) is met. The process related to the demonstration of whether the conditions for derogation set in Article 5(2) are met, is not in the remit of the BPC<sup>1</sup>. The detailed grounds for the overall conclusion are described in the assessment report.

### 2. BPC Opinion

#### 2.1. BPC Conclusions of the evaluation

##### a) Presentation of the active substance including the classification and labelling of the active substance

This evaluation covers the use of Reaction product of paraformaldehyde and 2-hydroxypropylamine (ratio 1:1), furthermore addressed as RP 1:1 in product type 6. RP 1:1 was originally notified as  $\alpha,\alpha',\alpha''$ -trimethyl-1,3,5-triazine-1,3,5(2H,4H,6H)-triethanol or HPT<sup>2</sup>. RP 1:1 is a formaldehyde-releaser. Specifications for the reference source are established.

The physico-chemical properties of the active substance and biocidal product have been evaluated and are deemed acceptable for the appropriate use and materials suitable for storage and transport of the active substance and biocidal product. Regarding the explosive properties the justification for non submission of data has not been accepted. An experimental test has to be conducted as the substance contains of unknown constituents and therefore the waiving cannot be justified by structural considerations.

Validated analytical methods are available for the active substance as manufactured and for the relevant and significant impurities. With regard to the methods submitted for determination of the hydrolysis product 2-hydroxypropylamin in water and soil the data has been considered as not sufficient.

The classification and labelling for RP 1:1 according to Regulation (EC) No 1272/2008 (CLP Regulation) as agreed by RAC-35 (December 2015) and published in Regulation (EC) No 2017/776 (10<sup>th</sup> adaption to technical progress):

Classification according to the CLP Regulation	
Hazard Class and Category Codes	Acute Tox. 4, H302 Acute Tox. 4, H332 Skin Corr. 1C, H314 Eye Dam. 1, H318 Skin Sens. 1A, H317 STOT RE 2, H373 Muta 2, H341* Carc. 1B, H350** Aquatic Chronic 2, H411
	* The classification as a mutagen need not apply if it can be shown that the maximum theoretical concentration of releasable formaldehyde, irrespective of the source, in the

<sup>1</sup> See document: Further guidance on the procedures related to the examination of the exclusion criteria and the conditions for derogation under Article 5(2) (CA-Nov14-Doc.4.5-Final).

<sup>2</sup> The renaming of  $\alpha,\alpha',\alpha''$ -trimethyl-1,3,5-triazine-1,3,5(2H,4H,6H)-triethanol - HPT into Reaction products of paraformaldehyde and 2-hydroxypropylamine (ratio 1:1) is not regarded as a redefinition according to Article 11 of Regulation (EU) No 1062/2014.

	mixture as placed on the market is less than 1%.” ** The classification as a carcinogen need not apply if it can be shown that the maximum theoretical concentration of releasable formaldehyde, irrespective of the source, in the mixture as placed on the market is less than 0.1%.
<b>Labelling</b>	
Pictograms	GHS 05, GHS 07, GHS 08, GHS 09
Signal Word	Danger
Hazard Statement Codes	H302: Harmful if swallowed H332: Harmful if inhaled H314: Causes severe skin burns and eye damage H317: May cause an allergic skin reaction H373: May cause damage to organs (gastrointestinal tract and respiratory tract) H341: Suspected of causing genetic defects H350: May cause cancer 411: Toxic to aquatic life with long lasting effects
Suppl. Hazard Statement Code	EUH071: Corrosive to the respiratory tract
<b>Specific Concentration limits, M-Factors</b>	M = not applicable

### b) Intended use, target species and effectiveness

RP 1:1 containing biocidal products are used as bactericides for the preservation of fuels (PT 6) which are prone to bacterial decay. The product is intended to be incorporated by industrial users into fuels during the formulation process, which is carried out automatically, to act as a preservative with bactericidal activity. Formulation is performed in closed systems with a high degree of automation resulting in a final concentration of the active substance of 0.005%.

The assessment of the biocidal activity of the active substance demonstrates that it has a sufficient level of efficacy against gram negative bacteria such as *Pseudomonas aeruginosa*, *Enterobacter aerogenes* and *Acinetobacter spec.*

The active substance is a formaldehyde-releaser. The biocidal activity of the active substance is due to the interaction of the released formaldehyde with protein, DNA and RNA. The interaction with protein results from a combination with the primary amide and the amino groups. It reacts with carboxyl, sulfhydryl and hydroxyl groups.

As formaldehyde is not specific for one cellular target, the development of resistance is unlikely, if sufficiently high formaldehyde concentrations are guaranteed that exceed the capacity of the innate detoxification systems.

### c) Overall conclusion of the evaluation including need for risk management measures

A common core dossier was developed for formaldehyde, which was agreed at a Biocides Technical Meeting. This core dossier forms the basis of the hazard assessment of formaldehyde for all formaldehyde releasing active substances.

## Human health

The toxicity of the active substances is dominated by skin sensitization and local irritation and local (in vitro) genotoxicity (but negative systemic in vivo genotoxicity) and the hydrolysis study and efficacy mode of action support that the equilibrium within the RP 1:1 quickly shifts towards formaldehyde and 2-hydroxypropylamine by dilution and by the reaction of formaldehyde with biological media. This is essentially the basis for reading across the classification of formaldehyde for germ cell mutagenicity category 2 and carcinogenicity category 1B. However the risk assessment provided below for local and for systemic effects includes also the potential for carcinogenic effects.

Risk for the application of RP 1:1 as PT 6 for the preventive as well as curative treatment of fuels is characterised in terms of application to fuel and use of treated fuel. 100% RP 1:1 as manufactured is added to fuels with a final concentrations of usually in the range of 0.005% (but also 0.1% was considered as reasonable worst case assumption for "shock doses" in analogy with the intended use of RP 3:2). Exposure to RP 1:1 has to be completely excluded due to the corrosive and sensitizing hazard. Exposure to treated fuel may happen due to sampling the mixing vessels and use of the fuel (tanking). Due to the low water content in fuel RP 1:1 is considered to be present largely in the non-hydrolysed state. Consequently the risk estimates are provided just for the situation of no hydrolysis. However it is recognised that toxicological reference values for the active substance are assured to be protective also for a release of formaldehyde at the site of contact since they are either read across from formaldehyde test values (local respiratory AEC) or are in the same range as the formaldehyde test values (systemic AEL). The toxicological reference values for the active substance are also assured to be protective for the hydrolysis product 2-hydroxypropylamine, since its AEL and AEC are higher compared to those of RP 1:1.

The table below summarises the exposure scenarios assessed.

Summary table: human health scenarios			
Scenario	Primary or secondary exposure and description of scenario	Exposed group	Conclusion
<b>PT 6:</b> Formulation of Diesel fuels	Primary exposure covering the following tasks mixing, sampling, filling  Exposure to active substance and fuels  Acceptable with local exhaust ventilation (LEV), automatic dosing system, technical and organisational RMM and PPE (like gloves, coveralls, masks) for high local hazard category (loading, mixing) and RMM for standard industrial workplace for other tasks.	Industrial workers	Acceptable with RMM and PPE.
<b>PT 6:</b> Refueling of engines	Primary exposure, refueling of engines at filling stations, risk management measures for preventing gaseous release  Exposure to fuels  Acceptable with vapour recovering which is standard at filling stations.	Professionals, general public	Acceptable with RMM.
<b>PT 6:</b> Refueling of engines	Secondary exposure (bystanders) during refueling of engines at filling stations, risk management measures for preventing gaseous release  Exposure to fuels  Acceptable with vapour recovering which is standard at filling stations.	General public	Acceptable with RMM.

Risk for loading of RP 1:1 to fuel mixing systems, sampling of treated fuels, cleaning of containers and maintenance of the system is considered. For the loading of the corrosive and sensitizing RP 1:1 to fuel mixing systems closed systems have to be used in order to allow concluding that the risk for local respiratory and local dermal effects is acceptable. Exposure to the fuel containing maximally 0.1% of RP 1:1 (below or borderline to classification limits for the mixture) shall be minimised with closed systems for the sampling procedure and appropriate ventilation according to actual technical standards at filling stations (Directive 2009/126/EC). In this case also risk for systemic effects is acceptable.

For PT 6 dermal and respiratory exposure to general public may occur via treated fuels, however due to the longer exposure intervals the professional exposure is considered worst case and resulted in acceptable risk ratios. No pets and dietary exposure are expected for PT6.

Considering that local irritation is a condition for the development of tumours and applying a deterministic threshold AEC and AEL, also the risk for potential carcinogenic effects appears acceptable.



## Environment

The risk characterization was based on the hydrolysis products 2-hydroxypropylamine and formaldehyde as during the disinfection and use of the metal working system RP 1:1 has almost completely hydrolysed. The parent compound itself is therefore not expected to reach any environmental compartment. 2-Hydroxypropylamine and formaldehyde are expected to be readily biodegradable in the environment and are unlikely to bioaccumulate in biota. For acute toxicity algae is the most sensitive species with a 72h-E<sub>c</sub>C<sub>50</sub> of 5.7 mg/L (geometric mean, *Desmodesmus subspicatus*) for formaldehyde and a 96h-E<sub>b</sub>C<sub>50</sub> of 118.4 mg/L (nominal, buffered, *Pseudokirchneriella subcapitata*) for 2-hydroxypropylamine. Both compounds are not classifiable towards environmental hazards based on the available data.

The table below summarises the exposure scenarios assessed.

Summary table: environment scenarios		
Scenario	Description of scenario including environmental compartments	Conclusion
<b>PT 6:</b> In-can preservatives (PT 6.5 Fuels)	Preservatives in fuels are used to prevent microbial growth in presence of water, the formation of slime and sludge and finally the deterioration of the fuels during storage in the tanks of machines. For fuel ending up in an engine, it is assumed that 100% of the substance will be burnt thus, emissions should not be considered. RP 1:1 is mainly used for diesel fuel and the storage of diesel; crude oil is not treated.  <b>Affected environmental compartments:</b> None.	Acceptable
<b>PT 6:</b> Storage of fuel in storage tanks	A scenario for emission of fuel preservatives from large oil storage tanks (for example refineries) along with the aqueous phase does not exist currently. Exposure estimates for PT 13 are expected to be similar to that of releases from large storage tanks and therefore may in principle cover the emissions for fuel preservatives as well.  <b>Affected environmental compartments:</b> STP, surface water, sediment, soil, groundwater	Acceptable for 2-hydroxypropylamine.  Not acceptable for formaldehyde for STP, surface water and soil.

No exposure and risks from the use of RP 1:1 in fuels ending up in machine tanks to the environment were identified.

Risks of formaldehyde for storage of fuel in storage tanks were acceptable only if formaldehyde concentrations in the water phase after on-site or off-site treatment is below 40 mg/L, unless further evidence is provided within product authorisation that measurements are not necessary.

Formaldehyde is a very reactive compound so it can be assumed that during and after application of the product the concentration of formaldehyde will decrease by a chain of chemical reactions. Because no scenario for storage of fuel in storage tanks exists currently, exposure estimates for PT13 are expected to cover the emissions for fuel preservatives as well. For PT13 exposure calculations degradation since last dosing, i.e. operating time since last biocide dosing, storage at end-user site or transport to waste management facility and

storage at waste management site was not taken into account (as there is no standard algorithm that can be used to derive such degradation rates). However, for an estimation of the exposure reduction, a supporting pilot study on the degradation of formaldehyde in used MWF was submitted. In this pilot study the formaldehyde concentration of a used emulsion of a metalworking company was measured, which was delivered to a waste treatment company. The measurements showed that after distillation, the content of formaldehyde in the distillate is below the limit of quantification (<40 mg/L). The pilot study results are considered as supportive information due to several limitations. Regarding a concentration of 40 mg formaldehyde per litre waste water in the influent of a municipal STP, no unacceptable risks concerning the affected environmental compartments are indicated. For a refinement of the exposure calculations additional robust information on degradation of formaldehyde between the last dosing and the start of waste treatment would be necessary. This additional information enables the use of an elimination fraction which includes degradation during use, storage at end-user site, transport to waste management facility and storage at waste management site.

It has to be highlighted, that the environmental exposure calculations are very conservative as no degradation of formaldehyde in the system during use was assumed. However the above mentioned pilot study provides evidence for high degradation of formaldehyde during use (83 % between dosing and removal of the metal working fluid). As the exceedance of the PEC/PNEC<sub>STP</sub> ratio regarding formaldehyde is around 2 (based on the risk characterisation of PT 13), the required new data will allow to refine the exposure calculations also in this respect, which will most likely result in an acceptable risk.

A risk management option for the terrestrial compartment is a restriction of the use of sewage sludge in agriculture depending on national legislations.

The measurements of formaldehyde in effluents take into consideration aggregate exposure from multiple uses and product types as well as exposure from other formaldehyde sources.

Concerning risks to surface water intended for the abstraction of drinking water the parametric value of 0.1 µg/L of the Drinking Water Directive 98/83/EC may be exceeded. This may be considered by the relevant national authorities when issuing permits for recovery plants.

## Overall conclusion

Overall a safe use has been identified for the use of RP 1:1 to preserve fuels in PT6 provided adequate RMM and PPE are considered for human health and environment.

## 2.2. Exclusion, substitution and POP criteria

### 2.2.1. Exclusion and substitution criteria

The table below summarises the relevant information with respect to the assessment of exclusion and substitution criteria:

Property		Conclusions	
CMR properties	Carcinogenicity (C)	Cat 1B	RP 1:1 does fulfil criterion (a) of Article 5(1)
	Mutagenicity (M)	Cat 2	
	Toxic for reproduction (R)	no classification required]	
PBT and vPvB properties	Persistent (P) or very Persistent (vP)	not P or vP	RP 1:1 does not fulfil

	Bioaccumulative (B) or very Bioaccumulative (vB)	not B or vB	criterion (e) of Article 5(1) and does not fulfil criterion (d) of Article 10(1)
	Toxic (T)	T	
Endocrine disrupting properties	RP 1:1 is not considered to have endocrine disrupting properties. RP 1:1 does not fulfil criterion (d) of Article 5(1).		
Respiratory sensitisation properties	No classification required. RP 1:1 does not fulfil criterion (b) of Article 10(1).		
Concerns linked to critical effects	RP 1:1 does not fulfil criterion (e) of Article 10(1)		
Proportion of non-active isomers or impurities	The substance does not contain a significant proportion of non-active isomers or impurities. RP 1:1 does not fulfil criterion (f) of Article 10(1).		

Consequently, the following is concluded:

Reaction products of paraformaldehyde and 2-hydroxypropylamine (ratio 1:1) does meet the exclusion criteria laid down in Article 5(1) of Regulation (EU) No 528/2012 by the released formaldehyde being a carcinogen Cat 1B.

Reaction products of paraformaldehyde and 2-hydroxypropylamine (ratio 1:1) does meet the conditions laid down in Article 10(1)(a) of Regulation (EU) No 528/2012, and is therefore considered as a candidate for substitution by meeting the exclusion criteria.

The exclusion and substitution criteria were assessed in line with the "Note on the principles for taking decisions on the approval of active substances under the BPR"<sup>3</sup> and in line with "Further guidance on the application of the substitution criteria set out under article 10(1) of the BPR"<sup>4</sup> agreed at the 54<sup>th</sup> and 58<sup>th</sup> meeting respectively, of the representatives of Member States Competent Authorities for the implementation of Regulation 528/2012 concerning the making available on the market and use of biocidal products. This implies that the assessment of the exclusion criteria is based on Article 5(1) and the assessment of substitution criteria is based on Article 10(1)(a, b, d, e and f).

### 2.2.2. POP criteria

A PBT assessment was performed for RP 1:1 and its hydrolysis products. Based on the available data RP 1:1, 2-hydroxypropylamine and formaldehyde are neither vPvB, nor PBT substances. Furthermore, none of the three substances meets two of the PBT criteria. Therefore, neither the parent nor its hydrolysis products meet the criteria for POPs either.

### 2.2.3. Public consultation for potential candidates for substitution and alternative substances or technologies

As RP 1:1 is considered a candidate for substitution ECHA launched the public consultation in accordance with Article 10(3) of Regulation (EU) No 528/2012. The public consultation took place from 4 November 2016 to 3 January 2017. Four contributions were submitted:

<sup>3</sup> See document: Note on the principles for taking decisions on the approval of active substances under the BPR (available from <https://circabc.europa.eu/d/a/workspace/SpacesStore/c41b4ad4-356c-4852-9512-62e72cc919df/CA-March14-Doc.4.1%20-%20Final%20-%20Principles%20for%20substance%20approval.doc>)

<sup>4</sup> See document: Further guidance on the application of the substitution criteria set out under article 10(1) of the BPR (available from [https://circabc.europa.eu/d/a/workspace/SpacesStore/dbac71e3-cd70-4ed7-bd40-fc1cb92cfe1c/CA-Nov14-Doc.4.4%20-%20Final%20-%20Further%20guidance%20on%20Art10\(1\).doc](https://circabc.europa.eu/d/a/workspace/SpacesStore/dbac71e3-cd70-4ed7-bd40-fc1cb92cfe1c/CA-Nov14-Doc.4.4%20-%20Final%20-%20Further%20guidance%20on%20Art10(1).doc))

one by an industry stakeholder association, two by individual companies and one by a member state. The same contributions were submitted in the consultation on the structurally and toxicologically related substance RP 3:2.

In the member state contribution it is stated that no information on alternatives is available as the product types are not covered by their national authorisation scheme. This may be the case for more member states.

In the three industry contributions information is submitted on the importance of formaldehyde releasers in the control of microbial growth in water-containing products or equipment. In addition, information on alternatives is submitted for all product types.

Three general observations are made in the industry contributions:

- First, it is stated that other formaldehyde releasers are not considered as alternatives as it can be foreseen that these will also be classified as carcinogen category 1B and subsequently meet the exclusion criteria. In total 10 other formaldehyde releasers are under evaluation and one (formaldehyde released from N,N-Methylenebismorpholine or MBM for PT6 and 13) is already approved.
- Second, it is stated that for an effective preservation of many water-based products a bactericide and fungicide is needed. Subsequently, fungicide active substances cannot be regarded as suitable alternatives.
- Last, it is stated that another class of bactericides are the isothiazolinones. Although these are not meeting the substitution criteria it should be considered that these are all classified as strong skin sensitisers. This triggers several obligations for the user making this class of active substances not suitable alternatives.

For PT 6 the only alternative mentioned in the industry contributions is CMI/MI. However, this active substance belongs to the class of isothiazolinones. In addition it is stated that CMI/MI is not soluble in fuel and contains halogen which is not allowed according to German Clean Air Act.

Several other active substances are already approved for PT 6. The only active substance approved for the same use is formaldehyde released from N,N-Methylenebismorpholine or MBM. However, this is a formaldehyde releaser also meeting the exclusion criteria.

The limited information available is insufficient to conclude on the availability of suitable alternatives for the intended uses assessed.

### **2.3 BPC opinion on the application for approval of the active substance RP 1:1 in product type 6**

As the exclusion criteria are met, RP 1:1 should normally not be approved unless one of the conditions for derogation set in Article 5(2) of Regulation (EU) No 528/2012 is met.

If RP 1:1 is approved and included in the Union list of approved active substances, the approval shall be subject to the following specific conditions:

1. Specification: the active substance has to be considered as substance of Unknown or Variable composition or Complex reaction products (UVC). Therefore the minimum purity is 1000 g/kg (100% by wt).
2. RP 1:1 is considered a candidate for substitution in accordance with Article 10(1)(a) of Regulation (EU) No 528/2012.
3. The authorisations of biocidal products are subject to the following condition:
  - a. Products shall only be authorised for use in Member States where at least one of the conditions set in Article 5(2) of Regulation (EU) No 528/2012 is met.

- b. The product assessment shall pay particular attention to the exposures, the risks and the efficacy linked to any uses covered by an application for authorisation, but not addressed in the Union level risk assessment of the active substance. In addition, pursuant to point 10 of Annex VI to Regulation (EU) No 528/2012, the product assessment shall include an evaluation as to whether the conditions of Article 5(2) of Regulation (EU) No 528/2012 can be satisfied.
  - c. In view of the risks identified for the uses assessed, the product assessment shall pay particular attention to:
    - i. Professionals and industrial workers;
    - ii. Sewage Treatment Plant, surface water and the terrestrial compartment.
4. The placing on the market of treated articles is subject to the following condition:
- a. The person responsible for the placing on the market of a treated article treated with or incorporating RP 1:1 shall ensure that the label of that treated article provides the information listed in the second subparagraph of Article 58(3) of the Regulation (EU) No 528/2012.

RP 1:1 does not fulfil the criteria according to Article 28(1) to enable inclusion in Annex I of Regulation (EU) 528/2012 as it is classified as Skin Corr. 1C, Skin Sens. 1A, STOT RE 2, Muta 2, Carc. 1B.

#### **2.4 Elements to be taken into account when authorising products**

1. The active substance RP 1:1 is considered as a candidate for substitution, and consequently the competent authority shall perform a comparative assessment as part of the evaluation of an application for national authorisation.
2. The following recommendations and risk mitigation measures have been identified for the uses assessed. Authorities should consider these risk mitigation measures when authorising products, together with possible other risk mitigation measures, and decide whether these measures are applicable for the concerned product:
  - a. If an unacceptable risk for professional users is identified for the product, safe operational procedures and appropriate organisational measures shall be established. Where exposure cannot be reduced to an acceptable level by other means, products should be used with appropriate personal protective equipment.
  - b. Surface water intended for the abstraction of drinking water may exceed the parametric value of 0.1 µg/L of the Drinking Water Directive 98/83/EC regarding the storage tank bottom water scenario, which is covered by the scenario of PT 13. This may be considered by the relevant national authorities when issuing permits for recovery plants.
  - c. Unacceptable risks are identified for the Sewage Treatment Plant, surface water and the terrestrial compartment regarding the storage tank bottom water scenario, which is covered by the scenario of PT 13. If the risk cannot be reduced to an acceptable level by appropriate risk mitigation measures or by other means, these uses should not be authorised.
  - d. Because no scenario for storage of fuel in storage tanks exists currently, environmental exposure estimates for PT 13 were used for the risk characterisation. Therefore refinements of PT 13 exposure calculations (information on degradation of the hydrolysis products between the last dosing and the start of waste treatment) provided at product authorisation stage are also relevant for the conclusions on the environmental risks for PT 6 (storage tank scenario).

## 2.5 Requirement for further information

Sufficient data have been provided to verify the conclusions on the active substance, permitting the proposal for the approval of RP 1:1. However, further data shall be required as detailed below. Data must be provided as soon as possible but not later than 6 months before the date of approval to the evaluating Competent Authority (Austria):

- a. Regarding the explosive properties of the active substance an experimental test has to be conducted as the substance contains unknown constituents and therefore the waiving cannot be justified by structural considerations.
- b. A specific or highly specific and fully validated analytical method for the determination of the hydrolysis product 2-hydroxypropylamin in water.
- c. For the refinement of the environmental risk assessment: monitoring of the concentration of formaldehyde in three representative spent fluids between the last dosing and the emulsion splitting (using techniques based on Kow and via evaporation) and before the biological waste treatment. If further refinement, is necessary may be an OECD STP simulation test 303 A or 314 should be conducted.