

CLH report

Proposal for Harmonised Classification and Labelling

Based on Regulation (EC) No 1272/2008 (CLP Regulation),
Annex VI, Part 2

Substance Name: S-Methoprene

EC Number: Not available

CAS Number: 65733-16-6

Index Number: Not available

Contact details for dossier submitter:

Pesticide Registration Division,
Department of Agriculture, Food and the Marine,
Backweston Laboratory Complex,
Celbridge,
Co. Kildare,
Ireland.

Version number: 1

Date: July 2015

CONTENTS

Part A.

1	PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING	7
1.1	SUBSTANCE.....	7
1.2	HARMONISED CLASSIFICATION AND LABELLING PROPOSAL	7
1.3	PROPOSED HARMONISED CLASSIFICATION AND LABELLING BASED ON CLP REGULATION	8
2	BACKGROUND TO THE CLH PROPOSAL	10
2.1	HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING	10
2.2	SHORT SUMMARY OF THE SCIENTIFIC JUSTIFICATION FOR THE CLH PROPOSAL	10
2.3	CURRENT HARMONISED CLASSIFICATION AND LABELLING.....	10
2.3.1	<i>Current classification and labelling in Annex VI, Table 3.1 in the CLP Regulation</i>	<i>10</i>
2.3.2	<i>Current classification and labelling in Annex VI, Table 3.2 in the CLP Regulation</i>	<i>10</i>
2.4	CURRENT SELF-CLASSIFICATION AND LABELLING	11
2.4.1	<i>Current self-classification and labelling based on the CLP Regulation criteria</i>	<i>11</i>
3	JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL	11
	SCIENTIFIC EVALUATION OF THE DATA	12
1	IDENTITY OF THE SUBSTANCE	12
1.1	NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE.....	12
1.2	COMPOSITION OF THE SUBSTANCE	13
1.2.1	<i>Composition of test material</i>	<i>13</i>
1.3	PHYSICO-CHEMICAL PROPERTIES	13
2	MANUFACTURE AND USES	15
2.1	MANUFACTURE	15
2.2	IDENTIFIED USES	15
3	CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES	16
3.1	<i>[INSERT HAZARD CLASS WHEN RELEVANT AND REPEAT SECTION IF NEEDED]</i>	<i>16</i>
3.1.1	<i>Summary and discussion of S-Methoprene</i>	<i>16</i>
3.1.2	<i>Comparison with criteria.....</i>	<i>16</i>
3.1.3	<i>Conclusions on classification and labelling</i>	<i>16</i>
4	HUMAN HEALTH HAZARD ASSESSMENT.....	16
4.1	TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)	16
4.1.1	<i>Non-human information.....</i>	<i>20</i>
4.1.2	<i>Human information.....</i>	<i>20</i>
4.1.3	<i>Summary and discussion on toxicokinetics</i>	<i>20</i>
4.2	ACUTE TOXICITY	21
4.2.1	<i>Non-human information.....</i>	<i>21</i>
4.2.1.1	<i>Acute toxicity: oral</i>	<i>21</i>
4.2.1.2	<i>Acute toxicity: inhalation.....</i>	<i>22</i>
4.2.1.3	<i>Acute toxicity: dermal.....</i>	<i>22</i>
4.2.1.4	<i>Acute toxicity: other routes</i>	<i>23</i>
4.2.2	<i>Human information.....</i>	<i>23</i>
4.2.3	<i>Summary and discussion of acute toxicity</i>	<i>23</i>
4.2.4	<i>Comparison with criteria.....</i>	<i>23</i>
4.2.5	<i>Conclusions on classification and labelling</i>	<i>23</i>
4.3	SPECIFIC TARGET ORGAN TOXICITY – SINGLE EXPOSURE (STOT SE).....	24
4.3.1	<i>Summary and discussion of Specific target organ toxicity – single exposure.....</i>	<i>24</i>
4.3.2	<i>Comparison with criteria.....</i>	<i>24</i>

4.3.3	<i>Conclusions on classification and labelling</i>	24
4.4	IRRITATION	24
	<i>Skin irritation</i>	24
4.4.1		24
4.4.1.1	Non-human information	24
4.4.1.2	Human information	25
4.4.1.3	Summary and discussion of skin irritation	25
4.4.1.4	Comparison with criteria	25
4.4.1.5	Conclusions on classification and labelling	25
4.4.2	<i>Eye irritation</i>	26
4.4.2.1	Non-human information	26
4.4.2.2	Human information	27
4.4.2.3	Summary and discussion of eye irritation	27
4.4.2.4	Comparison with criteria	27
4.4.2.5	Conclusions on classification and labelling	27
4.4.3	<i>Respiratory tract irritation</i>	27
4.4.3.1	Non-human information	27
4.4.3.2	Human information	27
4.4.3.3	Summary and discussion of respiratory tract irritation	27
4.4.3.4	Comparison with criteria	27
4.4.3.5	Conclusions on classification and labelling	27
4.5	CORROSIVITY	27
4.5.1	<i>Non-human information</i>	28
4.5.2	<i>Human information</i>	28
4.5.3	<i>Summary and discussion of corrosivity</i>	28
4.5.4	<i>Comparison with criteria</i>	28
4.5.5	<i>Conclusions on classification and labelling</i>	28
4.5.6	<i>Skin sensitisation</i>	29
4.5.6.1	Non-human information	29
4.5.6.2	Human information	29
4.5.6.3	Summary and discussion of skin sensitisation	30
4.5.6.4	Comparison with criteria	30
4.5.6.5	Conclusions on classification and labelling	30
4.5.7	<i>Respiratory sensitisation</i>	30
4.5.7.1	Non-human information	30
4.5.7.2	Human information	30
4.5.7.3	Summary and discussion of respiratory sensitisation	30
4.5.7.4	Comparison with criteria	30
4.5.7.5	Conclusions on classification and labelling	30
4.6	REPEATED DOSE TOXICITY	30
4.6.1	<i>Non-human information</i>	32
4.6.1.1	Repeated dose toxicity: oral	32
4.6.1.2	Repeated dose toxicity: inhalation	37
4.6.1.3	Repeated dose toxicity: dermal	37
4.6.1.4	Repeated dose toxicity: other routes	37
4.6.1.5	Human information	37
4.6.1.6	Other relevant information	37
4.6.1.7	Summary and discussion of repeated dose toxicity	37
4.7	SPECIFIC TARGET ORGAN TOXICITY (CLP REGULATION) – REPEATED EXPOSURE (STOT RE)	37
4.7.1	<i>Summary and discussion of repeated dose toxicity findings relevant for classification as STOT RE according to CLP Regulation</i>	37
4.7.2	<i>Comparison with criteria of repeated dose toxicity findings relevant for classification as STOT RE</i>	37
4.7.3	<i>Conclusions on classification and labelling of repeated dose toxicity findings relevant for classification as STOT RE</i>	38
4.8	GERM CELL MUTAGENICITY (MUTAGENICITY)	38
4.8.1	<i>Non-human information</i>	38
4.8.1.1	<i>In vitro</i> data	38
4.8.1.2	<i>In vivo</i> data	39
4.8.2	<i>Human information</i>	39
4.8.3	<i>Other relevant information</i>	39
4.8.4	<i>Summary and discussion of mutagenicity</i>	39
4.8.5	<i>Comparison with criteria</i>	39
4.8.6	<i>Conclusions on classification and labelling</i>	39
4.9	CARCINOGENICITY	40

4.9.1	<i>Non-human information</i>	40
4.9.1.1	Carcinogenicity: oral.....	40
Dosage level (ppm)	41
4.9.1.2	Carcinogenicity: inhalation.....	46
4.9.1.3	Carcinogenicity: dermal.....	46
4.9.2	<i>Human information</i>	46
4.9.3	<i>Other relevant information</i>	46
4.9.4	<i>Summary and discussion of carcinogenicity</i>	46
4.9.5	<i>Comparison with criteria</i>	46
4.9.6	<i>Conclusions on classification and labelling</i>	47
4.10	TOXICITY FOR REPRODUCTION	47
4.10.1	<i>Effects on fertility</i>	48
4.10.1.2	Human information.....	56
4.10.2	<i>Developmental toxicity</i>	56
4.10.2.1	Non-human information.....	56
4.10.2.2	Human information.....	59
4.10.3	<i>Other relevant information</i>	59
4.10.4	<i>Summary and discussion of reproductive toxicity</i>	59
4.10.5	<i>Comparison with criteria</i>	60
4.10.6	<i>Conclusions on classification and labelling</i>	60
4.11	OTHER EFFECTS	60
4.11.1	<i>Non-human information</i>	60
4.11.1.1	Neurotoxicity.....	60
4.11.1.2	Immunotoxicity.....	61
4.11.1.3	Specific investigations: other studies.....	61
4.11.1.4	Human information.....	61
4.11.2	<i>Summary and discussion</i>	61
4.11.3	<i>Comparison with criteria</i>	61
4.11.4	<i>Conclusions on classification and labelling</i>	61
5	ENVIRONMENTAL HAZARD ASSESSMENT	62
5.1	DEGRADATION	62
5.1.1	<i>Stability</i>	62
5.1.2	<i>Biodegradation</i>	63
	Biodegradation estimation.....	63
5.1.2.1	63
5.1.2.2	Screening tests.....	64
5.1.2.3	Simulation tests.....	64
5.1.3	<i>Summary and discussion of degradation</i>	64
5.2	ENVIRONMENTAL DISTRIBUTION	65
5.2.1	<i>Adsorption/Desorption</i>	65
5.2.2	<i>Volatilisation</i>	65
5.2.3	<i>Distribution modelling</i>	66
5.3	AQUATIC BIOACCUMULATION	66
5.3.1	<i>Aquatic bioaccumulation</i>	66
5.3.1.1	Bioaccumulation estimation.....	66
5.3.1.2	Measured bioaccumulation data.....	66
5.3.2	<i>Summary and discussion of aquatic bioaccumulation</i>	66
5.4	AQUATIC TOXICITY	67
5.4.1	<i>Fish</i>	68
5.4.1.1	Short-term toxicity to fish.....	68
5.4.1.2	Long-term toxicity to fish.....	69
5.4.2	<i>Aquatic invertebrates</i>	69
5.4.2.1	Short-term toxicity to aquatic invertebrates.....	69
5.4.2.2	Long-term toxicity to aquatic invertebrates.....	69
5.4.3	<i>Algae and aquatic plants</i>	70
5.4.3.1	Short-term toxicity to algae and aquatic plants.....	70
5.4.3.2	Long-term toxicity to algae and aquatic plants.....	70
5.4.4	<i>Other aquatic organisms (including sediment)</i>	70
5.4.4.1	Short-term toxicity to other aquatic organisms (including sediment).....	70
5.4.4.2	Long-term toxicity to other aquatic organisms (including sediment).....	70
5.5	COMPARISON WITH CRITERIA FOR ENVIRONMENTAL HAZARDS (SECTIONS 5.1 – 5.4)	70
5.6	CONCLUSIONS ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS (SECTIONS 5.1 – 5.4)	71

6 OTHER INFORMATION..... 71
7 REFERENCES..... 71
8 ANNEXES..... 71

Part A.

1 PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING

1.1 Substance

Table 1: Substance identity

Substance name:	S-Methoprene
EC number:	Not available
CAS number:	65733-16-6
Annex VI Index number:	Not available
Degree of purity:	Min. 95% w/w
Impurities:	Confidential Information. See Confidential Data & Information, S-Methoprene CAR. (See Technical dossier in IUCLID 5, section 1.2)

1.2 Harmonised classification and labelling proposal

S-Methoprene is not classified according to CLP Regulation (EC) No. 1272/2008.

Table 2: The current Annex VI entry and the proposed harmonised classification

	CLP Regulation
Current entry in Annex VI, CLP Regulation	No current entry
Current proposal for consideration by RAC	Aquatic Acute 1 H400 Aquatic Chronic 1 H410 Acute M-factor of 1 and Chronic M-factor of 1
Resulting harmonised classification (future entry in Annex VI, CLP Regulation)	Aquatic Acute 1 H400 Aquatic Chronic 1 H410 Acute M-factor of 1 and Chronic M-factor of 1

1.3 Proposed harmonised classification and labelling based on CLP Regulation

Table 3: Proposed classification according to the CLP Regulation

CLP Annex I ref	Hazard class	Proposed classification	Proposed SCLs and/or M-factors	Current classification ¹⁾	Reason for no classification ²⁾
2.1.	Explosives	-	-	-	Conclusive but not sufficient for classification
2.2.	Flammable gases	-	-	-	Not applicable to liquids
2.3.	Flammable aerosols	-	-	-	Not applicable to liquids
2.4.	Oxidizing gases	-	-	-	Not applicable to liquids
2.5.	Gases under pressure	-	-	-	Not applicable to liquids
2.6.	Flammable liquids	-	-	-	Conclusive but not sufficient for classification
2.7.	Flammable solids	-	-	-	Not applicable to liquids
2.8.	Self-reactive substances and mixtures	-	-	-	Conclusive but not sufficient for classification
2.9.	Pyrophoric liquids	-	-	-	Conclusive but not sufficient for classification
2.10.	Pyrophoric solids	-	-	-	Not applicable to liquids
2.11.	Self-heating substances and mixtures	-	-	-	Conclusive but not sufficient for classification
2.12.	Substances and mixtures which in contact with water emit flammable gases	-	-	-	Conclusive but not sufficient for classification
2.13.	Oxidizing liquids	-	-	-	Conclusive but not sufficient for classification
2.14.	Oxidizing solids	-	-	-	Not applicable to liquids
2.15.	Organic peroxides	-	-	-	Conclusive but not sufficient for classification
2.16.	Substance and mixtures corrosive to metals	-	-	-	Data lacking

3.1.	Acute toxicity - oral	-	-	-	-
	Acute toxicity - dermal	-	-	-	Conclusive but not sufficient for classification
	Acute toxicity - inhalation	-	-	-	Conclusive but not sufficient for classification
3.2.	Skin corrosion / irritation	-	-	-	Conclusive but not sufficient for classification
3.3.	Serious eye damage / eye irritation	-	-	-	Conclusive but not sufficient for classification
3.4.	Respiratory sensitisation	-	-	-	Data lacking
3.4.	Skin sensitisation	-	-	-	Conclusive but not sufficient for classification
3.5.	Germ cell mutagenicity	-	-	-	Conclusive but not sufficient for classification
3.6.	Carcinogenicity	-	-	-	-
3.7.	Reproductive toxicity	-	-	-	-
3.8.	Specific target organ toxicity –single exposure	-	-	-	Conclusive but not sufficient for classification
3.9.	Specific target organ toxicity – repeated exposure	-	-	-	Conclusive but not sufficient for classification
3.10.	Aspiration hazard	-	-	-	Data lacking
4.1.	Hazardous to the aquatic environment	H400 H410	Acute M-factor.=.1 Chronic M-factor = 1	none	
5.1.	Hazardous to the ozone layer	-	-	-	

¹⁾Including specific concentration limits (SCLs) and M-factors

²⁾Data lacking, inconclusive, or conclusive but not sufficient for classification

Labelling: Pictogram



GHS09

Signal word:

Warning

Hazard statements:

H410 Very toxic to aquatic life with long lasting effects.

Precautionary statements:

P273 Avoid release to the environment

P391 Collect spillage

P501 Dispose of contents/ container in accordance with applicable regulations

Acute M-factor of 1 and Chronic M-factor of 1

BACKGROUND TO THE CLH PROPOSAL

2.1 History of the previous classification and labelling

S-Methoprene has no previous human health or environmental classification and labeling elements.

2.2 Short summary of the scientific justification for the CLH proposal

Physical Effects CLH proposal

No physical effects classification proposal is required for S-Methoprene.

Health Effects CLH proposal

No health effects classification proposal is required for S-Methoprene.

Environmental CLH proposal

- H400 (which is implicit in the H410 labelling) follows from the acute toxicity of the active substance to *Daphnia magna*: $LC_{50} < 1$ mg a.s./L (48 hour $LC_{50} = 0.22$ mg a.s./L). An M-factor of 1 is applicable based on $0.1 < LC_{50} \leq 1$ mg a.s./l.
- H410 follows from the chronic toxicity of the active substance to *Daphnia magna*: $NOEC \leq 1$ mg a.s./L ($NOEC = 0.019$ mg/L) and the fact that the active substance is not rapidly degradable. Additionally, the $\log K_{ow} > 6$. An M-factor of 1 is applicable based on $0.01 < NOEC \leq 0.1$ mg/l.
- GHS09 Pictogram is required for ‘Aquatic acute 1’ and ‘Aquatic chronic 1’ category substance.
- Signal word ‘Warning’ is required for ‘Aquatic acute 1’ and ‘Aquatic chronic 1’ category substance.
- The statements P273, P391 and P501 follow a general precautionary approach for dangerous substances.

2.3 Current harmonised classification and labelling

2.3.1 Current classification and labelling in Annex VI, Table 3.1 in the CLP Regulation

There is no current harmonised classification and labelling for S-Methoprene in Annex VI, Table 3.1 under the CLP Regulation.

2.3.2 Current classification and labelling in Annex VI, Table 3.2 in the CLP Regulation

There is no current harmonised classification and labelling for S-Methoprene in Annex VI, Table 3.1 under the CLP Regulation.

2.4 Current self-classification and labelling

Not available - no information regarding S-Methoprene was found in the C&L Inventory database (database contains classification and labelling information on notified and registered substances received from manufacturers and importers; <http://www.echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database>).

2.4.1 Current self-classification and labelling based on the CLP Regulation criteria

Not available.

3 JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL

S-Methoprene is a biocidal active substance approved for use in biocidal products for product type 18 from 1 September 2015 as stated in Regulation (EC) 91/2014 of 31 January 2014. The classification and labelling proposal includes environmental toxicity endpoints and needs to be evaluated under the CLP Regulation (EC) No. 1272/2008

Part B.

SCIENTIFIC EVALUATION OF THE DATA


1 IDENTITY OF THE SUBSTANCE

1.1 Name and other identifiers of the substance

Table 5: Substance identity

EC number:	Not available
EC name:	Not available
CAS number (EC inventory):	Not available
CAS number:	65733-16-6
CAS name:	Not available
IUPAC name:	Isopropyl (2E,4E,7S)-11-methoxy-3,7,11-trimethyldodeca- 2,4-dienoate
CA Index Name	2,4-Dodecadienoic acid, 11-methoxy-3,7,11-trimethyl-1-methylethyl ester, (2E,4E,7S)-
CLP Annex VI Index number:	Not available
Molecular formula:	C ₁₉ H ₃₄ O ₃
Molecular weight range:	310.48 g/mol

Structural formula:



1.2 Composition of the substance

Table 6: Constituents (non-confidential information)

Constituent	Typical concentration	Concentration range	Remarks
S-Methoprene	95%	-	-

Current Annex VI entry:

Table 7: Impurities (non-confidential information)

Impurity	Typical concentration	Concentration range	Remarks
-	-	-	-
All impurities have been claimed confidential			

Current Annex VI entry:

Table 8: Additives (non-confidential information)

Additive	Function	Typical concentration	Concentration range	Remarks
-	-	-	-	-

Current Annex VI entry:

1.2.1 Composition of test material

1.3 Physico-chemical properties

Table 9: Summary of physico-chemical properties

Property	Value	Reference	Comment (e.g. measured or estimated)
Freezing point (state purity)	Purity: 98.3% < -22°C	Laky, V., 2006a	
Boiling point (state purity)	Purity: 99.6% 279.9 °C	Laky, V., 2006a	
Temperature of decomposition	Not applicable as the boiling point was estimated.		
Appearance (state purity)	Purity: > 95% A transparent pale yellow liquid at 24°C	Anderson, W., 1999	

	with a faint, fruity, waxy odour.		
Relative density (state purity)	Purity: > 95% 0.924 g/ml at 20°C	Anderson, W., 1999	
Surface tension	Purity: 98.3% 50.1 mN/m at 20°C (1mg/l)	Laky, V., 2006f	
Vapour pressure (in Pa, state temperature)	Purity: 98.1 % 0.623 mPa at 20°C 1.08 mPa at 25°C	Bates, M., 2007	
Henry's law constant (Pa m ³ mol ⁻¹)	0.0306 Pa x m ³ /mol at 20°C	Bates, M., 2007	
Solubility in water (g/l or mg/l, state temperature)	Purity: > 95% 6.85 mg/l at 20 °C	Anderson, W., 1999	
Solubility in organic solvents (in g/l or mg/l, state temperature) (Annex IIIA, point III.1)	Purity: 98.1% Hexane: > 5 10 ⁵ mg/l Methanol: > 4.5 10 ⁵ mg/l Acetone: > 5 10 ⁵ mg/l Temperature: 20 ± 1 °C	Laky, V., 2006a	
Stability in organic solvents used in biocidal products including relevant breakdown products (IIIA, point III.2)	Not required as no organic solvents are present in the technical.	Laky, V., 2006a	
Partition coefficient (log P _{OW}) (state temperature)	LogKow = 6.34	Rivendell International 2012	Calculated
Hydrolytic stability (DT ₅₀) (state pH and temperature) (point VII.7.6.2.1)	pH 1.2: 17 hours at 37 ± 0.5°C	Anderson, W., 1999	
	pH 4: Stable at 25 ± 0.5°C, 37 ± 0.5°C and 50 ± 0.5°C	Laky, V. (2002a),	
	pH 7: Stable at 25 ± 0.5°C, 37 ± 0.5°C and 50 ± 0.5°C		
	pH 9: Stable at 25 ± 0.5°C, 37 ± 0.5°C and 50 ± 0.5°C		
Dissociation constant (not stated in Annex IIA or IIIA; additional data requirement from TNsG)	Not required as S-Methoprene does not dissociate in water.		
UV/VIS absorption (max.) (if absorption > 290 nm state ε at	Purity: 95% <u>90% Neutral Methanol:</u> λ _{max} 264 nm; ε 26,700 <u>90% Acidified</u>	Anderson, W., 1999	

wavelength)	<u>Methanol:</u> λ_{\max} 264 nm; ϵ 26,600 90% Alkalinized <u>Methanol:</u> λ_{\max} 266 nm; ϵ 27,450		
Flammability	Purity: 98.3% 263 °C	Laky, V., 2006d	
Explosive properties	The molecular structure of S-Methoprene indicates that the substance has little or no explosive properties.		

2 MANUFACTURE AND USES

2.1 Manufacture

Not relevant for this dossier.

2.2 Identified uses

Insecticide (PT 18) – Biocide.

3 CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES

Table 10: Summary table for relevant physico-chemical studies

Method	Results	Remarks	Reference
-	-	-	-
-	-	-	-
-	-	-	-
-	-	-	-

3.1 *[Insert hazard class when relevant and repeat section if needed]*

3.1.1 Summary and discussion of S-Methoprene

Not applicable. S-Methoprene does not classify with respect to physical chemistry.

3.1.2 Comparison with criteria

No classification is warranted for S-Methoprene regarding physico-chemical hazardous properties based on study results summarised in Table 9 above.

3.1.3 Conclusions on classification and labelling

No classification is warranted for S-Methoprene with respect to physical chemistry.

4 HUMAN HEALTH HAZARD ASSESSMENT

4.1 Toxicokinetics (absorption, metabolism, distribution and elimination)

In a toxicokinetic study, [¹⁴C]-S-Methoprene was administered to male and female Sprague-Dawley rats, and its excretion, absorption, distribution and metabolism were examined. The study consisted of four groups: a single low level dose group (25 mg/kg bw), a single high level dose group (250 mg/kg bw), a low level repeat dose group (seven daily oral doses of 25 mg/kg bw), and a bile duct cannulated group (single administration at 25 mg/kg bw).

All rats survived the study and no signs of toxicity were observed. Measurement of radioactivity in the plasma revealed that in the low dose group the peak plasma concentrations for males and females were at 6 and 12 hours respectively. In the high dose group, the peak plasma concentrations were seen at 4 hours in males and 6 hours in females.

The majority of [¹⁴C]-S-Methoprene was excreted within 24 to 48 hours of administration, indicating that it is rapidly eliminated from the body. The primary routes of excretion of the

compound were in the faeces and expired air, with a lesser amount recovered in urine and cage rinses.

For all dosing regimens, the tissue radioactivity was negligible at 96 hours after administration, with the exception of the white fat, which contained up to 4.633 % of the administered radioactivity. This is in line with previous reports suggesting that S-Methoprene is a lipophilic compound. However, in all other tissues the levels of radioactivity decreased between the 6 and 96 hour time points, indicating that neither the compound nor its metabolites accumulate in these tissues.

In all dosing regimens, males excreted a higher percentage of the [¹⁴C]-S-Methoprene-derived radioactivity in the faeces than females. Sex differences in the tissue distribution of radioactivity were observed at the low level dose of [¹⁴C]-S-Methoprene. Female tissue contained higher radioactivity than male tissue in this group.

The recovery of radioactivity following administration of [14C]-S-Methoprene to rats indicated that the majority of radioactivity (40-60 % AR) was excreted via the faecal route and the majority of this was the unchanged parent compound [¹⁴C]-S-Methoprene (Quotient Bioresearch Report LIH/02). Of the remaining radioactivity 14-28 % AR was excreted via expired air indicating extensive metabolism of the parent molecule and incorporation of the radiolabel into endogenous components.

Based on the available evidence it is concluded that the metabolism of S-Methoprene proceeds via incorporation into natural products. Definitive identification of radioactive metabolites is rendered almost impossible by the high background observed from unlabelled natural components already in the circulation. The presence of S-Methoprene was confirmed using mass spectrometric techniques, whilst the presence of Methoprene acid, acetate and glycine were determined using co-chromatography with unlabelled or radiolabelled reference materials. The only component in rat excreta samples that represented > 10 % of administered radioactivity was RF23 and this component was shown to be the parent compound S-Methoprene.


Previous studies summarised by the World Health Organisation (WHO) (1998 and 1999) reached similar conclusions regarding the metabolism of Methoprene in animals. In the current study both S-Methoprene and S-Methoprene acid could be detected using GC-MS analysis. S-Methoprene was detected in faeces, whilst no other metabolites were detected. It would be expected that any components with structures similar to S-Methoprene would have been detected if present in the samples. The WHO document references metabolism in large animals (Cow, Steer and Hen) whereas the current study was performed in the rat. Generally, the smaller the animal species the higher the rate of metabolic turnover. The results are therefore consistent with rapid degradation/metabolism of Methoprene into C-1 and C-2 units and fully consistent with previously published data on the metabolism of Methoprene in the rat. In the current study there was one component (RF23, identified as S-Methoprene) which was greater than 10 % dose and four components (RF21, RF20, RF13 and RF7) that were greater than 5 %. These components were not identified in the current study but it should be noted that these were present in the faeces and may arise from unabsorbed material travelling through the GI tract following an oral dose and will therefore be of no toxicological concern.

It is concluded that the metabolism of S-Methoprene proceeds via extensive degradation of the aliphatic chains and this may lead to production of saturated and/or unsaturated aliphatic, alicyclic linear primary alcohols, aldehydes and acids as intermediates to the ultimate production of acetate.

In addition, a literature search has been conducted to provide basic toxicokinetic data for S-Methoprene. The bio-kinetics and metabolism of Methoprene were investigated in various mammalian species such as rat and guinea pig following a single oral administration showing that Methoprene is rapidly absorbed and excreted in urine, faeces and expired air. S-Methoprene is metabolised as a methyl branched fatty acid food and also via conjugation (O-dealkylation). Following treatment with glucuronidase, the two major metabolites in urine were 11-methoxy-3,7,11-trimethyldodeca-2,4-dienoic acid and 11-hydroxy-3,7,11-trimethyldodeca-2,4-dienoic acid and accounted for 75% of the radiolabel in urine. 77% of the recovered radiolabel represented intact Methoprene in the faeces.

An overall value for oral absorption to be used in the risk assessment (35%) was derived from the basic toxicokinetics studies.

Proposed metabolic pathway for [¹⁴C]-S-Methoprene



Dermal Absorption

A study investigating the rate and extent of absorption of S-Methoprene following topical application of a single formulation of the test substance to human skin was conducted. The mean percentage recovery of the applied test substance was $100 \pm 10\%$. The mean percentage found in the receptor fluid was 0.04%, the mean percentage found remaining in the skin was 1.61%, the mean percentage found in the stratum corneum (tape strips 6-20) was 1.21% and the mean percentage found remaining unabsorbed was 97.14% at 24 hours. As strips 3-5 were also considered part of the potentially absorbed dose an additional 0.58% was added to the value calculated by the Notifier

(2.86%). From this study it was determined the dermal absorption of [¹⁴C]-S-Methoprene is 3.44 ≈ 3.5%.

4.1.1 Non-human information

4.1.2 Human information

None available.

4.1.3 Summary and discussion on toxicokinetics

[¹⁴C]-S-Methoprene was administered to male and female Sprague-Dawley rats, and its excretion, absorption, distribution and metabolism were examined. Measurement of radioactivity in the plasma revealed that in the low dose group the peak plasma concentrations for males and females were at 6 and 12 hours respectively. In the high dose group, the peak plasma concentrations were seen at 4 hours in males and 6 hours in females.

The majority of [¹⁴C]-S-Methoprene was excreted within 24 to 48 hours of administration, indicating that it is rapidly eliminated from the body. The primary routes of excretion of the compound were in the faeces and expired air, with a lesser amount recovered in urine and cage rinses.

For all dosing regimens, the tissue radioactivity was negligible at 96 hours after administration, with the exception of the white fat, which contained up to 4.633 % of the administered radioactivity.

S-Methoprene is metabolised as a methyl branched fatty acid food and also via conjugation (O-dealkylation). Following treatment with glucuronidase, the two major metabolites in urine were 11-methoxy-3,7,11-trimethyldodeca-2,4-dienoic acid and 11-hydroxy-3,7,11-trimethyldodeca-2,4-dienoic acid and accounted for 75% of the radiolabel in urine. 77% of the recovered radiolabel represented intact Methoprene in the faeces.

An overall value for oral absorption to be used in the risk assessment (35%) was derived from the basic toxicokinetics studies.

The dermal absorption of [¹⁴C]-S-Methoprene was determined in an in-vitro dermal absorption study to be 3.44 ≈ 3.5%.

4.2 Acute toxicity

Table 11: Summary table of relevant acute toxicity studies

Method	Results	Remarks	Reference
OECD 401/US EPA 81-1. Rat HSD;SD 5050 mg/kg bw Single oral dose)	Value LD₅₀ / LC₅₀ > 5050 mg/kg. S-Methoprene does not require classification.	No deaths occurred at the 5050 mg/kg dose level. Clinical observations included crust around the nose, piloerection, diarrhoea, activity decrease and an oily yellow substance at the base of the tail, all of which were no longer evident by Day 6 of the study. Body weight gain remained unaffected. Terminal necropsy revealed no abnormalities	Kuhn, J. O. (1999a) S-Methoprene CAR IIIA 6.1.1
OECD 402/US EPA 81-1.(Rabbit Albino New Zealand White Male / Female 5/sex 5050mg/kg bw Single dermal dose)	Value LD₅₀ / LC₅₀ > 5050 mg/kg. S-Methoprene does not require classification.	Erythema was observed on Day 1, and no longer evident on Day 4. Signs of dermal irritation including erythema and desquamation were observed on Day 1 but were no longer evident on Day 4. Body weight gain was unaffected by the administration of S-Methoprene. Terminal necropsy revealed no observable abnormalities	Kuhn, J. O. (1999b) S-Methoprene CAR IIIA 6.1.2
OECD 403/US EPA 81-3. Rat 2.38 mg/l	Value LD₅₀ / LC₅₀ > 2.38 mg/L S-Methoprene does not require classification.	Clinical signs of toxicity included activity decrease and piloerection in both sexes. Red staining around the nose in males. Animals were asymptomatic by Day 1. No mortalities were recorded. Terminal necropsy revealed no observable abnormalities with the exception of discoloured lungs and a swollen large intestine in one male and discoloured lungs in two females. Body weight gain was generally unaffected by administration of the test substance however, one male failed to gain weight and one female lost weight during the first week.	Leeper, L. (1999) S-Methoprene CAR IIIA 6.1.3

*New studies submitted for the Biocides Review (2013).

4.2.1 Non-human information

4.2.1.1 Acute toxicity: oral

Kuhn, J.O. (1999a), Acute oral toxicity study in rats: The acute oral toxicity of S-Methoprene was investigated in Harlan Sprague Dawley rats. S-Methoprene was administered by oral gavage to

male and female rats (5/sex) at a dose level of 5050 mg/kg. The study is comparable to OECD guideline 401. No deaths occurred at the 5050 mg/kg dose level. Clinical observations included crust around the nose, piloerection, diarrhoea, activity decrease and an oily yellow substance at the base of the tail, all of which were no longer evident by Day 6 of the study. Body weight gain remained unaffected. Terminal necropsy revealed no abnormalities.

Based on the results from the acute exposure oral toxicity study in rats, the acute oral LD₅₀ for S-Methoprene was determined to be > 5050 mg/kg. The acute oral LD₅₀ of S-Methoprene in rats was determined to be greater than 5050 mg/kg.

In accordance with the provisions of CLP Regulation (EC) No. 1272/2008, S-Methoprene does not require classification for acute oral toxicity.

4.2.1.2 Acute toxicity: inhalation

Leeper, L. (1999), Acute inhalation toxicity study in rats. The acute inhalation toxicity of S-Methoprene in Sprague Dawley rats was investigated by administering a single dose of S-Methoprene to one group of 5/sex rats as a liquid aerosol by the inhalation route for an exposure duration of four hours and at a concentration of 2.38 mg/L. The study is comparable to OECD 403.

Clinical signs of toxicity included activity decrease and piloerection in both sexes and red staining around the nose in males. Animals were asymptomatic by Day 1. No mortalities were recorded throughout the study period. Terminal necropsy revealed no observable abnormalities with the exception of discoloured lungs and a swollen large intestine in one male and discoloured lungs in two females. Body weight gain was generally unaffected by administration of the test substance however, one male failed to gain weight and one female lost weight during the first week.

Table 12: Mortality data following inhalation of S-Methoprene technical

Dose (mg/L)	Males Mortality	Time of death – hrs (no of animals)	Dose (mg/L)	Females Mortality	Time of death – hrs (no of animals)
2.38	0/5	-	2.38	0/5	-

The acute inhalation LC₅₀ of S-Methoprene in male and female albino SD rats was determined to be greater than 2.38 mg/L. In accordance with the provisions of CLP Regulation (EC) No. 1272/2008, S-Methoprene does not require classification for acute inhalation toxicity.

4.2.1.3 Acute toxicity: dermal

Kuhn, J. O. (1999b), Acute dermal toxicity study in rabbits. The acute dermal toxicity of S-Methoprene was investigated by applying a single dose of S-Methoprene at a concentration of 5050 mg/kg to the skin of 5 albino New Zealand white rabbits/sex. The study is comparable to OECD guideline 402 and is described. Clinical observations revealed soft faeces at 1 and 4 hrs exposure to S-Methoprene in the male rabbit but this reaction was no longer evident by Day 1 of the examination period. Erythema was observed on Day 1, and no longer evident on Day 4. Signs of dermal irritation including erythema and desquamation were observed on Day 1 but were no longer evident on Day 4. Body weight gain was unaffected by the administration of S-Methoprene. Terminal necropsy revealed no observable abnormalities.

Table 13: Acute Dermal Toxicity in the Rabbit (Limit Test): Mortality Data

Dose (mg/ kg)	Males Mortality	Time of death – hrs (no of animals)	Dose (mg/kg)	Females Mortality	Time of death – hrs (no of animals)
5050	0/5	-	5050	0/5	-

The dermal LD₅₀ of S-Methoprene in rabbits was determined to be greater than 5050 mg/kg. In accordance with the provisions of CLP Regulation (EC) No. 1272/2008, S-Methoprene does not require classification for acute inhalation toxicity.

4.2.1.4 Acute toxicity: other routes

No data.

4.2.2 Human information

No data.

4.2.3 Summary and discussion of acute toxicity

Conclusion: The acute oral LD₅₀ and dermal LD₅₀ of S-Methoprene in rats were determined to be greater than 5050 mg/kg. The acute inhalation LC₅₀ of S-Methoprene in rats was found to be greater than 2.38mg/l. Clinical signs of intoxication were non-specific and reversible (piloerection, diarrhoea, decreased activity and some dermal irritation in the dermal study). All studies were carried out using protocols comparable to accepted guidelines and according to GLP (self-certified). In accordance with the provisions of CLP Regulation (EC) No. 1272/2008, S-Methoprene remains unclassified and requires no symbols or risk phrases for acute oral, dermal and inhalation toxicity.

4.2.4 Comparison with criteria

The classification criteria for the acute oral, dermal or inhalation toxicity of S-methoprene are not met.

4.2.5 Conclusions on classification and labelling

Taking the rat oral data, (greater than 5050 mg/kg bw/day) S-Methoprene does not classify for acute oral toxicity when compared to the requirements of the CLP Regulation (EC) No. 1272/2008. Equally, dermal or inhalation exposure study results do not warrant classification under CLP Regulation (EC) No. 1272/2008.

4.3 Specific target organ toxicity – single exposure (STOT SE)

4.3.1 Summary and discussion of Specific target organ toxicity – single exposure

There is no indication from the dossier presented that specific target organ toxicity will result from a single exposure.

4.3.2 Comparison with criteria

Not applicable.

4.3.3 Conclusions on classification and labelling

There is no indication from the dossier presented that specific target organ toxicity will result from a single exposure.

4.4 Irritation

4.4.1 Skin irritation

Table 14: Summary table of relevant skin irritation studies

Method	Results	Remarks	Reference
Skin irritation in NZW rabbits. S-Methoprene OECD 404	Non-irritating	-	Kuhn, J.O. (1999c) S-Methoprene CAR IIIA 6.1.4/1

4.4.1.1 Non-human information

Kuhn, J.O. (1999c), Primary Dermal Irritation Study In Rabbits. In a primary dermal irritation study, 3 young adult (2 males and 1 female) white albino New Zealand rabbits were dermally exposed to 0.5 mL S-Methoprene technical (LX 125-03) for a single 4-hour application to one intact site on each animal. The trunk of the animals was wrapped with a semi-permeable dressing and secured with strips of tape to retard evaporation and prevent possible ingestion of the test substance. At the end of the exposure period, the wrappings were removed and the skin was gently wiped with water and a clean cloth to remove any residual test substance. The acute dermal irritation index was calculated and S-Methoprene was classified according to the Draize method.

Erythema was observed in all rabbits at the 1 hr observation period and in one male and female at the 24 hr observation period. All rabbits returned to normal at the 48 hour observation period. No oedema was observed in any of the animals at throughout the study. The average score for erythema and oedema for all animals at 24-72 hours was 0.11 and 0, respectively.

Table 15: Dermal Irritation scores following exposure to S-Methoprene

Animal no.	Erythema						Oedema					
	1	2	3	4	5	6	1	2	3	4	5	6
After 1 hr	1	1	1	1	1	1	0	0	0	0	0	0

After 24 hr	0	1	0	1	0	0	0	0	0	0	0	0
After 48 hr	0	0	0	0	0	0	0	0	0	0	0	0
After 72 hr	0	0	0	0	0	0	0	0	0	0	0	0
Mean score 24-72 hr	0.11						0.0					

On the basis of reactions observed in this study and the criteria defined in CLP Regulation (EC) No. 1272/2008, S-Methoprene does not require classification for dermal irritation.

4.4.1.2 Human information

Not available.

4.4.1.3 Summary and discussion of skin irritation

S-Methoprene was not irritating to the skin in the studies presented.

4.4.1.4 Comparison with criteria

The criteria for skin irritation are not met.

4.4.1.5 Conclusions on classification and labelling

S-Methoprene does not classify for skin irritation when compared to the requirements of the CLP Regulation (EC) No. 1272/2008.

4.4.2 Eye irritation

Table 16: Summary table of relevant eye irritation studies

Method	Results	Remarks	Reference
Eye irritation in NZW rabbits S-Methoprene OECD 405	Non-irritant	Slight reversible erythema.	Kuhn, J.O. (1999d) S-Methoprene CAR IIIA 6.1.4/2

4.4.2.1 Non-human information

Kuhn, J.O. (1999d), Primary eye irritation study in rabbits. The acute eye irritation of S-Methoprene was investigated by instilling a single dose (0.1 mL) of S-Methoprene technical (LX 125-03) into the conjunctival sac of one eye of a group of six New Zealand White rabbits (3/sex). The lids were thereafter gently held together for one second and then released. The left eyes served as controls. Following installation of S-Methoprene, the eyes of all animals were observed for signs of ocular irritation at 1, 24, 48 and 72 hours after treatment. The grades of ocular reaction were recorded at each examination period.

Table 17: Eye Irritation Scores following exposure to S-Methoprene

Time/Rabbit	Corneal Opacity						Iridial Inflammation					
	9224	9226	9230	9225	9227	9229	9224	9226	9230	9225	9227	9229
	M	M	M	F	F	F	M	M	M	F	F	F
1h	0	0	0	0	+	+	0	0	0	0	0	0
24h	0	0	0	0	0	0	0	0	0	0	0	0
48h	0	0	0	0	0	0	0	0	0	0	0	0
72h	0	0	0	0	0	0	0	0	0	0	0	0
Mean Score (24-72 hr)	0						0					

+ slightly dulling of normal luster

Time/Rabbit	Conjunctival Redness						Conjunctival Chemosis					
	9224	9226	9230	9225	9227	9229	9224	9226	9230	9225	9227	9229
	M	M	M	F	F	F	M	M	M	F	F	F
1h	1	1	1	2	1	2	1	1	1	1	1	1
24h	0	0	0	1	0	1	0	0	0	0	0	0
48h	0	0	0	0	0	0	0	0	0	0	0	0
72h	0	0	0	0	0	0	0	0	0	0	0	0
Mean Score (24-72 hr)	0.11						0					

Average scores (24-72 hours) of 0 for cornea, 0 for Iris and 0 for Chemosis were recorded. Conjunctiva redness was scored at 0.11 for 24-72 hours. All effects had reversed at 48 hours and on the basis of reactions observed in this study and the criteria defined in CLP Regulation (EC) No. 1272/2008, S-Methoprene does not require classification for acute eye irritation.

4.4.2.2 Human information

Not available.

4.4.2.3 Summary and discussion of eye irritation

Slight reversible Conjunctiva redness was seen in the study presented. This was fully reversible within 48 hours.

4.4.2.4 Comparison with criteria

The criteria for eye irritation are not met.

4.4.2.5 Conclusions on classification and labelling

S-Methoprene does not classify for eye irritation when compared to the requirements of the CLP Regulation (EC) No. 1272/2008. Classification not warranted.

4.4.3 Respiratory tract irritation

4.4.3.1 Non-human information

S-Methoprene is not classifiable for skin or eye irritation and respiratory tract irritation is not anticipated. There was no evidence of respiratory tract irritation in the acute studies provided.

4.4.3.2 Human information

Not available.

4.4.3.3 Summary and discussion of respiratory tract irritation

Not indicated.

4.4.3.4 Comparison with criteria

Not applicable.

4.4.3.5 Conclusions on classification and labelling

Not applicable.

4.5 Corrosivity

Table 18: Summary table of relevant corrosivity studies

Method	Results	Remarks	Reference

4.5.1 Non-human information

S-Methoprene is not irritating to the skin or eyes in the rat. Corrosivity not relevant.

4.5.2 Human information

Not available.

4.5.3 Summary and discussion of corrosivity**4.5.4 Comparison with criteria**

Based upon the irritation studies submitted for S-Methoprene, discussed in Section 4.4.1 and 4.4.2 above, corrosivity is not anticipated.

4.5.5 Conclusions on classification and labelling

No classification.

4.5.6 Skin sensitisation

Table 19: Summary table of relevant skin sensitisation studies

Method	Results	Remarks	Reference
Beuhler Method. (OECD 406). Hartley guinea pigs	No sensitisation reaction	-	Kuhn, J. O. (1999e) CAR IIIA 6.1.5

4.5.6.1 Non-human information

Kuhn, J.O. (1999e), Dermal Sensitization Study In Guinea Pigs. In a dermal sensitization study with S-Methoprene technical (LX 125-03; 97.2%), 10 hartley-albino guinea pigs (5 animals/sex) were tested using the method of Buehler for a total of three six-hour insult periods. Induction treatments were on Days 1, 8 and 15. Observations for skin reactions were made approximately 24 hours and 48 hours after the first induction treatment and 48 hours after each subsequent treatment. Skin reactions were graded according to the Buehler scale. In a preliminary dose-range-finding study, 4 animals each (2/sex) were exposed to four different concentrations of S-Methoprene to determine the mildly irritating and highest non-irritating dose (100% v/v, 75% v/v, 50% v/v, 25% v/v in acetone). Based upon the results of the dose-range-finding studies, the test article was dosed as received for induction and challenge

Fourteen days after the last induction period, all test animals were challenged with 0.4 mL of undiluted test substance in the same manner as the induction treatments at a virgin test site on the right rear quadrant of the animal. A group of 10 naïve control (5/sex) animals were treated with S-Methoprene in the same manner. This group served as the control challenge group. Challenge application lasted 6 hours, as before. Observations were performed 24 hours and 48 hours after the challenge treatments.

Table 20: Summary of Skin Sensitization Results following Challenge Phase

	Number of animals with signs of allergic reactions / number of animals in group		
	Naive control	Test group (S-Methoprene)	Positive control (2-Mercapto- Benzothiazole)
Scored after 24h	0/10	0/10	9/10
Scored after 48h	0/10	0/10	8/10

The test substance, S-Methoprene, produced no irritation in naive control group animals after the single treatment at challenge. Similarly, the test substance produced no signs of irritation in test group animals after the challenge treatment and therefore did not elicit a sensitising reaction in guinea pigs. In accordance with the criteria defined in CLP Regulation (EC) No. 1272/2008, S-Methoprene does not require classification for skin sensitization.

4.5.6.2 Human information

Not available.

4.5.6.3 Summary and discussion of skin sensitisation

Dermal sensitisation was investigated in an acceptable test the “Beuhler test”. There was no response consistent with dermal sensitisation.

4.5.6.4 Comparison with criteria

The criteria for skin sensitisation are not met.

4.5.6.5 Conclusions on classification and labelling

No sensitisation effects were detected in the Buehler test and therefore classification of S-Methoprene for skin sensitisation is not warranted.

4.5.7 Respiratory sensitisation

No data.

Table 21: Summary table of relevant respiratory sensitisation studies

Method	Results	Remarks	Reference
-	-	-	-
-	-	-	-
-	-	-	-

4.5.7.1 Non-human information

No data.

4.5.7.2 Human information

No data.

4.5.7.3 Summary and discussion of respiratory sensitisation

No data.

4.5.7.4 Comparison with criteria

No data.

4.5.7.5 Conclusions on classification and labelling

Not applicable.

4.6 Repeated dose toxicity

Note: Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene as a juvenile hormone is restricted to the S enantiomer (S-Methoprene). S-Methoprene is the active isomer in the racemic mixture and is present in a 1:1 ratio;

the results obtained in the studies with Methoprene are therefore adjusted to give the S-Methoprene equivalent. The NOEL from studies carried out with Methoprene has been adjusted and the NOEL for S-Methoprene included below:

Table 22: Summary table of relevant repeated dose toxicity studies

Species/study/dose	Findings at LOAEL	GV (extrap.) CLP	Reference
Rat, Oral, 90 days, S-Methoprene technical (96%): 0, 200, 400 and 1000 mg/kg/day	200 mg/kg bw/day. Statistically significant changes in organ weight such as liver and kidneys NOAEL was not determined. NOAEL: < 200 mg/kg bw/day		Szakonyi, I.P. (2002) CAR IIIA 6.1.4/2
Dog, Oral 90 days, S-Methoprene technical (96%): 0, 100, 300 and 1000 mg/kg bw/day Daily (Capsule)	300 mg/kg bw/day in males and females. Gastrointestinal signs including thin faeces and diarrhoea and increase in liver weight in males and females and in ALP activity in females NOAEL: 100 mg/kg bw/day in males and females		Török, T. (2007)
Rat, Oral, 104 weeks, S-Methoprene technical (96%): 0, 10.9, 43.4 and 217 mg/kg/day	5000 ppm (equivalent to 217 mg/kg bw/day) for Methoprene 2500 ppm (equivalent to 108.5 mg/kg bw/day) for S-Methoprene - 5000 ppm (equivalent to 217 mg/kg bw/day) increased incidence of hepatic lesions (bile-duct proliferation and portal lymphocyte infiltration) in males and increased absolute and relative weights of the liver in females. NOAEL: 1000 ppm (equivalent to 43.4 mg/kg bw/day) for Methoprene 500 ppm (equivalent to 21.7 mg/kg bw/day) for S-Methoprene		Wazeter, F.X., Goldenthal E.I., Geil, R.G., Benson, B.W., Keller W.F. and Blanchard, G.L. (1975)
Mouse, Oral, 18 month Methoprene technical: 250, 1000 and 2500 ppm (equivalent to 32.7, 130.8 and 327 mg/kg bw/day, respectively) (Dietary)	2500 ppm (equivalent to 327 mg/kg bw/day) for S-Methoprene. Increases in focal accumulations of macrophages with brownish foamy cytoplasm in the liver, often associated with small necrotic foci and mononuclear inflammatory cells. NOAEL: Carcinogenicity; 2500 ppm (equivalent to 327 mg/kg bw/day) for Methoprene 1250 ppm (equivalent to 163.5 mg/kg bw/day) for S-Methoprene Toxicity; 1000ppm (equivalent to 130.8 mg/kg bw/day) for Methoprene 500 ppm (equivalent to 65.4mg/kg		Wazeter, F.X., Goldenthal, E.I., Geil, R.G. and Benson B.W. (1975)

Species/study/dose	Findings at LOAEL	GV (extrap.) CLP	Reference
	bw/day) for S-Methoprene 125 ppm (equivalent to 16.35 mg/kg bw/day) for S-Methoprene.		

Note: Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene to S-Methoprene

4.6.1 Non-human information

4.6.1.1 Repeated dose toxicity: oral

Study 1. Szakonyi, I.P. (2002) 90-day rat feeding study (CAR IIIA.6.4.1)

Table 23: Summary of main findings

Study	Main findings	
Rat, Oral, 90 days, S-Methoprene technical (96%): 0, 200, 400 and 1000 mg/kg/day	<p>Bodyweight gain was reduced compared to controls in male rats at 1000 mg/kg/day. A statistically significant difference was noted between weeks 7 and 13. However, this difference was reversible as the values of treated group exceeded the controls at the end of the recovery period. Sporadic statistically significant decreases were noted at 400 mg/kg/day.</p> <p>From day 63 until the end of the recovery period, the mean body weight in males at 1000 mg/kg/day was statistically significantly lower than controls.</p> <p>No treatment related changes in haematological parameters were recorded. At 400 mg/kg/day, increased HTC was noted in both males and females along with decreased MCHC in females. These variations were not considered to be treatment related. At 1000 mg/kg/day, slight increases in RBC and HTC in males and increase HGB in both males and females were recorded. Decreases in MCHC were elicited in both sexes at 1000 mg/kg/day. However, these variations were not biologically significant and were within the physiological range.</p> <p>Increase in the glucose concentration was recorded in all female dose groups. However, this increase was only deemed to be relevant at 1000 mg/kg/day. A treatment related effect was proposed. Other changes such as decrease in bilirubin in both sexes at 200 mg/kg/day and in males at 400 mg/kg/day, decrease in carbamide and creatinine in females at 200 and 1000 mg/kg/day, increase in cholesterol in females at 200, 400 and 1000 mg/kg/day were reported. At 400 and 1000 mg/kg/day, decreased AST activity in males and females was observed. During the recovery period, increase in albumin concentration in males at 1000 mg/kg/day was noted. However, these variations were not biologically significant and were within the physiological range.</p> <p>At necropsy, the female that died on day 77 displayed reddish mottled lungs and congestive liver. However this was deemed to be a consequence of the paragastric treatment. Macroscopic findings in the lungs (emphysema, pinprick-sized and point-like haemorrhages, reddish mottled colour) noted in the control and all treated groups were correlated to the extermination process or considered to be incidental. Other findings such as enlarged, pale</p>	Szakonyi, I.P. (2002)

	<p>or nutmeg-like liver, pale or enlarged kidneys, pyelectasis, enlarged testes, smaller than normal thymus, haemorrhages in the urinary bladder, ovarian cyst, hydrometra) are commonly reported in rats of this age.</p> <p>A treatment related increase in liver weight (absolute and relative to bodyweight and brain weight) was recorded in all treated groups in both males and females. Increased kidney weight (absolute and relative to the body and brain weight) was observed in all male treated groups and was considered to be dose related. Increase in absolute kidney weight was reported in female at 400 and 1000 mg/kg/day. A dose related increase in kidney weight relative to body weight was noted in all female treated groups. The kidney referred to the brain weight increased only in females at 1000 mg/kg/day. At the end of the recovery period, the relative kidney weight referred to the body weight remained significantly higher than in the control in male animals. Other changes in the organ weights including brain, spleen, adrenals, testes, epididymides, ovaries and uterus were considered to be individual or incidental findings. The increase of organ weight issues originated from the increased number of healthy active cells of the organ could be considered as an adaptational process.</p> <p>LO(A)EL: 200 mg/kg bw/day NO(A)EL: No value determined. The value is somewhere between 0 and 200 mg/kg bw/day but has not been determined.</p>	
--	--	--

*P<0.05, ** p<0.01, ns- not significant.

Table 24: Summary of body weight data in males and females (g)

Days of Study	Male				Female			
	0 mg/kg/day (weight ± SD)	200 mg/kg/day (weight ± SD)	400 mg/kg/day (weight ± SD)	1000 mg/kg/day (weight ± SD)	0 mg/kg/day (weight ± SD)	200 mg/kg/day (weight ± SD)	400 mg/kg/day (weight ± SD)	1000 mg/kg/day (weight ± SD)
1	169.15 ± 5.43	170.30 ± 5.54	169.10 ± 5.53	165.25 ± 7.47	146.30 ± 5.52	144.80 ± 5.81	146.60 ± 5.44	146.15 ± 6.48
7	229.95 ± 9.51	232.10 ± 10.61	231.00 ± 8.34	226.00 ± 7.40	172.35 ± 7.19	169.30 ± 7.13	167.30 ± 7.73	170.15 ± 9.86
14	286.30 ± 14.02	292.30 ± 17.86	290.50 ± 11.31	283.15 ± 11.37	195.45 ± 9.71	191.50 ± 7.52	191.50 ± 13.39	197.25 ± 13.19
21	329.95 ± 18.81	340.60 ± 25.78	335.30 ± 16.81	327.35 ± 17.59	218.60 ± 11.86	212.90 ± 8.06	210.30 ± 16.10	219.60 ± 15.01
28	367.25 ± 23.09	376.00 ± 34.42	367.90 ± 21.48	360.40 ± 20.93	232.20 ± 13.04	228.80 ± 9.54	226.50 ± 17.84	235.55 ± 17.83
35	396.40 ± 27.03	403.80 ± 40.07	397.00 ± 26.12	387.40 ± 24.13	243.00 ± 12.37	239.80 ± 14.38	237.60 ± 14.83	246.15 ± 19.98
42	423.65 ± 30.43	439.00 ± 43.15	422.80 ± 28.40	411.25 ± 27.52	254.00 ± 14.90	251.80 ± 16.02	252.20 ± 19.63	260.15 ± 20.78
49	437.95 ± 33.37	456.20 ± 47.87 ^c	433.70 ± 29.04	419.90 ± 30.75	260.70 ± 13.40	257.70 ± 13.12	256.50 ± 17.08	263.10 ± 20.93
56	457.05 ± 37.71	479.50 ± 49.36 ^d	450.10 ± 32.92	432.70 ± 29.74	268.05 ± 14.10	267.00 ± 15.03	265.90 ± 14.67	272.20 ± 23.82
63	471.80 ± 39.81 ^a	499.50 ± 54.18 ^d	464.20 ± 38.44	441.80 ± 30.46	274.90 ± 15.82	273.60 ± 16.10	272.10 ± 15.95	279.00 ± 23.19
70	485.05 ± 39.83 ^a	515.70 ± 59.45 ^{e, d}	474.90 ± 39.36	451.30 ± 33.37	277.20 ± 15.98	276.50 ± 18.79	274.50 ± 15.34	281.75 ± 24.19
77	498.95 ± 40.02 ^a	525.50 ± 58.84 ^{e, d}	485.40 ± 39.72	458.10 ± 33.27	285.05 ± 16.65	285.50 ± 17.02	282.20 ± 18.81	288.15 ± 25.17

84	497.65 ± 39.69 ^b	523.20 ± 58.19 ^d	484.50 ± 38.07	456.95 ± 34.81	275.80 ± 16.19	273.30 ± 16.49	271.70 ± 15.59	275.53 ± 26.77
89	508.65 ± 40.07 ^b	531.30 ± 60.31 ^{e,d}	490.30 ± 40.23	461.00 ± 34.97	281.40 ± 17.81	275.40 ± 17.70	273.60 ± 15.72	284.68 ± 23.25
96	527.6 ± 37.00	-	-	481.7 ± 38.59	284.80 ± 15.25	-	-	296.3 ± 21.85
103	537.8 ± 36.21	-	-	486.6 ± 33.14	290.4 ± 15.83	-	-	294.4 ± 26.20
110	550.6 ± 40.29	-	-	505.5 ± 38.02	298.3 ± 16.35	-	-	303.8 ± 30.43
117	551.4 ±	-	-	511.3 ± 41.70	300.8 ±	-	-	310.0 ± 33.74

^a p<0.05 and (group 4 - group 1) means group mean4 < group mean1

^b p<0.01 and (group 4 - group 1) means group mean4 < group mean1

^c p<0.05 and (group 4 - group 2) means group mean4 < group mean2

^d p<0.01 and (group 4 - group 2) means

Table 25: Summary of terminal organ and body weights (g), organ/body weight and organ/brain weight ratios

Data collection	Sex	Group	Dose	Liver weight	Liver/body weight	Liver/brain weight	Kidney weight	Kidney/body weight	Kidney/brain weight
At study termination	Male	1M	0	11.50 ± 1.93	2.330 ± 0.216	526.09 ± 83.14	2.57 ± 0.24	0.524 ± 0.051	117.39 ± 7.85
		2M	200	14.39 ± 2.32 ^a	2.751 ± 0.188 ^a	631.23 ± 90.34 ^a	3.11 ± 0.46 ^a	0.598 ± 0.073 ^d	136.22 ± 16.62 ^a
		3M	400	14.37 ± 1.28 ^b	3.016 ± 0.203 ^{b, f}	669.69 ± 66.77 ^b	3.25 ± 0.30 ^b	0.683 ± 0.071 ^{b, g}	151.12 ± 13.10 ^{b, g}
		4M	1000	16.94 ± 1.57 ^{c, d, e}	3.786 ± 0.373 ^{c, d, e}	781.38 ± 70.63 ^{c, d, e}	3.29 ± 0.23 ^c	0.733 ± 0.046 ^{c, d}	151.50 ± 9.34 ^{c, p}
At the end of the recovery period	Male	1M	0	10.88 ± 1.24	2.058 ± 0.260	481.94 ± 49.98	2.85 ± 0.24	0.537 ± 0.028	126.35 ± 9.78
		4M	1000	12.77 ± 1.13 ^{**}	2.619 ± 0.225 ^{**}	571.28 ± 46.26 ^{**}	3.07 ± 0.30	0.627 ± 0.029 ^{**}	137.23 ± 13.39
At study termination	Female	1F	0	6.87 ± 0.66	2.443 ± 0.186	341.16 ± 38.59	1.63 ± 0.17	0.581 ± 0.049	81.18 ± 10.15
		2F	200	7.90 ± 0.5 ¹	2.885 ± 0.129 ^{**}	390.54 ± 23.68 ^{**}	1.75 ± 0.16	0.638 ± 0.044 [*]	86.38 ± 7.01
		3F	400	9.71 ± 0.83 ^{j, 1}	3.572 ± 0.242 ^{**}	464.35 ± 40.31 ^{**}	1.86 ± 0.18 ^k	0.683 ± 0.047 ^{**}	88.81 ± 8.31
		4F	1000	13.45 ± 1.21 ^{m, n, o}	4.800 ± 0.364 ^{**}	663.94 ± 72.02 ^{**}	2.20 ± 0.26 ^{m, n, o}	0.789 ± 0.132 ^{**}	108.19 ± 12.80 ^{m, n, o}
At the end of the recovery period	Female	1F	0	6.58 ± 0.72	2.322 ± 0.199	318.17 ± 35.01	1.72 ± 0.11	0.608 ± 0.028	83.09 ± 4.14
		4F	1000	7.32 ± 0.80	2.524 ± 0.178 [*]	353.54 ± 35.94 [*]	1.78 ± 0.15	0.614 ± 0.052	85.88 ± 7.86

* Statistically different from control, p<0.05

** Statistically different from control, p<0.01

Conclusion:

S-Methoprene treatment at 200, 400 and 1000 mg/kg/day resulted in a treatment related increase in liver weight (absolute and relative to bodyweight and brain weight) in all treated groups in both males and females. Increased kidney weight (absolute and relative to the body and brain weight) was observed in all male treated groups and was considered to be dose related. Increase in absolute kidney weight was reported in female at 400 and 1000 mg/kg/day. A dose related increase in kidney weight relative to body weight was noted in all female treated groups. At the end of the recovery period, the relative kidney weight referred to the body weight remained significantly higher than in the control in male animals. However, there was no evidence of specific organ toxicity outside increase in weight. Indeed, relative liver weights were 30% and 60% increased at the 400 mg/kg/day and 1000 mg/kg/day respectively without any noticeable impact on clinical chemistry or histological findings.

Study 2. Török, T. (2007) 90-day dog feeding study (CAR IIIA.6.4.1)

<p>Dog 90-day, S-Methoprene technical (96%): 0, 100, 300 and 1000 mg/kg bw/day</p> <p>Daily (Capsule)</p>	<p>Gastrointestinal signs such as thin faeces and diarrhoea were observed in both sexes mostly in the 300 and 1000 mg/kg bw/day groups in a dose-related manner.</p> <p>There were no mortalities at any dose level during the study.</p> <p>No significant treatment related differences were found in body weights, food consumption, ophthalmoscopic examinations, haematology, urinalysis and macroscopic examinations.</p> <p>A statistically and biologically significant increase in ALKP activity was observed at the end of the midway and terminal period in males in the high dose group and in females in all treated groups. Enzyme activities were increased in a dose dependent manner but only exceeded historical control ranges in the higher dose groups. Increased ALKP activity is common although not specific in liver damage. The histopathological examination confirmed the suspicion of liver damage in the high dose groups.</p> <p>In both sexes in the 1000-mg/kg bw/day groups the weight of the livers showed a significant increase when compared to the control.</p> <p>In the 1000-mg/kg bw/day dose animals, a slight or medium degree of zonal vacuolisation of hepatocytes in the liver was observed. No other test material related lesions were detectable during the histopathology</p>
---	--

Table 26: Summary of body weight data in males and females (kg)

Weeks of Study	Male				Female			
	0 mg/kg bw/day (weight ± SD)	100 mg/kg bw/day (weight ± SD)	300 mg/kg bw/day (weight ± SD)	1000 mg/kg bw/day (weight ± SD)	0 mg/kg bw/day (weight ± SD)	100 mg/kg bw/day (weight ± SD)	300 mg/kg bw/day (weight ± SD)	1000 mg/kg bw/day (weight ± SD)
1	7.93 ±	8.05 ± 0.37	8.05 ±	7.90 ±	8.73 ± 0.56	8.80 ± 0.47	8.73 ± 0.87	8.75 ±
2	8.33 ±	8.45 ± 0.33	8.33 ±	8.25 ±	9.03 ± 0.49	8.93 ± 0.44	9.08 ± 0.84	8.98 ±
3	8.65 ±	8.78 ± 0.25	8.63 ±	8.35 ±	9.18 ± 0.48	9.20 ± 0.22	9.30 ± 0.65	9.18 ±
4	8.95 ±	9.05 ± 0.19	8.90 ±	8.85 ±	9.33 ± 0.34	9.35 ± 0.17	9.48 ± 0.65	9.30 ±
5	9.20 ±	9.33 ± 0.26	9.10 ±	9.13 ±	9.68 ± 0.41	9.60 ± 0.16	9.73 ± 0.75	9.48 ±
6	9.45 ±	9.58 ± 0.13	9.55 ±	9.28 ±	9.98 ± 0.51	9.78 ± 0.19	9.90 ± 0.85	9.63 ±
7	9.50 ±	9.78 ± 0.22	9.53 ±	9.50 ±	10.28 ±	9.95 ± 0.21	9.90 ± 0.75	9.75 ±
8	9.63 ±	10.00 ± 0.28	9.58 ±	9.68 ±	10.38 ±	10.10 ±	10.03 ±	9.83 ±

9	9.88 ±	10.23 ± 0.31	9.80 ±	10.05 ±	10.53 ±	10.28 ±	10.15 ±	9.93 ±
10	9.98 ±	10.30 ± 0.29	9.95 ±	10.18 ±	10.60 ±	10.15 ±	10.25 ±	9.95 ±
11	10.00 ± 0.75	10.45 ± 0.31	10.18 ± 0.68	10.30 ± 0.83	10.73 ± 0.15	10.43 ± 0.15	10.40 ± 0.91	10.13 ± 1.00
12	10.15 ± 0.87	10.65 ± 0.40	10.33 ± 0.56	10.33 ± 0.92	10.90 ± 0.24	10.50 ± 0.22	10.53 ± 0.86	10.28 ± 1.00
13	10.20 ± 0.88	10.65 ± 0.52	10.33 ± 0.57	10.35 ± 0.84	10.93 ± 0.15	10.63 ± 0.33	10.53 ± 0.99	10.23 ± 0.93

Table 27: Summary of terminal organ and body weights (g), organ/body weight and organ/brain weight ratios

Data collection	Sex	Dose (mg/kg bw/day)	Liver weight (g)	Liver/ body weight (%)	Liver/ brain weight (%)
Terminal	Male	0	510.17 ± 111.65	5.090 ± 0.951	563.50 ± 95.26
		100	617.37 ± 111.37	5.836 ± 1.280	723.56 ± 154.51
		300	569.81 ± 95.34	5.690 ± 0.843	704.70 ± 114.96
		1000	731.21* ± 82.75	7.158* ± 0.595	860.99** ± 66.32
Terminal	Female	0	487.14 ± 64.64	4.451 ± 0.603	584.30 ± 101.87
		100	562.99 ± 83.26	5.304 ± 0.678	661.51 ± 92.66
		300	595.68 ± 116.92	5.704* ± 0.549	729.61 ± 152.62
		1000	678.19* ± 96.88	6.837** ± 0.913	836.46 ± 169.69

* Significantly different at 0.05 using Duncan's multiple range test

** Significantly different at 0.01 using Duncan's multiple range test

Conclusion:

S-Methoprene treatment at 100, 300 and 1000 mg/kg/day produced clinical signs such as thin faeces and diarrhea and increased liver weight in males at 1000 mg/kg (140% of controls) and females from 300 mg/kg (128% and 153%) and biologically and statistically significant increase in alkaline phosphatase (ALP) values in females from 300mg/kg bw/day (>10%). In addition, in the 1000-mg/kg bw/day dose animals, a slight or medium degree of zonal vacuolisation of hepatocytes in the liver was observed in both sexes. No other test material related lesions were detectable during the histopathology. Based on the findings found under the conditions of this study, the LOAEL was established to be 300-mg/kg bw/day and the NOAEL was established to be 100-mg/kg bw/day in beagle dogs.

Note: Study 3. Wazeter, F.X., Goldenthal E.I., Geil, R.G., Benson, B.W., Keller W.F. and Blanchard, G.L. (1975) 104 week rat oral study (CAR IIIA.6.7.1) and Study 4 Wazeter, F.X., Goldenthal, E.I., Geil, R.G. and Benson B.W. (1975) 18-month mouse oral study are summarized in Section 4.9 of this document.

4.6.1.2 Repeated dose toxicity: inhalation

None available

4.6.1.3 Repeated dose toxicity: dermal

None available

4.6.1.4 Repeated dose toxicity: other routes

None available

4.6.1.5 Human information

None available

4.6.1.6 Other relevant information

None available

4.6.1.7 Summary and discussion of repeated dose toxicity

The toxicological properties of S-Methoprene upon short-term treatment were investigated in rat and dog. A sub chronic 90-day rat study (Szakonyi, I.P., 2002) did not provide an NOAEL value, and only an LOAEL of < 200 mg/kg was determined. The 90-day dog study (Torok, T., 2007) produced an NOAEL of 100mg/kg bw/day. The effects noted included clinical signs such as thin faeces and diarrhoea and increased liver weight in males at 1000 mg/kg (140% of controls) and females from 300 mg/kg (128% and 153%) and biologically and statistically significant increase in alkaline phosphatase (ALP) values in females from 300mg/kg bw/day (>10%). In addition vacuolization of hepatocytes were noted in both sexes at 1000mg/kg bw/day.

4.7 Specific target organ toxicity (CLP Regulation) – repeated exposure (STOT RE)

4.7.1 Summary and discussion of repeated dose toxicity findings relevant for classification as STOT RE according to CLP Regulation

4.7.2 Comparison with criteria of repeated dose toxicity findings relevant for classification as STOT RE

Classification for STOT Cat 2 is required when:

....on the basis of evidence from studies in experimental animals can be presumed to have the potential to be harmful to human health following repeated exposure. Substances are classified in category 2 for target organ toxicity (repeat exposure) on the basis of observations from appropriate studies in experimental animals in which significant toxic effects, of relevance to human health, were produced at generally moderate exposure concentrations.

According to the CLP Regulation (EC) No. 1272/2008 S-Methoprene should not be considered for classification because clinical signs in the dog and liver effects in the dog and rat were seen at doses

of 200 to 300 mg/kg bw/day. These doses are in excess of the cut-off value of $10 \leq 100$ mg/kg bw/day for Category 2 classification.

4.7.3 Conclusions on classification and labelling of repeated dose toxicity findings relevant for classification as STOT RE

S-Methoprene does not require classification for STOT RE according to the CLP Regulation (EC) No. 1272/2008.

4.8 Germ cell mutagenicity (Mutagenicity)

Table 28: Summary table of relevant *in vitro* and *in vivo* mutagenicity studies

Method	Results	Reference
<i>In vitro</i>		
Prokaryote gene mutation		
<i>Salmonella typhimurium</i> : TA 1535, TA 1537, TA 98 with pKM101, TA 100 with pKM101	Negative +/- S9	Hernádi, D. (2002) CAR IIIA 6.6.1
<i>Escherichia Coli</i> WP2 uvrA OECD 471		
Mammalian gene mutation		
None available		
Chromosomal aberrations		
Subline (KI) of Chinese hamster ovary cell line (CHO) OECD 473	Negative +/- S9	Béres, E. (2003) CAR IIIA 6.6.2
Subline (KI) of Chinese hamster ovary cell line (CHO) OECD 473	Negative +/- S9	Béres, E. (2002) CAR IIIA 6.6.3
DNA repair <i>in vitro</i>/cell transformation assay-mammalian		
None available		
<i>In vivo</i>		
Micronucleus assay		
None available		
Germ cell assays		
None available		

4.8.1 Non-human information

4.8.1.1 *In vitro* data

S-Methoprene did not induce gene mutations in bacterial cells *in vitro*. The *in vitro* chromosome aberration test in CHO cells revealed no evidence for clastogenic potential of S-Methoprene.

4.8.1.2 *In vivo* data

No *in vivo* data was available.

4.8.2 Human information

None available.

4.8.3 Other relevant information

None relevant.

4.8.4 Summary and discussion of mutagenicity

S-Methoprene was negative in both available *in vitro* studies.

4.8.5 Comparison with criteria

Not relevant.

4.8.6 Conclusions on classification and labelling

No classification.

4.9 Carcinogenicity

Table 29: Summary table of relevant carcinogenicity studies

Method	Results	Target organ/ principal effect at LOAEL	Reference
Rat, Charles River CD, Male/Female 50/sex/group Methoprene technical: 250, 1000 and 5000 ppm (equivalent to 10.9, 43.5 and 217.25 mg/kg bw/day) 104 weeks/daily	No treatment related tumours		Wazeter, F.X., Goldenthal, E.I., Geil, R.G., Benson, B.W., Keller, W.F. and Blanchard, G.L. (1975) CAR IIIA 6.4.1
Mouse, Charles River CD-1, Male/Female 50/sex/group Methoprene technical: 250, 1000 and 2500 ppm (equivalent to 0, 32.3, 130.4 and 325.9 mg/kg bw/day) 72- 78 weeks/daily	No treatment related tumours		Wazeter, F.X., Goldenthal, E.I., Geil, R.G., and Benson, B.W., (1975) CAR IIIA 6.4.1-2

Note: Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene to S-Methoprene.

4.9.1 Non-human information

4.9.1.1 Carcinogenicity: oral

Study 1: Oral carcinogenicity study in the rat

Methoprene technical (86.9%) was administered at target dietary doses of 0, 250, 1000 and 5000 ppm daily over a period of 104 weeks to 4 groups of 50 Charles River CD rats/sex/group.

No changes in general behavior or appearance was deemed to be treatment related. Incidental findings noted such as occasional soft stool, slight alopecia, ocular or nasal porphyrin discharge and nodules (usually on the abdomen, thorax, or sides of the rats) were recorded in all groups.

Survival at 250 and 1000 ppm exceeded that of controls. 38% of males survived in the control group and at 5000 ppm. 52% and 48% of females survived in the control group and at 5000 ppm, respectively.

No treatment related changes were noted in bodyweight at any dose levels. Few statistically significant changes were reported between control and treated groups. However these were not deemed to be treatment related. Results of the statistical analysis are summarized in the following Table.

Table 30: Summary of body weights

Dosage level (ppm)	Number surviving/Number initiated	
	Male	Female
Control	19/50	26/50
250	27/50	30/50
1000	23/50	29/50
5000	19/50	24/50

Table 31: Summary of body weights

Week of Study	Male (g)				Female (g)			
	0 ppm	250 ppm	1000 ppm	5000 ppm	0 ppm	250 ppm	1000 ppm	5000 ppm
13	479	470	463	463	261	263	258	250
26	546	535	534	532	277	285	277	269
39	599	578	584	571	303	309	302	290
52	636	623	623	626	327	335	337	318
65	685	674	676	669	375	378	365	344 ^a
78	675	662	656	677	377	394	371	344 ^a
91	651	639	628	665	390	403	386	369
104	599	597	584	616	379	365	364	370

^aSignificantly different from control group mean, p<0.05

No treatment related changes were reported for food consumption at any dose level. No treatment related changes were recorded for haematology, biochemistry and urinalysis test. No ophthalmologic changes were reported during the study at any dose levels.

No treatment related gross pathologic lesions were noted at any dose levels. Few lesions such as mass skin, white yellow foci lung, abscess lung, enlarged spleen, hemorrhage erosion, ulcerations stomach and chronic nephritis were reported in all groups including the control. However they are commonly reported in rats of this age.

No treatment related effect on organ weight was reported. The absolute and relative weights of the liver were elevated in females at 5000 ppm dose level. Results are summarized in the following table.

Table 32: Summary of absolute (g) and relative (% body weight) organ weights

Group Sex	Body weight	Spleen		Liver		Adrenals		Kidneys		Testis		Ovaries		Heart		Thyroid/parat hyroid		Brain		Pituitary	
		g	% x 10 ²	g	%	g	% x 10 ²	g	%	g	%	g	% x 10 ²	g	%	g	% x 10 ²	g	%	g	% x 10 ²
Control																					
male	599	1.15	0.20	21.27	3.65	0.126	2.22	5.67	0.99	3.25	0.54	-	-	1.81	0.31	0.059	1.00	2.25	0.39	0.030	0.51
female	379	0.79	0.21	12.78	3.42	0.154	4.18	3.16	0.84	-	-	0.145	0.39	1.29	0.34	0.045	1.18	1.94	0.53	0.101	2.73
250 ppm																					
male	597	1.15	0.19	17.13 _b	2.90 ^a	0.097 _a	1.69	4.70	0.81	3.70	0.62	-	-	1.93	0.33	0.047 _a	0.81	2.22	0.38	0.031	0.52
female	376	1.11	0.30	11.37	3.18	0.187	5.23	3.03	0.86	-	-	0.176	0.50 ^a	1.36	0.38	0.048	1.35	1.96	0.55	0.151	4.69
1000 ppm																					
male	581	1.17	0.21	18.95	3.38	0.107	1.90	5.56	0.98	3.45	0.61	-	-	1.82	0.32	0.055	0.95	2.19	0.38	0.032	0.53
female	364	0.75	0.21	11.99	3.35	0.153	4.24	3.00	0.85	-	-	0.141	0.39	1.19	0.34	0.049	1.44	1.85	0.53	0.120	3.95
5000 ppm																					
male	616	1.10	0.18	21.10	3.45	0.089 _a	1.49 ^a	5.47	0.91	3.23	0.53	-	-	1.73	0.29	0.063	1.05	2.07 ^b	0.34	0.027	0.45
female	370	0.65 ^a	0.18	15.41	4.13 ^b	0.143	4.08	3.14	0.86	-	-	0.149	0.43	1.25	0.34	0.051	1.47	1.86	0.52	0.110	3.22

Group mean relative organ weights shown in this table were calculated by averaging the individually calculated relative organ weights

^aSignificantly different from control group mean, p<0.05

^bSignificantly different from control group mean, p<0.01

No treatment related histopathologic lesions were noted at any dose levels. Few lesions such as chronic nephritis and adenoma pituitary were noted in control and high dose groups. However they are commonly reported in rats of this age. Histopathological evaluation showed an increased incidence of hepatic lesions, such as bile-duct proliferation and portal lymphocyte infiltration in males at 5000ppm dose level.

No treatment related tumours were reported. There were no statistically significant changes at any dose levels.

Based on the increased incidence of hepatic lesions such as bile-duct proliferation and portal lymphocyte infiltration in males at 5000 ppm dose level and the increased absolute and relative weights of the liver in females at 5000 ppm dose level, the NOEL was established to be 1000 ppm (equivalent to 43.45 mg/kg bw/day). Methoprene is concluded to be non-carcinogenic at dose levels up to 5000 ppm (equivalent to 217 mg/kg bw/day).

Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene as a juvenile hormone is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene to S-Methoprene. Therefore, the observed NOAEL in this study has been revised and a NOAEL for S-Methoprene is proposed:

- NOAEL for S-Methoprene = **21.72 mg/kg bw/day**

In general the level of reporting in this study is poor. There is no reference to findings at the high dose which may have been attributable to treatment, e.g., relative liver weight increase in high dose females; an apparent increase in liver histopathological findings in high dose animals (portal lymphocytic infiltration, bile duct proliferation, small groups of vacuolated hepatocytes).

Survival was lower than average for some male groups including controls (19/50, 27/50, 23/50, 19/50, i.e., 38%, 54%, 46%, 38% respectively). There does not seem to be a relationship with treatment. In order for a test to be considered negative for carcinogenicity, survival should be no less than 50%. Survival was less than 50% in the male controls, 1000, and 5000ppm groups.

It is noted that there was a very high incidence of lung disease in all groups.

Some data entries were missing from the Table 24 (no thymus input for 30/43 animals).

There was no discussion of these results in the light of the findings of the 90-day rat study (A6.4.1) where a statistically significant ($p < 0.01$) and treatment-related increase in absolute and relative liver weight was recorded from 200 mg/kg bw/day (lowest dose). Relative kidney weight was also significantly elevated in both sexes.

Study 2: Oral carcinogenicity study in the mouse

Methoprene technical (86.9 and 87.5%) was administered at target dietary doses of 0, 250, 1000 and 2500 ppm daily over a period of 72 weeks to 4 groups of 50 Charles River CD-1 mice/sex/group. Duration of treatment: 72 weeks for females treated at 1000 ppm and 78 weeks for all other groups. At study termination, all surviving mice were sacrificed, necropsied and fixed *in toto* in buffered neutral 10% formalin. Mice that died during the study were also necropsied and their tissues were saved if autolytic changes were not advanced.

No changes were observed in the mouse carcinogenicity study considered to be treatment related were seen in the general behavior and appearance of the animals. Females of the 1000ppm group were sacrificed at 72 weeks. No reason is given. It can be assumed that this was because survival had fallen to 20/50 (40%). The OECD criteria for a negative (carcinogenicity) test include the

stipulation that survival of all groups should be no less than 50%. Survival was lower than 50% in all treated groups of this study. Mortality rates are described in the following table.

Table 33: Summary of mortality results

Dosage level (ppm)	Number surviving/number initiated	
	Male	Female
Control 1	32/50	25/50
250	28/50	22/50
1000	27/50	20/50 ^a
2500	30/50	24/50

^a prior to sacrifice, week 72

All animals gained weight throughout the study. Increases in body weight were similar for control and treated mice. No changes in food consumption were reported in both sexes at any dose level.

It was noted that the treatment-related incidence of liver pathology (dark brown, finely granular pigment were observed in the cytoplasm of liver parenchymal cells of most male and female mice sacrificed at study termination at 2500 ppm. Many mice also displayed focal accumulations of macrophages with brownish, foamy cytoplasm in their livers. These changes were associated with small necrotic foci and mononuclear inflammatory cells. Less frequently, intracytoplasmic brown pigment was detected in Kupffer cells) was greater and more marked in the 2500-ppm females and in the 10 females from the 1000 ppm survivors. These pigmentary changes were not noted in mice at 250 ppm.

The intracytoplasmic pigment in the hepatocytes was reported to be PAS negative, negative for oil red-o stain and negative for iron. The macrophages stained positive. The composition of the treatment-related pigment accumulation is not known, therefore. It is likely to reflect some toxicity related endpoint and leads to the conclusion that following long-term exposure in the diet, methoprene resulted in probably toxicity-related liver pathology in the mouse, with females being more sensitive. The NOEL for this effect was 250 ppm.

Based on the presence of focal accumulations of macrophages with brownish foamy cytoplasm in the liver, often associated with small necrotic foci and mononuclear inflammatory cells, the NOEL for Methoprene technical under the conditions of this study was established as 1000 ppm (equivalent to 130.8 mg/kg bw/day).

No compound related tumourigenic effect was observed at any dose level. Incidence of the more commonly occurring tumours was similar between the control and the treated groups. There were no statistical differences reported. Furthermore overall tumour incidence was similar between this study and other 18-month studies conducted in this laboratory in this strain of mouse

Methoprene is not concluded to be carcinogenic at dose levels up to 2500 ppm (equivalent to 327 mg/kg bw/day).

Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene as a juvenile hormone is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene

to S-Methoprene. Therefore, the observed NOEL in this study has been revised and a NOEL for S-Methoprene is proposed:

4.9.1.2 Carcinogenicity: inhalation

No data

4.9.1.3 Carcinogenicity: dermal

No data

4.9.2 Human information

No data

4.9.3 Other relevant information

4.9.4 Summary and discussion of carcinogenicity

Technical Methoprene (86.9%) was administered to rats in the diet, in a combined carcinogenicity and chronic toxicity study, at nominal concentrations of up to 5000 ppm (equivalent to 108.6 mg/kg bw/day S-Methoprene) for 24 months. There was no evidence of carcinogenicity in this study. Similarly, there was no evidence of carcinogenicity in the mouse following 72-78 weeks of dietary administration of technical Methoprene (86.9% and 87.5%) at nominal concentrations of up to 2500 ppm (equivalent to 163.5 mg/kg bw/day S-Methoprene).

4.9.5 Comparison with criteria

The CLP criteria for classification as a category 1 Carcinogen are as follows:

“Known or presumed human carcinogens. A substance is classified in Category 1 for carcinogenicity on the basis of epidemiological and/or animal data. A substance may be further distinguished as: Category 1A: Category 1A, known to have carcinogenic potential for humans, classification is largely based on human evidence, or Category 1B: Category 1B, presumed to have carcinogenic potential for humans, classification is largely based on animal evidence. The classification in Category 1A and 1B is based on strength of evidence together with additional considerations. Such evidence may be derived from:

— human studies that establish a causal relationship between human exposure to a substance and the development of cancer (known human carcinogen); or

— animal experiments for which there is sufficient (1) evidence to demonstrate animal carcinogenicity (presumed human carcinogen).

In addition, on a case-by-case basis, scientific judgement may warrant a decision of presumed human carcinogenicity derived from studies showing limited evidence of carcinogenicity in humans together with limited evidence of carcinogenicity in experimental animals”.

The CLP criteria for classification as a category 2 Carcinogen are as follows:

“Substances are classified as a category 2 Carcinogen when evidence is obtained from human and/or animal studies, but which is not sufficiently convincing to place the substance in Category

*1A or 1B, based on strength of evidence together with additional considerations. Such evidence may be derived either from limited evidence of carcinogenicity in human studies or from **limited evidence** of carcinogenicity in animal studies.”*

There are two studies, one rat, and one mouse available for S-Methoprene. Carcinogenic potential is not demonstrated in either. Based on the finding of the aforementioned studies and the lack of any other data suggesting carcinogenicity no carcinogenicity classification is required.

4.9.6 Conclusions on classification and labelling

Based upon the findings of the submitted studies classification of S-Methoprene for carcinogenicity is not required.

4.10 Toxicity for reproduction

Table 34: Summary table of relevant reproductive toxicity studies

Method	Results	Remarks	Reference
*Oral, Rat, Charles River albino female rats, 19-22/group. Day 6-15 of gestation. Methoprene technical: 500 and 1000 mg/kg/day. The study was carried out before the availability of US EPA and OECD guidelines.	Maternal toxicity NOAEL Methoprene technical: 1000 mg/kg bw/day. S-Methoprene: 500 mg/kg/day Bifurcated sternum, cleft sternum and sternal asymmetry at 1000 mg/kg bw/day. Teratogenicity Embryotoxicity NOAEL: Methoprene 500 mg/kg bw/day and S-Methoprene 250 mg/kg/day.		Haley, S. (1972)
OECD Guideline 414, Oral, Rabbit, New Zealand White , Female, 17-25/group. Day 6-27 of gestation. S-Methoprene technical: 25, 100 and 1000 mg/kg/day	Maternal toxicity NOAEL S-Methoprene: 100 mg/kg bw/day. NOAEL Growth retarded foetuses, treatment related maternal death, abortions and vaginal bleeding at 1000 mg/kg bw/day. Teratogenicity Embryotoxicity : 100 mg/kg bw/day		Kolep Csete, K. (2008) CAR IIIA 6.8.1-1
OECD Guideline 414, Oral, Rat , Hsd. Brl. Han: WIST Rats, Female, 22-26/group. Day 5-19 of gestation. S-Methoprene technical: 60, 250 and 1000 mg/kg/day	Maternal toxicity NOAEL S-Methoprene: 250 mg/kg bw/day. NOAEL ↓food consumption and mean body weight gain. ↑post-implantation loss. Teratogenicity Embryotoxicity : 250 mg/kg bw/day		Kolep Csete, K. (2009) CAR IIIA 6.8.1-2

* The rat gavage study of *S. Haley* (1972) was considered invalid because of serious methodological and reporting inadequacies and was not considered in the assessment.

Note: Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene to S-Methoprene.

4.10.1 Effects on fertility

4.10.1.1 Non-human information

Study 1: Teratogenic study with Altosid technical in albino rats. Haley, S., (1972) - CAR IIIA 6.8.1(1)

Conclusion: The rat gavage study of *S. Haley (1972)* was considered invalid because of serious methodological and reporting inadequacies and therefore was not used in the risk assessment.

Study 2: Teratology study of test item S-Methoprene technical in rabbits. Kolep, C.K. (2008) - CAR IIIA 6.8.1(2)

Four groups of 25 inseminated female New Zealand albino white rabbits were dosed, once daily between days 6 to 27 of presumed gestation, by the oral route with S-Methoprene at dose levels of 0, 25, 100 and 1000 mg/kg bw/day. On gestation day 28, euthanasia of the animals was carried out. The internal organs were examined macroscopically and all changes were recorded. The number of corpora lutea in each ovary, the number of implantation sites in each uterine horn, the number of live foetuses, early and late embryonic deaths and dead foetuses were determined. The uterus with cervix and the left ovary were removed and weighed. Live foetuses were individually weighed. The crown rump length of the foetuses and the litter mean was calculated. All foetuses were examined externally and viscerally and the sex of the foetuses was determined.

Maternal effects:

Body weight

Clear maternal toxicity was evidenced by marked adverse effects on mean body weight, body weight gains and corrected body weights and some clinical signs observed at the top dose. Treatment related changes in mean maternal body weight, body weight gain and corrected body weight were seen 1000 mg/kg bw/day dams. A treatment-related and statistically significant ($p < 0.01$) reduction in bodyweight from was seen from day 18 in the 1000 mg/kg bw/day dose group. Body weight gain was statistically significantly ($p < 0.01$) reduced throughout the treatment period. The gravid uterine weight ($p < 0.05$), corrected body weight ($p < 0.01$) and corrected body weight gain ($p < 0.01$) were also statistically significantly lower for the 1000 mg/kg bw/day dose group.

Table 35: Mean maternal body weights (g)

Treatment group	Gestation day (TIME)					
	0	6	12	18	24	28
Control	4009.7 ±236.18	4221.1 ±348.34	4258.3 ±418.72	4433.4 ±387.19	4529.4 ±371.65	4600.0 ±345.08
25 mg/kg/day	3990.0 ±210.69	4242.0 ±223.67	4323.4 ±209.57	4472.5 ±250.09	4530.3 ±271.85	4543.9 ±248.51
100 mg/kg/day	4053.9 ±198.08	4309.2 ±212.84	4408.3 ±216.45	4560.3 ±246.25	4634.3 ±279.61	4659.2 ±302.37

1000 mg/kg/day	4022.8 ±194.26	4267.8 ±207.81	4146.4 ±252.90	4104.2** ±369.64	4097.2** ±416.93	4126.4** ±503.51
-----------------------	-------------------	-------------------	-------------------	---------------------	---------------------	---------------------

*= Statistically significant at $p < 0.05$

**= Statistically significant at $p < 0.01$

Table 36: Mean maternal body weights gain (g)

Treatment group	Gestation day (TIME)					
	0-6	6-12	12-18	18-24	24-28	0-28
Control	211.4 ±171.16	37.2 ±146.93	175.1 ±114.72	96.1 ±94.52	70.6 ±139.65	590.3 ±217.08
25 mg/kg/day	252.0 ±79.04	81.3 ±116.13	149.2 ±126.40	57.7 ±125.07	13.6 ±155.58	553.9 ±180.43
100 mg/kg/day	255.3 ±119.91	99.1 ±112.55	152.0 ±95.30	74.0 ±84.94	25.0 ±128.05	605.4 ±233.68
1000 mg/kg/day	245.0 ±69.65	-111.3** ±132.98	-42.2** ±222.91	-53.7* ±215.57	-14.8 ±237.31	141.2** ±412.06

*= Statistically significant at $p < 0.05$

**= Statistically significant at $p < 0.01$

Mortality

Three does died during the study, two in the control and one at 1000 mg/kg bw/day group. The top dose doe died following abortion and this was considered treatment-related. The two control animals were considered to have died of internal bacterial infection, therefore these deaths were unrelated to treatment.

Abortions and clinical signs

One animal aborted and was sacrificed in the low dose group on gestation day 21. Six animals aborted in the high dose group, (one died following abortion and five were sacrificed after abortion). Except for the doe, which died following abortion on gestation day 17, all the abortions in the high dose group were after the organogenetic period (from gestation day 18) and these rabbits had marked weight loss by the end of the first week of the treatment period. Vaginal bleeding was reported in one control animal, one animal dosed at 25 mg/kg bw/day and two animals dosed at 1000 mg/kg bw/day. One doe at 1000 mg/kg bw/day had bloody urine and blood was found in the cage. A high incidence in the late embryonic death was reported in this doe. Total intrauterine mortality was also noted in the doe that showed vaginal bleeding. The incidences of abortions and vaginal bleeding in the other treatment groups were within the normal range.

Clinical signs included diarrhoea in one animal at 25 mg/kg bw/day and soft faeces in 3, 4 and 6 animals in the 25, 100 and 1000 mg/kg bw/day dose groups, respectively. This was accompanied by reduced body weight in the 1000 mg/kg bw/day dose group. In addition, reduced activity noted in five rabbits in the high dose group was considered related to treatment.

Necropsy and pathology

No treatment related changes were reported at necropsy. One animal that died in the 1000 mg/kg bw/day dose group following abortion showed dark reddish mottled lungs, bloody discharge in the thoracic cavity and nutmeg like patterned lungs. These findings were considered related to abortion and physiological stress. Dead animals in the control group displayed necropsy findings such as blood around nasal orifice, dark reddish mottled lungs and inflammation exudates in the thoracic cavity. These deaths were not treatment-related and were as a result of inter-current or accidental bacterial infection. The maternal reproductive parameters, overall pregnancy rate, number of

corpora lutea, number of implantations were similar in all groups. There were no treatment related changes at histopathological examination.

In summary, there was evidence of maternal toxicity characterized by a significantly reduced mean body weights, body weight gains and corrected body weights at the high dose. Increased instances of treatment-related abortions and vaginal bleeding in the 1000 mg/kg/bw/day dose group were considered treatment-related also. The overly severe effects on the maternal animal do not allow discrimination between the endpoints of maternal and developmental toxicity at the top dose, if such exists with this substance.

Embryo and foetal effects:

The number of viable foetuses in the treated groups was comparable to controls. The number of female foetuses in the 1000 mg/kg bw/day dose group was statistically significantly higher than males (60 v's 40%). However, this incidence was not deemed to be treatment related.

Table 37: Intrauterine mortality, viable foetuses and their sex distribution

Group		Autopsy findings (Mean/Female)			
		Control	25 mg/kg bw/day	100 mg/kg bw/day	1000 mg/kg bw/day
No. Foetuses examined		19	22	21	18
Corpora Lutea	Mean	11.5	10.7	11.1	11.1
	SD	3.47	2.16	2.10	2.74
Pre-implantation sites	Mean	11.2	18.3	14.9	13.1
	SD	11.83	17.49	19.49	14.97
Implantation	Mean	10.4	8.7	9.5	9.8
	SD	3.47	2.41	2.69	3.30
Early Embryonic Death %	Mean	11.5	1.1	2.1	10.9
	SD	25.10	3.85	4.55	23.62
Late Embryonic Death %	Mean	4.6	3.3	3.5	7.4
	SD	5.96	5.60	5.90	16.79
Dead Foetuses %	Mean	1.6	0.5	5.1	2.0
	SD	3.19	2.35	21.80	6.07
Post-implantation Loss %	Mean	17.7	4.9	10.8	20.4
	SD	24.5	7.00	21.82	28.25
Total intrauterine mortality %	Mean	27.3	22.6	24.6	29.6
	SD	24.54	16.68	26.62	27.01
Viable Foetuses	Mean	8.5	8.2	8.5	7.7

	SD	3.45	2.02	3.30	2.97
Male Foetuses % (DN)	Mean	57.1	47.0	53.7	40.2*
	SD	18.58	20.69	16.19	17.82
Female Foetuses % (DN)	Mean	43.0	53.2	46.5	60.0*
	SD	18.56	20.72	16.05	17.85

*= p <0.05

DN= Duncan's Multiple Range Test

Table 38: Intrauterine mortality, viable foetuses and their sex distribution

Group		Autopsy findings (Mean/Female)			
		Control	25 mg/kg bw/day	100 mg/kg bw/day	1000 mg/kg bw/day
No. Foetuses examined		19	22	21	18
Corpora Lutea	Sum	218	236	234	200
Pre-implantation sites	Sum	24	45*	37	24
(Data compared to no. of corpora lutea)	%	11	19	16	12
Implantation	Sum	198	192	199	176
Early Embryonic Death %	Sum	23	3**	4**	16
(Data compared to no. of implantations)	%	12	2	2	9
Late Embryonic Death %	Sum	10	7	7	17
(Data compared to no. of implantations)	%	5	4	4	10
Dead Foetuses %	Sum	4	1	9	4
(Data compared to no. of implantations)	%	2	1	5	2
Post-implantation Loss %	Sum	37	11**	20*	37
(Data compared to no. of implantations)	%	19	6	6	21
Total intrauterine mortality %	Sum	61	56	57	61
(Data compared to no. of corpora lutea)	%	28	24	24	31
Viable Foetuses	Sum	161	181	179	139
Male Foetuses	Sum	89	86	96	55**
(Data compared to no. of viable foetuses)	%	55	48	54	40
Female Foetuses	Sum	72	95	83	84**
(Data compared to no. of viable foetuses)	%	45	52	46	60

foetuses)					
-----------	--	--	--	--	--

Remarks:

*= Statistically significant at $p < 0.05$ Chi Square

**= Statistically significant at $p < 0.01$ Chi Square

External findings

A biologically and statistically significantly lower foetal body weight ($p < 0.05$) and crown-rump length averages ($p < 0.01$) were found in the 1000 mg/kg bw/day dose group. Significant maternal toxicity was demonstrated at this dose level. The number of foetuses with abnormalities was significantly higher ($p < 0.01$) in the 1000 mg/kg bw/day dose group, The overall incidence of variations was statistically significantly higher in the 1000 mg/kg bw/day dose group when compared to the control. This difference was due to the higher incidence of growth retarded fetuses (body weight and crown-rump length evaluated as variations). A number of external variations (misaligned palatine rugae, protruding tongue and bent tail) were reported in the control and treated groups, without dose relationship.

Table 39: Summary of gravid uterine weight, corrected body weight and body weight gain of does

		Dose Groups				
		Control	25 mg/kg	100 mg/kg	1000 mg/kg	
gravid uterine Weight (g)	Mean	542.7	491.3	525.3	437.6	U
	SD	163.81	97.80	125.53	84.08	
	n	18	22	20	17*	
corrected body weight (g)	Mean	4092.1	4052.6	4152.1	3784.3	DN
	SD	285.79	275.83	235.31	271.48	
	n	18	22	20	17**	
corrected body weight gain (g)	Mean	66.1	62.6	88.7	-219.1	DN
	SD	188.89	178.09	182.73	227.36	
	n	18	22	20	17**	

Remarks:

*= Statistically significant at $p < 0.05$

**= Statistically significant at $p < 0.01$

U=Mann-Whitney U-Test Versus Control

DN= Duncan's Multiple Range Test

Visceral examination.

The incidence of rudimentary lung-lobe was significantly higher in the 1000 mg/kg bw/day dose group. The incidence of absent lung-lobe was significantly higher in the 25 mg/kg bw/day and in the 1000 mg/kg bw/day dose group. However, these values were within the historical control range. None of these differences were considered treatment related.

Skeletal examination.

Significantly higher incidences of enlarged anterior fontanelle and un-ossified proximal phalanges of pellex (bilateral) were noted. These incidences resulted in a significant increase in the incidence of foetuses with abnormalities and with variations in the 1000 mg/kg bw/day dose group. These changes were correlated to the intrauterine foetal growth retardation and were deemed to be treatment related.

The average incidence of skeletal malformations was lower in the treated groups than in the control groups. A number of skeletal malformations were recorded in all study groups including controls: (fused sternal bodies, wide sternal bodies, branched and /or fused rib cartilages or branched rib).

However, the skeletal malformations in all groups represented a normal background incidence and a statistical evaluation of skeletal parameters showed that all parameters were within the expected range.

Table 40: Results of external, visceral and skeletal examinations

Dose Groups					
		Control	25 mg/kg	100 mg/kg	1000 mg/kg
Litters examined	N	18	22	20	17
EXTERNAL EXAMINATION					
Foetuses examined	N	161	181	179	139
Foetuses with abnormalities	N	13	8	16	29**
	%	8	4	9	21
Variation	N	12	5*	13	27**
	%	7	3	7	19
Malformation	N	1	3	3	2
	%	1	2	2	1
Retarded in body weight	N	7	4	11	15*
	%	4	2	6	11
Retarded in crown-rump weight	N	6	4	11	22**
	%	4	2	6	16
VISCERAL EXAMINATION					
Foetuses examined	N	161	181	179	139
Foetuses with abnormalities	N	31	37	33	39
	%	19	20	18	28
Variation	N	28	32	31	37
	%	17	18	17	27
Malformation	N	3	5	2	2
	%	2	3	1	1
SKELETAL EXAMINATION					
Foetuses examined	N	152	181	171	139
Foetuses with abnormalities	N	19	18	19	30*
	%	13	10	11	22
Variation	N	11	14	14	26**
	%	7	8	8	19
Malformation	N	8	4	5	4
	%	5	2	3	3
NO. of FOETUSES WITH VARIATION					
	N	43	48	48	68**
	%	27	27	27	49
No. FOETUSES WITH MALFORMATION					
	N	10	8	9	7
	%	6	4	5	5

Remarks:

*= Statistically significant at $p < 0.05$ Chi Square

**= Statistically significant at $p < 0.01$ Chi Square

Conclusion:

Severe maternal toxicity (death, weight loss) was accompanied by significant foetolethality (abortions) and foetotoxicity (runts and retarded ossification) at the high dose of 1000 mg/kg bw/day. However, an NOEL was established for maternal and foetal toxicity at 100-mg/kg bw/day.

Ultimately, the dosing in the rabbit developmental study is considered inadequate. The top dose is considered to be inappropriately high and the mid-range dose provides an NOEL value. However, this NOEL value of 100mg/kg bw/day is used as the NOAEL value and is used to establish a systemic AEL acute reference value even though it may be a more conservative value than what may have been achieved if the study dosing was more appropriately considered.

Study 3: Teratology study of test item S-Methoprene technical in rats. Kolep, C.K. (2009) - CAR IIIA 6.8.1(2)

In a new study, the teratogenicity of S-Methoprene Technical was investigated by oral administration to 4 groups of pregnant female rats at the following concentrations: 0, 60, 250 and 1000 mg/kg bw/day followed by full examination of the dams and the fetuses. The number of sperm positive females in the study was 97. None of the females displayed clinical signs and all females survived until necropsy on gestation day 20.

Maternal Effects:

1000 mg/kg bw/day dose group: There was a statistically significant reduction in body weight gain between gestational days 17 to 20 ($p < 0.01$) and 0 to 20 ($p < 0.05$) compared to the vehicle control level. Corrected body weight was lower in this treatment group than in other experimental groups. However, this was not statistically significant due to the high standard deviation, but was considered biologically relevant. Food consumption of the dams was statistically significantly reduced ($p < 0.05$) during the whole treatment period. This was considered to be treatment related. Mottled reddish lungs were observed at necropsy for four dams in the 60 and 1000 mg/kg bw/day groups. This finding was attributed to euthanasia and was not considered to be treatment related.

250 mg/kg bw/day dose group: There were four non-pregnant animals in this treatment group. At termination on gestation day 20, there were 18 dams with live fetuses and more than five implantations. There were no changes observed in body weight, body weight gain or food consumption. At necropsy, there were no treatment related effects noted.

60 mg/kg bw/day dose group: There were two non-pregnant animals in this treatment group. At termination on gestation day 20, there were 19 dams with live fetuses and more than five implantations. In the 60 mg/kg bw/day treatment group, on necropsy, reddish mottled lungs were observed in four dams in this treatment group. This effect was attributed to euthanasia and was not considered to be treatment related. There were no changes observed in body weight, body weight gain or food consumption. At necropsy, there were no treatment related effects noted.

Table 41: Mean maternal body weights and body weight changes in the teratogenicity study with S-Methoprene Technical

Day of gestation	Dosage level (mg/kg bw/day)			
	Group mean maternal body weights (g)			
	0 (Control)	60	250	1000
	Mean \pm SD	Mean \pm SD	Mean \pm SD	Mean \pm SD
0	205.7 \pm 16.10	198.4 \pm 15.30	196.8 \pm 11.49	201.8 \pm 20.99
5	224.7 \pm 15.74	219.9 \pm 16.72	217.4 \pm 11.89	221.9 \pm 20.36
11	245.5 \pm 15.49	243.1 \pm 18.54	240.0 \pm 16.86	239.8 \pm 21.94
17	280.5 \pm 20.93	278.1 \pm 21.58	275.8 \pm 19.93	273.3 \pm 28.77
20	313.2 \pm 22.56	309.1 \pm 22.47	303.3 \pm 25.23	291.7 \pm 35.41

Days of gestation	Group mean maternal body weight gain (g)			
0 to 5	19.0 ± 6.01	21.5 ± 6.00	20.7 ± 5.12	20.1 ± 6.81
5 to 11	20.8 ± 5.10	23.2 ± 5.73	22.6 ± 7.33	17.8 ± 6.50
11 to 17	34.6 ± 10.44	34.9 ± 7.04	35.8 ± 10.06	33.5 ± 10.91
17 to 20	33.1 ± 9.81	31.1 ± 4.39	27.6 ± 14.73	18.4** ± 17.98
0 to 20	107.5 ± 16.39	110.7 ± 17.46	106.6 ± 21.40	89.9* ± 28.36

Significantly different from control:

* p<0.05

** p<0.01

Table 42: Mean maternal food consumption in the teratogenicity study with S-Methoprene Technical

Day of gestation	Dosage level (mg/kg bw/day)			
	Food consumption (g)			
	0 (Control)	60	250	1000
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD
0 to 5	20.7 ± 3.15	21.1 ± 2.79	20.9 ± 1.21	20.5 ± 3.46
5 to 11	20.0 ± 1.25	20.5 ± 1.72	20.9 ± 1.70	18.1* ± 3.06
11 to 17	21.4 ± 1.15	21.6 ± 1.94	21.9 ± 1.74	19.5* ± 2.48
17 to 20	22.6 ± 1.99	23.3 ± 2.48	21.9 ± 2.16	20.6* ± 2.35

Significantly different from control:

* p<0.05

Embryo and foetal effects:

Foetal body weight was not influenced by treatment with the test substance.

1000 mg/kg bw/day dose group: There was a statistically significant reduction in post-implantation loss at this dose level. There was no treatment related effect observed on either the type or incidence of malformations or variations during skeletal examination of the foetuses.

250 mg/kg bw/day dose group: Pre-implantation loss increased statistically significantly (13%, p<0.05) in this treatment group, however, it was below the historical control level (20%). No external or visceral malformations were noted in the 250 mg/kg bw/day treatment group. One foetus in the 250 mg/kg bw/day group displayed signs of vertebral abnormalities, having lumbar vertebrae malformations. However, this was not considered to be treatment related.

60 mg/kg bw/day dose group: In the 60 mg/kg mw/day treatment group, incomplete ossification of the skull was noted in three foetuses, while marked wavy ribs with or without complete ossification was found in six foetuses in the control and two in this treatment group. Vertebral abnormalities were found in three foetuses in the control and 60 mg/kg bw/day treatment group. There were no visceral malformations in this treatment group. None of the skeletal malformations were increased significantly in the test substance treatment groups

No treatment related foetal external abnormalities or visceral examinations were observed in the foetal litter. None of the skeletal malformations were increased significantly in the treatment groups.

Table 43: Pregnancy data in the teratogenicity study with S-Methoprene Technical

Dose groups	Dosage level (mg/kg bw/day)			
	0 (Control)	60	250	1000
No. of sperm positive females	26	22	22	27
Percentage of pregnant females	89 %	91 %	82 %	70 %
No. of non-pregnant females	3	2	4	8
No. of pregnant females with no implantation but corpora lutea	0	0	0	1
No. of dams with total intrauterine death	0	0	0	1
No. of pregnant females with viable foetuses, but with less than or equal to 5 implantations	0	1	0	1
Percentage of evaluated dams with malformed foetuses	39 %	37 %	11 %	25 %

Conclusion:

Maternal toxicity was evidenced at the high dose by a statistically significant reduction in food consumption and mean weight gain. There was also a statistically significant reduction in post-implantation loss at this dose level. A NOEL of 250 mg/kg/day was established for both maternal and developmental toxicity.

4.10.1.2**4.10.1.3 Human information**

No data.

4.10.2 Developmental toxicity**4.10.2.1 Non-human information****Table 44: Summary table of relevant reproductive toxicity studies**

Method	Results	Remarks	Reference
Rat, Long-Evans, Male and female, 20 animals /sex/ dose. Over a three generation period Methoprene technical : 500 and 2500 ppm equivalent to 0, 16.3 and 261.6 mg/kg bw/day.	Overall, minimal and insufficient parental toxicity was demonstrated in this study. The only effects were slight reduction in mean pup weights seen at day 21 of lactation in the F2 generation and throughout lactation in the F3 generation.	The study was carried out before the availability of US EPA and OECD guidelines. The dosing parameters for testing the substance were not extended sufficiently to	Killeen, J.C., Rapp, W.R. (1974) CAR IIIA 6.8.2

		produce the required parental toxicity and as a consequence this substance was not assessed to its full extent	
--	--	--	--

Note: Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene to S-Methoprene.

Study 1: A three generation reproduction study of Altosid™ in rats. Killeen, J.C. (1974) - CAR IIIA 6.8.2

Methoprene technical was administered to 3 groups of 20 Long-Evans rats sex/group at concentrations of 0, 500 and 2500 ppm (equivalent to 0, 16.3 and 261.6 mg/kg bw/day Methoprene technical and 0, 8.15 and 130.8 mg/kg bw/day S-Methoprene) over a three generation period. This study was not performed according to guidance but has been compared to OECD Guideline 416.

During the growth period, animals from both sexes in the F₀ and F₁ generations showed lower mean body weights and mean weight gains than controls at 2500-ppm dose level. In the F₂ generation, mean body weights were lower than control at the 500 and 2500-ppm dose level groups. However, mean weight gains were comparable. Results are summarised in Table 45.

Table 45: Summary Mean Body Weights – growth periods (g)

Dose level ppm	Growth period												Weight gain Initial-Week 9/10
	Initial	Baseline	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6	Week 7	Week 8	Week 9	Week 10	
F₀ generation - males													
0	79.0	113.3	149.9	188.0	227.8	257.4	286.4	308.6	328.6	345.4	361.9	-	282.9
500	79.1	115.9	152.8	196.1	238.8	264.8	287.1	309.7	329.3	345.9	360.7	-	281.6
2500	79.1	116.5	151.4	192.0	233.6	259.9	282.2	301.3	321.9	338.9	351.4	-	272.3
F₁ generation - males													
0	127.5	-	169.3	211.7	254.3	290.2	315.5	312.0	300.3	346.1	371.9	384.6	257.1
500	121.0	-	159.5	200.2	240.0	276.0	301.9	304.9	294.7	330.5	355.1	371.2	250.2
2500	122.0	-	159.8	203.2	241.3	276.8	302.5	308.2	293.9	325.9	346.3	365.9	243.9
F₂ generation - males													
0	155.5	-	192.6	244.3	281.6	315.2	337.3	367.1	383.5	396.2	404.5	392.0	236.5
500	131.9	-	172.2	211.7	252.2	281.4	298.8	325.7	344.0	357.3	366.1	366.9	235.0
2500	139.2	-	a	222.3	257.7	285.6	299.2	323.5	349.2	364.5	369.6	376.8	237.6
F₀ generation - females													
0	71.1	103.6	131.4	155.9	180.1	193.6	201.6	212.4	223.6	229.2	238.0	-	166.9
500	71.1	100.6	128.8	154.6	178.1	188.9	196.7	207.5	219.5	225.2	233.2	-	162.1
2500	71.1	99.7	126.7	149.9	174.4	185.7	194.6	203.6	215.2	220.6	227.1	-	156.0
F₁ generation - females													
0	113.2	-	134.7	158.7	182.2	200.0	211.9	213.7	209.3	224.9	231.5	243.8	130.6
500	110.5	-	131.0	156.3	175.9	193.9	205.0	209.8	203.1	223.3	226.9	241.1	130.6
2500	108.6	-	131.5	155.4	174.8	192.4	204.8	207.3	214.3	223.9	228.1	235.4	126.8
F₂ generation - females													
0	126.1	-	148.6	170.0	186.0	201.8	213.0	226.3	234.1	239.1	244.6	247.0	120.9
500	121.6	-	146.9	165.6	182.6	196.8	202.8	218.8	228.6	232.7	237.5	237.5	115.9
2500	114.2	-	137.8	156.8	172.4	188.6	195.5	208.6	215.8	222.0	225.5	232.2	118.0

^aNot recorded

During gestation and lactation, decreased maternal mean body weights were observed at 2500-ppm dose level throughout all generations. In the F₂ generation decreased maternal body weights were also observed at 500-ppm dose level. These effects were not considered to be treatment related. Results are summarised in Table 46.

Table 46: Summary Mean Maternal Body Weights (g)

Dose level (ppm)	Premating	Gestation				Weight gain	Lactation				Weight gain
		Day 0	Day 6	Day 15	Day 20	Gestation 0-20	Day 0	Day 4	Day 14	Day 21	Lactation 0-21
F₀ Mating											
0	238.0	240.3	260.9	301.1	358.3	118.1	286.7	292.8	301.9	293.2	6.5
500	233.2	239.5	261.6	303.8	363.3	123.8	295.9	302.6	311.8	303.9	8.1
2500	227.1	235.3	258.2	297.7	348.0	112.7	286.1	290.4	301.5	296.5	10.4
F₁ Mating											
0	243.8	251.2	279.1	309.8	368.8	117.6	301.3	307.1	308.1	300.3	-0.1
500	241.1	251.2	274.7	311.3	368.8	117.6	297.1	300.3	302.0	301.9	4.8
2500	235.4	244.0	269.4	304.2	357.6	113.6	292.7	307.3	313.3	291.4	-1.3
F₂ Mating											
0	247.0	252.4	282.4	316.4	370.9	118.1	299.0	310.4	310.4	311.6	12.6
500	237.5	245.7	269.0	303.5	356.1	110.4	296.8	298.6	306.3	309.7	12.9
2500	232.2	236.4	262.2	293.8	343.4	107.0	280.1	290.3	291.9	296.0	15.9

In the F₂ mating (F₃ generation), there was a statistically significant decrease in the percentage of live pups born at 2500-ppm dose level. This difference was due to two litters, which contained large numbers of stillborn pups and was not considered to be treatment related. Results are summarised in Table 47. Slightly decreased mean offspring weights were observed at 2500-ppm dose level in the F₁ mating (F₂ generation) on Day 21 of lactation and in the F₂ mating (F₃ generation) throughout the lactation period. Results are summarised in Table 47.

Table 47: Summary of Survival and Growth of Offspring

Dose level ppm	Mean gestation length	Mean No. born/litter		% Gestation survival	Mean No. weaned/litter	% Postnatal offspring survival Days			% Litters with offspring deaths	% Litters weaned	Mean live offspring weights (g)			
		Days	Alive			Dead	Alive/total born	0-4			4-14	4-21	Days 0-21	Day 0
F₀ Mating (F₁)														
0	22.0	11.3	0.1	99.4	9.6	92.8	98.0	98.0	25.0	93.8	5.988	9.070	25.168	36.486
500	21.9	10.9	0.1	99.4	8.4	96.0	97.1	97.1	31.3	100.0	5.838	8.726	26.546	38.319
2500	22.1	10.0	0.3	97.4	8.6	95.3	86.5**	86.5**	31.6	89.5	6.061	8.995	24.345	35.501
F₁ Mating (F₂)														
0	22.3	10.7	0.1	99.0	8.5	89.6	91.9	91.9	72.2	88.9	6.214	9.091	25.788	39.536
500	22.1	11.3	0.5	96.0	8.1	78.1**	89.6	89.6	70.6	88.2	5.785	7.946	23.484	37.101
2500	22.4	10.9	0.3	97.4	8.7	85.4	93.1	90.3	58.8	88.2	6.125	8.666	24.116	35.885
F₂ Mating (F₃)														
0	22.1	10.2	0.4	96.0	8.7	97.2	99.2	99.2	21.4	100.0	6.184	10.421	29.044	39.635
500	21.9	10.4	0.1	99.4	8.8	99.4	100.0	99.3	12.5	100.0	6.051	9.858	27.784	39.477
2500	22.1	9.7	1.2	88.8*	9.0	99.4	97.0	97.0	15.8	94.7	5.886	9.412	25.457	35.967

Significantly different from control:

* (p<0.05)

** (p<0.01)

At necropsy, no treatment related findings were reported in the F₃ generation. Based on the effects observed on the body weights in adult animals and offspring at 2500-ppm dose level, the NOEL for reproductive toxicity was established to be 500 ppm (equivalent to 16.3 mg/kg bw/day). An observed NOEL for S-Methoprene of 8.15 mg/kg bw/day is established.

Conclusion: Methoprene technical (86.9 and 87.5%) was administered in the diet to two generations of Long-Evans rats at two dose levels of 500 and 2500 ppm (equivalent to 8.15 mg/kg bw/day and 130.8 mg/kg bw/day S-Methoprene).

Overall, minimal and insufficient parental toxicity was demonstrated in this study. The only effects were slight reduction in mean pup weights seen at day 21 of lactation in the F₂ generation and

throughout lactation in the F3 generation. In conclusion, a clear NOEL of 500 ppm (equivalent to 8.15 mg/kg bw/day S-Methoprene) was established.

The study demonstrated there was no evidence of an adverse effect on reproduction. However it should be noted that insufficient parental effects and toxicity were demonstrated and only two doses were used. Ultimately, it is felt that the dosing parameters for testing the substance were not extended sufficiently to produce the required parental toxicity and as a consequence this substance was not assessed to its full extent.

4.10.2.2 Human information

None available.

4.10.3 Other relevant information

4.10.4 Summary and discussion of reproductive toxicity

Developmental Toxicity: Rat

The teratogenicity of S-Methoprene Technical was investigated by oral administration to 4 groups of pregnant female rats at the following concentrations: 0, 60, 250 and 1000 mg/kg bw/day. Maternal toxicity was evidenced at the high dose by a statistically significant reduction in food consumption and mean weight gain. There was also a statistically significant reduction in post-implantation loss at this dose level. A NOEL of 250 mg/kg/day was established for both maternal and developmental toxicity.

Developmental Toxicity: Rabbit

In the rabbit developmental study severe maternal toxicity (death, weight loss) was accompanied by significant foetolethality (abortions) and foetotoxicity (runts and retarded ossification) at the high dose of 1000 mg/kg bw/day. An NOEL was established for maternal and foetal toxicity at 100-mg/kg bw/day. Ultimately, the dosing in the rabbit developmental study is considered inadequate. The top dose is considered to be inappropriately high and the mid-range dose provides an NOEL value. However, this NOEL value of 100mg/kg bw/day is used as the NOAEL value and is used to establish a systemic AEL acute reference value even though it may be a more conservative value than what may have been achieved if the study dosing was more appropriately considered.

Fertility: Rat

Methoprene technical was administered in the diet to two generations of Long-Evans rats at two dose levels of 500 and 2500 ppm (equivalent to 8.15 mg/kg bw/day and 130.8 mg/kg bw/day S-Methoprene). Overall, minimal and insufficient parental toxicity was demonstrated in this study. The only effects were slight reduction in mean pup weights seen at day 21 of lactation in the F2 generation and throughout lactation in the F3 generation. In conclusion, 500 ppm (8.15 mg/kg bw/day) was a clear NOEL.

The study demonstrated there was no evidence of an adverse effect on reproduction. However it should be noted that insufficient parental effects and toxicity were demonstrated and only two doses were used. Ultimately, it is felt that the dosing parameters for testing the substance were not extended sufficiently to produce the required parental toxicity and as a consequence this substance was not assessed to its full extent.

4.10.5 Comparison with criteria

The criteria for classification as Cat 1A (H360D May damage the unborn child) are as follows:

‘...known to have produced an adverse effect on sexual function and fertility, or on development in humans or when there is evidence from animal studies, possibly supplemented with other information, to provide a strong presumption that the substance has the capacity to interfere with reproduction in humans. The classification of a substance in Category 1A is largely based on evidence from humans.’

The criteria for classification as Cat 1B (H360D May damage the unborn child) are as follows:

‘...clear evidence of an adverse effect on (sexual function and fertility or on) development in the absence of other toxic effects, or if occurring together with other toxic effects the adverse effect on reproduction is considered not to be a secondary non-specific consequence of other toxic effects’.

The criteria for classification as Cat 2 (H361D Suspected of damaging the unborn child) are as follows:

‘...some evidence from humans or experimental animals, possibly supplemented with other information, of an adverse effect on sexual function and fertility, or on development, and where the evidence is not sufficiently convincing to place the substance in Category 1. If deficiencies in the study make the quality of evidence less convincing, Category 2 could be the more appropriate classification. Such effects shall have been observed in the absence of other toxic effects, or if occurring together with other toxic effects the adverse effect on reproduction is considered not to be a secondary non-specific consequence of the other toxic effects.’

In accordance with the CLP Regulation (EC) No. 1272/2008 classification criteria for reproductive toxicants, S-Methoprene does not classify for developmental toxicity or teratogenicity. S-Methoprene was determined not to affect fertility in the rat in a 2-generation study.

4.10.6 Conclusions on classification and labelling

In accordance with the classification criteria of the CLP Regulation (EC) No. 1272/2008 and the Guidance to Regulation (EC) No. 1272/2008 on Classification, Labelling and Packaging of substances and mixtures), S-Methoprene does not warrant classification as a reproductive or developmental toxicant.

4.11 Other effects

4.11.1 Non-human information

4.11.1.1 Neurotoxicity

There was no indication of neurotoxicity such as behavioural changes or neurological disturbances from the standard systemic toxicity studies conducted. This includes any single or repeat dose studies conducted. In addition, neurotoxicity studies are not a standard requirement as they are only required for substances of similar or related structures to those capable of inducing delayed neurotoxicity such as organophosphates.

4.11.1.2 Immunotoxicity

No data.

4.11.1.3 Specific investigations: other studies

No data.

4.11.1.4 Human information

No data.

4.11.2 Summary and discussion

No data.

4.11.3 Comparison with criteria

No data.

4.11.4 Conclusions on classification and labelling

No data.

5 ENVIRONMENTAL HAZARD ASSESSMENT

5.1 Degradation

5.1.1 Stability

Table 48: Summary of relevant information on degradation

Method	Results	Remarks	Reference
Hydrolysis			
OECD 111	S-Methoprene technical was found to be hydrolytically stable at pH 4, 7 and 9 at all temperatures examined. (25, 37 and 50). At pH 1.2 hydrolysis of the test material was rapid with a DT ₅₀ value of 17 hours.	The study was considered acceptable with a Reliability score of 2.	Laky, V. (2002a), Hydrolysis of S-Methoprene technical as a function of pH. Toxicological Research Centre Ltd., H-8200 Veszprém-Szbadápuszta, P.O.Box 348, 8201, Hungary, unpublished report no.: 01/616-336AN. CAR IIIA 7.1.1.1.1
Photolysis in water			
OECD Draft Guideline: Phototransformation of Chemicals in Water-Direct and Indirect Photolysis (August 2000)	The aqueous photolysis study performed under lab conditions is not representative of natural conditions and highly overestimates the degradation potential of S-Methoprene and the levels of metabolites (15 d continuous irradiation with a Xe lamp, pH 7, sterilised, 22 ± 2 °C). Under field conditions, photolysis in water may only be relevant in the upper few centimetres of a water body.	The study was considered to be acceptable but yielded uncertain results. Reliability score of 2.	McCorquodale, G. (2009), Photodegradation of [¹⁴ C]-S Methoprene, Charles River, Tranent, Edinburgh, EH33 2NE, UK. Unpublished report no: 807299. CAR IIIA 7.1.1.1.2-1

5.1.2 Biodegradation

5.1.2.1 Biodegradation estimation

Table 49: Summary of relevant information on biodegradation

Method	Results	Remarks	Reference
Biodegradation			
OECD 301D	<p>S-Methoprene technical, at a concentration of 2 mg/l and 8 mg/l, attained 49.45% and 20.99% degradation, respectively, after 28 days.</p> <p>The concentration of 8 mg/L is exceeds the water solubility of S-Methoprene (6.85 mg/L).</p> <p>The results observed at 2 mg/L are considered reliable. S-Methoprene is not readily biodegradable.</p>	The study was considered acceptable with a Reliability score of 2.	<p>Gáty, S. (2002a), Determination of biodegradability of S-Methoprene Technical test item with closed bottle test.</p> <p>Toxicological Research centre Ltd., H-8201 Veszprém, Szabadságpuszta, P.O. Box 348, Hungary, unpublished report no.:01/616-322AN.</p> <p>CAR IIIA 7.1.1.2.1</p>
OECD 302C	The percentage biodegradation of S-Methoprene reached a mean of 4.2 % after 7 days, 24.5 % after 14 days, 77.5 % after 21 days and 85.8 % after 28 days.	The study was considered acceptable with a Reliability score of 1.	<p>Dr. Vértesi, A. (2014); Inherent Biodegradability of S-Methoprene In Modified MITI Test (II). TOXI-COOP ZRT, 8230 Balatonfüred, Arácsi út 97, Hungary , unpublished report No.: 484.462.3617</p>

This closed bottle test concluded that S-Methoprene is not readily biodegradable. The OECD guideline states “*If in a toxicity test, containing both the test substance and a reference compound, less than 35% degradation (based on total DOC) or less than 25% (based on total ThOD or ThCO₂) occurred within 14 days, the test substance can be assumed to be inhibitory (see Annex II for other toxicity tests). The test series should be repeated, using a lower concentration of test substance (if this can be done without seriously impairing the accuracy of the DOC determination) and/or a higher concentration of inoculum, but not greater than 30 mg solids/l.*”. At the higher test concentration of 8 mg/L less degradation took place relative to tests performed at 2 mg/L. This suggests a concentration effect. The study does not report the concentration of test item used in the toxicity test. If the test was performed at a low concentration of test item, negligible inhibition may take place resulting in the observed result in the tox control. The CA notes the OECD 301

guideline states “*If inhibition due to toxicity is to be avoided, it is suggested that the test substance concentrations used in ready biodegradability testing should be less than 1/10 of the EC₅₀ values (or less than EC₂₀ values) obtained in toxicity testing*”. For S-Methoprene the EC₅₀ for activated sludge is reported as >100 mg/L (3 hr). The test concentrations used in the experiment were 2 mg/L and 8 mg/L. The higher test concentration of 8 mg/L lies above the water solubility. The results at 2 mg/L maybe more reliable than the results at 8 mg/L. The validity criteria for the test were fulfilled.

The modified MITI (II) test showed >70% degradation within 28 days. This represents inherent biodegradability (as specified in TGD). The failure to reach 70% within 14 days means that the specific inherent biodegradability criteria were not met and therefore that extrapolation of the results for use in STP models is not possible.

5.1.2.2 Screening tests

Not relevant to this dossier

5.1.2.3 Simulation tests

Not relevant to this dossier

5.1.3 Summary and discussion of degradation

S-Methoprene is not considered to be readily biodegradable and it is hydrolytically stable at environmentally relevant pH values. Photolysis may only be relevant in the upper few centimeters of a water body. These findings are appropriately reflected in the classification and labelling of S-Methoprene.

5.2 Environmental distribution

5.2.1 Adsorption/Desorption

Table 50: Summary of adsorption/desorption

Method	Results	Remarks	Reference
Adsorption and mobility in soil			
OECD 106	S-Methoprene is readily adsorbed to and desorbed from the soil. With K_{oc} values of 537, 684 and 1407 in three soil types and an average of 876, S-Methoprene is classified as being of low mobility according to the McCall and UK Soil Survey and Land Research Centre Pesticide Mobility classification systems.	The study was considered acceptable with a Reliability score of 1.	Laky, V. (2002b), Adsorption/desorption test of S-Methoprene technical. Toxicological Research Centre Ltd., H8200 Veszprém-Szabadságpuszta, P.O.Box 378, 8201, Hungary, unpublished report no.: 01/616-331TL. CAR IIIA 7.2.3.1

No further studies were submitted due to the use pattern proposed for S-Methoprene and the indication that based on the above study that S-Methoprene exhibits low mobility in soil. The three soil types used in the study with the corresponding distribution coefficients are described below.

Table 51: Summary of soil types used in study

Soil type	Type 1	Type 2	Type 3
Sand (%)	14.7	48.9	88.96
Clay (%)	22.2	18.1	5
Organic carbon (%)	1.16	1.21	-
pH	7.31	5.65	4.64
K_d (L/kg)	7.9	6.5	5.5
K_{oc} (L/kg)	684	537	1407
1/n	-	-	-
r^2	-	-	-

5.2.2 Volatilisation

Not relevant for this dossier. The vapour pressure of (3.15 mPa) and molecular weight (310.5) of S-Methoprene allow that it will not readily volatilise into the atmosphere at ambient temperature and pressure.

5.2.3 Distribution modelling

Table 52: Summary of distribution modeling

Method	Results	Remarks	Reference
Mackay Level I fugacity model applying Type 1 type chemical partitioning	S-Methoprene was predicted to partition predominantly to soil (97.8%) and to a much lesser extent to sediment (2.2%). Insignificant amounts are anticipated to be distributed to suspended sediment (0.07%), to air (0.000001%), to fish (0.006%), to water (0.00003%) and to aerosols in the atmosphere (0.0000004%). S-Methoprene has a fugacity value of 3.03×10^{-11} Pa and is found to partition predominantly to soil.	The modeling was considered acceptable with a Reliability score of 1.	Laky, V. (2002b), Adsorption/desorption test of S-Methoprene technical. Toxicological Research Centre Ltd., H8200 Veszprém-Szabadságpuszta, P.O.Box 378, 8201, Hungary, unpublished report no.: 01/616-331TL. CAR IIIA 7.2.3.1

5.3 Aquatic Bioaccumulation

5.3.1 Aquatic bioaccumulation

5.3.1.1 Bioaccumulation estimation

For S-Methoprene (CAS: 65733-16-6) calculations using BCFBAF, the CAS number is provided to the software and the SMILES structure are generated automatically. Also the partition coefficient value is also provided, i.e. Log Kow 6.34. For this model other physical-chemical properties are not necessary as they are not part of this specific model calculation. The calculated BCF is 516. This result is consistent with literature values of the BCF of S-Methoprene. The UK Pesticide Database and the US EPA Integrated Pest Management Plan, 2006, both report a BCF of 457 for S-Methoprene. According to CLP Regulation (EC) No. 1272/2008 the bioconcentration factor (BCF) threshold limit for classification purposes is 500. From the calculation the BCF of S-Methoprene is 516, therefore, based on these results, S-Methoprene meets B criterion (CAR IIIA7.4.2.1). Please note that according to the criteria for the PBT assessment for Annex XIII to Regulation (EC) 1907/2006 S-Methoprene does not meet the B criterion (>2000).

5.3.1.2 Measured bioaccumulation data

No studies were performed on bioaccumulation.

5.3.2 Summary and discussion of aquatic bioaccumulation

The calculated bioaccumulation factor (BCF) of 516 marginally exceeds the threshold limit for classification purposes under Regulation (EC) No. 1272/2008. Therefore S-Methoprene is considered to have potential to bioaccumulate.

5.4 Aquatic toxicity

Table 53: Summary of relevant information of S-Methoprene technical on aquatic toxicity (text highlighted in bold indicates key study)

Method	Results	Remarks	Reference
OECD 203	96 hr LC ₅₀ =4.26 mg/L (measured)	Lowest endpoint in this study	CAR, IIIA7.4.1.1/01. Gáty, S (2002a). Fish acute toxicity study S-Methoprene technical test item on Zebrafish, Toxicological Research Centre Ltd., Report No. 01/616-009H, GLP (unpublished).
OECD 202 ; Comm Reg. (EC) No 440/2008 ; EPA Guideline 712-C-96-114	48 hr EC₅₀=0.22 mg/L (measured)	24 hr EC₁₀₀=0.66 mg/L (measured)	CAR, IIIA7.4.1.2/02. Istvan, A.(2012) Acute Toxicity of S-Methoprene on <i>Daphnia magna</i> in a 48-hour Acute Immobilisation Test, TOXI-COOP ZRT., 8230 Balatonfüred, Arácsi út 97 , Hungary, report no.: 484.441.3614 (unpublished).
OECD 202	48 hr LC ₅₀ =0.38 mg/L (nominal)	Supporting study	CAR, IIIA7.4.1.2/01. Gáty, S. (2002d) Acute immobilisation test with S-Methoprene technical in <i>Daphnia magna</i> , Toxicological Research Centre Ltd., Report No. 01/616-023DA, GLP (unpublished).
OECD 211	21 d NOEC=0.019 mg/L (measured)	Most sensitive long-term endpoint	CAR, IIIA7.4.3.4/01. Istavan, A. (2012) Chronic Toxicity of S-Methoprene

			to <i>Daphnia magna</i> in a 21-day Reproduction Test, TOXI-COOP ZRT., 8230 Balatonfüred, Arácsi út 97 , Hungary report no.: 484.447.3615 (unpublished).
OECD 201	72 hr E _r C ₅₀ =2.264 mg/L (nominal)	No remarks	CAR III, 7.4.1.3/01. Hernádi, D. (2002) Algal growth inhibition test with S-Methoprene technical. Toxicological Research Centre Ltd., Report No. 01/616-022AL, GLP (unpublished).
OECD 209	3 hr EC ₅₀ >100mg/L	Static respiration inhibition test	CAR, IIIA7.4.1.4/01 Gáty, S. (2002c) Activated sludge, respiration inhibition test with S-Methoprene technical test item. Toxicological Research Centre Ltd., Report No. 01/616-027AS, GLP (unpublished).

Note: Methoprene is a terpenoid consisting of a racemic mixture of two enantiomers (R and S in a ratio of 1:1). The activity of Methoprene is restricted to the S enantiomer (S-Methoprene). Since S-Methoprene is the active isomer in the racemic mixture and as they are present in a 1:1 ratio it is possible to extrapolate the results obtained in the studies with Methoprene to S-Methoprene.

5.4.1 Fish

5.4.1.1 Short-term toxicity to fish

Nominal test concentrations in the static study (A7.4.1.1) were 0.63, 1.25, 2.50, 5.00 and 10.00 mg/L. The concentration of the test material was not measured, before, after or during the course of the experiment. Observations of fish were carried out at 3, 6, 24, 48, 72, and 96 hr. Mortality and

sub-lethal effects (fast motility of operculum, decreased activity, localisation near bottom of the aquarium, darkening of body colour and eyes) were recorded at sampling times. No adverse effects were noted in the controls or the two lowest test concentrations. The LC₅₀ values were calculated by probit analysis with 95 % confidence limits using TOXSTAT 3.5 statistical software. At 2.50 mg/L fast motility of operculum, decreased activity, localisation near bottom of the aquarium were recorded. At the highest two concentrations these effects in addition to the darkening of body colour were recorded. The highest concentration used producing no mortality was 1.25 mg/L and the lowest concentration producing 100 % mortality was 10.00 mg/L. The study has a reliability of 2 due to the minor deviation from the OECD Guideline 203, whereby test concentrations were not measured. S-Methoprene is acutely toxic to fish, with the LC₅₀ at 96h of 4.26 mg/L.

5.4.1.2 Long-term toxicity to fish

No long-term toxicity study was submitted. Not relevant for this dossier.

5.4.2 Aquatic invertebrates

5.4.2.1 Short-term toxicity to aquatic invertebrates

The key study on *Daphnia magna* (A7.4.1.2.2) was conducted in semi-static conditions because it is an adsorbing substance, of the test item. All vessels were preconditioned for at least two days prior to test initiation using solutions of the test substance in order to minimise the concentration decline during the experiment. Water renewal periods were 24 hours. The measured test concentrations (geometric mean of four replicates) were 0.07, 0.12, 0.22, 0.40 and 0.66 mg/L. Immobilisation of the test animals was recorded at 24 and 48 hr after treatment. At 24 and 48 hr EC₅₀ values of the test item and their confidence limits were calculated using Probit analysis by SPSS PC+ software. The 24 hr EC₅₀ was > 0.66 mg/L and the 48 hr EC₅₀ was calculated to be 0.22 mg/L. The corresponding 48 hr NOEC and EC₁₀₀ were 0.12 and 0.66 mg/L respectively. This study has a reliability of 1. S-Methoprene is acutely toxic to the aquatic invertebrate, *Daphnia magna*, with the EC₅₀ at 48h of 0.22 mg/L.

The supporting study on *Daphnia magna* (A7.4.1.2.1) had five nominal concentrations of S-Methoprene: 0.15, 0.24, 0.39, 0.63 and 1.00 mg/L. There were two deviations from the OECD Guideline 202: the concentration of the test substance was not measured at the highest and lowest concentrations at the beginning and end of the test and the hardness of the dilution water was not documented. These deviations were not considered to have affected the scientific validity of the study or the interpretation of the results, however, the study has a reliability of 2. The 24 hr EC₅₀ of S-Methoprene for *Daphnia magna* was calculated to be 0.48 mg/L and 48 hrEC₅₀ value was calculated to be 0.38 mg/L.

5.4.2.2 Long-term toxicity to aquatic invertebrates

In this semi-static study (A7.4.3.4), the nominal test item concentrations were 0.01, 0.02, 0.04, 0.07, 0.12 and 0.20 mg/L. Test solutions were renewed three times per week. The test solutions were prepared using water miscible solvent (acetone). An untreated control and an additional solvent control group were investigated concurrently. There were 10 replicates per treatment. The concentration of the solvent (acetone) was 0.1 mL/L in each test concentration and in the additional solvent control. The measured test item concentrations deviated more than 20 % from the nominal during the experiment therefore the time-weighted mean of the measured start and end concentrations at each water renewal period were calculated in order to determine exposure

concentrations. The calculated time-weighted mean concentrations were the followings: 0.009, 0.019, 0.030, 0.049, 0.074 and 0.131 mg/L. All biological results are based on these measured test item concentrations. The offspring produced by each parent animal were removed and counted daily from the appearance of the first brood. Mortality was measured daily. Additionally, the length of the parent animals was measured at the end of the test. The validity criteria of the study were fulfilled. The long-term NOEC at 21 d, based on offspring, was statistically determined to be 0.019 mg/L (Bonferroni t-Test, $\alpha=0.05$). This was also the NOEC for growth measurement. Mortality results did not follow a dose response pattern so these results were excluded from the reproductive output analysis. S-Methoprene is chronically toxic to the aquatic invertebrate, *Daphnia magna*.

5.4.3 Algae and aquatic plants

5.4.3.1 Short-term toxicity to algae and aquatic plants

In the study on algal growth (A7.4.1.3) nominal test concentrations were 0.0625, 0.1250, 0.5000, 1.000 and 2.000mg/L. There were three vessels per test concentration, six per control group and three per reference concentration. The S-Methoprene concentrations were not analysed throughout the study, however this deviation was not considered to affect the scientific validity of the study. Cell concentrations were determined at 0, 24, 48, and 72 hours. The 0 – 72 hour average specific growth of S-Methoprene at the following concentrations: 0.125, 0.25, 0.5, 1 and 2 mg/L were significantly different from that of the control group. The ErC_{50} was the only endpoint that was documented: $ErC_{50} = 2.264$ mg/L.

5.4.3.2 Long-term toxicity to algae and aquatic plants

Not relevant for this dossier.

5.4.4 Other aquatic organisms (including sediment)

5.4.4.1 Short-term toxicity to other aquatic organisms (including sediment)

In the study on the inhibition to aquatic microbial activity (A7.4.1.4) the initial test concentrations were nominally 6.3, 12.5, 25, 50 and 100 mg/L. Actual concentrations were not documented. The test parameter was respiration inhibition, based on oxygen consumption rate, in activated sludge. Measurements of oxygen uptake of the activated sludge were performed 3 hours after initiating aeration. The validity criteria for the control respiration rates and reference material EC_{50} values were satisfied. The effect of the test material on the respiration of activated sewage sludge micro-organisms gave a 3 hour EC_{50} of greater than 100 mg/L. Therefore, it can be assumed that S-Methoprene is non-toxic to sewage treatment microbes.

5.4.4.2 Long-term toxicity to other aquatic organisms (including sediment)

Not relevant for this dossier.

5.5 Comparison with criteria for environmental hazards (sections 5.1 – 5.4)

CLP

S-Methoprene classifies as Acute Category 1 based on the lowest endpoint from acute studies on all three trophic levels: *Daphnia magna*: 48 hr EC₅₀=0.22 (0.1 < LC₅₀ ≤ 1 mg a.s./L). Chronic toxicity was only assessed for aquatic invertebrates, however since this can be expected to be the most sensitive level the resulting hazard category accurately reflects the chronic risk to aquatic organisms. Therefore, S-Methoprene also classifies as Chronic Category 1 classification: *Daphnia magna*: NOEC = 0.019 mg/L (measured) (0.01 < NOEC ≤ 0.1 mg/L), and is not readily degradable.

5.6 Conclusions on classification and labelling for environmental hazards (sections 5.1 – 5.4)

Based on the requirements of the CLP Regulation (EC) No. 1272/2008; S-Methoprene should be Classified:

Aquatic Chronic 1

H400 Very toxic to aquatic life

H410 Very toxic to aquatic life with long lasting effects

Signal Word: Warning

Pictogram:



The environmental hazard pictogram is required

P273 Avoid release to the environment

P391 Collect spillage

P501 Dispose of contents/container in accordance with applicable regulations

Acute M-factor of 1 is applicable based on $0.1 < LC_{50} \leq 1$ mg/L.

Chronic M-factor of 1 is applicable based on $0.01 < NOEC \leq 0.1$ mg/L.

6 OTHER INFORMATION

Not applicable.

7 REFERENCES

8 ANNEXES