

Summary of product characteristics for a biocidal product

Product name: STORM PASTE

Product type(s): PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

Authorisation number: NL-0000040-0000

R4BP 3 asset reference number: NL-0000040-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	3
4. Authorised use(s)	3
5. General directions for use	14
5.1. Instructions for use	14
5.2. Risk mitigation measures	15
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	15
5.4. Instructions for safe disposal of the product and its packaging	16
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	16
6. Other information	16

Administrative information

1.1. Trade names of the product

STORM Paste
Storm Paste
Storm Paste
STORM PASTE
Storm Pasta
STORM PASTE
STORM PASTE
Storm Pasta
STORM PASTE
Storm Pasta
STORM PASTE

1.2. Authorisation holder

Name and address of the authorisation holder	Name	BASF Nederland B.V.
	Address	Divisio Agro Groningensingel 1 6835 EA Arnhem Netherlands
Authorisation number	NL-0000040-0000	
R4BP 3 asset reference number	NL-0000040-0000	
Date of the authorisation	05/07/2013	
Expiry date of the authorisation	08/03/2023	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	BASF Agro B.V., Arnhem (NL) – Freienbach Branch
Address of the manufacturer	Huobstrasse 3 8808 Pfäffikon SZ Switzerland
Location of manufacturing sites	St. Michaels Industrial Estate, WA8 8TJ Widnes, United Kingdom

1.4. Manufacturer(s) of the active substance(s)

Active substance	34 - Flocoumafen
Name of the manufacturer	BASF Agro B.V., Arnhem (NL) – Freienbach Branch
Address of the manufacturer	Huobstrasse 3 8808 Pfäffikon SZ Switzerland
Location of manufacturing sites	Lower Road, WA8 8 NS Halebank Widnes, United Kingdom

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Flocoumafen	4-hydroxy-3- [(1RS,3RS;1RS,3RS)- 1,2,3,4-tetrahydro-3- [4- (4- trifluoromethylbenzyloxy) phenyl]-1-	Active Substance	90035-08-8	421-960-0	0,005

2.2. Type of formulation

RB - Bait (ready for use)

3. Hazard and precautionary statements

Hazard statements

May damage the unborn child.
May cause damage to organs (blood) through prolonged or repeated exposure .

Precautionary statements

If medical advice is needed, have product container or label at hand.
Keep out of reach of children.
Read carefully and follow all instructions.
Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Wear protective gloves.
IF exposed or concerned: Call a POISON CENTER or doctor/physician.
Store locked up.
Dispose of contents to comply with local requirements.
Get medical advice if you feel unwell.

4. Authorised use(s)

4.1 Use description

Use 1 - House mice and rats - trained professionals - indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides.

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mouse
Development stage: Juveniles to adults

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: Juveniles to adults

Scientific name: *Rattus rattus*
Common name: Black or roof rat
Development stage: Juveniles to adults

Field(s) of use

	Indoor
Application method(s)	<p>Bait application - Pulsed baiting</p> <p>Bait formulation:</p> <ul style="list-style-type: none"> - Ready-to-use bait to be used in tamper resistant bait stations - Covered and protected baiting points (as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations).
Application rate(s) and frequencies	<p>For mice: 15-30 g of bait (1-2 bait units) per bait station; For rats: 45-60 g of bait (3-4 bait units) per bait station. - -</p> <p>Mice: 15-30 g of bait (1-2 bait units) per bait station . If more than one bait station is needed, the minimum distance between bait stations should be of 1-2 meters. Rats: 45-60 g of bait (3-4 bait units) per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters</p> <p>Pulsed baiting technique: Replace eaten bait only after 3 days and then at maximum 7 day intervals. Move any untouched bait to active areas.</p> <p>At each visit, replace eaten bait. Place bait throughout the infested area, in tamper-resistant bait boxes or at covered and protected bait points.</p> <p>If control is not achieved, survey site again to establish reasons and consider re-siting or increasing the number of bait points.</p>
Category(ies) of users	Trained professional
Pack sizes and packaging material	<p>Sachet of paper / plastic laminate (PE and/or PP) containing 15 g bait, further packed in:</p> <p>Plastic container: 3 kg – 10 kg, PP or HDPE or PET, closure: lid of PP or PE</p> <p>Reclosable bag: 3 kg – 10 kg, paper laminate, re-closable flexible bags</p> <p>Lined reclosable cardboard box : 3 kg – 10 kg, re-closable LDPE lined carton</p>

4.1.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.

- Baits must be placed to minimise the exposure to non-target species and children.
 - [When available] Follow the specific instructions for pulsed baiting provided by the applicable code of good practice at national level.
 - Replace eaten bait only after 3 days and then at maximum 7 day intervals.
 - Collect any spilled bait and dead rodents.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

4.1.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- The baits must be securely deposited in a way so as to minimise the risk of consumption by other animals or children. Where possible secure baits so that they cannot be dragged away.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- See general directions for use.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- See general directions for use.

4.2 Use description

Use 2 - House mice and rats- trained professionals- outdoor around buildings

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	<p>Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: Juveniles to adults.</p> <p>Scientific name: <i>Rattus norvegicus</i> Common name: Brown rat Development stage: Juveniles to adults.</p> <p>Scientific name: <i>Rattus rattus</i> Common name: Black or roof rat Development stage: Juveniles to adults.</p>
Field(s) of use	<p>Outdoor</p> <p>outdoor around buildings</p>
Application method(s)	<p>Bait application -</p> <p>Pulsed baiting</p> <p>Bait formulation: -Ready-to-use bait to be used in tamper-resistant bait stations -Covered and protected baiting points(as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations) -Direct application of ready-to-use bait into the burrow – only if authorised; see the corresponding additional elements to be considered in addendum 4</p>
Application rate(s) and frequencies	<p>For mice: 15-30 g of bait (1-2 bait units) per bait station; For rats: 45-60 g of bait (3-4 bait units) per bait station. Bait products: 45-60 g of bait (3-4 bait units) per burrow. - - -</p> <p>Mice: 15-30 g of bait (1-2 bait units) per bait station . If more than one bait station is needed, the minimum distance between bait stations should be of 1-2 meters. Rats: 45-60 g of bait (3-4 bait units) per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters Pulsed baiting technique:</p>

Replace eaten bait only after 3 days and then at maximum 7 day intervals. Move any untouched bait to active areas.
At each visit, replace eaten bait. Place bait throughout the infested area, in tamper-resistant bait boxes or at covered and protected bait points.

If control is not achieved, survey site again to establish reasons and consider re-siting or increasing the number of bait points.

Category(ies) of users

Trained professional

Pack sizes and packaging material

Sachet of paper / plastic laminate (PE and/or PP) containing 15 g bait, further packed in:

Plastic container: 3 kg – 10 kg, PP or HDPE or PET, closure: lid of PP or PE
Reclosable bag: 3 kg – 10 kg, paper laminate, re-closable flexible bags

Lined reclosable cardboard box : 3 kg – 10 kg, re-closable LDPE lined carton

4.2.1 Use-specific instructions for use

- Protect the bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which baits has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- Baits must be placed to minimise the exposure to non-target species and children.

- [When available] Follow the specific instructions for pulsed baiting provided by the applicable code of good practice at national level.
- Replace eaten bait only after 3 days and then at maximum 7 day intervals. Collect any spilled bait and dead rodents.
- Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- See general directions for use

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- See general directions for use

4.3 Use description

Use 3 - House mice - professionals - indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: Juveniles to adults
Field(s) of use	Indoor
Application method(s)	Bait application - Description: - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	15-30 g of bait (1-2 bait units) per bait station - - Place bait throughout the infested area, in tamper-resistant bait boxes. 15-30 g of bait (1-2 bait units) per bait station . If more than one bait station is needed, the minimum distance between bait stations should be of 1-2 meters.
Category(ies) of users	Professional
Pack sizes and packaging material	Sachet of paper / plastic laminate (PE and/or PP) containing 15 g bait, further packed in: Plastic container: 3 kg – 10 kg, PP or HDPE or PET, closure: lid of PP or PE Reclosable bag: 3 kg – 10 kg, paper laminate, re-closable flexible bags Lined reclosable cardboard box : 3 kg – 10 kg, re-closable LDPE lined carton

4.3.1 Use-specific instructions for use

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.3.2 Use-specific risk mitigation measures

- see general directions for use.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- See general directions for use.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- See general directions for use.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- See general directions for use.

4.4 Use description

Use 4 - Rats - professionals - indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides.

Target organism(s) (including development stage)

Scientific name: Rattus norvegicus
Common name: Brown rat
Development stage: Juveniles to adults

Scientific name: Rattus rattus
Common name: Black or roof rat
Development stage: Juveniles to adults

Field(s) of use

	Indoor
Application method(s)	Bait application - Description: - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	45-60 g of bait (3-4 bait units) per bait station. - - - Place bait throughout the infested area, in tamper-resistant bait boxes. 45-60 g of bait (3-4 bait units) per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters.
Category(ies) of users	Professional
Pack sizes and packaging material	Sachet of paper / plastic laminate (PE and/or PP) containing 15 g bait, further packed in: Plastic container: 3 kg – 10 kg, PP or HDPE or PET, closure: lid of PP or PE Reclosable bag: 3 kg – 10 kg, paper laminate, re-closable flexible bags Lined reclosable cardboard box : 3 kg – 10 kg, re-closable LDPE lined carton

4.4.1 Use-specific instructions for use

<p>- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</p> <p>- [When available] Follow any additional instructions provided by the relevant code of best practice.</p>
--

4.4.2 Use-specific risk mitigation measures

- see general directions for use

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- See general directions for use.

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- See general directions for use.

4.5 Use description

Use 5 - House mice and rats- professionals- outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides.

Target organism(s) (including development stage)

Scientific name: *Mus musculus*
Common name: House mouse
Development stage: Juveniles to adults.

Scientific name: *Rattus norvegicus*
Common name: Brown rat
Development stage: Juveniles to adults.

Scientific name: *Rattus rattus*
Common name: Black or roof rat
Development stage: Juveniles to adults.

Field(s) of use	Outdoor outdoor around buildings
Application method(s)	Bait application - Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequencies	For mice: 15-30 g of bait (1-2 bait units) per bait station; For rats: 45-60 g of bait (3-4 bait units) per bait station. - - - Place bait throughout the infested area, in tamper-resistant bait boxes. Mice: 15-30 g of bait (1-2 bait units) per bait station . If more than one bait station is needed, the minimum distance between bait stations should be of 1-2 meters. Rats: 45-60 g of bait (3-4 bait units) per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5-10 meters.
Category(ies) of users	Professional
Pack sizes and packaging material	Sachet of paper / plastic laminate (PE and/or PP) containing 15 g bait, further packed in: Plastic container: 3 kg – 10 kg, PP or HDPE or PET, closure: lid of PP or PE Reclosable bag: 3 kg – 10 kg, paper laminate, re-closable flexible bags Lined reclosable cardboard box : 3 kg – 10 kg, re-closable LDPE lined carton

4.5.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited for mice at least every 2 to 3 days at the beginning and for rats only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- *[When available]* Follow any additional instructions provided by the relevant code of best practice.

4.5.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.
- Do not wash the bait stations with water between applications.

- The product information (i.e. label and/or leaflet) shall clearly show that:

- The product shall not be supplied to the general public (e.g. "for professionals only").
- The product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- Users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations")

-Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

- The baits must be securely deposited in a way so as to minimise the risk of consumption by other animals or children. Where possible secure baits so that they cannot be dragged away.

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- See general directions for use

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- See general directions for use

5. General directions for use

5.1. Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- *[If national policy or legislation require it]* When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Bait in sachets: do not open the sachets containing the bait

5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
 - To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment in line with the recommendations provided by the relevant code of practice.
 - Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
 - Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
the product shall not be supplied to the general public (e.g. "for professionals only").
the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
 - Do not wash the bait stations with water between applications.
 - Dispose dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting.

If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [insert national phone number] "

- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements:

Do not reuse the container for any other purpose.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.

- Store in places prevented from the access of children, birds, pets and farm animals.

- Shelf life: The shelf-life of the product is 5 years. Product is not to be stored at temperatures >30 °C.

6. Other information

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.

- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

- This product contains a bittering agent and a dye.