

# Summary of product characteristics for a biocidal product

**Product name:** FAAR BLE

**Product type(s):** PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

**Authorisation number:** FR-2014-0020

**R4BP 3 asset reference number:** FR-0001028-0000

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	4
2. Product composition and formulation	4
2.1. Qualitative and quantitative information on the composition of the biocidal product	4
2.2. Type of formulation	4
3. Hazard and precautionary statements	5
4. Authorised use(s)	5
5. General directions for use	18
5.1. Instructions for use	18
5.2. Risk mitigation measures	18
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	19
5.4. Instructions for safe disposal of the product and its packaging	19
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	19
6. Other information	20

## Administrative information

### 1.1. Trade names of the product

FAAR BLE  
BROFAR  
PARADE 51 B  
BROMABLE  
BROMAGRAIN RAT 50  
BROMAGRAIN RONGEUR 50  
SUPP' BROMAGRAIN 50  
STOP BROMAGRAIN 50  
BROMASEED RAT 50  
BROMASEED RODENT 50  
STOP BROMASEED 50

### 1.2. Authorisation holder

**Name and address of the  
authorisation holder**

Name

TRIPLAN SA

Address

BP 258 LA POSTE FRANCAISE AD500 ANDORRA LA VELLA  
France

**Authorisation number**

FR-2014-0020

**R4BP 3 asset reference  
number**

FR-0001028-0000

**Date of the authorisation**

18/06/2013

**Expiry date of the  
authorisation**

01/07/2024

### 1.3. Manufacturer(s) of the biocidal products

**Name of the manufacturer**

SOFAR France

**Address of the manufacturer**

ZA DU DREVERS BP02 29190 PLEYBEN France

**Location of manufacturing sites**

ZA DU DREVERS BP02 29190 PLEYBEN France

**Name of the manufacturer**

AEDES PROTECTA

**Address of the manufacturer**

75 rue d'Orgemont 95210 SAINT-GRATIEN France

**Location of manufacturing sites**

LIEU DIT DOUILLAC 81310 PARISOT France

**Name of the manufacturer**

IRIS

**Address of the manufacturer**

1126A, Avenue du Moulinas, Route de Saint-Privat 30340 SALINDRES France

**Location of manufacturing sites**

1126A, Avenue du Moulinas, Route de Saint-Privat 30340 SALINDRES France

**Name of the manufacturer**

HDA

**Address of the manufacturer**

ZA LA CHARME MENETROL 63200 RIOM France

**Location of manufacturing sites**

ZA LA CHARME MENETROL 63200 RIOM France

**Name of the manufacturer**

NOXIMA

**Address of the manufacturer**

CARREFOUR JEAN MONNET - LACROIX-SAINT OUEN 60201 COMPIEGNE France

**Location of manufacturing sites**

CARREFOUR JEAN MONNET - LACROIX SAINT OUEN 60201 COMPIEGNE France

**Name of the manufacturer**

INDUSTRIAL CHIMICA SRL

**Address of the manufacturer**

VIA SORGAGLIA 40 35020 ARRE France

**Location of manufacturing sites**

VIA SORGAGLIA 40 35020 ARRE Italy

<b>Name of the manufacturer</b>	RATOUCY SAS
<b>Address of the manufacturer</b>	29 avenue de la Forêt - BP145 89303 JOIGNY CEDEX France
<b>Location of manufacturing sites</b>	29 AVENUE DE LA FORET - LOOZE - BP145 89303 JOIGNY CEDEX France

<b>Name of the manufacturer</b>	FARMAVIT OOD
<b>Address of the manufacturer</b>	Bul Tsar Boris III, n° 62, Office n° 1 1612 SOFIA Bulgaria
<b>Location of manufacturing sites</b>	Indulstriana str 2. - Pleven District 5960 GULIANTSI Bulgaria

<b>Name of the manufacturer</b>	SALOMEZ
<b>Address of the manufacturer</b>	ZI AV. du Général de GAULLE 89130 TOUCY France
<b>Location of manufacturing sites</b>	ZI AV. du Général de GAULLE 89130 TOUCY France

<b>Name of the manufacturer</b>	FARMA-CHEM SA
<b>Address of the manufacturer</b>	Industrial Area of Sindos P.O. BOC 1026 Block 53 Zone C 57022 THESSALONIKI Greece
<b>Location of manufacturing sites</b>	Industrial Area of Sindos P.O. BOC 1026 Block 53 Zone C 57022 THESSALONIKI Greece

<b>Name of the manufacturer</b>	AGROZA AGROTECHNICAL LTD
<b>Address of the manufacturer</b>	AITOLIAS 2 & EK. VENIZELOU 158 16341 ATHENS Greece
<b>Location of manufacturing sites</b>	AITOLIAS 2 & EK. VENIZELOU 158 16341 ATHENS Greece

<b>Name of the manufacturer</b>	AGROLOGY SA
<b>Address of the manufacturer</b>	P.O. BOX 1089, Industrial Area of Thessaloniki 57022 THESSALONIKI Greece
<b>Location of manufacturing sites</b>	P.O. BOX 1089, Industrial Area of Thessaloniki 57022 THESSALONIKI France

#### 1.4. Manufacturer(s) of the active substance(s)

Active substance	12 - Bromadiolone
Name of the manufacturer	PM TEZZA SRL
Address of the manufacturer	Via tre ponti 22 37050 S. MARIA DI ZEVIO (VR) Italy
Location of manufacturing sites	Via tre ponti 22 37050 S. MARIA DI ZEVIO (VR) Italy

## 2. Product composition and formulation

#### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[3-(4'-Bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	Active Substance	28772-56-7	249-205-9	0,005

#### 2.2. Type of formulation

Grain bait, ready to use
--------------------------

### 3. Hazard and precautionary statements

#### Hazard statements

May damage the unborn child.

Causes damage to organs through prolonged or repeated exposure (blood).

#### Precautionary statements

Obtain special instructions before use.

Do not handle until all safety precautions have been read and understood.

Do not breathe dust.

Wash hands thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wear protective gloves.

IF exposed or concerned: Get medical advice.

Get medical attention if you feel unwell.

Store locked up.

Dispose of container to in accordance with national regulations.

### 4. Authorised use(s)

#### 4.1 Use description

##### Use 1 - HOUSE MICE AND/OR RATS - TRAINED PROFESSIONALS - INDOOR

#### Product type

PT14 - Rodenticides (Pest control)

#### Where relevant, an exact description of the authorised use

Not relevant for rodenticides

#### Target organism(s) (including development stage)

Scientific name: Rattus norvegicus  
Common name: Brown rat  
Development stage: Adults

Scientific name: Rattus norvegicus  
Common name: Brown rat  
Development stage: Juveniles

Scientific name: Rattus rattus  
Common name: Roof rat  
Development stage: Adults

Scientific name: Rattus rattus  
Common name: Roof rat  
Development stage: Juveniles

Scientific name: Mus musculus

Common name: House mouse  
Development stage: Adults

Scientific name: Mus musculus  
Common name: House mouse  
Development stage: Juveniles

#### Field(s) of use

Indoor

Indoor

#### Application method(s)

Method: Bait application

Detailed description:

- Ready-to-use bait to be used in tamper-resistant bait stations
- [Covered and protected baiting points]

#### Application rate(s) and frequencies

Application Rate: Rats: 200g of bait per bait station separated 5 to 10 meters Mice : 40g of bait per bait station separated by 1 to 2 meters

Dilution (%): 0

Number and timing of application:

Bait products:

Rats (*Rattus norvegicus* & *Rattus rattus*):

- High infestation: 200 g of bait per bait station separated by 5 meters.
- Low infestation: 200 g of bait per bait station separated by 10 meters.

House mice (*Mus musculus*):

- High infestation: 40 g of bait per bait station separated by 1 meter..
- Low infestation: 40 g of bait per bait station separated by 2 meters.

#### Category(ies) of users

Trained professional

#### Pack sizes and packaging material

Minimum pack size of 3 kg.

(In France only : minimum pack size of 5 kg)

Package is restricted to separately packed bags with a maximum of 10 kg per packed bag for loose grain baits.

PE/PP sachets (10-20-25-30-40-50-60-90-100-200g) packed in:

- Bags (paper/PE/PP) (5-10-15-20-25-30 kg)
- Bucket (PE) (5-10-15-18-20-25-30kg)
- Carton box (carton) (5-10-12-15-20-25-30-50 kg)
- Bait box (PET/PP/PE/PVC)
- Metal box (5-10-15-20-25kg)

Loose baits packed in:

- Bags (paper/PE/PP) (5-10 kg)
- Bucket (PE/PP) (5-10 kg)
- Carton box (carton) (5-10kg)
- Sachet PE or PP (100-200-300-400-500-600-700-800-900-1000g) packed in carton box (5-10 kg)
- Metal box (5-10kg)

#### 4.1.1 Use-specific instructions for use



- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

#### 4.1.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use the product in pulsed baiting treatments.

#### 4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

### 4.2 Use description

**Use 2 - HOUSE MICE AND/OR RATS - TRAINED PROFESSIONALS - OUTDOOR AROUND BUILDINGS**

**Product type**

PT14 - Rodenticides (Pest control)

**Where relevant, an exact description of the authorised use**

Not relevant for rodenticides

**Target organism(s) (including development stage)**

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: Adults

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: Juveniles

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: Adults

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: Juveniles

Scientific name: *Mus musculus*  
Common name: House mouse  
Development stage: Adults

Scientific name: *Mus musculus*  
Common name: House mouse  
Development stage: Juveniles

**Field(s) of use**

Outdoor

Outdoor around buildings

**Application method(s)**

Method: Bait application

Detailed description:

- Ready-to-use bait to be used in tamper-resistant bait stations.
- [Covered and protected baiting points]

**Application rate(s) and frequencies**

Application Rate: Rats 200 g of bait per bait station separated by 5 to 10 meters Mice : 40 g of bait per bait station separated by 1 to 2 meters

Dilution (%): 0

Number and timing of application:

Bait products:

Rats (*Rattus norvegicus* & *Rattus rattus*):

- High infestation: 200 g of bait per bait station separated by 5 meters.

- Low infestation: 200 g of bait per bait station separated by 10 meters.

House mice (*Mus musculus*):

- High infestation: 40 g of bait per bait station separated by 1 meter.

	<ul style="list-style-type: none"> <li>- Low infestation: 40 g of bait station separated by 2 meters.</li> </ul>
<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	<p>Minimum pack size of 3 kg. (In France only : minimum pack size of 5 kg) Package is restricted to separately packed bags with a maximum of 10 kg per packed bag for loose grain baits. PE/PP sachets (10-20-25-30-40-50-60-90-100-200g) packed in:</p> <ul style="list-style-type: none"> <li>- Bags (paper/PE) (5-10-15-20-25-30 kg)</li> <li>- Bucket (PE) (5-10-15-18-20-25-30kg)</li> <li>- Carton box (carton) (5-10-12-15-20-25-30-50 kg)</li> <li>- Bait box (PET/PP/PE/PVC)</li> <li>- Metal box (5-10-15-20-25kg)</li> </ul> <p>Loose baits packed in:</p> <ul style="list-style-type: none"> <li>- Bags (paper/PE/PP) (5-10 kg)</li> <li>- Bucket (PE) (5-10 kg)</li> <li>- Carton box (carton) (5-10 kg)</li> <li>- Sachet PE or PP (100-200-300-400-500-600-700-800-900-1000g) packed in carton box (5-10 kg)</li> <li>- Metal box (5-10kg)</li> </ul>

#### 4.2.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- [For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species].

#### 4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

### 4.3 Use description

#### Use 3 - RATS - TRAINED PROFESSIONALS - OUTDOOR OPEN AREAS & WASTE DUMPS

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Adults  Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles  Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults  Scientific name: Rattus rattus Common name: Roof rat Development stage: Juveniles
<b>Field(s) of use</b>	Outdoor  Other  Outdoor open areas Outdoor waste dumps
<b>Application method(s)</b>	Method: Bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations. - [Covered and protected baiting points]
<b>Application rate(s) and frequencies</b>	Application Rate: 200g of bait per bait station separated by 5 to 10 meters. Dilution (%): Number and timing of application: Bait products:

	<p>Rats (<i>Rattus norvegicus</i> &amp; <i>Rattus rattus</i>):</p> <ul style="list-style-type: none"> <li>- High infestation: 200 g of bait per bait station separated by 5 meters.</li> <li>- Low infestation: 200 g of bait per bait station separated by 10 meters.</li> </ul>
<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	<p>Minimum pack size of 3 kg. (In France only : minimum pack size of 5 kg) Package is restricted to separately packed bags with a maximum of 10 kg per packed bag for loose grain baits. PE/PP sachets (10-20-25-30-40-50-60-90-100-200g) packed in:</p> <ul style="list-style-type: none"> <li>- Bags (paper/PE) (5-10-15-20-25-30 kg)</li> <li>- Bucket (PE) (5-10-15-18-20-25-30kg)</li> <li>- Carton box (carton) (5-10-12-15-20-25-30-50 kg)</li> <li>- Bait box (PET/PP/PE/PVC)</li> <li>- Metal box (5-10-15-20-25kg)</li> </ul> <p>Loose baits packed in:</p> <ul style="list-style-type: none"> <li>- Bags (paper/PE/PP) (5-10 kg)</li> <li>- Bucket (PE) (5-10 kg)</li> <li>- Carton box (carton) (5-10 kg)</li> <li>- Sachet PE or PP (100-200-300-400-500-600-700-800-900-1000g) packed in carton box (5-10 kg)</li> <li>- Metal box (5-10kg)</li> </ul>

#### 4.3.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- [For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species].

#### 4.3.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not apply this product directly in the burrows.

#### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

--

#### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

--

#### 4.4 Use description

##### Use 4 - (not relevant in France) HOUSE MICE - PROFESSIONALS - INDOOR

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: House mouse Development stage: Adults  Scientific name: Mus musculus Common name: House mouse Development stage: Juveniles
Field(s) of use	Indoor  Indoor
Application method(s)	Method: Bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations - [Covered and protected baiting points]
Application rate(s) and frequencies	Application Rate: 40g of bait per bait station Dilution (%): Number and timing of application: House mice (Mus musculus): 40 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 1 meter.
Category(ies) of users	Professional
Pack sizes and packaging material	Minimum pack size of 3 kg. (In France only : minimum pack size of 5 kg) Package is restricted to separately packed bags with a maximum of 10 kg per packed bag for loose grain baits. PE/PP sachets (10-20-25-30-40-50-60-90-100-200g) packed in: - Bags (paper/PE) (5-10-15-20-25-30 kg) - Bucket (PE) (5-10-15-18-20-25-30kg) - Carton box (carton) (5-10-12-15-20-25-30-50 kg) - Bait box (PET/PP/PE/PVC) -Metal box (5-10-15-20-25kg)

Loose baits packed in:

- Bags (paper/PE/PP) (5-10 kg)
- Bucket (PE) (5-10 kg)
- Carton box (carton) (5-10 kg)
- Sachet PE or PP (100-200-300-400-500-600-700-800-900-1000g) packed in carton box (5-10 kg)

- Metal box (5-10kg)

#### 4.4.1 Use-specific instructions for use

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.  
- [When available] Follow any additional instructions provided by the relevant code of best practice.

#### 4.4.2 Use-specific risk mitigation measures

#### 4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

#### 4.5 Use description

Use 5 - (not relevant in France) - RATS - PROFESSIONALS - INDOOR

Product type

PT14 - Rodenticides (Pest control)

**Where relevant, an exact description of the authorised use**

**Target organism(s) (including development stage)**

Not relevant for rodenticides

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: Adults

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: Juveniles

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: Adults

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: Juveniles

**Field(s) of use**

Indoor

Indoor

**Application method(s)**

Method: Bait application

Detailed description:

- Ready-to-use bait to be used in tamper-resistant bait stations
- [Covered and protected baiting points]

**Application rate(s) and frequencies**

Application Rate: Rats : 200 g of bait per bait station.

Dilution (%):

Number and timing of application:

Rats (*Rattus norvegicus* & *Rattus rattus*): 200 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 4 meters.

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

Minimum pack size of 3 kg.

(In France only : minimum pack size of 5 kg)

Package is restricted to separately packed bags with a maximum of 10 kg per packed bag for loose grain baits.

PE/PP sachets (10-20-25-30-40-50-60-90-100-200g) packed in:

- Bags (paper/PE) (5-10-15-20-25-30 kg)
- Bucket (PE) (5-10-15-18-20-25-30kg)
- Carton box (carton) (5-10-12-15-20-25-30-50 kg)
- Bait box (PET/PP/PE/PVC)
- Metal box (5-10-15-20-25kg)

Loose baits packed in:

- Bags (paper/PE/PP) (5-10 kg)
- Bucket (PE) (5-10 kg)
- Carton box (carton) (5-10 kg)
- Sachet PE or PP (100-200-300-400-500-600-700-800-900-1000g) packed in carton box (5-10 kg)
- Metal box (5-10kg)



#### 4.5.1 Use-specific instructions for use

- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

#### 4.5.2 Use-specific risk mitigation measures

#### 4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

#### 4.6 Use description

##### Use 6 - (not relevant in France) - HOUSE MICE AND/OR RATS - PROFESSIONALS - OUTDOOR AROUND BUILDINGS

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	<div>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Adults</div> <div>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles</div> <div>Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults</div> <div>Scientific name: Rattus rattus Common name: Roof rat Development stage: Juveniles</div>

Scientific name: Mus musculus  
Common name: House mouse  
Development stage: Adults

Scientific name: Mus musculus  
Common name: House mouse  
Development stage: Juveniles

## Field(s) of use

Outdoor

Outdoor around buildings

## Application method(s)

Method: Bait application

Detailed description:

- Ready-to-use bait to be used in tamper-resistant bait stations
- [Covered and protected baiting points]

## Application rate(s) and frequencies

Application Rate: Rats : 200g of bait per bait station Mice :40 g of bait per bait station

Dilution (%):

Number and timing of application:

Rats (*Rattus norvegicus* & *Rattus rattus*): 200 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 4 meters.

House mice (*Mus musculus*): 40 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 1 meter.

## Category(ies) of users

Professional

## Pack sizes and packaging material

Minimum pack size of 3 kg.

(In France only : minimum pack size of 5 kg)

Package is restricted to separately packed bags with a maximum of 10 kg per packed bag for loose grain baits.

PE/PP sachets (10-20-25-30-40-50-60-90-100-200g) packed in:

- Bags (paper/PE) (5-10-15-20-25-30 kg)
- Bucket (PE) (5-10-15-18-20-25-30kg)
- Carton box (carton) (5-10-12-15-20-25-30-50 kg)
- Bait box (PET/PP/PE/PVC)
- Metal box (5-10-15-20-25kg)

Loose baits packed in:

- Bags (paper/PE/PP) (5-10 kg)

- Bucket (PE) (5-10 kg)
- Carton box (carton) (5-10 kg)
- Sachet PE or PP (100-200-300-400-500-600-700-800-900-1000g) packed in carton box (5-10 kg)
- Metal box (5-10kg)

#### 4.6.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

#### 4.6.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

#### 4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

## 5. General directions for use

### 5.1. Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
  - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
  - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
  - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
  - The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
  - Where possible, bait stations must be fixed to the ground or other structures.
  - Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
  - [If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
  - Bait should be secured so that it cannot be dragged away from the bait station.
  - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
  - Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
  - Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- FOR TRAINED PROFESSIONAL ONLY- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
  - If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- FOR PROFESSIONNALS ONLY Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- FOR PROFESSIONNALS ONLY Remove the remaining bait or the bait stations at the end of the treatment period.
- Instructions for use that are "bait-specific":
- Bait in sachets: [For non-emptiable sachets - Do not open the sachets containing the bait].
  - Loose pellets-granules, grains: Place the bait in the baiting point by using a dosage devise. Specify the methods to minimise dust (e.g. wet wiping).
  - Loose pellets-granules, grains: Decanting is to be avoided. In case decanting cannot be avoided, an RPE of APF 10 has to be used.

### 5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- FOR TRAINED PROFESSIONAL ONLY Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- FOR TRAINED PROFESSIONAL ONLY Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements
- FOR PROFESSIONAL ONLY To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week)..
- FOR PROFESSIONAL ONLY Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- FOR PROFESSIONAL ONLY. The product information (i.e. label and/or leaflet) shall clearly show that:  
the product shall not be supplied to the general public (e.g. "for professionals only").  
the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").  
users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- FOR PROFESSIONAL ONLY Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

### 5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:  
- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label [insert country specific information]. Contact a veterinary surgeon in case of ingestion by a pet [insert country specific information]
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre"
- Hazardous to wildlife.

### 5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

### 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Shelf-life : 2 year
- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.

## 6. Other information

- In France only : The authorisation holder has to monitor the resistance phenomenon of rodent populations toward the active substance bromadiolone. Results of the resistance monitoring must be submitted at the renewal of the product
- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.