

Summary of product characteristics for a biocidal product

Product name: JADE BLOCK

Product type(s): PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

Authorisation number: UK-2013-0711

R4BP 3 asset reference number: UK-0001718-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	1
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
4.1.1 Use-specific instructions for use	4
4.1.2 Use-specific risk mitigation measures	4
4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	4
4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	5
4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	5
4.2.1 Use-specific instructions for use	6
4.2.2 Use-specific risk mitigation measures	6
4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	6
4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	6
4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	7
4.3.1 Use-specific instructions for use	8
4.3.2 Use-specific risk mitigation measures	8
4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	9
4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	9
4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	9
4.4.1 Use-specific instructions for use	10
4.4.2 Use-specific risk mitigation measures	11

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	11
4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	11
4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	11
4.5.1 Use-specific instructions for use	13
4.5.2 Use-specific risk mitigation measures	13
4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	13
4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	14
4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	14
4.6.1 Use-specific instructions for use	15
4.6.2 Use-specific risk mitigation measures	16
4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	16
4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	16
4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	16
4.7.1 Use-specific instructions for use	18
4.7.2 Use-specific risk mitigation measures	18
4.7.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	18
4.7.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	18
4.7.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	19
5. General directions for use	19
5.1. Instructions for use	19
5.2. Risk mitigation measures	21
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	21
5.4. Instructions for safe disposal of the product and its packaging	21
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	22
6. Other information	22

Administrative information

1.1. Trade names of the product

JADE BLOCK
ENDORATS RAT KILLER BLOCKS

1.2. Authorisation holder

Name and address of the authorisation holder	Name	LODI S.A.S.
	Address	Parc d'Activités des Quatre Routes 35390 Grand Fougeray France
Authorisation number	UK-2013-0711	
R4BP 3 asset reference number	UK-0001718-0000	
Date of the authorisation	30/09/2012	
Expiry date of the authorisation	25/04/2023	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	Lodi SAS
Address of the manufacturer	Parc d'activités des quatre routes 35390 Grand Fougeray France
Location of manufacturing sites	Lodi SAS, Parc d'activités des quatre routes 35390 Grand Fougeray France
	Parc d'activités du Pays du Grand Fougeray Espace Nord 24 et 26 rue des Pionniers 35390 GRAND FOUGERAY France

1.4. Manufacturer(s) of the active substance(s)

Active substance	12 - Bromadiolone
Name of the manufacturer	PelGar International Ltd.
Address of the manufacturer	Unit 13 Newman Lane GU34 2QR Alton United Kingdom
Location of manufacturing sites	Via Tre Ponti 37050 S. Maria di Zevio Italy

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[3-(4'-Bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	Active Substance	28772-56-7	249-205-9	0.005

2.2. Type of formulation

RB - Bait (ready for use)

3. Hazard and precautionary statements

Hazard statements	<p>May damage the unborn child.</p> <p>Causes damage to organs (blood) through prolonged or repeated exposure .</p>
Precautionary statements	<p>Obtain special instructions before use.</p> <p>Do not handle until all safety precautions have been read and understood.</p> <p>Do not breathe dust.</p> <p>Wash hands thoroughly after handling.</p> <p>Do no eat, drink or smoke when using this product.</p> <p>Wear protective gloves.</p>

IF exposed or concerned: Get medical advice.

Get medical advice if you feel unwell.

Store locked up.

Dispose of contents to in accordance with the national regulation.

4. Authorised use(s)

4.1 Use description

Use 1 - House Mouse - Professionals - Indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Mus musculus-House mouse-Adults Mus musculus-House mouse-Juveniles
Field(s) of use	Indoor Indoor
Application method(s)	Bait application - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	20-30g bait in bait station - - 20-30 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be 5 or 10 meters depending of the infestation.
Category(ies) of users	Professional
Pack sizes and packaging material	Minimum pack size 2.5kg Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30 Packaging material and size: Bucket (PE or PP) :

20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20), 6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)

30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)

Cardboard box (with inner liner in PE if bait is unwrapped)

20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20), 6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)

30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)

Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:

- 2.5 kg (125*20 g) or (84*30 g)
- 3 kg(150*20g) or (100*30g)
- 4 kg (200*20g) or (134*30 g)
- 5 kg (250*20g) or (167*30g)

4.1.1 Use-specific instructions for use

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. [When available] Follow any additional instructions provided by the relevant code of best practice

4.1.2 Use-specific risk mitigation measures

None

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.2 Use description

Use 2 - Rat - Professionals- Indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticide
Target organism(s) (including development stage)	Rattus norvegicus-Brown rat-Adults Rattus norvegicus-Brown rat-Juveniles
Field(s) of use	Indoor Indoor
Application method(s)	Bait application - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	60-100 g of bait per bait station - 0 - 60-100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters (high infestation) and 10 meters (low infestation).
Category(ies) of users	Professional
Pack sizes and packaging material	Minimum pack size 2.5kg Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30 Packaging material and size:

Bucket (PE or PP) :

20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20), 6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)

30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)

Cardboard box (with inner liner in PE if bait is unwrapped)

20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20), 6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)

30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)

Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:

- 2.5 kg (125*20 g) or (84*30 g)
- 3 kg(150*20g) or (100*30g)
- 4 kg (200*20g) or (134*30 g)
- 5 kg (250*20g) or (167*30g)

4.2.1 Use-specific instructions for use

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

[When available] Follow any additional instructions provided by the relevant code of best practice

4.2.2 Use-specific risk mitigation measures

None

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.3 Use description

Use 3 - House mice and rats - Professionals -Outdoor around buildings

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Mus musculus-House mouse-Adults Mus musculus-House mouse-Juveniles Rattus norvegicus-Brown rat-Adults Rattus norvegicus-Brown rat-Juveniles
Field(s) of use	Outdoor Other Outdoor around buildings
Application method(s)	Bait application - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	20-30 g (for mice) and 60-100 g for rats - 0 - 20-30 g (for mice) and 60-100 g for rats in tamper-resistant bait stations. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters (high infestation) and 10 meters (low infestation)
Category(ies) of users	Professional

Pack sizes and packaging material

Minimum pack size 2.5kg
Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30
Packaging material and size:
Bucket (PE or PP) :
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Cardboard box (with inner liner in PE if bait is unwrapped)
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:
- 2.5 kg (125*20 g) or (84*30 g)
- 3 kg(150*20g) or (100*30g)
- 4 kg (200*20g) or (134*30 g)
- 5 kg (250*20g) or (167*30g)

4.3.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.3.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.4 Use description

Use 4 - House mice and/or rats - Trained professionals - Indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Mus musculus-House mouse-Adults Mus musculus-House mouse-Juveniles Rattus norvegicus-Brown rat-Adults Rattus norvegicus-Brown rat-Juveniles
Field(s) of use	Indoor Indoors
Application method(s)	Bait application - Ready-to-use bait to be used in covered bait points or in tamper-resistant bait stations
Application rate(s) and frequencies	20-30g (mice) / 60-100 g (rats) - 0 - 20-30g (mice) / 60-100 g (rats) Mice Low infestation : 20-30 g every 10 metres

High infestation : 20-30 g every 5 metres
Rats
Low infestation : 60-100 g every 10 metres
High infestation : 60-100 g every 5 metres

Permanent baiting –
Low infestation : 20-30 g every 10 metres
High infestation : 20-30 g every 5 metres
Rats
Low infestation : 60-100 g every 10 metres
High infestation : 60-100 g every 5 metres

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size 2.5kg
Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30
Packaging material and size:
Bucket (PE or PP) :
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Cardboard box (with inner liner in PE if bait is unwrapped)
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:
- 2.5 kg (125*20 g) or (84*30 g)
- 3 kg(150*20g) or (100*30g)
- 4 kg (200*20g) or (134*30 g)
- 5 kg (250*20g) or (167*30g)

4.4.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- For permanent baiting - Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.
[When available]

4.4.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.

The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

- Do not use this product in pulsed baiting treatments.

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.5 Use description

Use 5 - House mice and/or rats - Trained professionals - Outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Mus musculus-House mouse-Adults
Mus musculus-House mouse-Juveniles
Rattus norvegicus-Brown rat-Adults
Rattus norvegicus-Brown rat-Juveniles

Field(s) of use

Outdoor
Other
Outdoor around buildings

Application method(s)

Bait application -
Ready-to-use bait to be used in covered bait points or in tamper-resistant bait stations or direct application of ready-to-use bait into the burrow

Application rate(s) and frequencies

20-30g (mice) / 60-100 g (rats) - 0 -

20-30g (mice) / 60-100 g (rats)
Mice
Low infestation : 20-30 g every 10 metres
High infestation : 20-30 g every 5 metres
Rats
Low infestation : 60-100 g every 10 metres
High infestation : 60-100 g every 5 metres
- In burrows: 60-100g of bait per burrow.

Permanent baiting –
Low infestation : 20-30 g every 10 metres
High infestation : 20-30 g every 5 metres
Rats
Low infestation : 60-100 g every 10 metres
High infestation : 60-100 g every 5 metres

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size 2.5kg
Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30
Packaging material and size:
Bucket (PE or PP) :
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),

6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
 30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 kg (200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (317*30), 10 kg (334*30)
 Cardboard box (with inner liner in PE if bait is unwrapped)
 20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
 6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
 30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 kg (200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (317*30), 10 kg (334*30)
 Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:
 - 2.5 kg (125*20 g) or (84*30 g)
 - 3 kg(150*20g) or (100*30g)
 - 4 kg (200*20g) or (134*30 g)
 - 5 kg (250*20g) or (167*30g)

4.5.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- For permanent baiting –Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population. [When available] Follow any additional instructions provided by the relevant code of best practice.
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. [When available] Follow any additional instructions provided by the relevant code of best practice.
- When used in burrows: Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

4.5.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
 - Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
 - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice
 - Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.
- Do not use this product in pulsed baiting treatments

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.6 Use description

Use 6 - Rats - Trained professionals - Outdoor open areas & waste dumps

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticide
Target organism(s) (including development stage)	Rattus norvegicus-Brown rat-Adults Rattus norvegicus-Brown rat-Juveniles
Field(s) of use	Outdoor Other Outdoor open areas & waste dumps
Application method(s)	Bait application - Ready-to-use bait to be used in covered bait points or in tamper-resistant bait stations
Application rate(s) and frequencies	60-100 g of bait per bait station - 0 - 60-100 g of bait per bait station - - - High infestation: 60-100 g of bait per bait station every 5 meters - Low infestation: 60-100 g of bait per bait station every 10 meters

- In burrows: 60-100g of bait per burrow.

Permanent baiting –

Low infestation : 60-100 g every 10 metres

High infestation : 60-100 g every 5 metres

Category(ies) of users

Trained professional

Pack sizes and packaging material

Minimum pack size 2.5kg
Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30
Packaging material and size:
Bucket (PE or PP) :
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Cardboard box (with inner liner in PE if bait is unwrapped)
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:
- 2.5 kg (125*20 g) or (84*30 g)
- 3 kg(150*20g) or (100*30g)
- 4 kg (200*20g) or (134*30 g)
- 5 kg (250*20g) or (167*30g)

4.6.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- For permanent baiting - Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- [For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species].
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- When used in burrows: Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

4.6.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.
- Do not use this product for pulsed baiting.

4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.7 Use description

Use 7 - Rats - Trained professionals - Sewers

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticide
Target organism(s) (including development stage)	Rattus norvegicus-Brown rat-Adults Rattus norvegicus-Brown rat-Juveniles

Field(s) of use	Other Sewers
Application method(s)	Bait application - Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water.
Application rate(s) and frequencies	200-300 g - 0 - 200-300 g every 30-50 meters Bait products: - High infestation: (300) grams per manhole. - Low infestation: (200) grams per manhole. Permanent baiting – Low infestation : 20-30 g every 10 metres High infestation : 20-30 g every 5 metres Rats Low infestation : 60-100 g every 10 metres High infestation : 60-100 g every 5 metres
Category(ies) of users	Trained professional
Pack sizes and packaging material	Minimum pack size 2.5kg Grams of bait (individually wrapped in PE or PP sachet or unwrapped) : 20-30 Packaging material and size: Bucket (PE or PP) : 20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20), 6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20) 30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg

(250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Cardboard box (with inner liner in PE if bait is unwrapped)
20g: 2.5 kg (125*20), 3 kg(150*20), 3.5 kg (175*20), 4 kg (200*20), 4.5 kg (225*20), 5 kg (250 *20), 5.5 kg (275*20), 6 kg (300*20),
6.5 kg (325*20) , 7 kg (350*20), 7.5 kg (375*20), 8 kg (400*20), 8.5 kg (425*20), 9 kg (450*20), 9.5 kg (475*20), 10 kg (500*20)
30g: 2.5 kg (84*30), 3 kg (100*30), 3.5 kg (117*30), 4 kg (134*30), 4.5 kg (150*30), 5 kg (167*30), 5.5 kg (184*30), 6 k g(200*30), 6.5 kg (217*30), 7 kg (234*30), 7.5 kg (250*30), 8 kg (267*30), 8.5 kg (284*30), 9 kg (300*30), 9.5 kg (347*30), 10 kg (334*30)
Pre-baited station (PP, PVC,PS) in cardboard box containing each 1 block of 20 or 30 g:
- 2.5 kg (125*20 g) or (84*30 g)
- 3 kg(150*20g) or (100*30g)
- 4 kg (200*20g) or (134*30 g)
- 5 kg (250*20g) or (167*30g)

4.7.1 Use-specific instructions for use

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- For permanent baiting –Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population. [When available] Follow any additional instructions provided by the relevant code of best practice.

4.7.2 Use-specific risk mitigation measures

- [If national policy or legislation requires it] Place baits only in sewer systems which are connected to the sewage treatment plant.
- Do not use this product in pulsed baiting treatments.
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

4.7.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.7.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.7.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

5. General directions for use

5.1. Instructions for use

Instructions for Use - Professionals

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drink as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations/ points should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 2.5.3 for the information to be shown on the label).
- [If national policy or legislation require it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets, farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.
- Bait in sachets: Do not open the sachets containing the bait.

Instructions for Use – Trained Professionals

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 2.5.3 for the information to be shown on the label).
- [If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Bait in sachets: [For non-emptiable sachets - Do not open the sachets containing the bait].

IE Only: The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

5.2. Risk mitigation measures

Risk mitigation measures - Professionals

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week). [Where relevant, specify if more frequent or daily inspection is required].
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
 - -the product shall not be supplied to the general public (e.g. "for professionals only").
 - - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - -users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

Risk mitigation measures – Trained Professionals

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose of dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of: Dermal exposure, wash skin with water and then with water and soap.

Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [insert national phone number]".

Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Shelf-life: 24 months

Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.

Store in places prevented from the access of children, birds, pets and farm animals.

Keep only in original container.

6. Other information

Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after consumption of the bait.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

This product contains a bittering agent and a dye.