

Rezumatul caracteristicilor produsului pentru un produs biocid

Denumirea produsului: Roban Pasta Bait

Tipul (tipurile) de produs (produse): TP 14 - Rodenticide

TP 14 - Rodenticide

Numărul curent: RO/2014/0067/MRA/ UK-2011-0085

Numărul curent R4BP 3: RO-0006284-0000

Cuprins

| | |
|---|----|
| Informații administrative | 1 |
| 1.1. Denumirea comercială a produsului | 1 |
| 1.2. Detinatorul autorizației | 1 |
| 1.3. Producătorul (Producătorii) produselor biocide | 1 |
| 1.4. Producatorul (producatorii) substanței (substanțelor) active | 2 |
| 2. Compoziția și formularea produsului | 2 |
| 2.1. Informații calitative și cantitative despre compoziția produsului biocid | 2 |
| 2.2. Tipul preparatului | 2 |
| 3. Fraze de pericol și de precauție | 3 |
| 4. Utilizare (utilizări) autorizată (autorizate) | 3 |
| 5. Direcții generale de utilizare | 8 |
| 5.1. Instrucțiuni de utilizare | 8 |
| 5.2. Măsuri de reducere a riscurilor | 9 |
| 5.3. Particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului | 9 |
| 5.4. Instrucțiuni pentru eliminarea în siguranță a produsului și a ambalajului acestuia | 10 |
| 5.5. Condiții de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare | 10 |
| 6. Alte informații | 10 |

Informații administrative

1.1. Denumirea comercială a produsului

| |
|---|
| Roban Pasta Bait |
| Agriforce Wildcat Pasta Bait |
| Murin Dife Pasta Bait |
| Pest-Stop Super Mouse Killer |
| Romax Mouse DP |
| Nippon Rodenticide Fresh Bait |
| Wilko Mouse & Rat Killer |
| Rascal Difenacoum Pasta Bait |
| Carrs Billington Rat & Mouse Killer Difenacoum Pasta Bait |

1.2. Detinatorul autorizației

| | | |
|--|--------------------------------|---|
| Numele și adresa detinatorului autorizației | Numele | PelGar International Limited |
| | Adresa | 18 rue des Remparts d'Ainay 69002 Lyon Franța |
| Numărul curent | RO/2014/0067/MRA/ UK-2011-0085 | |
| Numărul curent R4BP 3 | RO-0006284-0000 | |
| Data autorizației | 29/09/2011 | |
| Data de expirare a autorizației | 01/07/2024 | |

1.3. Producătorul (Producătorii) produselor biocide

| | |
|--------------------------------------|---|
| Numele producătorului | PelGar International Limited |
| Adresa producătorului | Unit 13 Newman Lane GU34 2QR Alton Regatul Unit |
| Adresa locurilor de producție | Unit 13 Newman Lane GU34 2QR Alton Regatul Unit |

1.4. Producătorul (producătorii) substanței (substanțelor) active

| | |
|--------------------------------------|---|
| Substanța activă | 26 - Difenacum |
| Numele producătorului | PelGar International Limited |
| Adresa producătorului | Unit 13 Newman Lane GU34 2QR Alton Regatul Unit |
| Adresa locurilor de producție | Prazska 54 280 02 Kolin Republica Cehă |

2. Compoziția și formularea produsului

2.1. Informații calitative și cantitative despre compoziția produsului biocid

| Denumirea comună | Denumirea IUPAC | Funcția | Numărul CAS | Numărul CE | Continut (%) |
|------------------|---|------------------|-------------|------------|--------------|
| Difenacum | 3-(3-biphenyl-4-yl-1,2,3,4-tetrahydro-1-naphthyl)-4-hydroxycoumarin | Substanța activă | 56073-07-5 | 259-978-4 | 0,005 |

2.2. Tipul preparatului

| |
|-------------------------------|
| RB - Bait (gata de utilizare) |
|-------------------------------|

3. Fraze de pericol și de precauție

Fraze de pericol

Poate provoca leziuni ale organelor blood în caz de expunere prelungită sau repetată.
Poate dăuna fătului.

Fraze de precauție

Purtați mănuși de protecție.
ÎN CAZ DE expunere sau de posibilă expunere:Consultați medicul.
Consultați medicul, dacă nu vă simțiți bine.
A se depozita sub cheie.
Aruncați conținutul la in accordance with national regulations.
Procurați instrucțiuni speciale înainte de utilizare.
A nu se manipula decât după ce au fost citite și înțelese toate măsurile de securitate.

4. Utilizare (utilizări) autorizată (autorizate)

4.1 Modul de utilizare

Utilizarea 1 - Professional - indoors and outdoors

Tipul produsului

TP 14 - Rodenticide

Descrierea exactă a utilizării autorizate, după caz

Not relevant for rodenticides

Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)

denumirea științifică: Mus musculus
denumirea comună: House mouse
etapă de dezvoltare: Adults and juveniles

denumirea științifică: Rattus norvegicus
denumirea comună: Brown rat
etapă de dezvoltare: Adults and juveniles

Domeniul de utilizare

de interior
în aer liber
Indoors
Outdoors - around buildings

Metoda (metodele) de aplicare

Metoda: cerere momeală
Descriere detaliată:
Ready-to-use bait to be used in tamper-resistant bait stations

Doza (dozele) și frecvența de aplicare

Frecvența de aplicare: RATS: Up to 200 g of bait per bait station. MICE: Up to 40 g of bait per bait station.
Diluare (%): 100
Numărul și calendarul de aplicare:

Mice
High infestation – Up to 40g bait in bait stations every 2 metres
Low infestation – Up to 40g bait in bait stations every 5 metres
Rats
High infestation – Up to 200g bait in bait stations every 5 metres
Low infestation – Up to 200g bait in bait stations every 10 metres

Categoria (categoriile) de utilizatori

profesional

Dimensiunile ambalajului și materialul de ambalare

10 g or 15 g edible paper tea-bag sachets or 30 g polyethylene film 'sausages' in fibre-board carton with polyethylene liner – 3 to 20kg

10g or 15 g edible paper tea-bag sachets or 30 g polyethylene film 'sausages' in polyethylene or polypropylene packs – 3 to 10kg

10 g or 15 g edible paper tea-bag sachets or 30 g polyethylene film 'sausages' in paper/polyethylene or aluminium/polyethylene sachets with or without cardboard outer.. – 3 to 20 kg

10 g or 15 g edible paper tea-bag sachets in HDPE tub – 3 to 20 kg

15 g edible paper tea-bag sachets in polypropylene pouch – 3 to 20 kg

10 g or 15 g sachets in plastic (PE) bag within cardboard outer box – 3 to 20 kg

Pre-filled or refillable tamper proof HDPE or polypropylene mouse bait station containing 2 x 15 g sachets inside a tamper-proof polypropylene box. Bait stations packed in units of 1, 2 or 4 in a cardboard outer or plastic heat-sealed container then in multiples with cardboard outer – 3 to 20 kg.

Pre-filled or refillable tamper proof HDPE or polypropylene rat bait station containing 12 x 15 g sachets secured on a fixed metal bar inside a tamper-proof polypropylene box. Bait stations packed in units of 1 or 2 in cardboard outer or plastic heat-sealed container then in multiples with cardboard outer – 3 to 20 kg.

0.1, 0.15, 0.2 or 0.3 kg mastic tubes for application by caulking gun. Mastic tubes supplied singly or in multiples in cardboard outer – 0.3 to 3.6 kg

Loose paste in polypropylene or polyethylene packs for application by decorator's knife/spatula. – 3 to 5kg

200 g polythene film 'sausage' in a collagen outer casing for use in rat bait boxes, packed in fibreboard outer. – 3 to 10kg

30 g polythene film 'sausage' in a collagen outer casing for use in mouse bait boxes,

packed in fibreboard outer – 3 to 10kg

200 g polypropylene or polystyrene container with a heat-sealed plastic laminate lid in fibreboard cartons. – 3 to 20kg

4.1.1 Instrucțiuni specifice de utilizare

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas non-labile to flooding.
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.1.2 Măsurile specifice de reducere a riscului de utilizare

- Do not apply this product directly in burrows.

4.1.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.1.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.1.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

4.2 Modul de utilizare

Utilizarea 2 - Trained professional - indoors and outdoors (around buildings), in sewers, in open areas and in waste dumps

| | |
|---|--|
| Tipul produsului | TP 14 - Rodenticide |
| Descrierea exactă a utilizării autorizate, după caz | Not relevant for rodenticides |
| Organism(organisme)-țintă (inclusiv stadiul de dezvoltare) | denumirea științifică: Mus musculus denumirea comună: House mouse etapă de dezvoltare: Adults and juveniles denumirea științifică: Rattus norvegicus denumirea comună: Brown rat etapă de dezvoltare: Adults and juveniles |
| Domeniul de utilizare | de interior în aer liber Other Indoors Outdoors - around buildings Outdoors - in open areas Outdoors - in waste dumps In sewers |
| Metoda (metodele) de aplicare | Metoda: cerere momeală Descriere detaliată: Ready-to-use bait to be used in tamper-resistant bait stations, covered and protected baiting points. Direct application of ready-to-use bait into the burrow. Ready-to-use bait in sewers to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. |
| Doza (dozele) și frecvența de aplicare | Frecvența de aplicare: RATS: 20 to 200 g of bait per bait station. MICE: 10 to 40 g of bait per bait station. Diluare (%): 100 Numărul și calendarul de aplicare: Mice High infestation – 10 to 40g bait in bait stations every 2 metres Low infestation – 10 to 40g bait in bait stations every 5 metres Rats High infestation – 20 to 200g bait in bait stations every 5 metres Low infestation – 20 to 200g bait in bait stations every 10 metres |
| Categoria (categoriile) de utilizatori | instruiți profesional |
| Dimensiunile ambalajului și materialul de ambalare | 10 g or 15 g edible paper tea-bag sachets or 30 g polyethylene film 'sausages' in fibre-board carton with polyethylene liner – 3 to 20kg 10g or 15 g edible paper tea-bag sachets or 30 g polyethylene film 'sausages' in polyethylene or polypropylene packs – 3 to 10kg |

10 g or 15 g edible paper tea-bag sachets or 30 g polyethylene film 'sausages' in paper/polyethylene or aluminium/polyethylene sachets with or without cardboard outer.. – 3 to 20 kg

10 g or 15 g edible paper tea-bag sachets in HDPE tub – 3 to 20 kg

15 g edible paper tea-bag sachets in polypropylene pouch – 3 to 20 kg

10 g or 15 g sachets in plastic (PE) bag within cardboard outer box – 3 to 20 kg

Pre-filled or refillable tamper proof HDPE or polypropylene mouse bait station containing 2 x 15 g sachets inside a tamper-proof polypropylene box. Bait stations packed in units of 1, 2 or 4 in a cardboard outer or plastic heat-sealed container then in multiples with cardboard outer – 3 to 20 kg.

Pre-filled or refillable tamper proof HDPE or polypropylene rat bait station containing 12 x 15 g sachets secured on a fixed metal bar inside a tamper-proof polypropylene box. Bait stations packed in units of 1 or 2 in cardboard outer or plastic heat-sealed container then in multiples with cardboard outer – 3 to 20 kg.

0.1, 0.15, 0.2 or 0.3 kg mastic tubes for application by caulking gun. Mastic tubes supplied singly or in multiples in cardboard outer – 0.3 to 3.6 kg

Loose paste in polypropylene or polyethylene packs for application by decorator's knife/spatula. – 3 to 5kg

200 g polythene film 'sausage' in a collagen outer casing for use in rat bait boxes, packed in fibreboard outer. – 3 to 10kg

30 g polythene film 'sausage' in a collagen outer casing for use in mouse bait boxes, packed in fibreboard outer – 3 to 10kg

200 g polypropylene or polystyrene container with a heat-sealed plastic laminate lid in fibreboard cartons. – 3 to 20kg

4.2.1 Instrucțiuni specifice de utilizare

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
 - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
 - Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent)
 - Remove the remaining product at the end of treatment period (except when directly applied into burrows)
 - For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Use in sewers: - Baits must be applied in a way so that they do not come into contact with water and are not washed away.

4.2.2 Măsuri specifice de reducere a riscului de utilizare

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.

4.2.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.2.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.2.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

5. Direcții generale de utilizare

5.1. Instrucțiuni de utilizare

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
 - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
 - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
 - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
 - The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
 - Where possible, bait stations must be fixed to the ground or other structures.
 - Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
 - Bait should be secured so that it cannot be dragged away from the bait station.
 - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
 - Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
 - Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
-
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
 - The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the CRRU UK Code of Best Practice (or equivalent). Sites where burrow baiting is used should be visited more frequently than those where secure bait boxes are employed.
 - If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
 - If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- For non-emptiable sachets:- Do not open the sachets containing the bait.

5.2. Măsuri de reducere a riscurilor

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.
 - Do not use in areas where resistance to the active substance can be suspected.
 - Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
 - Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
 - Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
 - Dispose dead rodents in accordance with local requirements.
- UK only: In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).
- UK only: To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.
- UK only: Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.
- UK only: When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

5.3. Particularitati ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

UK only: UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice. Contact a veterinary surgeon in case of ingestion by a pet.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident call a poison center".

- Hazardous to wildlife.

5.4. Instrucțiuni pentru eliminarea în siguranță a produsului și a ambalajului acestuia

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

UK only: For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

5.5. Condiții de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 2 years

6. Alte informații

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing of them.
- This product contains a bittering agent and a dye.

